

ON GRADIENT RICCI SOLITONS WITH CONSTANT SCALAR CURVATURE

MANUEL FERNÁNDEZ-LÓPEZ AND EDUARDO GARCÍA-RÍO

(Communicated by Guofang Wei)

ABSTRACT. We use the theory of isoparametric functions to investigate gradient Ricci solitons with constant scalar curvature. We show rigidity of gradient Ricci solitons with constant scalar curvature under some conditions on the Ricci tensor, which are all satisfied if the manifold is curvature homogeneous. This leads to a complete description of four- and six-dimensional Kähler gradient Ricci solitons with constant scalar curvature.

1. INTRODUCTION AND MAIN RESULTS

A complete n -dimensional Riemannian manifold (M, g) is said to be a *gradient Ricci soliton* if there exists a smooth function f on M such that

$$(1.1) \quad Rc + H_f = \lambda g,$$

where Rc is the Ricci tensor, H_f denotes the Hessian of the function f , and λ is a real number. For $\lambda > 0$ the Ricci soliton is *shrinking*, for $\lambda = 0$ it is *steady*, and for $\lambda < 0$ it is *expanding*. The function f is called the *potential function* of the gradient Ricci soliton. Gradient Ricci solitons play an important role in Hamilton's Ricci flow as they correspond to self-similar solutions, and often arise as singularity models. Therefore it is important to classify gradient Ricci solitons and to understand their geometry. We refer to [3, 7, 8] and the references therein for background on Ricci solitons.

The Ricci soliton equation (1.1) links geometric information about the curvature of the manifold through the Ricci tensor and the geometry of the level sets of the potential function by means of their second fundamental form. Hence, classifying gradient Ricci solitons under some curvature conditions is a natural problem. Gradient Ricci solitons with constant scalar curvature were investigated by Petersen and Wylie in [15], who showed that constant scalar curvature is a very restrictive condition in the steady case, since it leads to Ricci flatness. Therefore we focus on the non-steady case in what follows.

If a non-steady gradient Ricci soliton has constant scalar curvature R , then it is bounded as $0 \leq R \leq n\lambda$ in the shrinking case, and $n\lambda \leq R \leq 0$ in the expanding case. Our first result shows that the possible values of the scalar curvature are quantified by the soliton constant λ as follows.

Received by the editors September 17, 2014 and, in revised form, December 12, 2014.

2010 *Mathematics Subject Classification*. Primary 53C25, 53C20, 53C44.

Key words and phrases. Gradient Ricci soliton, scalar curvature.

The authors were supported by projects GRC2013-045 and MTM2013-41335-P with FEDER funds (Spain).

Theorem 1. *Let (M^n, g) be an n -dimensional complete gradient Ricci soliton with constant scalar curvature R . Then $R \in \{0, \lambda, \dots, (n-1)\lambda, n\lambda\}$.*

Petersen and Wylie showed that the extremal values for the scalar curvature in Theorem 1 are achieved if and only if the underlying Riemannian structure is Einstein [15]. We investigate the realizability of the extremal values $R = \lambda$ and $R = (n-1)\lambda$, showing that the value $R = \lambda$ cannot occur in the shrinking case, while any complete gradient Ricci soliton with constant scalar curvature $R = (n-1)\lambda$ is necessarily rigid (see Theorem 10).

A gradient soliton is said to be *rigid* if it is isometric to a quotient of $N \times \mathbb{R}^k$, where N is an Einstein manifold and $f = \frac{\lambda}{2}|x|^2$ on the Euclidean factor. That is, the Riemannian manifold (M, g) is isometric to $N \times_{\Gamma} \mathbb{R}^k$, where Γ acts freely on N and by orthogonal transformations on \mathbb{R}^k . Clearly any rigid gradient Ricci soliton has constant scalar curvature. When M is compact, a Ricci soliton (M, g) is rigid if and only if it is Einstein, and the constancy of the scalar curvature characterizes rigidity as a consequence of the Hopf maximum principle (see, for example, [9]). In the more general setting of complete non-compact gradient Ricci solitons, Petersen and Wylie showed rigidity under some additional assumptions [15]. One such condition is the harmonicity of the Riemann curvature tensor. In the shrinking case, a gradient Ricci soliton is rigid if and only if the Weyl tensor is harmonic [11, 14].

Our main result shows rigidity of gradient Ricci solitons with constant scalar curvature under the assumption that the Ricci operator has constant rank. From this it follows that curvature homogeneous gradient Ricci solitons are rigid.

Theorem 2. *A complete gradient Ricci soliton with constant scalar curvature is rigid if and only if the Ricci operator has constant rank.*

Remark 3. A special family of rigid gradient Ricci solitons are the homogeneous ones. Indeed, it was shown in [16] that any homogeneous gradient Ricci soliton is rigid. An immediate application of Theorem 2 shows that the homogeneity condition can be replaced by the more general one of curvature homogeneity. A Riemannian manifold (M, g) is said to be *k -curvature homogeneous* if for any two points $p, q \in M$ there exists a linear isometry $\varphi_{pq} : T_p M \rightarrow T_q M$ which preserves the curvature tensor and its covariant derivatives up to order k , i.e., $\varphi_{pq}^* \nabla^l R(q) = \nabla^l R(p)$ for all $l = 0, \dots, k$. Clearly, any locally homogeneous Riemannian manifold is k -curvature homogeneous for all k , and conversely, if (M^n, g) is k -curvature homogeneous for sufficiently large k (for instance $k \geq \frac{1}{2}n(n-1)$), then it is locally homogeneous. However, there are plenty of examples of Riemannian manifolds which are 0-curvature homogeneous but not homogeneous (there are even many curvature homogeneous Riemannian manifolds whose curvature tensor does not correspond to any homogeneous space [1]). Since any curvature homogeneous Riemannian manifold has constant Ricci curvatures, the Ricci operator has constant rank, and thus it follows from Theorem 2 that any 0-curvature homogeneous complete gradient Ricci soliton is rigid, which generalizes [16, Theorem 1.1].

Gradient Ricci solitons with constant scalar curvature are rigid in dimension three [17] (in the shrinking case rigidity was shown without any additional assumption in [4]). We extend the above results to the case of Kähler Ricci solitons by showing the following.

Theorem 4. *Any Kähler gradient Ricci soliton with constant scalar curvature is rigid in dimension $n = 4, 6$.*

The previous result strongly depends on the number of different Ricci curvatures and does not hold necessarily in the general non-Kähler case, where we have:

Theorem 5. *Any four-dimensional complete gradient shrinking Ricci soliton with constant scalar curvature $R \neq 2\lambda$ is rigid. Moreover, any four-dimensional gradient shrinking (resp., expanding) Ricci soliton with constant scalar curvature $R = 2\lambda$ has non-negative (resp., non-positive) Ricci curvature.*

2. PROOF OF THE RESULTS

The results will be obtained by considering the geometric information underlying equation (1.1). The analysis of the level sets of the potential function plays a crucial role since our assumption of constant scalar curvature imposes some restrictions on their geometry. First of all we introduce some definitions to be used in what follows.

A non-constant C^2 function $f : M \rightarrow \mathbb{R}$ is said to be a *transnormal function* if

$$(2.1) \quad |\nabla f|^2 = b(f)$$

for some C^2 function b on the range of f in \mathbb{R} . The function f is said to be an *isoparametric function* if moreover it satisfies

$$(2.2) \quad \Delta f = a(f)$$

for some continuous function a on the range of f in \mathbb{R} [2, 12, 13, 18].

Equation (2.1) implies that the level set hypersurfaces of f are parallel hypersurfaces and it follows from equation (2.2) that these hypersurfaces have constant mean curvature (i.e., they are isoparametric hypersurfaces).

Given a function $f : M \rightarrow \mathbb{R}$ we denote $f_{min} = \min\{f(x) / x \in M\}$ and $f_{max} = \max\{f(x) / x \in M\}$, if they exist. If the function is transnormal we define the sets $M_- = \{x \in M / f(x) = f_{min}\}$ and $M_+ = \{x \in M / f(x) = f_{max}\}$, which are called the *focal varieties* of f . Note that M_- and/or M_+ could be the empty set.

If (M, g) is a gradient Ricci soliton, the potential function (after a possible rescaling) satisfies $|\nabla f|^2 = 2\lambda f - R$ (see, for example, [7, 15]) and $\Delta f = n\lambda - R$. Thus, if the Ricci soliton has constant scalar curvature, the potential function satisfies (2.1) and (2.2), thus being an isoparametric function on (M, g) . First of all we show that at least one of the focal varieties is necessarily non-empty.

Lemma 6. *Let (M, g) be a gradient Ricci soliton with constant scalar curvature. If it is shrinking, then $M_- \neq \emptyset$ and if it is expanding, then $M_+ \neq \emptyset$.*

Proof. We need to show that the potential function has a minimum (resp., maximum) if the Ricci soliton is shrinking (resp., expanding). After translating f by a constant we can suppose that $2\lambda f = |\nabla f|^2$. This shows that f has the same sign as λ , and moreover it vanishes at the same points where ∇f does so. Hence, if $p \in M$ is such that $\nabla f(p) = 0$, then $f(p) = 0$ and the potential function attains a minimum (resp., maximum) at $p \in M$.

We will argue by contradiction, assuming that $\nabla f \neq 0$ at any point to get a contradiction. Define the function $r : (M, g) \rightarrow \mathbb{R}$ by $r = \sqrt{\frac{2f}{\lambda}}$. Now it follows that $\nabla r = \frac{\sqrt{\lambda} \nabla f}{2|\nabla f|}$, from which we have that r satisfies the Eikonal equation $|\nabla r|^2 = 1$. Thus r is a distance function and the integral curves of ∇r are geodesics. Hence

∇r is a complete vector field, due to completeness of (M, g) . So, if $\nabla f \neq 0$, then the range of r must be \mathbb{R} , which contradicts the fact that r is non-negative. This finishes the proof. \square

Remark 7. Cao and Zhou [5] showed that if (M, g) is a shrinking gradient Ricci soliton, then the potential function satisfies $\frac{\lambda}{2}(r(p) - c_1)^2 \leq f(p) \leq \frac{\lambda}{2}(r(p) - c_1)^2$, where $r(x) = d(p_0, p)$ is the distance function from some fixed point $p_0 \in M$, and c_1 and c_2 are positive constants. Previous inequalities provide an alternative proof of the fact that f attains a global minimum in the shrinking case.

Proof of Theorem 1. Wang proved in [18] that the focal varieties M_- and M_+ of a transnormal function are smooth submanifolds of M (whenever they are not empty). Also it was shown in [18] that the restriction of the Hessian H_f of a transnormal function f satisfying (2.1) to M_\pm has only two eigenvalues, 0 and $\frac{1}{2}b'(f)$. Indeed, one has $H_f(X, Y) = 0$ for all $X, Y \in TM_\pm$, and $H_f(V, W) = \frac{1}{2}b'(f)g(V, W)$ for all $V, W \in TM_\pm^\perp$.

In our case, since $b(f) = 2\lambda f - R$, we have that $b'(f) = 2\lambda$. Now, it follows from (1.1) that the restriction of the Ricci tensor to the (non-empty) focal submanifolds M_\pm is of the form

$$Rc|_{M_\pm} = \begin{pmatrix} \lambda I_k & 0 \\ 0 & 0_{n-k} \end{pmatrix},$$

where $k = \text{rank } Rc|_{M_\pm}$. The previous lemma shows that at least one of the focal varieties is not empty, and hence taking traces one obtains that the possible values of the scalar curvature are quantified by the Ricci soliton constant λ , thus proving Theorem 1. \square

A basic fact in our analysis is the following [15, Proposition 1.3]: a gradient shrinking (resp., expanding) Ricci soliton is rigid if and only if it has constant scalar curvature and $0 \leq Rc \leq \lambda$ (resp., $\lambda \leq Rc \leq 0$).

Proof of Theorem 2. Recall that the f -Laplacian of the scalar curvature satisfies $\Delta_f R = \lambda R - |Rc|^2$, from where $|Rc|^2 = \lambda R$ in the constant scalar curvature setting (see, for example, [15]). Next, assuming that the rank of the Ricci operator is constant, say $\text{rank } Rc = \dim M_\pm = k$, one has that the scalar curvature is $R = k\lambda$, since it is constant. Let us denote by R_i , $i = 1, \dots, k$, the non-zero eigenvalues of the Ricci operator at any point of M . Then

$$\begin{aligned} \sum_{i=1}^k (R_i - \lambda)^2 &= |Rc|^2 - 2\lambda R + k\lambda^2 \\ &= \lambda R - 2\lambda R + \lambda R = 0. \end{aligned}$$

Hence all non-zero eigenvalues satisfy $R_1 = R_2 = \dots = R_k = \lambda$, and thus it follows that the Ricci tensor has only two eigenvalues, 0 and λ , from which it follows that $0 \leq Rc \leq \lambda$ and rigidity is a consequence of the results in [15].

The proof is analogous in the expanding case, using that $M_+ \neq \emptyset$ and $Rc|_{M_+} = \text{diag}[\lambda, \dots, \lambda, 0, \dots, 0]$ to obtain that $\lambda \leq Rc \leq 0$. \square

Remark 8. The proof of Theorem 2 relies on the fact that $|Rc|^2 = \lambda R$ in the constant scalar curvature setting, from which it follows that $|Rc|^2$ is constant provided that R is also. The converse is also true in the non-expanding setting. In fact, since $R^2 \leq n|Rc|^2$, one has that R is bounded if $|Rc|^2$ is constant. If a steady gradient Ricci

soliton has constant $|Rc|^2$, then it is Ricci flat since at the maximum/supremum of R , $\Delta_f R = -2|Rc|^2 \leq 0$ and at the minimum/infimum of R , $\Delta_f R = -2|Rc|^2 \geq 0$, from which it follows that $|Rc|^2 \equiv 0$ and the manifold is Ricci flat. Also, if (M, g) is a shrinking gradient Ricci soliton, then R is constant provided that $|Rc|^2$ is also. Indeed, at any point where R attains a maximum/supremum R^* one has that $\Delta_f R = \lambda R^* - 2|Rc|^2 \leq 0$, which implies $\lambda R^* \leq 2|Rc|^2$. Analogously, at any point where R attains a minimum/infimum R_* one has that $\Delta_f R = \lambda R_* - 2|Rc|^2 \geq 0$ and thus $\lambda R_* \geq 2|Rc|^2$. Hence $R^* \leq R_*$, which shows that R is constant.

It is shown in [16] that 3-dimensional gradient Ricci solitons with constant scalar curvature are rigid. Note that the Ricci tensor of such a soliton has at most three different eigenvalues. We extend this result in the following theorem.

Theorem 9. *Let (M, g) be a gradient Ricci soliton with constant scalar curvature. If Rc has at most three different eigenvalues, then it is rigid.*

Proof. Note that Rc has at least one zero eigenvalue due to the constancy of the scalar curvature. Indeed, the Ricci operator satisfies $Rc(\nabla f) = \frac{1}{2}\nabla R$ at any point $p \in M$ where $(\nabla f)(p) \neq 0$, which shows that Rc has a zero eigenvalue on $M \setminus M_{\pm}$ and, as shown in the proof of Theorem 1, it also has a zero eigenvalue on M_{\pm} .

Assume that it has two non-zero (possibly) different eigenvalues R_1, R_2 of multiplicities k_1 and k_2 . Since R and $|Rc|^2 = \lambda R$ are constant, R_1 and R_2 must be constant. Indeed, they solve the system of equations

$$\begin{cases} k_1 R_1 + k_2 R_2 = R, \\ k_1 R_1^2 + k_2 R_2^2 = \lambda R, \end{cases}$$

which only has two solutions if $R_1 \neq R_2$ and one solution if $R_1 = R_2$. Because of the continuity of the eigenvalues of the Ricci tensor it follows that they are constant, from which we get rigidity. \square

Proof of Theorem 4. Since for Kähler manifolds (M, g, J) the Ricci tensor is invariant under the action of the complex structure J (that is, it satisfies $Rc(J\cdot, J\cdot) = Rc(\cdot, \cdot)$), each Ricci curvature has even multiplicity. Since the Ricci tensor has a zero eigenvalue with multiplicity at least 2 it follows that Rc has at most two different eigenvalues if $n = 4$ and three different eigenvalues if $n = 6$. The result now follows from Theorem 9. \square

An alternative proof of the rigidity of four-dimensional Kähler Ricci solitons has been recently given by Wu, Wu, and Wylie in [19], who followed a different approach based on the harmonicity of the half-Weyl conformal curvature tensor.

Theorem 1 showed that the possible values of the scalar curvature of any gradient Ricci soliton of constant scalar curvature are $R \in \{0, \lambda, \dots, (n-1)\lambda, n\lambda\}$. Petersen and Wylie proved in [15] that the extremal values may occur only for Einstein metrics. Next we show that the soliton is necessarily rigid if $R = (n-1)\lambda$ and that $R \neq \lambda$ in the shrinking case.

Theorem 10. *Any complete gradient Ricci soliton with constant scalar curvature $R = (n-1)\lambda$ is rigid. Moreover, no complete gradient shrinking Ricci soliton may exist with $R = \lambda$.*

Proof. Let us denote by R_i ($i = 1, \dots, n - 1$) the non-zero eigenvalues of the Ricci operator at any point of M . It holds that

$$\begin{aligned} \sum_{i=1}^{n-1} (R_i - \lambda)^2 &= |Rc|^2 - 2\lambda R + (n-1)\lambda^2 \\ &= \lambda R - 2\lambda R + \lambda R = 0. \end{aligned}$$

So all non-zero eigenvalues satisfy $R_1 = R_2 = \dots = R_{n-1} = \lambda$. Thus, Rc has constant rank and the Ricci soliton is rigid.

To show the non-existence of complete shrinking solitons with $R = \lambda$ we argue by contradiction. If $R = \lambda$, then the focal variety $M_- \neq \emptyset$ is a minimal submanifold (see [12]). Since it has dimension one it must be totally geodesic. Moreover it is compact since f is proper. So we have that M_- is \mathbb{S}^1 . Now, M is diffeomorphic to a vector bundle over \mathbb{S}^1 (see [2, 13]). But this contradicts the fact that M has finite fundamental group [20]. \square

Remark 11. It is shown in [17, Corollary 2] that the existence of a non-zero eigenvalue of the Ricci operator of multiplicity $n - 1$ leads to rigidity. The assumption $R = (n - 1)\lambda$ in Theorem 10 is equivalent to the existence of an eigenvalue of multiplicity $n - 1$ along the focal varieties M_\pm .

Although our results are not so conclusive for other values of the scalar curvature, the Ricci curvature has sign as follows.

Proposition 12. *Let (M, g) be a gradient shrinking (resp., expanding) Ricci soliton with constant scalar curvature $R = k\lambda$. If $\text{rank } Rc \leq k + 1$, then the Ricci curvature is non-negative (resp., non-positive).*

Proof. Let us denote by R_i ($i = 1, \dots, k + 1$) the possibly non-zero eigenvalues of the Ricci operator at any point of M . We have that

$$\sum_{i=1}^{k+1} R_i^2 = |Rc|^2 = \lambda R = k\lambda^2 = \frac{1}{k} \left(\sum_{i=1}^{r+1} R_i \right)^2.$$

Now, adapting the proof of [6, Lemma 1], one has that for any reals α_i , $i = 1, \dots, r$ ($r \geq 2$), such that

$$\sum_{i=1}^r \alpha_i^2 \leq \frac{1}{r-1} \left(\sum_{i=1}^r \alpha_i \right)^2,$$

then all the α_i 's are non-negative or non-positive.

Hence all non-zero eigenvalues of the Ricci operator are positive or negative. In the shrinking case it is $R > 0$ from where we get $Rc \geq 0$, and in the expanding case it holds that $R < 0$, which gives $Rc \leq 0$. \square

Remark 13. If the scalar curvature of a gradient Ricci soliton satisfies $R = (n - 2)\lambda$, then $\text{rank } Rc \leq n - 1$ since Rc has at least a zero eigenvalue due to the constancy of the scalar curvature. Hence the Ricci curvature is non-negative (resp., non-positive) in the shrinking (resp., expanding) case.

Proof of Theorem 5. The proof immediately follows from Theorem 10 and Proposition 12. \square

Recall that a gradient shrinking (expanding) Ricci soliton is rigid if and only if it has constant scalar curvature and $0 \leq Rc \leq \lambda$ (resp., $\lambda \leq Rc \leq 0$) [15]. Next, we compute the f -Laplacian of the Ricci tensor acting on the gradient of the potential function in terms of the eigenvalues of the Ricci tensor aimed to show that if the scalar curvature is constant and $(\Delta_f Rc)(\nabla f, \nabla f) = 0$, then rigidity of gradient shrinking (resp., expanding) Ricci solitons is characterized by only one of the bounds: $Rc \geq 0$ or $Rc \leq \lambda$ (resp., $Rc \leq 0$ or $Rc \geq \lambda$).

Lemma 14. *Let (M, g) be a gradient Ricci soliton with constant scalar curvature. If R_i denote the eigenvalues of the Ricci tensor, then it holds that*

$$(\Delta_f Rc)(\nabla f, \nabla f) = 2 \sum_{i=1}^n (\lambda - R_i)^2 R_i.$$

Proof. Let us consider $p \in M$ such that $(\nabla f)(p) \neq 0$. Let $\{E_1, \dots, E_n\}$ be eigenvectors of Rc at p and take normal coordinates centered at p induced by the orthonormal basis $\{E_1, \dots, E_n\}$. Then, using that $Rc(\cdot, \nabla f) = 0$, one has

$$\begin{aligned} (\Delta_f Rc)(\nabla f, \nabla f) &= \sum_{i=1}^n (\nabla_{E_i, E_i}^2 Rc)(\nabla f, \nabla f) - (\nabla_{\nabla f} Rc)(\nabla f, \nabla f) \\ &= \sum_{i=1}^n (\nabla_{E_i}(\nabla_{E_i} Rc))(\nabla f, \nabla f) \\ &= \sum_{i=1}^n [\nabla_{E_i}(\nabla_{E_i} Rc)(\nabla f, \nabla f) - 2(\nabla_{E_i} Rc)(\nabla_{E_i} \nabla f, \nabla f)] \\ &= \sum_{i=1}^n \nabla_{E_i}(-2Rc(\nabla_{E_i} \nabla f, \nabla f)) - 2(\nabla_{E_i} Rc)(\nabla_{E_i} \nabla f, \nabla f) \\ &= -2 \sum_{i=1}^n [E_i Rc(\nabla_{E_i} \nabla f, \nabla f) - Rc(\nabla_{E_i} \nabla_{E_i} \nabla f, \nabla f) \\ &\quad - Rc(\nabla_{E_i} \nabla f, \nabla_{E_i} \nabla f)] \\ &= 2 \sum_{i=1}^n (\lambda - R_i)^2 R_i. \end{aligned}$$

□

Corollary 15. *Let (M, g) be a gradient shrinking (resp., expanding) Ricci soliton with constant scalar curvature. If $(\Delta_f Rc)(\nabla f, \nabla f) = 0$ and $Rc \geq 0$ or $Rc \leq \lambda$ (resp., $Rc \leq 0$ or $Rc \geq \lambda$), then the soliton is rigid.*

Proof. First of all note that, since $|Rc|^2 = \lambda R$, one has $\sum_{i=1}^n (\lambda - R_i)^2 R_i = \sum_{i=1}^n R_i^2 (R_i - \lambda)$. Now it follows from the constancy of the scalar curvature and $(\Delta_f Rc)(\nabla f, \nabla f) = 0$ that $\sum_{i=1}^n (\lambda - R_i)^2 R_i = 0$. Hence

$$\sum_{i=1}^n (\lambda - R_i)^2 R_i = \sum_{i=1}^n R_i^2 (R_i - \lambda) = 0.$$

Thus, under our assumptions on the bounds of the Ricci tensor we get that $R_i \in \{0, \lambda\}$, and thus the Ricci curvatures are constant, which shows rigidity. □

As shown in Theorem 5, four-dimensional complete shrinking gradient Ricci solitons with constant scalar curvature $R \neq 2\lambda$ are rigid. If the scalar curvature

satisfies $R = 2\lambda$ we need some additional assumptions to get rigidity. Before stating the next result, we recall some basic facts about scalar curvature invariants. The space of scalar curvature invariants of order two, $I(1, n)$, is one-dimensional and it is generated by the scalar curvature R . The space corresponding to four-order invariants $I(2, n)$ is generated by $\{R, \Delta R, |Rc|^2, |Rm|^2\}$. Some other contractions of the curvature and the Ricci tensors, giving rise to new $(0, 2)$ -symmetric tensors fields (\check{R} , $\check{R}c$ and $L(Rc)$), will be used in what follows. With respect to an orthonormal basis, set

$$\check{R}_{ij} = \sum_{abc} R_{abci} R_{abcj}, \quad \check{R}c_{ij} = \sum_a R_{ai} R_{aj}, \quad L(Rc)_{ij} = 2 \sum_{ab} R_{iabj} R_{cab}.$$

Considering the four-dimensional Gauss-Bonnet integrand $|Rm|^2 - 4|Rc|^2 + R^2$, the following curvature identity is shown in [10]:

$$\check{R} - 2\check{R}c - L(Rc) + RRc = \frac{1}{4}(|Rm|^2 - 4|Rc|^2 + R^2)g.$$

Next, observe that the f -Laplacian of the principal Ricci curvatures (see, for example, [9]) is given by

$$\Delta_f R_i = 2\lambda R_i - 2R_{ikis} R c^{ks} = 2\lambda R_i + L(Rc)_{ii}.$$

Theorem 16. *Let (M, g) be a four-dimensional complete gradient shrinking Ricci soliton with constant scalar curvature $R = 2\lambda$. Then it is rigid if and only if one of the following assumptions holds:*

- (i) $Rc \leq \lambda$,
- (ii) $(\Delta_f Rc)(\nabla f, \nabla f) = 0$,
- (iii) $\check{R}(\nabla f, \nabla f) \leq \frac{1}{4}(|Rm|^2 - 4\lambda^2)g(\nabla f, \nabla f)$.

Proof of Theorem 16. Under our first assumption, $Rc \leq \lambda$, we have that $0 \leq Rc \leq \lambda$, from which follows rigidity [15]. Now, assume that $(\Delta_f Rc)(\nabla f, \nabla f) = 0$. It follows from the previous lemma that $\sum_{i=1}^4 (\lambda - R_i)^2 R_i = 0$. Since $Rc \geq 0$ we have that $R_i \in \{0, \lambda\}$. Thus Rc has constant eigenvalues and the soliton is rigid.

Let $\{E_1, E_2, E_3, E_4\}$ be an orthonormal basis consisting of eigenvectors of the Ricci tensor with $E_1 = \frac{\nabla f}{|\nabla f|}$. Then $\Delta_f R_1 = L(Rc)_{11}$. Since $R = 2\lambda$, it follows from Proposition 12 and Remark 13 that $\Delta_f R_1 = \frac{1}{|\nabla f|^2(p)}(\Delta_f Rc)(\nabla f, \nabla f) \geq 0$.

Now, since the scalar curvature is constant, $R = 2\lambda$, $Rc(E_1, E_1) = \check{R}c(E_1, E_1) = 0$, and $|Rm|^2 - 4|Rc|^2 + R^2 = |Rm|^2 - 4\lambda^2$. Hence the curvature identity above becomes

$$L(Rc)(E_1, E_1) = \check{R}(E_1, E_1) - \frac{1}{4}|Rm|^2 + \lambda^2.$$

Now, it follows from assumption (iii) that $L(Rc)(E_1, E_1) \leq 0$, and hence $\Delta_f R_1 = 0$ and rigidity is a consequence of assumption (ii).

Finally observe that conditions (i)-(iii) are satisfied by any rigid four-dimensional gradient Ricci soliton. \square

Remark 17. As a final observation, note that a *complete gradient Ricci soliton is rigid if and only if the Ricci operator Rc and its powers Rc^2 , Rc^3 , and Rc^4 have constant traces*. In fact, proceeding as in the proof of Theorem 2 one has that the function

$$\sum_i R_i^2 (\lambda - R_i)^2 = \sum_i R_i^4 - 2\lambda \sum_i R_i^3 + \lambda^2 \sum_i R_i^2$$

is constant provided that the traces of the first powers of the Ricci operator are so. Since its value on the focal submanifold is zero we have that the function $\sum_i R_i^2(\lambda - R_i)^2$ vanishes on M . Thus $R_i \in \{0, \lambda\}$, which shows that the Ricci curvatures are constant, from which it follows that the soliton is rigid.

Further note that while the constancy of the scalar curvature implies the constancy of the trace of Rc^2 , the functions $\text{tr } Rc^3$ and $\text{tr } Rc^4$ are not yet completely understood.

REFERENCES

- [1] Eric Boeckx, Oldřich Kowalski, and Lieven Vanhecke, *Riemannian manifolds of conullity two*, World Scientific Publishing Co., Inc., River Edge, NJ, 1996. MR1462887 (98h:53075)
- [2] J. Bolton, *Transnormal systems*, Quart. J. Math. Oxford Ser. (2) **24** (1973), 385–395. MR0336645 (49 #1418)
- [3] Huai-Dong Cao, *Recent progress on Ricci solitons*, Recent advances in geometric analysis, Adv. Lect. Math. (ALM), vol. 11, Int. Press, Somerville, MA, 2010, pp. 1–38. MR2648937 (2011d:53061)
- [4] Huai-Dong Cao, Bing-Long Chen, and Xi-Ping Zhu, *Recent developments on Hamilton’s Ricci flow*, Surveys in differential geometry. Vol. XII. Geometric flows, Surv. Differ. Geom., vol. 12, Int. Press, Somerville, MA, 2008, pp. 47–112, DOI 10.4310/SDG.2007.v12.n1.a3. MR2488948 (2010c:53096)
- [5] Huai-Dong Cao and Detang Zhou, *On complete gradient shrinking Ricci solitons*, J. Differential Geom. **85** (2010), no. 2, 175–185. MR2732975 (2011k:53040)
- [6] Qing-Ming Cheng, *Compact locally conformally flat Riemannian manifolds*, Bull. London Math. Soc. **33** (2001), no. 4, 459–465, DOI 10.1017/S0024609301008074. MR1832558 (2002g:53045)
- [7] Bennett Chow, Sun-Chin Chu, David Glickenstein, Christine Guenther, James Isenberg, Tom Ivey, Dan Knopf, Peng Lu, Feng Luo, and Lei Ni, *The Ricci flow: techniques and applications. Part I*, Mathematical Surveys and Monographs, vol. 135, American Mathematical Society, Providence, RI, 2007. Geometric aspects. MR2302600 (2008f:53088)
- [8] Bennett Chow, Peng Lu, and Lei Ni, *Hamilton’s Ricci flow*, Graduate Studies in Mathematics, vol. 77, American Mathematical Society, Providence, RI; Science Press, New York, 2006. MR2274812 (2008a:53068)
- [9] Manolo Eminent, Gabriele La Nave, and Carlo Mantegazza, *Ricci solitons: the equation point of view*, Manuscripta Math. **127** (2008), no. 3, 345–367, DOI 10.1007/s00229-008-0210-y. MR2448435 (2009m:53088)
- [10] Yunhee Euh, JeongHyeong Park, and Kouei Sekigawa, *A curvature identity on a 4-dimensional Riemannian manifold*, Results Math. **63** (2013), no. 1-2, 107–114, DOI 10.1007/s00225-011-0164-3. MR3009674
- [11] Manuel Fernández-López and Eduardo García-Río, *Rigidity of shrinking Ricci solitons*, Math. Z. **269** (2011), no. 1-2, 461–466, DOI 10.1007/s00209-010-0745-y. MR2836079 (2012g:53072)
- [12] Jianquan Ge and Zizhou Tang, *Geometry of isoparametric hypersurfaces in Riemannian manifolds*, Asian J. Math. **18** (2014), no. 1, 117–125, DOI 10.4310/AJM.2014.v18.n1.a6. MR3215342
- [13] Reiko Miyaoka, *Transnormal functions on a Riemannian manifold*, Differential Geom. Appl. **31** (2013), no. 1, 130–139, DOI 10.1016/j.difgeo.2012.10.005. MR3010083
- [14] Ovidiu Munteanu and Natasa Sesum, *On gradient Ricci solitons*, J. Geom. Anal. **23** (2013), no. 2, 539–561, DOI 10.1007/s12220-011-9252-6. MR3023848
- [15] Peter Petersen and William Wylie, *Rigidity of gradient Ricci solitons*, Pacific J. Math. **241** (2009), no. 2, 329–345, DOI 10.2140/pjm.2009.241.329. MR2507581 (2010j:53071)
- [16] Peter Petersen and William Wylie, *On gradient Ricci solitons with symmetry*, Proc. Amer. Math. Soc. **137** (2009), no. 6, 2085–2092, DOI 10.1090/S0002-9939-09-09723-8. MR2480290 (2010a:53073)
- [17] Peter Petersen and William Wylie, *On the classification of gradient Ricci solitons*, Geom. Topol. **14** (2010), no. 4, 2277–2300, DOI 10.2140/gt.2010.14.2277. MR2740647 (2012a:53060)
- [18] Qi Ming Wang, *Isoparametric functions on Riemannian manifolds. I*, Math. Ann. **277** (1987), no. 4, 639–646, DOI 10.1007/BF01457863. MR901710 (88h:53056)

- [19] J.-Y. Wu, P. Wu and W. Wylie, Gradient shrinking Ricci solitons of half harmonic Weyl curvature, to appear.
- [20] William Wylie, *Complete shrinking Ricci solitons have finite fundamental group*, Proc. Amer. Math. Soc. **136** (2008), no. 5, 1803–1806, DOI 10.1090/S0002-9939-07-09174-5. MR2373611 (2009a:53059)

IES MARÍA SARMIENTO, CONSELLERÍA DE EDUCACIÓN, XUNTA DE GALICIA, LUGO, SPAIN
E-mail address: manufl@edu.xunta.es

FACULTY OF MATHEMATICS, UNIVERSITY OF SANTIAGO DE COMPOSTELA, SANTIAGO DE COMPOSTELA, GALICIA, SPAIN
E-mail address: eduardo.garcia.rio@usc.es