

CR STRUCTURES ON OPEN MANIFOLDS

HOWARD JACOBOWITZ AND PETER LANDWEBER

(Communicated by Franc Forstneric)

Dedicated with admiration to André Haefliger

ABSTRACT. We show that the vanishing of the higher dimensional homology groups of a manifold ensures that every almost CR structure of codimension k may be homotoped to a CR structure. This result is proved by adapting a method Haefliger used to study foliations (and previously applied to study the relation between almost complex and complex structures on manifolds) to the case of (almost) CR structures on open manifolds.

1. INTRODUCTION

An almost complex structure on an even dimensional smooth manifold M is a smooth linear vector bundle map J on the tangent bundle of M satisfying $J^2 = -I$. We pass to the complexified tangent bundle and denote by B the bundle of eigenspaces of J with eigenvalue $+i$. So the elements of B correspond to the anti-holomorphic vectors. We have

$$B \cap \overline{B} = \{0\} \text{ and } B \oplus \overline{B} = \mathbf{C} \otimes T(M).$$

These steps are reversible, and in this paper we shall view an almost complex structure as being such a subbundle of the complexified tangent bundle.

Given an almost complex structure B on M , one says that B defines a complex structure on M if in an open neighborhood of each point, complex coordinates may be introduced so that B is spanned by the set of vectors

$$\frac{\partial}{\partial \overline{z}_j}, \quad j = 1, \dots, \frac{1}{2} \dim M.$$

The Newlander-Nirenberg Theorem [21] asserts that B defines a complex structure if and only if B is involutive, by which we mean that the bracket of two smooth sections of B defined on an open subset of M is also a section of B , i.e., the space \mathcal{B} of smooth vector fields with values in B satisfies

$$[\mathcal{B}, \mathcal{B}] \subset \mathcal{B}.$$

More generally, any subbundle of $\mathbf{C} \otimes T(M)$ is said to be involutive if this condition is satisfied.

This, of course, is reminiscent of the Frobenius condition for foliations. It is to be expected that various techniques developed in the theory of foliations would have

Received by the editors September 3, 2014 and, in revised form, December 12, 2014.

2010 *Mathematics Subject Classification*. Primary 32V05; Secondary 55S35.

Key words and phrases. Almost complex, almost CR, Haefliger structure, Gromov's h-principle.

analogues whenever subbundles of $\mathbf{C} \otimes T(M)$ are involutive. In particular, we ask: When does M admit an involutive subbundle B which satisfies $B \cap \overline{B} = \{0\}$ and has a specified complex dimension? For almost complex structures these techniques do extend; see [1], [16]. Using somewhat different techniques from foliation theory, we provide an answer for almost CR structures on manifolds having vanishing homology in high dimensions.

There are two basic steps to our argument. Following Haefliger's work in foliation theory, we broaden the definition of an almost CR structure to a more flexible category (in Section 3) in order to apply (in Section 7) homotopy theory to study the existence of liftings of a continuous map $X \rightarrow B$ to a map $X \rightarrow E$, where $E \rightarrow B$ is a given fibration. This is used to construct a manifold X , foliated by complex manifolds, and a map $f : \mathbf{C} \otimes T(M) \rightarrow T^{1,0}(X)$ with appropriate properties. We then use Gromov's h-principle to prove an analogue of the Gromov-Phillips Theorem in order to replace f by a map $F : M \rightarrow X$ whose differential has these properties (specifically, conditions (1) and (2) of Theorem 4.1) and thereby induces on M the desired CR structure.

For convenience, here is the main result. Definitions are in the next section.

Theorem 1.1. *If*

$$H_p(M^{2n+k}; \mathbf{Z}) = 0 \text{ for } p \geq n + k + 1,$$

then every smooth almost CR structure of codimension k on M is homotopic to a C^ω CR structure of codimension k . In particular, every C^∞ CR structure may be deformed to a C^ω CR structure.

2. THE BASICS

Let M be a manifold of dimension $2n + k$ with $k > 0$. Let N stand for some unspecified manifold of the same dimension, which will often be just an open ball in \mathbf{R}^{2n+k} . All manifolds, bundles, and maps are of class C^∞ , and all manifolds are paracompact, unless otherwise indicated. (It is likely that the results discussed here also hold with only minimal smoothness assumptions.) All open sets which are introduced to state local results are taken to be connected and sufficiently small.

An almost CR structure of codimension k on M is a complex subbundle $B \subset \mathbf{C} \otimes T(M)$ of complex dimension n that satisfies $B \cap \overline{B} = \{0\}$.

A CR structure of codimension k is an almost CR structure B of codimension k that in addition is involutive.

A generic CR immersion is an immersion f of M into a complex manifold X of complex dimension $n + k$ such that $(\mathbf{C} \otimes T(M)) \cap f^*T^{0,1}(X)$ has complex dimension n at all points of M . We set $B = (\mathbf{C} \otimes T(M)) \cap f^*T^{0,1}(X)$ and observe that such an immersion induces a CR structure on M . Conversely, when the CR structure B is given we require that the immersion induces this CR structure.

Remark 1. Results similar to ours should hold for a large class of involutive systems.

Two things are well known:

- Not all CR structures can be obtained by immersions into complex manifolds. That is, there exist smooth CR structures that are not induced by local embeddings into \mathbf{C}^{n+k} . Nirenberg [22, p. 13] gave the first examples of such nonrealizable CR structures.

- Real analytic CR structures are obtained by immersions into \mathbf{C}^{n+k} and there is even an associated uniqueness result. For the convenience of the reader we now state and prove this result.

Lemma 2.1. *If M has a C^ω CR structure B of codimension k , then there exists an open covering*

$$M = \bigcup_j \mathcal{O}_j$$

and C^ω generic CR embeddings

$$f_j : \mathcal{O}_j \rightarrow \mathbf{C}^{n+k}.$$

Further, for each pair (i, j) with $\mathcal{O}_i \cap \mathcal{O}_j \neq \emptyset$ there exists an open set U_{ij} containing $f_i(\mathcal{O}_i \cap \mathcal{O}_j)$ and a biholomorphism $\gamma_{ij} : U_{ji} \rightarrow U_{ij}$ with

$$f_i = \gamma_{ij} \circ f_j \text{ on } \mathcal{O}_i \cap \mathcal{O}_j.$$

Proof. We reduce this lemma to the corresponding result for C^ω almost complex manifolds by applying the following theorem.

Theorem 2.1 ([6], [19]). *Let Q_1, \dots, Q_N be real analytic vector fields in a neighborhood of the origin in \mathbf{R}^{2N} satisfying*

- (1) $Q_1, \dots, Q_N, \overline{Q_1}, \dots, \overline{Q_N}$ span $\mathbf{C} \otimes \mathbf{R}^{2N}$,
- (2) $[Q_i, Q_j]$ is in the linear span of $\{Q_1, \dots, Q_N\}$ for $1 \leq i, j \leq N$.

Then there exist complex coordinates $\{z_1, \dots, z_N\}$ on a possibly smaller neighborhood of the origin such that

$$\{Q_1, \dots, Q_N\} = \{\partial_{z_1}, \dots, \partial_{z_N}\}.$$

To use this theorem to prove the lemma, let L_1, \dots, L_n be real analytic and span the CR bundle B near some point $p \in M$. Choose coordinates

$$x_1, \dots, x_{2n}, t_1, \dots, t_k$$

so that

$$\partial x_1, \dots, \partial x_{2n}$$

span $B \oplus \overline{B}$ at p . First write

$$L_j = \sum_{p=1}^{2n} \alpha_{jp}(x, t) \partial_{x_p} + \sum_{m=1}^k \beta_{jm}(x, t) \partial_{t_m}, \quad j = 1, \dots, n,$$

with coefficients α and β in C^ω and then extend each vector field to \mathbf{R}^{2n+2k} by

$$L_j = \sum_{p=1}^{2n} \alpha_{jp}(x, t + is) \partial_{x_p} + \sum_{m=1}^k \beta_{jm}(x, t + is) \partial_{t_m}.$$

Let \tilde{B} be the linear span of these extended vector fields together with the vector fields

$$\partial_{t_m} + i \partial_{s_m}, \quad m = 1, \dots, k.$$

Then, using the fact that the coefficients are holomorphic in $t + is$, \tilde{B} is involutive and so the above theorem applies:

$$\tilde{B} = \{\partial_{z_1}, \dots, \partial_{z_{n+k}}\}.$$

This gives us an embedding of a neighborhood of p in M into \mathbf{C}^{n+k} with

$$B = (\mathbf{C} \otimes T(M)) \cap T^{0,1}(\mathbf{C}^{n+k}).$$

Thus the embedding is a generic CR embedding.

To find γ_{ij} we assume that $\mathcal{O}_i \cap \mathcal{O}_j$ is not empty. Thus we have a diffeomorphism

$$\Phi : f_i(\mathcal{O}_i \cap \mathcal{O}_j) \rightarrow f_j(\mathcal{O}_i \cap \mathcal{O}_j)$$

with $\Phi_*B = B$. This implies that each component of Φ is a C^ω CR function. That is, for each local section L of B we have

$$L\Phi_j = 0.$$

Such a function on a generic submanifold of a complex manifold is the restriction of a holomorphic function. See, for instance [2, p. 29, Corollary 1.7.13]. We note for later use that the proof of this corollary also shows that the extension is unique.

Let γ_{ij} denote this extension. If γ_{ij} were not a biholomorphism in some neighborhood of $f_i(\mathcal{O}_i \cap \mathcal{O}_j)$, then there would be a nonzero vector

$$v = \sum_{m=1}^{n+k} \alpha_m \partial z_m$$

annihilated by $d\gamma_{ij}$ at some $p \in f_i(\mathcal{O}_i \cap \mathcal{O}_j)$. Since the submanifold is CR generic, $v = T - iJT$ for some $T \in TM|_p$. But since γ_{ij} has been extended holomorphically

$$\begin{aligned} d\gamma v &= d\gamma T - id\gamma JT \\ &= d\gamma T - iJ(d\gamma T). \end{aligned}$$

So if γ_{ij} is not a biholomorphism, $d\gamma_{ij}$ annihilates a nonzero vector in TM and this is impossible because γ_{ij} is a real diffeomorphism on $f_i(\mathcal{O}_i \cap \mathcal{O}_j)$. \square

We note that

$$\gamma_{ik} = \gamma_{ij} \circ \gamma_{jk}$$

on the domain $f_k(\mathcal{O}_i \cap \mathcal{O}_j \cap \mathcal{O}_k)$. To see this let $p \in f_k(\mathcal{O}_i \cap \mathcal{O}_j \cap \mathcal{O}_k)$. Then there exists some $x \in \mathcal{O}_i \cap \mathcal{O}_j \cap \mathcal{O}_k$ with $f_k(x) = p$. Thus

$$\gamma_{ik}(p) = \gamma_{ik} \circ f_k(x) = f_i(x).$$

Further,

$$\gamma_{ij} \circ \gamma_{jk}(p) = \gamma_{ij} \gamma_{jk} \circ f_k(x) = \gamma_{ij} \circ f_j(x) = f_i(x) = \gamma_{ik}(p).$$

By the uniqueness of extensions of holomorphic functions off of generic submanifolds, as referenced in the previous proof, the fact that

$$\gamma_{ik} = \gamma_{ij} \circ \gamma_{jk}$$

holds on $f_k(\mathcal{O}_i \cap \mathcal{O}_j \cap \mathcal{O}_k)$ implies that it holds wherever it makes sense.

3. HAEFLIGER STRUCTURES

Following a well-known procedure in foliation theory, as in [17] or [20] we generalize the objects in Lemma 2.1 by dropping the requirements that the maps are smooth (indeed analytic) and are embeddings. As a reminder, we consider only paracompact manifolds. Let A be some index set.

Definition 1. A Haefliger CR structure of codimension k on M^{2n+k} consists of

- an open covering $M = \bigcup_j \mathcal{O}_j$, $j \in A$,

- continuous maps $f_j : \mathcal{O}_j \rightarrow \mathbf{C}^{n+k}$,
- local biholomorphisms γ_{ij} of \mathbf{C}^{n+k} defined for each pair (i, j) such that $\mathcal{O}_i \cap \mathcal{O}_j \neq \emptyset$ satisfying:
 - (1) $\gamma_{ik} = \gamma_{ij} \circ \gamma_{jk}$ at all points where both sides are defined, and
 - (2) $f_i = \gamma_{ij} \circ f_j$ on $\mathcal{O}_i \cap \mathcal{O}_j$.

Lemma 2.1 shows that a C^ω CR structure admits a Haefliger CR structure of a special kind. Namely, each f_j is an embedding.

Let A_{ij} and B_{ij} denote, respectively, the domain and codomain of γ_{ij} . So γ_{ij} induces a bijection $\gamma_{ij*} : T^{1,0}(A_{ij}) \rightarrow T^{1,0}(B_{ij})$. We may restrict γ_{ij*} to the image of $\mathcal{O}_i \cap \mathcal{O}_j$ and then pull back to M to obtain transition functions g_{ij} that patch $\mathcal{O}_i \times \mathbf{C}^{n+k}$ to $\mathcal{O}_j \times \mathbf{C}^{n+k}$ over $\mathcal{O}_i \cap \mathcal{O}_j$, and so determine a vector bundle:

Definition 2. The normal bundle ν of a Haefliger CR structure is the complex $(n + k)$ -dimensional vector bundle over M with transition functions g_{ij} .

Theorem 3.1. *If B is a C^ω CR structure, then $(\mathbf{C} \otimes T(M))/B$ is isomorphic to the normal bundle of a Haefliger CR structure.*

Proof. By Lemma 2.1, M admits a Haefliger CR structure. Each f_i is a generic CR map, so we have over \mathcal{O}_i ,

$$B \cong f_{i*}B = (\mathbf{C} \otimes T(M)) \cap T^{0,1}.$$

We claim this implies

$$(3.1) \quad ((\mathbf{C} \otimes T(M))/B)|_{\mathcal{O}_i} \cong f_i^*T^{1,0}.$$

To see this, consider the map $\mathbf{C} \otimes T(M) \rightarrow T^{1,0}$ given by, again over \mathcal{O}_i ,

$$\Phi : Y \mapsto f_{i*}Y - iJf_{i*}Y,$$

where J is the standard anti-involution on \mathbf{C}^{n+k} . Note that $\Phi(Y) = 0$ if and only if

$$Y \in (\mathbf{C} \otimes T(M)) \cap T^{0,1} = f_{i*}B.$$

Thus the kernel of Φ has complex dimension n , which implies that Φ is surjective. Then (3.1) follows.

But the transition functions for $T^{1,0}|_{f_i(M)}$ are just the derivative maps g_{ij} . Thus $(\mathbf{C} \otimes T(M))/B$ is isomorphic to the normal bundle of this Haefliger CR structure. □

The same conclusion holds if B is homotopic (even through $2n$ -plane bundles which may not be almost CR) to a C^ω CR structure.

We can now state the first of two basic steps in the proof of our main theorem. Here there is no restriction on the manifold. But for the second step, Theorem 7.1, the topology of the manifold is restricted. In particular, M needs to be open.

Theorem 3.2. *Let B be a smooth almost CR structure of codimension k on M^{2n+k} . If $(\mathbf{C} \otimes T(M))/B$ is isomorphic to the normal bundle of a Haefliger CR structure, then B is homotopic through almost CR structures of codimension k to a C^ω CR structure.*

The next two sections contain the proof of this theorem.

Corollary 1. *Let B be a smooth CR structure. Then B is homotopic to a C^ω CR structure if and only if $(\mathbf{C} \otimes T(M))/B$ is isomorphic to the normal bundle of a Haefliger CR structure.*

It is not known if every smooth CR structure is homotopic to a C^ω CR structure. It is possible that this is true on some manifolds and false on others. (It is certainly true locally.) We shall see in Theorem 8.1 that it is true provided M satisfies some topological restrictions.

4. THE CR EMBEDDING

In this section we reduce the proof of Theorem 3.2 to an h-principle argument which is then provided in the next section.

Lemma 4.1. *Given a Haefliger CR structure on M^{2n+k} there exists a manifold X of real dimension $4n + 3k$ and an embedding $\iota : M \rightarrow X$ such that*

- (1) X is a fiber bundle over M with complex structure on the fibers.
- (2) X admits a foliation \mathcal{F}^{2n+k} transverse to the fibers.
- (3) The normal bundle ν of the Haefliger structure is isomorphic to

$$\iota^*T(X)/T(M).$$

Proof. Let M have local charts

$$\phi_i : \mathcal{O}_i \rightarrow \mathbf{R}^{2n+k}.$$

Set Ω equal to the disjoint union

$$\Omega = \bigsqcup_i (\phi_i(\mathcal{O}_i) \times \mathbf{C}^{n+k}),$$

and define an equivalence relation on Ω by setting

$$(x_i, y_i) \sim (x_j, y_j)$$

whenever

$$(4.1) \quad x_j = \phi_j \phi_i^{-1}(x_i) \quad \text{and}$$

$$(4.2) \quad y_j = \gamma_{ji}(y_i).$$

Here γ_{ji} are the local biholomorphisms of the Haefliger structure. Denote the quotient space, with the quotient topology, by Z . Note that the projection map

$$\Omega \rightarrow Z$$

is an open map and that the projection map

$$(4.3) \quad \pi : Z \rightarrow M$$

is well defined. The multi-valued map of M into Ω

$$p \mapsto \bigsqcup_i (p, f_i(p))$$

for $p \in \mathcal{O}_i$ passes down to a well-defined map

$$\iota : M \rightarrow Z.$$

Clearly, $\pi \circ \iota$ is the identity.

We shall see that Z is Hausdorff near $\iota(M)$. This provides the manifold X . To show that Z is Hausdorff near the image of M let $a = [(x_i, y_i)]$ and $b = [(x_j, y_j)]$ be points of Z . If $x_j \neq \phi_j \phi_i^{-1}(x_i)$, then we can separate a and b by disjoint open sets. So now let $a = [(x_i, y_i)]$ and $b = [(\phi_j \phi_i^{-1}(x_i), y_j)]$. We may assume (by staying close enough to $\iota(M)$) that y_i is in the domain of γ_{ji} . Thus if $y_j \neq \gamma_{ji}(y_i)$ we may again separate a and b while if $y_j = \gamma_{ji}(y_i)$, then $a = b$.

Note that the transition functions for X are

$$(4.4) \quad \psi_{ji}(x, y) = (\phi_j \phi_i^{-1}(x), \gamma_{ji}(y))$$

for $x \in \phi_i(\mathcal{O}_i \cap \mathcal{O}_j)$ and y close enough to $\gamma_{ji}(f_i)$. This implies that the two natural foliations on Ω corresponding to each factor in $\phi_i(\mathcal{O}_i) \times \mathbf{C}^{n+k}$ pass down to X . In particular, the complex structure on each fiber $\pi^{-1}(p)$ is well defined. (Here the map π of (4.3) is restricted to $X \subset Z$.) Thus (1) and (2) of the lemma hold. Further, the transition functions for $T(X)$ are

$$\psi_{ji*}(x, y) = ((\phi_j \phi_i^{-1})_*(x), \gamma_{ji*}(y)).$$

Restricting to $\iota(M)$ we have

$$\begin{aligned} \psi_{ji*}(x) &= ((\phi_j \phi_i^{-1})_*(x), \gamma_{ji*}(f_i(x))) \\ &= ((\phi_j \phi_i^{-1})_*, g_{ji}(f_i(x))). \end{aligned}$$

So the transition functions for the normal bundle of $\iota(M) \subset X$ are the same as those for ν , the normal bundle of the Haefliger structure. This establishes (3). \square

To explain what comes next, we review a well-known procedure in foliation theory. Let \mathcal{M} have a foliation \mathcal{F} of codimension q . If $\Phi : M \rightarrow \mathcal{M}$ is transverse to the leaves of \mathcal{F} , in the sense that $\Phi_*T(M)$ and $T(\mathcal{F})$ together span $T(\mathcal{M})$ at each point of the image of M , then, as is easily seen, \mathcal{F} induces a foliation on M , of the same codimension. By the Gromov-Phillips Theorem (see, for instance, the expository paper *Foliations and foliated vector bundle*, in the collection [20], especially pages 293–297) when M is open, it is enough to have a bundle map

$$\Psi : T(M) \rightarrow T(\mathcal{M})$$

satisfying the transversality condition, for then the base map of Ψ may be deformed to a transverse map $\tilde{\Phi}$.

We rephrase this as: An open manifold M admits a codimension- q foliation provided there exists a vector bundle map Ψ such that the composition

$$T(M) \xrightarrow{\Psi} T(\mathcal{M}) \longrightarrow T(\mathcal{M})/T(\mathcal{F})$$

is surjective, where \mathcal{F} is a codimension- q foliation of \mathcal{M} .

To state the modification of the Gromov-Phillips Theorem we use, we first need some definitions. Let $\pi : X \rightarrow M$ be the projection provided by the lemma. Recall that each fiber $\pi^{-1}(p)$ is a complex manifold. Set

$$T_f = \bigcup_{p \in M} T(\pi^{-1}(p)) \subset T(X)$$

and

$$T_f^{1,0} = \bigcup_{p \in M} T^{1,0}(\pi^{-1}(p)) \subset \mathbf{C} \otimes T(X).$$

The subscript is to remind us that we are looking at tangents to the fiber. The form of the transition functions for X (see (4.4)) shows that the map

$$\mu : \mathbf{C} \otimes T(X) \rightarrow T_f^{1,0}$$

is well defined.

Although the following is patterned upon the Gromov-Phillips Theorem, the restriction to open manifolds is not necessary here.

Theorem 4.1. *If there exists a surjective bundle map*

$$\Psi : \mathbf{C} \otimes T(M) \rightarrow T_f^{1,0}|_M,$$

then there exists a smooth map

$$F : M \rightarrow X$$

such that

(1) *The composition*

$$\mu F_* : \mathbf{C} \otimes T(M) \rightarrow \mathbf{C} \otimes T(X) \rightarrow T_f^{1,0}$$

is surjective.

(2) $\mu_* F_*$ *is injective when restricted to* $T(M)$.

(3) $B_1 = \ker \mu F_*$ *is homotopic to* $B = \ker \Psi$.

This is the heart of the proof of Theorem 3.2. We prove it in the next section by means of Gromov’s h-principle. Here let us first see how it implies the conclusion of Theorem 3.2 and then how we use the hypothesis

$$(\mathbf{C} \otimes T(M))/B \cong \nu$$

to obtain the surjective map

$$\Psi : \mathbf{C} \otimes T(M) \rightarrow T_f^{1,0}|_M.$$

We first show that B_1 defines a CR structure.

Lemma 4.2. *The bundle* B_1 *is involutive.*

Proof. It suffices to assume that a and b are local sections of B_1 and to show that

$$\mu F_*[a, b] = 0.$$

We work in local coordinates. So we write

$$F(x) = (f(x), y(x)) \in \phi_i(\mathcal{O}) \times \mathbf{C}^{n+k}.$$

For any $a \in \mathbf{C} \otimes T(M)$, we have (note that we now use $a(y)$ to mean the action of the vector a on the function y and we employ the summation convention)

$$F_*(a) = a(f_j)\partial_{x_j} + a(y_k)\partial_{z_k} + a(\overline{y_k})\partial_{\overline{z_k}}$$

and

$$\mu F_* = a(y_k)\partial_{z_k}.$$

It follows that

$$B_1 = \ker \mu F_* = \{a \in \mathbf{C} \otimes T(M) : a(y_k) = 0 \text{ for } k = 1, \dots, n + 1\}$$

and this is involutive. □

Lemma 4.3.

$$B_1 \cap \overline{B_1} = \{0\}.$$

Proof. This is a consequence of the second condition of the theorem. □

Thus B_1 is a CR structure homotopic to B . (The proof of Theorem 4.1 will show that this homotopy is through almost CR structures.)

Finally, we need the map Ψ .

Lemma 4.4. *If $(\mathbf{C} \otimes T(M))/B$ is isomorphic to the Haefliger normal bundle ν , then there exists a surjective bundle map*

$$\Psi : \mathbf{C} \otimes T(M) \rightarrow T_f^{1,0}|_M$$

with $B = \ker \Psi$.

Proof. We have

$$\mathbf{C} \otimes T(M) \rightarrow (\mathbf{C} \otimes T(M))/B \xrightarrow{\alpha} \nu \xrightarrow{\beta} T_f^{1,0}|_M.$$

Since α and β are isomorphisms, the composite map Ψ is surjective and has kernel B . □

5. AN APPLICATION OF THE H-PRINCIPLE

The general set-up is to define a subset $\mathcal{R} \subset J^1(M, X)$, called a relation, with the property that $j^1(F) \in \mathcal{R}$ implies that F satisfies the conclusions of Theorem 4.1. See [8], [11], [14] for discussions of the h-principle.

Again choosing local coordinates $x \in \mathbf{R}^{2n+k}$ for M , $y \in \mathbf{R}^{2n+k}$, and $z \in \mathbf{C}^{n+k}$ for ν we see that

$$\mu F_* : \mathbf{C} \otimes T(M) \rightarrow T_f^{1,0}$$

is surjective if the rank of the matrix

$$\left(\frac{\partial z_i}{\partial x_j} \right), \quad 1 \leq i \leq n+k \text{ and } 1 \leq j \leq 2n+k,$$

is maximal (i.e., equal to $n+k$) and that

$$\mu F_* : T(M) \rightarrow T_f^{1,0}$$

is injective if the rank of the $(2n+2k) \times (2n+k)$ matrix

$$\begin{pmatrix} \frac{\partial z_i}{\partial x_j} \\ \frac{\partial \bar{z}_i}{\partial x_j} \end{pmatrix}$$

is also maximal.

So we define our relation by

$$\mathcal{R} = \{(p, c, a_i^j) : \text{rank}(a) = n+k, \quad \text{rank} \begin{pmatrix} a \\ \bar{a} \end{pmatrix} = 2n+k\}$$

where, as above

$$1 \leq i \leq n+k, \quad 1 \leq j \leq 2n+k.$$

We claim that \mathcal{R} is an open and ample subset of $J^1(M, X)$. For the definition of *ample* see any of the above cited works or [15]; for the proof of the claim see [15, pp. 157–159]. It is a consequence of the h-principle that the desired F exists provided we can find a section

$$\sigma : M \rightarrow \mathcal{R}.$$

That is, the existence of a formal solution implies the existence of a genuine solution.

To see that we do have a formal solution, we identify a bundle map

$$\sigma : \mathbf{C} \otimes T(M) \rightarrow T_f^{1,0}$$

with a section

$$\tilde{\sigma} : M \rightarrow J^1(M, X)$$

by writing

$$\sigma(\partial_{x_j}|_p) = h_i^j \partial_{z_i}|_{\sigma(p)}$$

and then setting

$$\tilde{\sigma}(p) = (p, \sigma(p), h_i^j).$$

As long as σ is surjective and its kernel satisfies $B \cap \overline{B} = \{0\}$, so

$$\sigma : T(M) \rightarrow T_f^{1,0}$$

is injective, $\tilde{\sigma}$ is a formal solution,

$$\tilde{\sigma} : M \rightarrow \mathcal{R} \subset J^1(M, X).$$

In this context, Lemma 4.4 asserts the existence of a formal solution.

The h-principle asserts more than the existence of a genuine solution. In fact, the genuine solution is obtained by a deformation

$$\sigma_t : M \rightarrow \mathcal{R}, \quad 0 \leq t \leq 1,$$

such that $\sigma_0 = \tilde{\sigma}$ and $\sigma_1 = j^1(F)$. Now set

$$B_t = \ker \mu\sigma_t.$$

We have $\text{rank } B_t = n$ and $B_t \cap \overline{B_t} = \{0\}$ for all t , and moreover B_1 is involutive.

Thus B_t represents a deformation of B through almost CR structures to a true CR structure.

Finally, once we have $F : M \rightarrow X$ we may deform F , maintaining the open conditions on the ranks of the matrices, to a C^ω map.

The proof of Theorem 3.2 is complete.

6. CLASSIFYING SPACES

We first note that the definition of a Haefliger CR structure makes sense (for fixed n and k) if M is merely a topological space. To specify n and k we speak of an (n, k) Haefliger CR structure. (Actually, only $n + k$ appears in the definition, so we occasionally speak of an $(n + k)$ Haefliger structure.) Also if X and Y are topological spaces and Y has an (n, k) Haefliger CR structure, then any continuous map $\phi : X \rightarrow Y$ induces an (n, k) Haefliger CR structure on X .

Definition 3. Two (n, k) Haefliger CR structures on X are homotopic if there exists an (n, k) Haefliger CR structure on $X \times I$ which pulls back to the given structures on $X \times \{0\}$ and $X \times \{1\}$.

In the following, we assume that all Haefliger CR structures are on paracompact spaces, and thus the classifying spaces for these structures are also paracompact.

Definition 4. A topological space $\mathcal{B}_{n,k}$ with an (n, k) Haefliger CR structure is a classifying space for (n, k) Haefliger CR structures if:

- Every (n, k) Haefliger CR structure on a manifold N is induced by a continuous map $\phi : N \rightarrow \mathcal{B}_{n,k}$.
- Two (n, k) Haefliger CR structures on N are homotopic if and only if the corresponding continuous maps into $\mathcal{B}_{n,k}$ are homotopic.

The associated $(n + k)$ -dimensional normal bundle over $\mathcal{B}_{n,k}$ is denoted by $\nu_{n,k}$.

Theorem 6.1. *There exists a classifying space for (n, k) Haefliger CR structures.*

This result is a particular case of the existence of classifying spaces for topological groupoid structures (as proved by Haefliger [12], [13] and others—see [18, p. 48] or [20, p. 312] for expository accounts). Indeed, these are precisely the classifying spaces for complex structures studied by Landweber [16] and by Adachi [1] and usually denoted by $B\Gamma_q^{\mathbf{C}}$. In the present case, the codimension q is taken to be $n + k$. Here $\Gamma_q^{\mathbf{C}}$ denotes the topological groupoid of germs of biholomorphisms between open subsets of \mathbf{C}^q .

7. THE LIFTING PROBLEM

We have started with a complex subbundle $B \subset \mathbf{C} \otimes T(M)$ and have set $\nu = (\mathbf{C} \otimes T(M))/B$. Using a Hermitian metric on $\mathbf{C} \otimes T(M)$ we may also consider ν as a subbundle of $\mathbf{C} \otimes T(M)$ complementary to B in $\mathbf{C} \otimes T(M)$. Our aim is to show that the subbundle $\nu \subset \mathbf{C} \otimes T(M)$ of rank $n + k$ is the normal bundle of some $(n + k)$ Haefliger structure on M . Since bundles over M are classified by homotopy classes of maps of M into $BGL(n + k)$ (we denote by $GL(m)$ the complex general linear group $GL(m, \mathbf{C})$) and Haefliger structures are classified by homotopy classes of maps of M into $\mathcal{B}_{n,k}$, the following set-up is natural. For any complex vector bundle ξ of rank m over a paracompact space X , let $\text{cl}(\xi) : X \rightarrow BGL(m)$ denote a classifying map for ξ . Thus we have the commutative diagram

$$\begin{array}{ccc}
 M & \xrightarrow{\text{cl}(\nu) \times \text{cl}(B)} & BGL(n + k) \times BGL(n) \\
 & \searrow \text{cl}(\mathbf{C} \otimes T(M)) & \downarrow B(\iota) \\
 & & BGL(2n + k)
 \end{array}$$

where ι denotes the inclusion of $GL(n + k) \times GL(n)$ into $GL(2n + k)$ by means of block matrices and $B(\iota)$ is the map of classifying spaces induced by this inclusion.

As above, let $\nu_{n,k}$ denote the normal bundle on the classifying space $\mathcal{B}_{n,k}$. Note that $\nu_{n,k}$ defines a map $\text{cl}(\nu_{n,k}) : \mathcal{B}_{n,k} \rightarrow BGL(n + k)$, unique up to homotopy.

Theorem 7.1. *If*

$$(7.1) \quad H_p(M^{2n+k}; \mathbf{Z}) = 0 \text{ for } p \geq n + k + 1,$$

then there exists a lift

$$\begin{array}{ccc}
 & & \mathcal{B}_{n,k} \times BGL(n) \\
 & \nearrow & \downarrow \text{cl}(\nu_{n,k}) \times B(\text{id}) \\
 M & \xrightarrow[\text{cl}(\nu) \times \text{cl}(B)]{} & BGL(n + k) \times BGL(n).
 \end{array}$$

Proof. The homotopy fiber $\mathcal{F} = \mathcal{F}_{n,k}$ of the map $\text{cl}(\nu_{n,k}) : \mathcal{B}_{n,k} \rightarrow BGL(n + k)$ is $(n + k)$ -connected (cf. [1] and [16], although the latter paper needs to be supplemented by Gromov’s h-principle for totally real immersions, as given in [10] and [11]). Since the obstructions to lifting a map

$$(7.2) \quad M \rightarrow BGL(n + k) \times BGL(n)$$

lie in $H^{j+1}(M; \{\pi_j(\mathcal{F})\})$ for the local coefficient system $\{\pi_j(\mathcal{F})\}$, and since $\pi_j(\mathcal{F}) = 0$ for $0 \leq j \leq n + k$, the nonzero obstructions can lie only in $H^{j+1}(M; \{\pi_j(\mathcal{F})\})$

for $j \geq n + k + 1$. (For discussion of local coefficient systems, also known as bundles of coefficients or systems of local groups, we refer to Steenrod’s book [25] on fiber bundles, or to his paper [24]. For a more recent account dealing with fibrations, see Whitehead’s book [26, Chapter VI].) Since complex general linear groups are connected, their classifying spaces are simply connected, and so the map $\text{cl}(\nu_{n,k}) : \mathcal{B}_{n,k} \rightarrow BGL(n + k)$, which we view as a fibration, has simple local coefficient systems in the sense of [24]. This means that there is a unique isomorphism between the homotopy groups of any two fibers. It follows that the local coefficient systems over M appearing here are also simple. As a result, the cohomology groups here are ordinary cohomology groups and we can apply the usual universal coefficient theorem which expresses cohomology in terms of integral homology (note that homotopy groups beyond the first one are abelian and so the universal coefficient theorem applies). Hence if (7.1) holds, then by the universal coefficient theorem we also have

$$H^{j+1}(M; \{\pi_j(\mathcal{F})\}) = 0 \text{ for } j \geq n + k + 1,$$

and so every map (7.2) lifts to a map

$$M \rightarrow \mathcal{B}_{n,k} \times BGL(n).$$

□

Remark 2. The hypothesis of Theorem 7.1 can be slightly weakened, namely, it suffices to require that the integral homology groups in (7.1) vanish for $p \geq n + k + 2$, and be free abelian for $p = n + k + 1$. In particular, this weakened hypothesis is satisfied if the manifold M^{2n+k} has the homotopy type of a CW complex of dimension at most $n + k + 1$. This last condition is satisfied if M^{2n+k} has the homotopy type of an open $(n + k + 2)$ -manifold, which follows from [23, Lemma 1]. In short, if M^{2n+k} is homotopic to an open $(n + k + 2)$ -manifold, then the lift exists and Theorem 1.1 holds in this slightly more general case.

8. THE THEOREM

Theorem 8.1. *If M^{2n+k} satisfies the condition (7.1) of Theorem 7.1, then every smooth almost CR structure of codimension k on M is homotopic to a C^ω CR structure of codimension k . In particular, every C^∞ CR structure may be deformed to a C^ω CR structure.*

Proof. The lifting $M \rightarrow \mathcal{B}_{n,k} \times BGL(n)$ of Theorem 7.1 shows that $\nu \oplus B$ is isomorphic to some bundle $\nu^0 \oplus B^0$ with ν^0 the normal bundle of some (n, k) Haefliger structure on M and B^0 a complex bundle of rank n . The map $\text{cl}(\nu) \times \text{cl}(B)$ is homotopic to the map $\text{cl}(\mathbf{C} \otimes T(M))$ and so, using the fact that the diagram is commutative, we see that

$$\nu^0 \oplus B^0 \cong \mathbf{C} \otimes T(M).$$

In particular, ν^0 and B^0 can be identified with subbundles of $\mathbf{C} \otimes T(M)$. Further, since the map $\text{cl}(\nu_{n,k}) \times B(\text{id})$ is the identity on the $BGL(n)$ factor, we have that ν is isomorphic to ν^0 . Thus Theorem 3.2 may be applied to show that B may be deformed to some CR structure. Further, since this CR structure is induced by a map $F : M \rightarrow X$, as in Theorem 4.1, the map F itself may be deformed to yield a C^ω CR structure. This last step just uses that the real analytic functions are dense in the set of smooth functions, by the Whitney approximation theorem. □

Note that the requirement that the integral homology of M vanish in high dimensions excludes compact manifolds, provided that $n > 1$. In the case $n = 1$ the theorem holds for a trivial reason: Each almost CR structure with $n = 1$ is automatically involutive.

In view of the analogy between foliations and involutive structures it is natural to try to place into this new context Bott's necessary condition for a subbundle of TM to be homotopic to a foliation. See [3], [4], or [5] for Bott's original argument and some of its further consequences.

A version of Theorem 8.1 also holds for $k = 0$, as in [1] and [16], but our proof does not. The reason for this, in a very similar case, is explained on page 160 of [15].

Actually, for almost complex structures the conclusion of Theorem 8.1 may be strengthened with a slight weakening of the hypothesis. The existence of a J -convex proper Morse function on M^{2n} , $n > 2$, with no critical point of index greater than n implies that the almost complex structure may be deformed in such a way that the Morse function becomes a plurisubharmonic exhaustion function, and so the new structure is Stein. See [7] and [9]. (Analogous results in the CR case, at least for $k = 1$, might be possible.) Note that the existence of a proper Morse function with no critical points of index greater than q implies $H_p(M, \mathbf{Z}) = 0$ for $p > q$. So the existence of the postulated Morse function in [7] and [9] implies (7.1). The reverse implication is in general false; the vanishing of higher homology groups does not imply the existence of a Morse function without critical points of high index. We are unable to determine whether in the present case, that is in the presence of an almost complex structure (even one homotopic to an integrable structure) and with q equal to half the dimension of M , this reverse implication does hold.

REFERENCES

- [1] Masahisa Adachi, *Embeddings and immersions*, Translations of Mathematical Monographs, vol. 124, American Mathematical Society, Providence, RI, 1993. Translated from the 1984 Japanese original by Kiki Hudson. MR1225100 (95a:57039)
- [2] M. Salah Baouendi, Peter Ebenfelt, and Linda Preiss Rothschild, *Real submanifolds in complex space and their mappings*, Princeton Mathematical Series, vol. 47, Princeton University Press, Princeton, NJ, 1999. MR1668103 (2000b:32066)
- [3] Raoul Bott, *On a topological obstruction to integrability*, Global Analysis (Proc. Sympos. Pure Math., Vol. XVI, Berkeley, Calif., 1968), Amer. Math. Soc., Providence, R.I., 1970, pp. 127–131. MR0266248 (42 #1155)
- [4] Raoul Bott, *On topological obstructions to integrability*, Actes du Congrès International des Mathématiciens (Nice, 1970), Gauthier-Villars, Paris, 1971, pp. 27–36. MR0425983 (54 #13932)
- [5] Raoul Bott, *Lectures on characteristic classes and foliations*, Lectures on algebraic and differential topology (Second Latin American School in Math., Mexico City, 1971), Springer, Berlin, 1972, pp. 1–94. Lecture Notes in Math., Vol. 279. Notes by Lawrence Conlon, with two appendices by J. Stasheff. MR0362335 (50 #14777)
- [6] Beno Eckmann and Alfred Frölicher, *Sur l'intégrabilité des structures presque complexes* (French), C. R. Acad. Sci. Paris **232** (1951), 2284–2286. MR0042202 (13,75f)
- [7] Yakov Eliashberg, *Topological characterization of Stein manifolds of dimension > 2* , Internat. J. Math. **1** (1990), no. 1, 29–46, DOI 10.1142/S0129167X90000034. MR1044658 (91k:32012)
- [8] Y. Eliashberg and N. Mishachev, *Introduction to the h -principle*, Graduate Studies in Mathematics, vol. 48, American Mathematical Society, Providence, RI, 2002. MR1909245 (2003g:53164)
- [9] Franc Forstnerič and Marko Slapar, *Stein structures and holomorphic mappings*, Math. Z. **256** (2007), no. 3, 615–646, DOI 10.1007/s00209-006-0093-0. MR2299574 (2008g:32036)

- [10] M. L. Gromov, *Convex integration of differential relations. I* (Russian), *Izv. Akad. Nauk SSSR Ser. Mat.* **37** (1973), 329–343. MR0413206 (54 #1323)
- [11] Mikhael Gromov, *Partial differential relations*, *Ergebnisse der Mathematik und ihrer Grenzgebiete (3)* [Results in Mathematics and Related Areas (3)], vol. 9, Springer-Verlag, Berlin, 1986. MR864505 (90a:58201)
- [12] A. Haefliger, *Feuilletages sur les variétés ouvertes* (French), *Topology* **9** (1970), 183–194. MR0263104 (41 #7709)
- [13] André Haefliger, *Homotopy and integrability*, *Manifolds—Amsterdam 1970* (Proc. Nuffic Summer School), *Lecture Notes in Mathematics*, Vol. 197, Springer, Berlin, 1971, pp. 133–163. MR0285027 (44 #2251)
- [14] H. Jacobowitz, *Convex integration and the h-principle*. *Lecture Notes Series*, 55. Seoul National University, Seoul, 2011.
- [15] Howard Jacobowitz and Peter Landweber, *Manifolds admitting generic immersions into \mathbb{C}^N* , *Asian J. Math.* **11** (2007), no. 1, 151–165, DOI 10.4310/AJM.2007.v11.n1.a14. MR2304588 (2008f:32044)
- [16] Peter S. Landweber, *Complex structures on open manifolds*, *Topology* **13** (1974), 69–75. MR0339210 (49 #3973)
- [17] H. Blaine Lawson Jr., *Foliations*, *Bull. Amer. Math. Soc.* **80** (1974), 369–418. MR0343289 (49 #8031)
- [18] H. Blaine Lawson Jr., *The quantitative theory of foliations*, American Mathematical Society, Providence, R. I., 1977. Expository lectures from the CBMS Regional Conference held at Washington University, St. Louis, MO, January 6–10, 1975; Conference Board of the Mathematical Sciences Regional Conference Series in Mathematics, No. 27. MR0448368 (56 #6675)
- [19] Paulette Libermann, *Problèmes d'équivalence relatifs à une structure presque complexe sur une variété à quatre dimensions* (French), *Acad. Roy. Belgique. Bull. Cl. Sci. (5)* **36** (1950), 742–755. MR0040791 (12,749i)
- [20] John Milnor, *Collected papers of John Milnor. IV*, American Mathematical Society, Providence, RI, 2009. Homotopy, homology and manifolds; Edited by John McCleary. MR2590677 (2011g:01020)
- [21] A. Newlander and L. Nirenberg, *Complex analytic coordinates in almost complex manifolds*, *Ann. of Math. (2)* **65** (1957), 391–404. MR0088770 (19,577a)
- [22] Louis Nirenberg, *Lectures on linear partial differential equations*, American Mathematical Society, Providence, R.I., 1973. Expository Lectures from the CBMS Regional Conference held at the Texas Technological University, Lubbock, TX, May 22–26, 1972; Conference Board of the Mathematical Sciences Regional Conference Series in Mathematics, No. 17. MR0450755 (56 #9048)
- [23] Anthony Phillips, *Submersions of open manifolds*, *Topology* **6** (1967), 171–206. MR0208611 (34 #8420)
- [24] N. E. Steenrod, *Homology with local coefficients*, *Ann. of Math. (2)* **44** (1943), 610–627. MR0009114 (5,104f)
- [25] Norman Steenrod, *The Topology of Fibre Bundles*, *Princeton Mathematical Series*, vol. 14, Princeton University Press, Princeton, N. J., 1951. MR0039258 (12,522b)
- [26] George W. Whitehead, *Elements of homotopy theory*, *Graduate Texts in Mathematics*, vol. 61, Springer-Verlag, New York-Berlin, 1978. MR516508 (80b:55001)

DEPARTMENT OF MATHEMATICAL SCIENCES, RUTGERS UNIVERSITY, CAMDEN, NEW JERSEY 08102

E-mail address: jacobowi@camden.rutgers.edu

DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, PISCATAWAY, NEW JERSEY 08854

E-mail address: landwebe@math.rutgers.edu