

CLOSED IDEALS OF OPERATORS ON AND COMPLEMENTED SUBSPACES OF BANACH SPACES OF FUNCTIONS WITH COUNTABLE SUPPORT

WILLIAM B. JOHNSON, TOMASZ KANIA, AND GIDEON SCHECHTMAN

(Communicated by Thomas Schlumprecht)

ABSTRACT. Let λ be an infinite cardinal number and let $\ell_\infty^c(\lambda)$ denote the subspace of $\ell_\infty(\lambda)$ consisting of all functions that assume at most countably many non-zero values. We classify all infinite-dimensional complemented subspaces of $\ell_\infty^c(\lambda)$, proving that they are isomorphic to $\ell_\infty^c(\kappa)$ for some cardinal number κ . Then we show that the Banach algebra of all bounded linear operators on $\ell_\infty^c(\lambda)$ or $\ell_\infty(\lambda)$ has the unique maximal ideal consisting of operators through which the identity operator does not factor. Using similar techniques, we obtain an alternative to Daws' approach description of the lattice of all closed ideals of $\mathcal{B}(X)$, where $X = c_0(\lambda)$ or $X = \ell_p(\lambda)$ for some $p \in [1, \infty)$, and we classify the closed ideals of $\mathcal{B}(\ell_\infty^c(\lambda))$ that contains the ideal of weakly compact operators.

1. INTRODUCTION AND THE STATEMENTS OF THE MAIN RESULTS

The aim of this paper is to contribute to the two closely related programs of classifying complemented subspaces of Banach spaces and classifying the maximal ideals of (bounded, linear) operators acting thereon. Our results constitute a natural continuation of the research undertaken by the authors in [7, 14–16] concerning the maximal ideals of the Banach algebras $\mathcal{B}(X)$ for certain classical Banach spaces X . From this perspective, we complement results ([9, 22]) of Gramsch and Luft, who classified all closed ideals of operators acting on Hilbert spaces, and a result ([4]) of Daws, who classified all closed ideals of operators acting on $c_0(\lambda)$ and $\ell_p(\lambda)$ for λ uncountable and $p \in [1, \infty)$. Nevertheless, bearing in mind that the similar problem of classifying all closed ideals of operators acting on $\ell_\infty(\lambda)$ is most likely intractable (as this would require, in particular, understanding *all* injective Banach spaces), we offer instead a complete description of the maximal ideals for the space

Received by the editors February 11, 2015 and, in revised form, December 10, 2015 and January 5, 2016.

2010 *Mathematics Subject Classification*. Primary 46H10, 47B38, 47L10; Secondary 06F30, 46B26, 47L20.

The first-named author was supported in part by NSF DMS-1301604 and the U.S.-Israel Binational Science Foundation.

The second-named author was supported by the Warsaw Centre of Mathematics and Computer Science.

The third-named author was supported in part by the U.S.-Israel Binational Science Foundation.

Part of this work was done when the third-named author was participating in the NSF Workshop in Analysis and Probability held at Texas A&M University.

$\ell_\infty(\lambda)$ and its closed subspace

$$\ell_\infty^c(\lambda) = \{z \in \ell_\infty(\lambda) : z(\alpha) \neq 0 \text{ for at most countably many } \alpha < \lambda\}.$$

(For consistency of notation we sometimes denote ℓ_∞ by $\ell_\infty^c(\omega)$.) Our first result then reads as follows.

Theorem 1.1. *Let λ be an infinite cardinal number. Suppose that X is either $\ell_\infty(\lambda)$ or $\ell_\infty^c(\lambda)$. Then*

$$\begin{aligned} \mathcal{I}_X(X) &= \{T \in \mathcal{B}(X) : T \text{ is not bounded below on any copy of } X\} \\ &= \{T \in \mathcal{B}(X) : I_X \neq ATB \text{ for all } A, B \in \mathcal{B}(X)\} \end{aligned}$$

is the unique maximal ideal of $\mathcal{B}(X)$.

(We explain the required notation and terminology in the next section, while proofs are postponed to the final section.)

Actually, we are able to describe all the closed ideals of $\ell_\infty^c(\lambda)$ containing the ideal of weakly compact operators—they consist precisely of operators that do not preserve copies of $\ell_\infty^c(\kappa)$ for some κ not exceeding the cardinal successor of λ .

Theorem 1.2. *Let λ be an infinite cardinal number. Then every closed ideal of $\mathcal{B}(\ell_\infty^c(\lambda))$ that contains the ideal of weakly compact operators is equal to $\mathcal{I}_{\ell_\infty^c(\kappa)}(\ell_\infty^c(\lambda))$ for some infinite cardinal number $\kappa \leq \lambda^+$.*

Remark. One cannot hope that a statement similar to Theorems 1.5 or 1.2 holds for the lattice of closed ideals of $\mathcal{B}(\ell_\infty(\lambda))$. Indeed, in general the lattice of closed ideals of $\mathcal{B}(\ell_\infty(\lambda))$ need not be linearly ordered. Let $E_1 = L_\infty(\{0, 1\}^{\omega_2})$ and $E_2 = \ell_\infty(\omega_1)$. Then both E_1 and E_2 embed into $\ell_\infty(\omega_2)$, but E_1 does not embed into E_2 and E_2 does not embed into E_1 ([28, Theorem 1.7]). Consider the (not necessarily closed) ideals \mathcal{I}_i consisting of operators that admit a factorisation through E_i ($i = 1, 2$). These sets are obviously closed under compositions from left and right, and, since $E_i \cong E_i \oplus E_i$ ($i = 1, 2$), they are also closed under addition, so they are indeed ideals of $\mathcal{B}(\ell_\infty(\omega_2))$. We claim that the closures of \mathcal{I}_i ($i = 1, 2$) are incomparable. Indeed, let P_i be a projection on $\ell_\infty(\omega_2)$ with range isomorphic to E_i ($i = 1, 2$), and suppose that $P_1 \in \overline{\mathcal{I}_2}$. Since $P_1^2 = P_1$, we actually have $P_1 \in \mathcal{I}_2$ (see [19, Lemma 2.7]). This, however, is impossible because it would imply that the range of P_1 is isomorphic to a subspace of E_2 . The other case is symmetric.

This observation shows even more: there are 1-injective Banach spaces E of the form $L_\infty(\mu)$ for which $\mathcal{B}(E)$ has more than one maximal ideal. It is readily seen that $E = E_1 \oplus E_2$ is such an example. Here $E = L_\infty(\mu)$, where μ is the direct product measure of the Haar measure on $\{0, 1\}^{\omega_2}$ with the counting measure on ω_1 .

A key result in the proof of Theorem 1.1 is the following theorem, which, we believe, is of interest in itself. In order to state it, we require a piece of terminology.

A Banach space X is *complementably homogeneous* if for each closed subspace Y of X such that Y is isomorphic to X , there exists a closed, complemented subspace Z of X such that Z is contained in Y and Z is isomorphic to X . That $\ell_1(\lambda)$ is complementably homogeneous follows directly from Lemma 2.1(ii). The Banach spaces $c_0(\lambda)$ have actually a stronger property: every isomorphic copy of $c_0(\lambda)$ in $c_0(\lambda)$ is complemented ([2, Proposition 2.8]). For $p \in (1, \infty)$ complementable homogeneity of $\ell_p(\lambda)$ is also well known and easy to prove; we shall include a proof of that fact (Proposition 3.10) for the sake of completeness.

Theorem 1.3. *For each infinite cardinal number λ , the Banach space $\ell_\infty^c(\lambda)$ is complementably homogeneous.*

We then employ Theorem 1.3 to characterise, up to isomorphism, all infinite-dimensional complemented subspaces of $\ell_\infty^c(\lambda)$, where λ is any uncountable cardinal number.

Theorem 1.4. *Let λ be an infinite cardinal number. Then every infinite-dimensional, complemented subspace of $\ell_\infty^c(\lambda)$ is isomorphic either to ℓ_∞ or to $\ell_\infty^c(\kappa)$ for some cardinal $\kappa \leq \lambda$. Consequently, $\ell_\infty^c(\lambda)$ is a primary Banach space.*

Let us remark that $\ell_\infty^c(\lambda)$ is a hyperplane in the space $E := \text{span}\{\mathbf{1}_\lambda, \ell_\infty^c(\lambda)\}$. The space E in turn is easily seen to be isomorphic to its hyperplanes as it contains ℓ_∞ . More importantly, E is a unital sub-C*-algebra of $\ell_\infty(\lambda)$ (the operations are pointwise), hence by the Gelfand–Naimark duality, $E \cong C(K)$ for some compact Hausdorff space K . The space E has one more incarnation—it is isometrically isomorphic to the space $L_\infty(\mu)$, where μ is the counting measure on λ restricted to the σ -field of sets that are countable or have countable complement. The space $L_\infty(\mu)$ is not isomorphic to a complemented subspace of a dual space. Indeed, if it were isomorphic to a complemented subspace of a dual space, then it would be complemented in its second dual, which is an injective space, so $L_\infty(\mu)$ would be injective. However, it is an observation of Pełczyński and Sudakov ([24]) that $\ell_\infty^c(\lambda)$ (λ uncountable) is not complemented in $\ell_\infty(\lambda)$, so it is not injective.

It is noteworthy that there are very few kinds of $C(K)$ -spaces with completely understood complemented subspaces. To the best of our knowledge, these are:

- $c_0(\lambda)$ for any set λ ; these spaces are isomorphic to $C(\lambda^1)$ where λ^1 is the one-point compactification of the discrete set λ . Every infinite-dimensional, complemented subspace of $c_0(\lambda)$ is isomorphic to $c_0(\kappa)$ for some cardinal number $\kappa \leq \lambda$ (see [10] or [2, Proposition 2.8]).
- $C[0, \omega^\omega]$, where $[0, \omega^\omega]$ carries the order topology. Every infinite-dimensional, complemented subspace of $C[0, \omega^\omega]$ is isomorphic either to c_0 or to $C[0, \omega^\omega]$ ([3, Theorem 3]).

No further separable examples—apart from c_0 and $C[0, \omega^\omega]$ —of infinite-dimensional $C(K)$ -spaces with completely understood complemented subspaces are known.

- $C(\beta\mathbb{N}) \cong \ell_\infty$. Every infinite-dimensional, complemented subspace of ℓ_∞ is isomorphic to ℓ_∞ ; this is Lindenstrauss' theorem ([20]).
- $C(K)$, where K is the compact scattered space constructed by Koszmider under the Continuum Hypothesis ([17]); it is the Stone space of a Boolean algebra generated by a certain almost disjoint family of subsets of \mathbb{N} . Every infinite-dimensional, complemented subspace of $C(K)$ is isomorphic either to c_0 or $C(K)$.
- Spaces of the form $C(K \sqcup L) \cong C(K) \oplus C(L)$, where K is scattered, $C(L)$ is a Grothendieck space (equivalently, every operator from $C(L)$ to a separable space is weakly compact; for the definition and basic properties of Grothendieck spaces see, e.g., [5, p. 150]), and all the complemented subspaces of $C(K)$ and $C(L)$ are classified. (So far, for K we can take the one-point compactification of a discrete set, $[0, \omega^\omega]$, or the above-mentioned space constructed by Koszmider; and for L we can take $\beta\mathbb{N}$, so that $C(L)$ is isometrically isomorphic to ℓ_∞ , a prototypical example of a Grothendieck

space; see [11].) Each complemented subspace of $C(K) \oplus C(L)$ is isomorphic to a space of the form $X \oplus Y$, where X is a complemented subspace of $C(K)$ and Y is complemented in $C(L)$.

Indeed, if K is scattered, then $C(K)^* \cong \ell_1(|K|)$, so it follows from [12, Corollary 2] that the unit ball of $C(K)^*$ is weak*-sequentially compact. Applying [5, Corollary 5 on p. 150], we infer that every operator $T: C(L) \rightarrow C(K)$ is weakly compact and hence strictly singular by [23]. It follows then from [30, Theorem 1.1] that every complemented subspace of $C(K) \oplus C(L)$ is of the desired form.

We therefore extend the above list by adding the following classes of examples:

- $\ell_\infty^c(\lambda)$, for any uncountable cardinal number λ ,
- $\ell_\infty^c(\lambda) \oplus C(K)$, where $C(K)$ is isomorphic to one of the spaces $c_0(\kappa)$, $C[0, \omega^\omega]$ or the above-mentioned space constructed by Koszmider. Indeed, for any cardinal number λ , $\ell_\infty^c(\lambda)$ is a Grothendieck space (see Proposition 3.7 below), so we can use the antepenultimate clause to deduce the claim.

As a by-product of our investigations, we give an alternative approach to Daws' description of all closed ideals of $\mathcal{B}(c_0(\lambda))$ and $\mathcal{B}(\ell_p(\lambda))$ for λ uncountable and $p \in [1, \infty)$.

Theorem 1.5. *Let λ be an infinite cardinal number and let $p \in [1, \infty)$. Then every non-zero, proper closed ideal of $\mathcal{B}(c_0(\lambda))$ and $\mathcal{B}(\ell_p(\lambda))$ is of the form $\mathcal{S}_{c_0(\kappa)}(c_0(\lambda))$ and $\mathcal{S}_{\ell_p(\kappa)}(\ell_p(\lambda))$, respectively, for some infinite cardinal number $\kappa \leq \lambda$.*

2. PRELIMINARIES

In this paper, we consider Banach spaces over the fixed scalar field either of real or of complex numbers. Our terminology is standard and follows mainly [1] and [21]. By *operator*, we mean a bounded, linear operator acting between Banach spaces. Let X and Y be Banach spaces. We denote by $\mathcal{B}(X, Y)$ the Banach space of all operators from X to Y and write $\mathcal{B}(X)$ for $\mathcal{B}(X, X)$. An operator $P \in \mathcal{B}(X)$ is a *projection* if $P^2 = P$.

Let $T: X \rightarrow Y$ be an operator. We say that T is *bounded below* if there is a constant $c > 0$ such that $\|Tx\| \geq c\|x\|$ for all $x \in X$. It is clear an operator is bounded below if and only if it is injective and has closed range, in which case it is an isomorphism onto its range.

Let E be a Banach space. We say that an operator T acting between Banach spaces is *bounded below on a copy of E* if there exists a subspace E_0 of the domain of T such that E_0 is isomorphic to E and the operator $T|_{E_0}$ is bounded below. For each pair of Banach spaces X, Y we set

$$\mathcal{S}_E(X, Y) = \{T \in \mathcal{B}(X, Y) : T \text{ is not bounded below on any copy of } E\},$$

and write $\mathcal{S}_E(X) = \mathcal{S}_E(X, X)$. We call operators which belong to $\mathcal{S}_E(X, Y)$ *E -singular operators*. We denote by \mathcal{S}_E the class of all E -singular operators acting between arbitrary Banach spaces. In general, the set $\mathcal{S}_E(X)$ need not be closed under addition. This is readily seen in the case where $E = X = \ell_p \oplus \ell_q$ for $1 \leq p < q < \infty$.

We use von Neumann's definition of an ordinal number. We identify cardinal numbers with initial ordinal numbers. For instance, ω_1 is the first uncountable

ordinal, ω_2 is the second one and so on. If λ is a cardinal and $\alpha < \lambda$ we often write $[0, \alpha)$ and (α, λ) for the set of ordinals less than λ which are, respectively, less than α and greater than α . Even though α and $[0, \alpha)$ are equal, we distinguish these symbols to avoid confusion in the situation when we think of α as an element of λ and in the situation when it is considered a subset of it. The *cofinality* of a cardinal number λ is the minimal cardinality κ of a family of cardinals strictly less than λ , say $\{\lambda_\alpha : \alpha < \kappa\}$, such that $\lambda = \sum_{\alpha < \kappa} \lambda_\alpha$. The cofinality of ω_1 is ω_1 while the cofinality of ω_ω is countable. If Λ is a set, we denote by $|\Lambda|$ the cardinality of Λ . For a cardinal number λ , we denote by λ^+ its cardinal successor; for instance $\omega_1^+ = \omega_2$, etc.

For a subset Y of a topological space X we define the *density* of Y , $\text{dens } Y$, to be the minimal cardinality of a dense subset of Y .

Let λ be an infinite cardinal and let E_λ denote one of the spaces: $c_0(\lambda)$, $\ell_p(\lambda)$ ($p \in [1, \infty]$) or $\ell_\infty^c(\lambda)$. For each subset $\Lambda \subseteq \lambda$ the following formula defines a contractive projection P_Λ on E_λ :

$$(2.1) \quad (P_\Lambda f)(\alpha) = \begin{cases} f(\alpha), & \text{if } \alpha \in \Lambda \\ 0, & \text{if } \alpha \notin \Lambda \end{cases} \quad (f \in E_\lambda).$$

We then denote the range of P_Λ by E_Λ . Of course, the range of P_Λ is isometrically isomorphic to $E_{|\Lambda|}$. We denote by $R_\Lambda : E_\lambda \rightarrow E_\Lambda$ the *restriction operator*, which identifies an element f in the range of P_Λ with the restriction of f , $R_\Lambda f$, to the set Λ . For $f \in E_\lambda$ we define the *support* of f by $\text{supp } f = \{\alpha < \lambda : f(\alpha) \neq 0\}$. A function f is *supported* in a set Λ if $\text{supp } f \subseteq \Lambda$.

We shall also require the following lemma due to Rosenthal ([26, Proposition 1.2, Corollary on p. 29 and Remark 1 on p. 30]).

Lemma 2.1. *Let λ be an infinite cardinal number and let X be a Banach space.*

- (i) *Suppose that $T : \ell_\infty(\lambda) \rightarrow X$ is an operator such that $T|_{c_0(\lambda)}$ is bounded below. Then there exists a set $\Lambda \subset \lambda$ of cardinality λ such that $T|_{\ell_\infty(\Lambda)}$ is bounded below. Consequently, if Z is an injective Banach space and $T : Z \rightarrow X$ is an operator that is bounded below on a copy of $c_0(\lambda)$, then T is bounded below on a copy of $\ell_\infty(\lambda)$.*
- (ii) *Suppose that $A \subseteq X$ is a closed subspace and $T : X \rightarrow \ell_1(\lambda)$ is an operator such that for some $\delta > 0$ and some bounded set $\{y_\alpha : \alpha < \lambda\} \subset A$ we have*

$$\delta \leq \|Ty_\alpha - Ty_\beta\|, \quad (\alpha, \beta < \lambda, \alpha \neq \beta).$$

Then T is bounded below on some subspace of A that is isomorphic to $\ell_1(\lambda)$ and complemented in X .

- (iii) *Suppose that $T : c_0(\lambda) \rightarrow X$ is an operator such that*

$$\inf\{\|Te_\alpha\| : \alpha < \lambda\} > 0,$$

where $\{e_\alpha : \alpha < \lambda\}$ is the canonical basis of $c_0(\lambda)$. Then there exists a set $\Lambda \subset \lambda$ of cardinality λ such that $T|_{c_0(\Lambda)}$ is bounded below.

We say that a Banach space is *primary* if whenever X is isomorphic to $Y \oplus Z$, then at least one of the spaces Y or Z is isomorphic to X .

3. AUXILIARY RESULTS

Proposition 3.1. *Let λ and τ be infinite cardinal numbers and suppose that λ is uncountable. Let $T: \ell_\infty^c(\tau) \rightarrow Y$ be an operator that is not bounded below on any sublattice isometric to $c_0(\lambda)$. Then for every $\varepsilon > 0$ there is a subset Λ of τ so that $|\Lambda| < \lambda$ and*

$$\|TR_{\tau \setminus \Lambda}\| \leq \varepsilon.$$

Consequently, if $Y = \ell_\infty^c(\tau)$ and T is a projection onto a subspace X , then X is isomorphic to a complemented subspace of $\ell_\infty^c(\kappa)$ for some cardinal number $\kappa < \lambda$.

Proof. Let $(f_\alpha)_{\alpha \in \Gamma}$ be a collection of disjointly supported unit vectors in $\ell_\infty^c(\tau)$ that is maximal with respect to the property that $\|Tf_\alpha\| \geq \varepsilon$ for each α . Let Λ be the union of the supports of the f_α ($\alpha \in \Gamma$). In the case where Γ is finite, $|\Lambda| < \lambda$ as Λ is countable. In the case where Γ is infinite, by the hypothesis on T and Lemma 2.1(iii), we have $|\Gamma| < \lambda$, so that also $|\Lambda| < \lambda$. By the maximality of the collection $(f_\alpha)_{\alpha \in \Gamma}$, if f is a unit vector whose support is contained in $\lambda \setminus \Lambda$, then $\|Tf\| < \varepsilon$, which implies that $\|TR_{\tau \setminus \Lambda}\| \leq \varepsilon$.

For the “consequently” statement, suppose that T is a projection onto a subspace X . Then

$$I_X = (TR_{\tau \setminus \Lambda} + TR_\Lambda)|_X,$$

so

$$\|I_X - (TR_\Lambda)|_X\| \leq \varepsilon.$$

Hence if $\varepsilon < 1$, there is an operator U on X so that

$$I_X = (UTR_\Lambda)|_X.$$

Thus I_X factors through $\ell_\infty^c(|\Lambda|)$. □

The next proposition, which is surely known, puts the hypothesis of Proposition 3.1 into perspective.

Proposition 3.2. *Let τ and λ be cardinal numbers and suppose that λ is uncountable. Let $E = \ell_\infty^c(\tau)$ and $E_\lambda = c_0(\lambda)$ or let $E = \ell_p(\tau)$ and $E_\lambda = \ell_p(\lambda)$ for some $1 < p < \infty$. Suppose that $(x_\alpha)_{\alpha \in \lambda}$ is a transfinite sequence in E that is equivalent to the unit vector basis of E_λ . Then there is a set $\Lambda \subset \lambda$ with $|\Lambda| = \lambda$ such that $(x_\alpha)_{\alpha \in \Lambda}$ have disjoint supports.*

Consequently,

- (i) *if $T: E \rightarrow E_\tau$ is a bounded, linear operator, then there is a set $\Lambda \subset \lambda$ with $|\Lambda| = \lambda$ such that $(Te_\alpha)_{\alpha \in \Lambda}$ have disjoint supports;*
- (ii) *if $\tau < \lambda$, then $c_0(\lambda)$ is not isomorphic to a subspace of $\ell_\infty^c(\tau)$.*

Proof. Take $\Lambda \subset \lambda$ maximal with respect to the property that the vectors $(x_\alpha)_{\alpha \in \Lambda}$ have disjoint supports. Let Γ be the union of the supports of $(x_\alpha)_{\alpha \in \Lambda}$. Suppose, for contradiction, that $|\Lambda| < |\lambda|$. Then also $|\Gamma| < |\lambda|$ since λ is uncountable. By the maximality of $(x_\alpha)_{\alpha \in \Lambda}$, each x_α for $\alpha \in \lambda \setminus \Lambda$ is non-zero at some coordinate in Γ . Consequently, uncountably many x_α are non-zero at the same coordinate $\gamma \in \Gamma$. So, without loss of generality, we have for some $\varepsilon > 0$ a sequence of distinct x_{α_n} so that for all n the real part of $x_{\alpha_n}(\gamma)$ is larger than ε . Then for every N ,

$$\left\| \sum_{n=1}^N x_{\alpha_n} \right\| \geq \begin{cases} \varepsilon \cdot N, & \text{if } E = \ell_\infty^c(\tau), \\ \frac{\varepsilon \cdot N}{N^{1/p}}, & \text{if } E = \ell_p(\tau), \end{cases}$$

contradicting the fact that $(x_\alpha)_{\alpha \in \lambda}$ is equivalent to the unit vector basis for E_λ . □

We thus obtain an analogue of Lemma 2.1(iii) in the case of $\ell_p(\lambda)$ for $p \in (1, \infty)$.

Corollary 3.3. *Let λ be an uncountable cardinal number and let $p \in (1, \infty)$. Suppose that $T: \ell_p(\lambda) \rightarrow \ell_p(\lambda)$ is a bounded, linear operator such that*

$$\inf\{\|Te_\alpha\|: \alpha < \lambda\} > 0.$$

Then there is a set $\Lambda \subset \lambda$ with $|\Lambda| = \lambda$ such that TR_Λ is bounded below.

Although not needed for the present paper, it is worth remarking that the “consequently” statement in Proposition 3.2 can be improved, as is shown by the following (probably known) simple proposition.

Proposition 3.4. *Let $\tau < \lambda$ be cardinal numbers. Then $c_0(\lambda)$ is not isomorphic to a subspace of $\ell_\infty(\tau)$.*

Proof. Since the other cases are clear, we assume that τ is infinite and hence that λ is uncountable. Suppose, for contradiction, that $(x_\alpha)_{\alpha \in \lambda}$ is a set of vectors in $\ell_\infty(\tau)$ that is equivalent to the unit vector basis of $c_0(\lambda)$. For each $t \in \tau$, the set A_t of all $\alpha \in \lambda$ for which $x_\alpha(t)$ is not zero must be countable (this was already used in the proof of Proposition 3.2). But then the cardinality of $\bigcup_{t \in \tau} A_t$ is at most $\omega \cdot \tau = \tau < \lambda$. □

We also obtain a characterisation of complemented subspaces of $\ell_\infty^c(\omega_1)$ that do not embed into ℓ_∞ .

Corollary 3.5. *Every infinite-dimensional, complemented subspace of $\ell_\infty^c(\omega_1)$ that does not contain a subspace isomorphic to $c_0(\omega_1)$ is isomorphic to ℓ_∞ .*

Proof. Let X be an infinite-dimensional, complemented subspace of $\ell_\infty^c(\omega_1)$ that does not contain a subspace isomorphic to $c_0(\omega_1)$. Let P be any projection onto X . By Proposition 3.1, X is isomorphic to a complemented subspace of $\ell_\infty^c(\omega) = \ell_\infty$ and hence is isomorphic to ℓ_∞ . □

In view of Corollary 3.5, in order to complete the classification of the complemented subspaces of $\ell_\infty^c(\lambda)$, that is, to prove Theorem 1.3, we need to show that a complemented subspace X of $\ell_\infty^c(\lambda)$ that contains an isomorphic copy of $c_0(\lambda)$ must be isomorphic to $\ell_\infty^c(\lambda)$. We begin by showing that such an X must contain a subspace isomorphic to $\ell_\infty^c(\lambda)$.

Proposition 3.6. *Let κ and λ be uncountable cardinal numbers and let $T: \ell_\infty^c(\lambda) \rightarrow \ell_\infty^c(\lambda)$ be an operator. Assume that T is bounded below on a subspace of $\ell_\infty^c(\lambda)$ that is isomorphic to $c_0(\kappa)$. Then there is a subspace Y of $\ell_\infty^c(\lambda)$ that is isometric to $\ell_\infty^c(\kappa)$ such that $T|_Y$ is bounded below.*

Proof. By Proposition 3.2 there is a transfinite sequence $(x_\beta)_{\beta < \kappa}$ of disjointly supported unit vectors in $\ell_\infty^c(\lambda)$ so that the restriction of T to $\overline{\text{span}}\{x_\beta: \beta < \kappa\}$ is bounded below. Define an isometry $S: \ell_\infty^c(\kappa) \rightarrow \ell_\infty^c(\lambda)$ by

$$(Sz)(\gamma) = \begin{cases} z(\beta)x_\beta(\gamma), & \gamma \in \text{supp } x_\beta, \\ 0, & \text{otherwise} \end{cases} \quad (z \in \ell_\infty^c(\kappa), \beta < \kappa).$$

The condition on T says that the operator $TS: \ell_\infty^c(\kappa) \rightarrow \ell_\infty^c(\lambda)$ is bounded below on $c_0(\kappa)$. Regard $\ell_\infty^c(\kappa)$ as the subspace $\ell_\infty^c([0, \kappa))$ of $\ell_\infty(\lambda)$. By injectivity of $\ell_\infty(\lambda)$, the operator TS has a norm-preserving extension to an operator $U: \ell_\infty(\lambda) \rightarrow \ell_\infty(\lambda)$ [21, Proposition 2.f.2]. The operator U is an isomorphism on $c_0(\kappa)$, hence by

Lemma 2.1(i), there is a subset Γ of κ with cardinality κ such that the restriction of U to $\ell_\infty(\Gamma)$ is an isomorphism. But U maps $\ell_\infty^c(\Gamma)$ into $\ell_\infty^c(\lambda)$, which is to say that TS is an isomorphism on $\ell_\infty^c(\Gamma)$, that $Y := S[\ell_\infty^c(\Gamma)]$ is isometric to $\ell_\infty^c(\kappa)$ and $T|_Y$ is bounded below. \square

The following proposition is essentially known in the sense that it can be easily deduced from a string of standard results concerning Grothendieck spaces; however, we offer here a direct proof.

Proposition 3.7. *For every cardinal number λ , $\ell_\infty^c(\lambda)$ is a Grothendieck space.*

Proof. If $\ell_\infty^c(\lambda)$ is not a Grothendieck space there exists an operator $T: \ell_\infty^c(\lambda) \rightarrow c_0$ that is not weakly compact (see [5, p. 150]). Then by [23] there is a subspace Y of $\ell_\infty^c(\lambda)$ isomorphic to c_0 such that $T|_Y$ is bounded below. Then Y is contained in E_Λ for some countable set $\Lambda \subset \lambda$. In particular, $T|_{E_\Lambda}$ can be regarded as an operator from ℓ_∞ into c_0 that is bounded below on a copy of c_0 . By Lemma 2.1(i), T must be bounded below on a copy of ℓ_∞ ; a contradiction as T maps into c_0 . \square

We need a consequence of [26, Lemma 1.1].

Lemma 3.8. *Let Γ be uncountable and let T be an operator on $\ell_\infty(\Gamma)$ such that T is the identity on $c_0(\Gamma)$. Then for every $\varepsilon > 0$ there is $\Gamma' \subset \Gamma$ with $|\Gamma'| = |\Gamma|$ so that*

$$\|(R_{\Gamma'}T)|_{\ell_\infty(\Gamma')} - I_{\ell_\infty(\Gamma')}\| < \varepsilon,$$

where $R_{\Gamma'}: \ell_\infty(\Gamma) \rightarrow \ell_\infty(\Gamma')$ is the restriction operator.

Proof. We identify $\ell_\infty(\Gamma)^*$ with the finitely additive, scalar-valued measures on the σ -algebra of all subsets of Γ . For $\gamma \in \Gamma$, set $\mu_\gamma = T^*\delta_\gamma$, where δ_γ is the point mass measure at γ . By [26, Lemma 1.1] there is $\Gamma' \subset \Gamma$ with $|\Gamma'| = |\Gamma|$ so that for all $\gamma \in \Gamma'$ we have $|\mu_\gamma|(\Gamma' \setminus \{\gamma\}) < \varepsilon$. \square

Proof of Theorem 1.3. We need to show that if X is a subspace of $\ell_\infty^c(\lambda)$ isomorphic to $\ell_\infty^c(\lambda)$, then X contains a subspace isomorphic to $\ell_\infty^c(\lambda)$ that is complemented in $\ell_\infty^c(\lambda)$. The case where λ is countable follows from injectivity of ℓ_∞ , so let us suppose that λ is uncountable. Let \leq denote the order on X that is induced from its isomorphism with $\ell_\infty^c(\lambda)$. Applying Proposition 3.2 twice in succession, we get a transfinite sequence $\{f_\gamma: \gamma < \lambda\}$ of unit vectors in X that are disjoint both relative to the order structure on X given by \leq and the order structure on $\ell_\infty^c(\lambda)$ given by its pointwise ordering $<$. Let Z be the lattice closure in $(\ell_\infty^c(\lambda), <)$ of the linear span of $\{f_\gamma: \gamma < \lambda\}$ (so that Z is isometrically isomorphic to $\ell_\infty^c(\lambda)$ and there is a norm-one projection Q from $\ell_\infty^c(\lambda)$ onto Z). By replacing X itself with a subspace, we might as well assume that in (X, \leq) the f_γ form the unit vector basis for $c_0(\lambda)$ and hence that in (X, \leq) the space X is the pointwise closure of the f_γ . Let $J: (Z, <) \rightarrow (X, \leq)$ denote the natural surjective order isomorphism that is the identity on the set $\{f_\gamma: \gamma < \lambda\}$. Now

$$(JQ)|_X: X \rightarrow X$$

is an operator on X that is the identity on $\{f_\gamma: \gamma < \lambda\}$, so by Lemma 3.8 (applied to any extension of the operator to an operator on the $\ell_\infty(\lambda)$ lattice generated by (X, \leq)) we have a subset Λ of λ of cardinality λ so that

$$(3.1) \quad \|(R_\Lambda JQ)|_{X_\Lambda} - I_{X_\Lambda}\| < \varepsilon$$

where X_Λ is the pointwise closure of $\{f_\gamma : \gamma \in \Lambda\}$ in (X, \triangleleft) , DR_Λ is the restriction mapping on (X, \triangleleft) , and $\varepsilon > 0$ can be as small as we want; in particular, $\varepsilon < 1$. So there is an automorphism U on X_Λ so that $UR_\Lambda JQ$ is the identity on X_Λ and hence $UR_\Lambda JQ$ is a projection from $\ell_\infty^c(\lambda)$ onto X_Λ . Since X_Λ is isomorphic to $\ell_\infty^c(\lambda)$, this completes the proof. \square

The above proof of Theorem 1.3 can be adjusted to get the following stronger result.

Proposition 3.9. *Let κ and λ be infinite cardinal numbers. If X is a subspace of $\ell_\infty^c(\lambda)$ that is isomorphic to $\ell_\infty^c(\kappa)$, then there is a subspace Y of X that is isomorphic to $\ell_\infty^c(\kappa)$ and complemented in $\ell_\infty^c(\lambda)$.*

Proposition 3.10. *For each uncountable cardinal number λ and $p \in [1, \infty)$ the Banach space $\ell_p(\lambda)$ is complementably homogeneous.*

Proof. The case $p = 1$ is covered by Lemma 2.1(ii). When $p \in (1, \infty)$, the result follows from Proposition 3.2. \square

Remark 3.11. It is perhaps worthwhile to note that an analogue of Proposition 3.9 holds also for $E_\lambda = \ell_p(\lambda)$ ($p \in [1, \infty)$); that is, if Y is a subspace of E_λ isomorphic to E_κ for some cardinal number κ , then there is a subspace $X \subseteq Y$ isomorphic to E_κ that is complemented in E_λ . This is because every copy of E_κ in E_λ is contained in E_Λ for some set $\Lambda \subset \lambda$ with $|\Lambda| = \kappa$.

Proposition 3.12. *Let λ and κ be infinite cardinal numbers, $p \in [1, \infty)$ and let X be a Banach space. Then the sets*

$$\mathcal{S}_{c_0(\lambda)}(X, \ell_\infty^c(\kappa)) \quad \text{and} \quad \mathcal{S}_{\ell_p(\lambda)}(X, \ell_p(\kappa))$$

are closed under addition.

Proof. The case $\lambda = \omega$ follows from [16, Proposition 2.5]. Let us consider the case where λ is uncountable.

Suppose that T, S are in $\mathcal{S}_{c_0(\lambda)}(X, \ell_\infty^c(\kappa))$ and assume, in search of contradiction, that $T + S$ is not in $\mathcal{S}_{c_0(\lambda)}(X, \ell_\infty^c(\kappa))$. Then there exists a subspace $X_1 \subseteq X$ isomorphic to $c_0(\lambda)$ and $\varepsilon > 0$ such that $\varepsilon\|z\| \leq \|(T + S)(z)\|$ for all $z \in X_1$. Arguing similarly as in the proof of Proposition 3.1, we infer that there exists a subspace $Z_1 \subset X_1$ isomorphic to $c_0(\lambda)$ such that $\|T|_{Z_1}\| \leq \frac{\varepsilon}{3}$. Repeating the above argument, we conclude that there is a subspace $Z_2 \subseteq Z_1$ isomorphic to $c_0(\lambda)$ such that $\|S|_{Z_2}\| \leq \frac{\varepsilon}{3}$. Consequently, for all $z \in Z_2$ we have

$$\varepsilon\|z\| \leq \|Tz + Sz\| \leq \frac{\varepsilon}{3}\|z\| + \frac{\varepsilon}{3}\|z\| = \frac{2}{3}\varepsilon\|z\|;$$

a contradiction.

The proof in the case of $\ell_p(\lambda)$ for $p \in (1, \infty)$ is analogous (here we use Corollary 3.3 instead of Lemma 2.1(iii) when arguing as in the proof of Proposition 3.1).

The proof of the case of $p = 1$ is actually contained in [26]. Indeed, suppose contrapositively that $T, S: X \rightarrow \ell_1(\kappa)$ are operators such that $T + S$ is not in $\mathcal{S}_{\ell_1(\lambda)}(X, \ell_1(\kappa))$. Let Y be a copy of $\ell_1(\lambda)$ in X such that $(T + S)|_Y$ is an isomorphism. By [26, Theorem 3.3], there is a set of unit vectors $\{x_\alpha : \alpha < \lambda\}$ in Y and disjoint sets $E_\alpha \subset \kappa$ ($\alpha < \lambda$) such that for some $\delta > 0$ $\|((T + S)x_\alpha)|_{E_\alpha}\| \geq \delta$. It follows that for some subset $\Lambda \subseteq \lambda$ of cardinality λ and for either T or S , say T , $\|(Tx_\alpha)|_{E_\alpha}\| \geq \delta/2$ for all $\alpha \in \Lambda$. It now follows from [26, Propositions 3.2 and 3.1]

that Λ contains a subset Λ' of the same cardinality such that T is bounded below on the closed linear span of $\{x_\alpha : \alpha \in \Lambda'\}$. □

Remark. One can improve a bit the ℓ_1 -version of the previous proposition: If X contains an isomorphic copy of $\ell_1(\lambda)$, then for each $\varepsilon > 0$ it contains a $(1 + \varepsilon)$ -isomorphic copy of $\ell_1(\lambda)$ that is complemented by means of a projection of norm at most $1 + \varepsilon$.

This follows from a simple combination of the non-separable counterpart of the James distortion theorem for ℓ_1 ([13], cf. [8]), which provides the $(1 + \varepsilon)$ -isomorphic copy of $\ell_1(\lambda)$ together with Dor’s result [6, Theorem A], which provides the needed projection.

This remark is provided here because it may be useful in questions related to the classification of commutators. Similar facts were useful when classifying the commutators in, say, the algebra of bounded operators on ℓ_1 .

Proposition 3.12 can be strengthened in the case of $c_0(\lambda)$:

Proposition 3.13. *For each infinite cardinal number λ , $\mathcal{S}_{c_0(\lambda)}$ forms a closed operator ideal.*

Proof. Let X and Y be Banach spaces. The set $\mathcal{S}_{c_0(\lambda)}(X, Y)$ is closed in $\mathcal{B}(X, Y)$ and obviously is closed under taking compositions with other operators (whenever these make sense). The only non-trivial thing is to verify that it is indeed closed under addition.

Let $T, S \in \mathcal{B}(X, Y)$ be operators such that $T + S$ is bounded below by $\delta > 0$ on some subspace $X_0 \subseteq X$ isomorphic to $c_0(\lambda)$. Let $\{e_\alpha : \alpha < \lambda\}$ be a transfinite sequence in X_0 equivalent to the canonical basis of $c_0(\lambda)$. Since

$$\delta \leq \|Te_\alpha + Se_\alpha\|$$

for all $\alpha < \lambda$, there is a set Λ of cardinality λ such that $\|Te_\alpha\| \geq \frac{\delta}{2}$ or $\|Se_\alpha\| \geq \frac{\delta}{2}$ for all $\alpha \in \Lambda$. It follows then from Lemma 2.1(iii) that at least one of the operators T or S does not belong to $\mathcal{S}_{c_0(\lambda)}(X, Y)$. □

Theorem 3.14. *Let κ and λ be infinite cardinals and let $p \in [1, \infty]$. Then \mathcal{A} is a closed ideal of \mathcal{B} if*

- $\mathcal{A} = \mathcal{S}_{c_0(\kappa)}(c_0(\lambda))$ and $\mathcal{B} = \mathcal{B}(c_0(\lambda))$;
- $\mathcal{A} = \mathcal{S}_{\ell_p(\kappa)}(\ell_p(\lambda))$ and $\mathcal{B} = \mathcal{B}(\ell_p(\lambda))$;
- $\mathcal{A} = \mathcal{S}_{\ell_\infty^c(\kappa)}(\ell_\infty^c(\lambda))$ and $\mathcal{B} = \mathcal{B}(\ell_\infty^c(\lambda))$.

Moreover,

$$(3.2) \quad \mathcal{S}_{\ell_\infty^c(\kappa)}(\ell_\infty^c(\lambda)) = \mathcal{S}_{c_0(\kappa)}(\ell_\infty^c(\lambda)).$$

Proof. The first clause as well as the second one in the case of $p \in [1, \infty)$ follow directly from Proposition 3.12.

We claim that $\mathcal{S}_{\ell_\infty^c(\kappa)}(\ell_\infty^c(\lambda)) = \mathcal{S}_{c_0(\kappa)}(\ell_\infty^c(\lambda))$, which is plainly enough as Proposition 3.12 asserts that $\mathcal{S}_{c_0(\kappa)}(\ell_\infty^c(\lambda))$ is a closed ideal of $\mathcal{B}(\ell_\infty^c(\lambda))$. Let us suppose that $T \notin \mathcal{S}_{c_0(\kappa)}(\ell_\infty^c(\lambda))$. If κ is countable, then arguing as in the proof of Proposition 3.7, we infer that T preserves a copy of ℓ_∞ . In the case where κ is uncountable, Proposition 3.2 gives us a transfinite sequence of disjointly supported unit vectors $(x_\beta)_{\beta < \kappa}$ such that T is bounded below on $\overline{\text{span}}\{x_\beta : \beta < \kappa\}$. By Proposition 3.6, T is bounded below on some copy of $\ell_\infty^c(\kappa)$, hence $T \notin \mathcal{S}_{\ell_\infty^c(\kappa)}(\ell_\infty^c(\lambda))$.

On the other hand, $\mathcal{S}_{\ell_\infty^c(\kappa)}(\ell_\infty^c(\lambda)) \supseteq \mathcal{S}_{c_0(\kappa)}(\ell_\infty^c(\lambda))$ trivially, so the proof of the claim is complete.

Let us consider the remaining case of $\mathcal{A} = \mathcal{S}_{\ell_\infty(\kappa)}(\ell_\infty(\lambda))$ and $\mathcal{B} = \mathcal{B}(\ell_\infty(\lambda))$. Suppose contrapositively that T, S are operators on $\ell_\infty(\lambda)$ such that $T + S \notin \mathcal{S}_{\ell_\infty(\kappa)}(\ell_\infty(\lambda))$. Let $X \subseteq \ell_\infty(\lambda)$ be a copy of $c_0(\kappa)$ on which $T + S$ is bounded below. Since $\mathcal{S}_{c_0(\lambda)}$ is an operator ideal (Proposition 3.13), either T or S is bounded below on some copy of $c_0(\lambda)$. By Lemma 2.1(i), at least one of those operators is bounded below on a copy of $\ell_\infty(\lambda)$, which means that either T or S is not in $\mathcal{S}_{\ell_\infty(\kappa)}(\ell_\infty(\lambda))$. \square

The next lemma is a counterpart of [4, Proposition 5.1].

Lemma 3.15. *Let κ and λ be infinite cardinal numbers and suppose that κ is uncountable and is not a successor of any cardinal number. If E_λ is one of the spaces $c_0(\lambda)$, $\ell_\infty^c(\lambda)$ or $\ell_p(\lambda)$ for some $p \in [1, \infty)$, then*

$$\overline{\bigcup_{\rho < \kappa} \mathcal{S}_{E_\rho}(E_\lambda)} = \mathcal{S}_{E_\kappa}(E_\lambda).$$

Consequently, if the cofinality of κ is uncountable

$$\bigcup_{\rho < \kappa} \mathcal{S}_{E_\rho}(E_\lambda) = \mathcal{S}_{E_\kappa}(E_\lambda).$$

Proof. Only the inclusion from right to left requires justification. Certainly we may suppose that κ is uncountable as otherwise the left hand side is trivially equal to the right hand side.

Take $T \in \mathcal{S}_{E_\kappa}(E_\lambda)$. We claim that for each $\varepsilon > 0$ there is a set Λ with $|\Lambda| < \kappa$ such that $\|P_\Lambda T - T\| < \varepsilon$. We split the proof of this claim into two subcases.

- Let us consider first the case where $E_\lambda = \ell_1(\lambda)$. Assume that the assertion fails for a certain $\varepsilon > 0$. Then there is a set of unit vectors $\{x_\alpha : \alpha < \kappa\}$ in Y and finite disjoint sets $E_\alpha \subset \lambda$ ($\alpha < \kappa$) such that for some $\delta \in (0, \varepsilon)$ we have $\|Tx_\alpha|_{E_\alpha}\| \geq \delta$ ($\alpha < \kappa$). It now follows from [26, Propositions 3.2 and 3.1] that κ contains a subset of the same cardinality, K say, such that T is bounded below on $\overline{\text{span}}\{x_\alpha : \alpha \in K\} \cong \ell_1(\kappa)$.
- Assume that our assertion is not true in the remaining cases, which means that for some $\varepsilon_0 > 0$ and all sets Λ with cardinality less than κ we have

$$\|P_{\lambda \setminus \Lambda} T\| = \|P_\Lambda T - T\| \geq \varepsilon_0.$$

If $E_\lambda = c_0(\lambda)$ or $E_\lambda = \ell_p(\lambda)$ for some $p \in (1, \infty)$, this is a contradiction. Indeed, otherwise by Proposition 3.2(i), we would have a family of disjointly supported unit vectors $\{z_\gamma : \gamma < \kappa\}$ such that $\{Tz_\gamma : \gamma < \kappa\}$ are also disjointly supported and have norm at least $\varepsilon_0/2$. In particular, T would be bounded below on a copy of E_κ against the assumption. For $E_\lambda = \ell_\infty^c(\lambda)$, as in the case of $c_0(\lambda)$, there would exist a family of disjointly supported unit vectors $\{z_\gamma : \gamma < \kappa\}$ such that $\{Tz_\gamma : \gamma < \kappa\}$ are also disjointly supported and have norm at least $\varepsilon_0/2$. Thus T would preserve a copy of $c_0(\lambda)$, hence by (3.2), also a copy of $\ell_\infty^c(\kappa)$.

Since $P_\Lambda T \in \mathcal{S}_{E_\rho}(E_\lambda)$ ($n \in \mathbb{N}$), we conclude that T belongs to the closure of the ideal $\bigcup_{\rho < \kappa} \mathcal{S}_{E_\rho}(E_\lambda)$ as desired.

The final assertion follows from the well-known fact that in first-countable spaces increasing unions of closed sets that are well ordered by inclusion are closed when κ has uncountable cofinality. \square

Lemma 3.16. *For each infinite cardinal number λ we have*

$$\mathcal{W}(\ell_\infty^c(\lambda)) = \mathcal{S}_{\ell_\infty}(\ell_\infty^c(\lambda)),$$

where \mathcal{W} denotes the ideal of weakly compact operators.

Proof. Let T be an operator on $\ell_\infty^c(\lambda)$. Then by [23], T is not weakly compact if and only if T is bounded below on some copy of c_0 . Suppose that T is not weakly compact and let F be a copy of c_0 that is preserved by T . Then there exists a countable set Λ such that $F \subseteq E_\Lambda$. Since E_Λ is isomorphic to ℓ_∞ , by Lemma 2.1(i), T preserves a copy of ℓ_∞ . \square

Lemma 3.17. *Let κ and λ be infinite cardinal numbers and let $T \in \mathcal{B}(\ell_\infty^c(\lambda))$. Suppose that $T \notin \mathcal{S}_{\ell_\infty^c(\kappa)}(\ell_\infty^c(\lambda))$. Then $\mathcal{S}_{\ell_\infty^c(\kappa^+)}(\ell_\infty^c(\lambda))$ is contained in the ideal generated by T .*

Proof. Note that $\mathcal{B}(\ell_\infty^c(\lambda))$ is nothing but $\mathcal{S}_{\ell_\infty^c(\lambda^+)}(\ell_\infty^c(\lambda))$. (Recall that λ^+ denotes the immediate cardinal successor of λ .) It is then enough to consider only the case where $\kappa \leq \lambda$. For simplicity of notation, for any cardinal number κ set $E_\kappa = \ell_\infty^c(\kappa)$.

Fix $T \notin \mathcal{S}_{E_\kappa}(E_\lambda)$ and let E be a subspace of E_λ isomorphic to E_κ on which T is bounded below and such that $F = T[E]$ is complemented (such E exists by Proposition 3.9).

Let $S \in \mathcal{S}_{E_{\kappa^+}}(E_\lambda)$. By Proposition 3.6, S is not bounded below on any sublattice isometric to $c_0(\kappa^+)$. By Proposition 3.1, for each $\varepsilon > 0$, there exists a subset $\Lambda_\varepsilon \subset \lambda$ such that $|\Lambda_\varepsilon| \leq \kappa$ and $\|SR_{\lambda \setminus \Lambda_\varepsilon}\| \leq \varepsilon$. Putting $\Lambda = \bigcup_{n=1}^\infty \Lambda_{1/n}$, we have $S = SR_\Lambda$. Thus S factors through $\ell_\infty^c(\kappa)$, and hence through $T|_E$. Since F is complemented, S factors through T as well. \square

4. PROOFS OF THEOREMS 1.1, 1.2, 1.4 AND 1.5

We are now in a position to prove Theorems 1.1, 1.2, 1.4 and 1.5 (we have already proved Theorem 1.3 in the previous section).

Proof of Theorem 1.1. The case of $\lambda = \omega$ follows from [21, Proposition 2.f.4] as explained in [18, p. 253]. Suppose then that λ is uncountable. The set $\mathcal{S}_X(X)$ is indeed a closed ideal of $\mathcal{B}(X)$ by Theorem 3.14. To show that it is the unique maximal ideal, take $T \notin \mathcal{S}_X(X)$. Then there are subspaces E and F of X , both isomorphic to X such that $T_E: E \rightarrow F$ is an isomorphism and F is complemented (if $X = \ell_\infty(\lambda)$, then it follows directly from injectivity of $\ell_\infty(\lambda)$; if $X = \ell_\infty^c(\lambda)$, then we apply Theorem 1.3).

Let $W = (T|_E)^{-1}$ be regarded as an operator into X . Furthermore, let $U: X \rightarrow F$ be an isomorphism. Since F is complemented, U^{-1} can be extended to an operator on X , say V . Consequently,

$$I_X = VT(WU)$$

belongs to the ideal generated by T . For this reason, $I_X = ATB$ for some operators $A, B \in \mathcal{B}(X)$ if and only if $T \notin \mathcal{S}_X(X)$. \square

Proof of Theorem 1.2. We keep the notation of the proof of Lemma 3.17.

Let κ be an infinite cardinal. By Lemma 3.17, if $T \notin \mathcal{S}_{E_\kappa}(E_\lambda)$, then $\mathcal{S}_{E_{\kappa^+}}(E_\lambda)$ is contained in the ideal generated by T . We will show that if \mathcal{J} is a closed ideal of $\mathcal{B}(E_\lambda)$, then $\mathcal{J} = \mathcal{S}_{E_\kappa}(E_\lambda)$ for some cardinal κ .

Set

$$\tau = \sup\{\rho: \rho \text{ is a cardinal number and } \mathcal{S}_{E_\rho}(E_\lambda) \subseteq \mathcal{J}\}.$$

This number is well defined as, by Lemma 3.16, \mathcal{J} contains $\mathcal{W}(E_\lambda) = \mathcal{S}_{E_\omega}(E_\lambda)$. If the supremum τ is attained and $S \in \mathcal{J} \setminus \mathcal{S}_{E_\tau}(E_\lambda)$, then the ideal generated by S , hence \mathcal{J} as well, would contain $\mathcal{S}_{E_\tau^+}(E_\lambda)$, which is impossible by the definition of τ . Consequently, $\mathcal{J} = \mathcal{S}_{E_\tau}(E_\lambda)$. Consider now the case where $\tau > \rho$ for each ρ such that $\mathcal{S}_{E_\rho}(E_\lambda)$ is contained in \mathcal{J} . Since \mathcal{J} is closed it contains the closure of $\bigcup_{\rho < \kappa} \mathcal{S}_{E_\rho}(E_\lambda)$, which is to say that $\mathcal{S}_{E_\kappa}(E_\lambda) \subseteq \mathcal{J}$ by Lemma 3.15. However, if this inclusion were strict $\mathcal{S}_{E_{\kappa^+}}(E_\lambda)$ would be contained in \mathcal{J} , which would again contradict the definition of τ . □

Proof of Theorem 1.4. Suppose that X is an infinite-dimensional complemented subspace of $\ell_\infty^c(\lambda)$ that is not isomorphic to ℓ_∞ . Then there is a cardinal number $\kappa \leq \lambda$ such that X is isomorphic to a complemented subspace of $\ell_\infty^c(\kappa)$ and $c_0(\kappa)$ embeds into X . Consequently, Proposition 3.6 applies and so X contains a subspace isomorphic to $\ell_\infty^c(\kappa)$. By Theorem 1.3, X contains a complemented subspace isomorphic to $\ell_\infty^c(\kappa)$. Since $\ell_\infty^c(\kappa)$ is isomorphic to $\ell_\infty(\ell_\infty^c(\kappa))$, the ℓ_∞ -sum of countably many copies of itself, the Pełczyński decomposition method (cf. [1, Theorem 2.2.3]) yields that X is isomorphic to $\ell_\infty^c(\kappa)$. □

Proof of Theorem 1.5. Let E_λ denote one of the spaces $c_0(\lambda)$ or $\ell_p(\lambda)$ for some $p \in [1, \infty)$. For $\lambda = \omega$ the result is well known (see, e.g., [25, Theorem on p. 82]). Suppose then that λ is uncountable.

Since E_λ has the approximation property, the ideal of compact operators $\mathcal{K}(E_\lambda)$ is the smallest closed non-zero ideal of $\mathcal{B}(E_\lambda)$ and $\mathcal{K}(E_\lambda) = \mathcal{S}_{E_\omega}(E_\lambda)$ ([25, Theorem on p. 82]). On the other hand $\mathcal{B}(E_\lambda) = \mathcal{S}_{E_{\lambda^+}}(E_\lambda)$.

Let κ be an infinite cardinal. All we need to do is to show that if

$$T \notin \mathcal{S}_{\ell_\infty^c(\kappa)}(\ell_\infty^c(\lambda)),$$

then $\mathcal{S}_{\ell_\infty^c(\kappa^+)}(\ell_\infty^c(\lambda))$ is contained in the ideal generated by T . However, the remainder of the proof is completely analogous to the proof of Theorem 1.2, so we leave the details to the reader. □

ACKNOWLEDGMENTS

We thank the referee for an unusually detailed and helpful report.

REFERENCES

- [1] Fernando Albiac and Nigel J. Kalton, *Topics in Banach space theory*, Graduate Texts in Mathematics, vol. 233, Springer, New York, 2006. MR2192298 (2006h:46005)
- [2] Spiros A. Argyros, Jesús F. Castillo, Antonio S. Granero, Mar Jiménez, and José P. Moreno, *Complementation and embeddings of $c_0(I)$ in Banach spaces*, Proc. London Math. Soc. (3) **85** (2002), no. 3, 742–768, DOI 10.1112/S0024611502013618. MR1936819 (2003k:46022)
- [3] Y. Benyamini, *An extension theorem for separable Banach spaces*, Israel J. Math. **29** (1978), no. 1, 24–30. MR0482075 (58 #2163)
- [4] Matthew Daws, *Closed ideals in the Banach algebra of operators on classical non-separable spaces*, Math. Proc. Cambridge Philos. Soc. **140** (2006), no. 2, 317–332, DOI 10.1017/S0305004105009102. MR2212284 (2007c:46049)

- [5] Joseph Diestel, *Geometry of Banach spaces—selected topics*, Lecture Notes in Mathematics, Vol. 485, Springer-Verlag, Berlin-New York, 1975. MR0461094 (57 #1079)
- [6] Leonard E. Dor, *On projections in L_1* , Ann. of Math. (2) **102** (1975), no. 3, 463–474. MR0420244 (54 #8258)
- [7] Detelin Dosev, William B. Johnson, and Gideon Schechtman, *Commutators on L_p , $1 \leq p < \infty$* , J. Amer. Math. Soc. **26** (2013), no. 1, 101–127, DOI 10.1090/S0894-0347-2012-00748-6. MR2983007
- [8] Daniel P. Giesy, *On a convexity condition in normed linear spaces*, Trans. Amer. Math. Soc. **125** (1966), 114–146. MR0205031 (34 #4866)
- [9] Bernhard Gramsch, *Eine Idealstruktur Banachscher Operatoralgebren* (German), J. Reine Angew. Math. **225** (1967), 97–115. MR0206727 (34 #6544)
- [10] Antonio S. Granero, *On the complemented subspaces of $c_0(I)$* , Atti Sem. Mat. Fis. Univ. Modena **46** (1998), no. 1, 35–36. MR1628581 (99c:46016)
- [11] A. Grothendieck, *Sur les applications linéaires faiblement compactes d’espaces du type $C(K)$* (French), Canadian J. Math. **5** (1953), 129–173. MR0058866 (15,438b)
- [12] J. Hagler and W. B. Johnson, *On Banach spaces whose dual balls are not weak* sequentially compact*, Israel J. Math. **28** (1977), no. 4, 325–330. MR0482086 (58 #2173)
- [13] Robert C. James, *Uniformly non-square Banach spaces*, Ann. of Math. (2) **80** (1964), 542–550. MR0173932 (30 #4139)
- [14] Tomasz Kania and Niels Jakob Laustsen, *Uniqueness of the maximal ideal of the Banach algebra of bounded operators on $C([0, \omega_1])$* , J. Funct. Anal. **262** (2012), no. 11, 4831–4850, DOI 10.1016/j.jfa.2012.03.011. MR2913688
- [15] Tomasz Kania and Niels Jakob Laustsen, *Operators on two Banach spaces of continuous functions on locally compact spaces of ordinals*, Proc. Amer. Math. Soc. **143** (2015), no. 6, 2585–2596, DOI 10.1090/S0002-9939-2015-12480-X. MR3326039
- [16] Tomasz Kania and Niels Jakob Laustsen, *Uniqueness of the maximal ideal of operators on the ℓ_p -sum of ℓ_∞^n ($n \in \mathbb{N}$) for $1 < p < \infty$* , Math. Proc. Cambridge Philos. Soc. **160** (2016), no. 3, 413–421, DOI 10.1017/S0305004115000766. MR3479542
- [17] Piotr Koszmider, *On decompositions of Banach spaces of continuous functions on Mrówka’s spaces*, Proc. Amer. Math. Soc. **133** (2005), no. 7, 2137–2146 (electronic), DOI 10.1090/S0002-9939-05-07799-3. MR2137881 (2006a:46030)
- [18] Niels Jakob Laustsen and Richard J. Loy, *Closed ideals in the Banach algebra of operators on a Banach space*, Topological algebras, their applications, and related topics, Banach Center Publ., vol. 67, Polish Acad. Sci., Warsaw, 2005, pp. 245–264, DOI 10.4064/bc67-0-20. MR2143930 (2006g:47115)
- [19] Niels Jakob Laustsen, Thomas Schlumprecht, and András Zsák, *The lattice of closed ideals in the Banach algebra of operators on a certain dual Banach space*, J. Operator Theory **56** (2006), no. 2, 391–402. MR2282689 (2008a:47119)
- [20] Joram Lindenstrauss, *On complemented subspaces of m* , Israel J. Math. **5** (1967), 153–156. MR0222616 (36 #5666)
- [21] Joram Lindenstrauss and Lior Tzafriri, *Classical Banach spaces*, Lecture Notes in Mathematics, Vol. 338, Springer-Verlag, Berlin-New York, 1973. MR0415253 (54 #3344)
- [22] Erhard Luft, *The two-sided closed ideals of the algebra of bounded linear operators of a Hilbert space*, Czechoslovak Math. J. **18 (93)** (1968), 595–605. MR0238086 (38 #6362)
- [23] A. Pełczyński, *Banach spaces on which every unconditionally converging operator is weakly compact*, Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys. **10** (1962), 641–648. MR0149295 (26 #6785)
- [24] A. Pełczyński and V. N. Sudakov, *Remark on non-complemented subspaces of the space $m(S)$* , Colloq. Math. **9** (1962), 85–88. MR0139001 (25 #2441)
- [25] Albrecht Pietsch, *Operator ideals*, North-Holland Mathematical Library, vol. 20, North-Holland Publishing Co., Amsterdam-New York, 1980. Translated from German by the author. MR582655 (81j:47001)
- [26] Haskell P. Rosenthal, *On relatively disjoint families of measures, with some applications to Banach space theory*, Studia Math. **37** (1970), 13–36. MR0270122 (42 #5015)
- [27] Haskell P. Rosenthal, *Correction to the paper: “On relatively disjoint families of measures, with some applications to Banach space theory”*, Studia Math. **37** (1970/71), 311–313. MR0305040 (46 #4170)

- [28] Haskell P. Rosenthal, *On injective Banach spaces and the spaces $C(S)$* , Bull. Amer. Math. Soc. **75** (1969), 824–828. MR0243335 (39 #4657)
- [29] Haskell P. Rosenthal, *On injective Banach spaces and the spaces $L^\infty(\mu)$ for finite measure μ* , Acta Math. **124** (1970), 205–248. MR0257721 (41 #2370)
- [30] P. Wojtaszczyk, *On projections and unconditional bases in direct sums of Banach spaces. II*, Studia Math. **62** (1978), no. 2, 193–201. MR0500084 (58 #17791)

DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TEXAS 77843
E-mail address: johnson@math.tamu.edu

INSTITUTE OF MATHEMATICS, POLISH ACADEMY OF SCIENCES, ŚNIADECKICH 8, 00-956
WARSZAWA, POLAND; DEPARTMENT OF MATHEMATICS AND STATISTICS, FYLDE COLLEGE, LAN-
CASTER UNIVERSITY, LANCASTER LA1 4YF, UNITED KINGDOM —AND — SCHOOL OF MATHEMAT-
ICAL SCIENCES, WESTERN GATEWAY BUILDING, UNIVERSITY COLLEGE CORK, CORK, IRELAND
E-mail address: tomasz.marcin.kania@gmail.com

DEPARTMENT OF MATHEMATICS, WEIZMANN INSTITUTE OF SCIENCE, REHOVOT, ISRAEL
E-mail address: gideon@weizmann.ac.il