

QUOTIENTS OF MAPPING CLASS GROUPS FROM $\text{Out}(F_n)$

KHALID BOU-RABEE AND CHRISTOPHER J. LEININGER

(Communicated by David Futer)

ABSTRACT. We give a short proof of Masbaum and Reid’s result that mapping class groups involve any finite group, appealing to free quotients of surface groups and a result of Gilman, following Dunfield–Thurston.

Let Σ_g be a closed oriented surface of genus g and let F_n be a nonabelian free group of rank n . The fundamental group, $\pi_1(\Sigma_g)$, is residually free [Bau62], and F_n has a wealth of finite index subgroups [MKS04, p. 116]. In [DT06], N. Dunfield and W. Thurston consider the action of the mapping class group $\text{Mod}(\Sigma_g)$ on the set of finite index normal subgroups of $\pi_1(\Sigma_g)$ with finite simple quotients and in particular those containing the kernel of an epimorphism $\pi_1(\Sigma_g) \rightarrow F_g$. Their observations relating to work of R. Gilman [Gil77] give rise to finite index subgroups of $\text{Mod}(\Sigma_g)$ that surject symmetric groups of arbitrarily large order; see the discussion following the proof of [DT06, Theorem 7.4].

Theorem 1 (Dunfield–Thurston). *For all $g \geq 3$, $r \geq 1$, there exists an epimorphism $\phi : \pi_1(\Sigma_g) \rightarrow F_g$ and a prime q , so that*

$$\{N \triangleleft \pi_1(\Sigma_g) \mid \ker \phi < N \text{ and } \pi_1(\Sigma_g)/N \cong \text{PSL}(2, q)\}$$

has at least r elements, and its (finite index) stabilizer in $\text{Mod}(\Sigma_g)$ acts as the full symmetric group on this set.

We explain the proof of this in Section 1.2. In this note, we observe that since every finite group embeds in some finite symmetric group, Theorem 1 provides a new elementary proof of a result of G. Masbaum and A. Reid [MR12]. Recall that a group G involves a group H if there exist a finite index subgroup $L \leq G$ and a surjective map $\phi : L \rightarrow H$.

Corollary 2 (Masbaum–Reid). *Let $\Sigma_{g,m}$ be a surface of genus g with m punctures. If $3g - 3 + m \geq 1$ (or $g = 1$ and $m = 0$), then $\text{Mod}(\Sigma_{g,m})$ involves any finite group.*

The few mapping class groups not covered by the corollary are finite groups; see e.g. [FM12]. Corollary 2 is also proved using arithmetic methods by F. Grunewald, M. Larsen, A. Lubotzky, and J. Malestein [GLLM15].

Further applications of the quotients from Theorem 1 include new proofs of residual finiteness and separability of handlebody groups; see Section 3 for theorem statements and proofs.

Received by the editors December 5, 2017, and, in revised form, March 27, 2018.

2010 *Mathematics Subject Classification.* Primary 20E26, 20F65.

Key words and phrases. Mapping class groups, involve, finite groups.

The first author was supported in part by NSF grant DMS-1405609.

The second author was supported in part by NSF grant DMS-1510034.

1. PRELIMINARIES

1.1. G -defining subgroups. Here we collect some results surrounding definitions and discussions in Gilman [Gil77]. Let G and F be groups. A G -defining subgroup of F is a normal subgroup N of F such that F/N is isomorphic to G . Let $X(F, G)$ denote the set of all G -defining subgroups of F . The automorphism group $\text{Aut}(F)$ acts on normal subgroups of F while preserving their quotients and hence on the set $X(F, G)$ of G -defining subgroups of F . Since inner automorphisms act trivially, the action descends to an action of the outer automorphism group of F , $\text{Out}(F)$, on $X(F, G)$. If G is finite and F is finitely generated, one obtains a finite permutation representation of $\text{Out}(F)$. Let F_n be the free group of rank n . The following is Theorem 1 of [Gil77].

Theorem 3 (Gilman). *For any $n \geq 3$ and prime $p \geq 5$, $\text{Out}(F_n)$ acts on the $\text{PSL}(2, p)$ -defining subgroups of F_n as the alternating or symmetric group, and both cases occur for infinitely many primes.*

From the proof, Gilman obtains the following strengthened form of residual finiteness for $\text{Out}(F_n)$.

Corollary 4 (Gilman). *For any $n \geq 3$, the group $\text{Out}(F_n)$ is residually finite alternating and residually finite symmetric via the quotients from Theorem 3.*

This means that for any $\phi \in \text{Out}(F_n) - \{1\}$, there exist primes p so that the action of $\text{Out}(F_n)$ on $X(F_n, \text{PSL}(2, p))$ is alternating (and also primes p so that the action is symmetric), and ϕ acts nontrivially.

We will also need the following well-known fact, obtained from the classical embedding of a free group into $\text{PSL}(2, \mathbb{Z})$ as a subgroup of finite index (cf. A. Peluso [Pel66]).

Lemma 5. *For any $n \geq 2$, any element $\alpha \in F_n - \{1\}$, and all but finitely many primes p , there exists a $\text{PSL}(2, p)$ -defining subgroup of F_n not containing α .*

Proof. Let F_n be a finite index, free subgroup of rank n in the free group $F_2 := \langle a, b \rangle$. Identify F_n with its image in $\text{PSL}(2, \mathbb{Z})$ under the injective homomorphism $F_2 \rightarrow \text{PSL}(2, \mathbb{Z})$ given by

$$a \mapsto \begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix} \quad \text{and} \quad b \mapsto \begin{pmatrix} 1 & 0 \\ 2 & 1 \end{pmatrix}.$$

Let $\alpha \in F_n - \{1\}$ be given and let $A \in \text{SL}(2, \mathbb{Z})$ be a matrix representing α . Since $\alpha \neq 1$, we may assume that A has either a nonzero off-diagonal entry $d \neq 0$ or a diagonal entry $d > 1$. Then for any prime p not dividing d in the former case or $d \pm 1$ in the latter, we have that $\pi_p(\alpha)$ is nontrivial in the quotient $\pi_p : \text{PSL}(2, \mathbb{Z}) \rightarrow \text{PSL}(2, p)$; that is, $\alpha \notin \ker \pi_p$.

Since F_n has finite index in F_2 , there exists $m \geq 1$ so that the matrices

$$\begin{pmatrix} 1 & m \\ 0 & 1 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 1 & 0 \\ m & 1 \end{pmatrix}$$

represent elements of F_n in $\text{PSL}(2, \mathbb{Z})$. For any prime p not dividing m , the π_p -image of these elements generates $\text{PSL}(2, p)$. Thus, for all but finitely many primes p , $\ker \pi_p \cap F_n$ is a $\text{PSL}(2, p)$ -defining subgroup not containing α . \square

1.2. Handlebody subgroups and maps to free groups. Let $\Sigma = \Sigma_g$ be a closed surface of genus $g \geq 2$ and let $H = H_g$ be a handlebody of genus g . Given a homeomorphism $\phi: \Sigma \rightarrow \partial H \subset H$, the induced homomorphism is a surjection $\phi_*: \pi_1(\Sigma) \rightarrow \pi_1(H) \cong F_g$. As is well-known, every epimorphism $\pi_1(\Sigma) \rightarrow F_g$ arises in this way (see e.g. [LR02, Lemma 2.2]). The kernel $\Delta_\phi = \ker(\phi_*)$ is an F_g -defining subgroup and is the subgroup generated by the simple closed curves on Σ whose ϕ -images bound disks in H . We write H_ϕ for the handlebody H , equipped with the homeomorphism $\phi: \Sigma \rightarrow \partial H$.

Let $\text{Mod}(H_\phi)$ denote the subgroup of the mapping class group $\text{Mod}(\Sigma)$ consisting of the isotopy classes of homeomorphisms that extend over H_ϕ (via the identification $\phi: \Sigma \rightarrow \partial H$). Equivalently, $\text{Mod}(H_\phi)$ consists of those mapping classes $[f]$ such that $f_*(\Delta_\phi) = \Delta_\phi$; that is, $\text{Mod}(H_\phi)$ is the stabilizer in $\text{Mod}(\Sigma)$ of Δ_ϕ . Any element $[f] \in \text{Mod}(H_\phi)$ induces an automorphism we denote $\Phi_*([f]) \in \text{Out}(F_g)$, which defines a homomorphism $\Phi_*: \text{Mod}(H_\phi) \rightarrow \text{Out}(F_g)$. The main result of [Gri64] implies the next proposition.

Proposition 6. *For any $g \geq 0$ and homeomorphism*

$$\phi: \Sigma \rightarrow \partial H, \quad \Phi_*: \text{Mod}(H_\phi) \rightarrow \text{Out}(F_g)$$

is surjective.

The kernel of Φ_* , the set of mapping classes in $\text{Mod}(H_\phi)$ that act trivially on $\pi_1(H)$, is also a well-studied subgroup denoted $\text{Mod}_0(H_\phi)$.

Recall that $X(F_g, G)$ and $X(\pi_1(\Sigma), G)$ are the sets of G -defining subgroups of F_g and $\pi_1(\Sigma)$, respectively. Define

$$X^\phi(\pi_1(\Sigma), G) := \{\phi_*^{-1}(N) \mid N \in X(F_g, G)\} \subset X(\pi_1(\Sigma), G).$$

Alternatively, this is precisely the set of G -defining subgroups containing Δ_ϕ :

$$X^\phi(\pi_1(\Sigma), G) = \{N \in X(\pi_1(\Sigma), G) \mid \Delta_\phi < N\}.$$

Lemma 7. *The handlebody subgroup is contained in the stabilizer*

$$\text{Mod}(H_\phi) < \text{stab } X^\phi(\pi_1(\Sigma), G).$$

Moreover, if $\text{Out}(F_g)$ acts on $X(F_g, G)$ as the full symmetric group, then $\text{Mod}(H_\phi)$ (and hence $\text{stab } X^\phi(\pi_1(\Sigma), G)$) acts on $X^\phi(\pi_1(\Sigma), G)$ as the full symmetric group.

Proof. Let $N \in X^\phi(\pi_1(\Sigma), G)$ and let $[f] \in \text{Mod}(H_\phi)$, where f is a representative homeomorphism. Since $f_*(\Delta_\phi) = \Delta_\phi$, we have that $\Delta_\phi < f_*(N)$ and $f_*(N) \in X^\phi(\pi_1(\Sigma), G)$. Thus, f_* preserves $X^\phi(\pi_1(\Sigma), G)$, as required.

The last statement follows immediately from Proposition 6 and the fact that the bijection from the correspondence theorem $X^\phi(\pi_1(\Sigma), G) \rightarrow X(F_g, G)$ is Φ_* -equivariant. \square

2. MAPPING CLASS GROUPS INVOLVE ANY FINITE GROUP:
THE PROOFS OF THEOREM 1 AND COROLLARY 2

Here we give the proof of Theorem 1, following Dunfield–Thurston (see [DT06, pp. 505–506]).

Proof of Theorem 1. Fix $g \geq 3$ and let Π be the infinitely many primes for which $\text{Out}(F_g)$ acts on the $\text{PSL}(2, p)$ -defining subgroups as the symmetric group, guaranteed by Theorem 3. As a consequence of Corollary 4, the cardinality of

$X(F_g, \text{PSL}(2, p))$ is unbounded over any infinite set of primes p , and hence there exists a prime $p \in \Pi$ where the number of $\text{PSL}(2, p)$ -defining subgroups is $R \geq r$.

By Lemma 7, $\text{stab } X^\phi(\pi_1(\Sigma), \text{PSL}(2, p))$ acts on $X^\phi(\pi_1(\Sigma), \text{PSL}(2, p))$ as the symmetric group, defining a surjective homomorphism

$$\text{stab } X^\phi(\pi_1(\Sigma), \text{PSL}(2, p)) \rightarrow \text{Sym}(X^\phi(\pi_1(\Sigma), \text{PSL}(2, p))) \cong S_R.$$

Since $X(\pi_1(\Sigma), \text{PSL}(2, p))$ is a finite set, $\text{stab } X^\phi(\pi_1(\Sigma), \text{PSL}(2, p)) < \text{Mod}(\Sigma)$ has finite index, completing the proof. \square

Proof of Corollary 2. For g, m as in the statement and any $r \in \mathbb{N}$, we show that there is a finite index subgroup of $\text{Mod}(\Sigma_{g,m})$ that surjects a symmetric group on at least r elements. Since any finite subgroup is isomorphic to a subgroup of some such symmetric group, this will prove the theorem.

First observe that for any $m, g \geq 0$, the kernel of the action of $\text{Mod}(\Sigma_{g,m})$ on the m punctures of $\Sigma_{g,m}$ is a finite index subgroup $\text{Mod}'(\Sigma_{g,m}) < \text{Mod}(\Sigma_{g,m})$. Furthermore, if $0 \leq m < m'$, there is a surjective homomorphism

$$\text{Mod}'(\Sigma_{g,m'}) \rightarrow \text{Mod}'(\Sigma_{g,m})$$

obtained by “filling in” $m' - m$ of the punctures; see [FM12].

Now, because $\text{Mod}'(\Sigma_{0,4}) \cong F_2$ and the symmetric group on r elements is 2-generated, it follows that $\text{Mod}'(\Sigma_{0,4})$ surjects S_r . From the previous paragraph, it follows that $\text{Mod}'(\Sigma_{0,m})$ surjects S_r for all $m \geq 4$. Similarly, $\text{Mod}(\Sigma_{1,0}) \cong \text{SL}(2, \mathbb{Z})$, which has a finite index subgroup isomorphic to F_2 , and so there is a finite index subgroup of $\text{Mod}(\Sigma_{1,m})$ that surjects S_r for all $m \geq 0$. As shown in [BH71], there is a surjective homomorphism $\text{Mod}(\Sigma_{2,0}) \rightarrow \text{Mod}(\Sigma_{0,6})$, and consequently, we may find surjective homomorphisms from finite index subgroups of $\text{Mod}(\Sigma_{2,m})$ to S_r for all $m \geq 0$. From this and the previous paragraph, it suffices to assume $g \geq 3$ and $m = 0$. The required surjective homomorphism to a symmetric group in this case follows from Theorem 1, completing the proof. \square

3. SEPARATING HANDLEBODY SUBGROUPS AND RESIDUAL FINITENESS

The finite quotients of $\text{Mod}(\Sigma)$ coming from surjective homomorphisms $\pi_1(\Sigma_g) \rightarrow F_g$ also allow us to deduce the following result of [LM07]. Recall that a subgroup $K < F$ is said to be separable in F if for any $\alpha \in F - K$, there exists a finite index subgroup $G < F$ containing K and not containing α .

Theorem 8 (Leininger-McReynolds). *For any $g \geq 2$ and homeomorphism to the boundary of a handlebody, $\phi: \Sigma \rightarrow \partial H$, the groups $\text{Mod}(H_\phi)$ and $\text{Mod}_0(H_\phi)$ are separable in $\text{Mod}(\Sigma_g)$.*

While the proof of separability of $\text{Mod}(H_\phi)$ in $\text{Mod}(\Sigma_g)$ works for all $g \geq 2$, separability of $\text{Mod}_0(\Sigma_g)$ follows from the discussion here only when $g \geq 3$.

Proof. Let $\Sigma = \Sigma_g, g \geq 2$. For any p , any $[h] \in \text{stab } X^\phi(\pi_1(\Sigma), \text{PSL}(2, p))$, and any $\alpha \in \Delta_\phi$, we have $h_*(\alpha) \in K$ for all $K \in X^\phi(\pi_1(\Sigma), \text{PSL}(2, p))$. By Lemma 7 this is true for all $[h] \in \text{Mod}(H_\phi)$.

Now let $[f] \in \text{Mod}(\Sigma) - \text{Mod}(H_\phi)$, so that $f_*(\Delta_\phi) \not\subset \Delta_\phi$. Let $\gamma \in \Delta_\phi$ be an element such that (the conjugacy class of) $f_*(\gamma)$ is not in Δ_ϕ . (In fact, well-defining $f_*(\gamma)$ requires a choice of basepoint preserving representative homeomorphisms for the mapping class of f , which we make arbitrarily.) Then $\phi_*(f_*(\gamma)) \in F_g - \{1\}$,

and so by Lemma 5, we can find a prime p and a $\text{PSL}(2, p)$ -defining subgroup $N \in X(F_g, \text{PSL}(2, p))$ so that $\phi_*(f_*(\gamma)) \notin N$. Therefore,

$$f_*(\gamma) \notin \phi_*^{-1}(N) \in X^\phi(\pi_1(\Sigma), \text{PSL}(2, p)),$$

and hence $[f] \notin \text{stab } X^\phi(\pi_1(\Sigma), \text{PSL}(2, p))$. Since $\text{stab } X^\phi(\pi_1(\Sigma), \text{PSL}(2, p))$ is a finite index subgroup containing $\text{Mod}(H_\phi)$ (by Lemma 7) and not containing $[f]$, and since $[f]$ was arbitrary, it follows that $\text{Mod}(H_\phi)$ is separable.

Since $\text{Mod}_0(H_\phi) < \text{Mod}(H_\phi)$ and since $\text{Mod}(H_\phi)$ is separable, it suffices to consider an element $[f] \in \text{Mod}(H_\phi) \setminus \text{Mod}_0(H_\phi)$ and produce a finite index subgroup of $\text{Mod}(\Sigma)$ containing $\text{Mod}_0(H_\phi)$ and not containing $[f]$. For all p , $\text{Mod}_0(H_\phi)$ is contained in the subgroup of $\text{stab } X^\phi(\pi_1(\Sigma), \text{PSL}(2, p))$ consisting of those mapping classes that act trivially on $X^\phi(\pi_1(\Sigma), \text{PSL}(2, p))$. Since $[f] \notin \text{Mod}_0(H_\phi)$, $\Phi_*([f]) \neq 1$ in $\text{Out}(F_g)$. For $g \geq 3$, Corollary 4 implies that for some p , $\Phi_*([f])$ acts nontrivially on $X(F_g, \text{PSL}(2, p))$. Therefore, $[f]$ acts nontrivially on $X^\phi(\pi_1(\Sigma), \text{PSL}(2, p))$, and so the finite index subgroup $G < \text{Mod}(\Sigma)$ consisting of those mapping classes preserving the subset $X^\phi(\pi_1(\Sigma), \text{PSL}(2, p))$ and acting trivially on this does not contain $[f]$, proving that $\text{Mod}_0(H_\phi)$ is separable. \square

Mapping class groups were shown to be residually finite by Grossman as a consequence of the fact that surface groups are conjugacy separable; see [Gro75].

Theorem 9 (Grossman). *Mapping class groups are residually finite.*

Residual finiteness of $\text{Mod}(\Sigma_g)$ follows immediately from separability of the handlebody subgroups $\text{Mod}(H_\phi)$ and the following.

Lemma 10. *The intersection of all handlebody subgroups $\text{Mod}(H_\phi)$ over all homeomorphisms $\phi: \Sigma_g \rightarrow \partial H$ is trivial if $g \geq 3$ and isomorphic to $\mathbb{Z}/2\mathbb{Z}$ if $g = 2$. The intersection of handlebody subgroups $\text{Mod}_0(H_\phi)$ is trivial for all $g \geq 2$.*

Proof. In [Mas86], Masur proved that the limit set of the handlebody subgroup in the Thurston boundary of Teichmüller space is a nowhere dense subset. The intersection of all handlebody subgroups is a normal subgroup and so is either finite or else has limit set equal to the entire Thurston boundary. By Masur’s result, we must be in the former case, and hence the intersection of handlebody subgroups is finite. But $\text{Mod}(\Sigma_g)$ has no nontrivial finite, normal subgroups if $g \geq 3$, while for $g = 2$, the only nontrivial, finite normal subgroup is the order-two subgroup generated by the hyperelliptic involution. This proves the first statement. The second follows from the first and the fact that the hyperelliptic involution of Σ_2 induces a nontrivial automorphism of $F_2 \cong \pi_1(H)$, for any homeomorphism $\phi: \Sigma_2 \rightarrow H$. \square

Proof of Theorem 9 for $\text{Mod}(\Sigma_g)$, with $g \geq 2$. An equivalent formulation of residual finiteness is that the intersection of all finite index subgroups is trivial. Therefore Theorem 8 and Lemma 10 immediately imply the result. \square

ACKNOWLEDGMENTS

The authors would like to thank Alan Reid and Alex Lubotzky for helpful conversations and Benson Farb for suggesting improvements on an earlier draft.

REFERENCES

- [Bau62] Gilbert Baumslag, *On generalised free products*, Math. Z. **78** (1962), 423–438, DOI 10.1007/BF01195185. MR0140562
- [BH71] Joan S. Birman and Hugh M. Hilden, *On the mapping class groups of closed surfaces as covering spaces*, Advances in the theory of Riemann surfaces (Proc. Conf., Stony Brook, N.Y., 1969), Ann. of Math. Studies, No. 66, Princeton Univ. Press, Princeton, N.J., 1971, pp. 81–115. MR0292082
- [DT06] Nathan M. Dunfield and William P. Thurston, *Finite covers of random 3-manifolds*, Invent. Math. **166** (2006), no. 3, 457–521, DOI 10.1007/s00222-006-0001-6. MR2257389
- [FM12] Benson Farb and Dan Margalit, *A primer on mapping class groups*, Princeton Mathematical Series, vol. 49, Princeton University Press, Princeton, NJ, 2012. MR2850125
- [Gil77] Robert Gilman, *Finite quotients of the automorphism group of a free group*, Canad. J. Math. **29** (1977), no. 3, 541–551, DOI 10.4153/CJM-1977-056-3. MR0435226
- [GLLM15] Fritz Grunewald, Michael Larsen, Alexander Lubotzky, and Justin Malestein, *Arithmetic quotients of the mapping class group*, Geom. Funct. Anal. **25** (2015), no. 5, 1493–1542, DOI 10.1007/s00039-015-0352-5. MR3426060
- [Gri64] H. B. Griffiths, *Automorphisms of a 3-dimensional handlebody*, Abh. Math. Sem. Univ. Hamburg **26** (1963/1964), 191–210, DOI 10.1007/BF02992786. MR0159313
- [Gro75] Edna K. Grossman, *On the residual finiteness of certain mapping class groups*, J. London Math. Soc. (2) **9** (1974/75), 160–164, DOI 10.1112/jlms/s2-9.1.160. MR0405423
- [LM07] Christopher J. Leininger and D. B. McReynolds, *Separable subgroups of mapping class groups*, Topology Appl. **154** (2007), no. 1, 1–10, DOI 10.1016/j.topol.2006.03.013. MR2271769
- [LR02] Christopher J. Leininger and Alan W. Reid, *The co-rank conjecture for 3-manifold groups*, Algebr. Geom. Topol. **2** (2002), 37–50, DOI 10.2140/agt.2002.2.37. MR1885215
- [Mas86] Howard Masur, *Measured foliations and handlebodies*, Ergodic Theory Dynam. Systems **6** (1986), no. 1, 99–116, DOI 10.1017/S014338570000331X. MR837978
- [MKS04] Wilhelm Magnus, Abraham Karrass, and Donald Solitar, *Combinatorial group theory: Presentations of groups in terms of generators and relations*, 2nd ed., Dover Publications, Inc., Mineola, NY, 2004. MR2109550
- [MR12] Gregor Masbaum and Alan W. Reid, *All finite groups are involved in the mapping class group*, Geom. Topol. **16** (2012), no. 3, 1393–1411, DOI 10.2140/gt.2012.16.1393. MR2967055
- [Pel66] Ada Peluso, *A residual property of free groups*, Comm. Pure Appl. Math. **19** (1966), 435–437, DOI 10.1002/cpa.3160190406. MR0199245

DEPARTMENT OF MATHEMATICS, CITY COLLEGE OF NEW YORK, CUNY, NEW YORK, NEW YORK 10031

Email address: khalid.math@gmail.com

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS URBANA-CHAMPAIGN, URBANA, ILLINOIS 61801

Email address: c.j.leininger95@gmail.com