

ON THE FEJES TÓTH PROBLEM ABOUT THE SUM OF ANGLES BETWEEN LINES

DMITRIY BILYK AND RYAN W. MATZKE

(Communicated by Alexander Iosevich)

ABSTRACT. In 1959 Fejes Tóth posed a conjecture that the sum of pairwise nonobtuse angles between N unit vectors in \mathbb{S}^d is maximized by periodically repeated elements of the standard orthonormal basis. We obtain new improved upper bounds for this sum, as well as for the corresponding energy integral. We also provide several new approaches to the only settled case of the conjecture: $d = 1$.

1. INTRODUCTION

Fejes Tóth has formulated a variety of problems and conjectures about point distributions on the sphere. In particular, in 1959 he posed the following question [7]: what is the maximal value of the sum of pairwise acute angles defined by N vectors in the sphere \mathbb{S}^d ? More precisely, determine which N -element point sets $Z = \{z_1, \dots, z_N\} \subset \mathbb{S}^d$ maximize the discrete energy

$$(1.1) \quad E(Z) = \frac{1}{N^2} \sum_{i,j=1}^N \arccos |z_i \cdot z_j| = \frac{1}{N^2} \sum_{i,j=1}^N \min \{ \arccos(z_i \cdot z_j), \pi - \arccos(z_i \cdot z_j) \}.$$

He conjectured that this sum is maximized by the periodically repeated copies of the standard orthonormal basis.

Conjecture 1.1 (Fejes Tóth, 1959 [7]). *Let $d \geq 1$ and $N = m(d+1) + k$ with $m \in \mathbb{N}_0$ and $0 \leq k \leq d$. Then the discrete energy (1.1) on the d -dimensional sphere \mathbb{S}^d is maximized by the point set $Z = \{z_1, \dots, z_N\} \subset \mathbb{S}^d$ with $z_{p(d+1)+i} = e_i$, where $\{e_i\}_{i=1}^{d+1}$ is the standard orthonormal basis of \mathbb{R}^{d+1} , i.e., Z consists of $m+1$ copies of k elements of the orthonormal basis of \mathbb{R}^{d+1} and m copies of the remaining $d+1-k$ basis elements. In this case,*

$$E(Z) = \frac{\pi}{2N^2} (k(k-1)(m+1)^2 + 2km(d+1-k)(m+1) + (d-k)(d+1-k)m^2).$$

In particular, if $N = m(d+1)$, the sum is maximized by m copies of the orthonormal basis:

$$(1.2) \quad \max_{\substack{Z \subset \mathbb{S}^d \\ \#Z=N}} E(Z) = \frac{\pi}{2} \cdot \frac{d}{d+1}.$$

Received by the editors January 23, 2018.

2010 *Mathematics Subject Classification*. Primary 11K38, 52C35.

This work is supported by the NSF grant DMS 1665007 (the first author) and the NSF Graduate Research Fellowship 00039202 (the second author).

This conjecture has been independently stated in [9] for all $d \geq 1$ (Fejes Tóth originally formulated it just for \mathbb{S}^2).

We also formulate a continuous version of the conjecture. Let $\mathcal{B}(\mathbb{S}^d)$ denote the set of Borel probability measures on the sphere, i.e., $\mu \geq 0$ and $\mu(\mathbb{S}^d) = 1$, and let $F : [-1, 1] \rightarrow \mathbb{R}$ be a (bounded or positive) measurable function. For $\mu \in \mathcal{B}(\mathbb{S}^d)$ we define the energy integral of μ with respect to the potential F as

$$I_F(\mu) = \int_{\mathbb{S}^d} \int_{\mathbb{S}^d} F(x \cdot y) d\mu(x) d\mu(y).$$

In this notation, the discrete energy of a set $Z = \{z_1, \dots, z_N\}$ is

$$E_F(Z) = \frac{1}{N^2} \sum_{i,j=1}^N F(z_i \cdot z_j) = I_F \left(\frac{1}{N} \sum_{k=1}^N \delta_{z_k} \right).$$

In our case, $F(t) = \arccos |t|$ and we suppress the dependence on F in the subscript, as we did in (1.1), i.e.,

$$I(\mu) = \int_{\mathbb{S}^d} \int_{\mathbb{S}^d} \arccos |x \cdot y| d\mu(x) d\mu(y).$$

In accordance with Conjecture 1.1 it is natural to guess that $I(\mu)$ should be maximized by the measure whose mass is equally distributed between the elements of the orthonormal basis:

$$\nu_{ONB} = \frac{1}{d+1} \sum_{i=1}^{d+1} \delta_{e_i}.$$

Conjecture 1.2. *The energy integral $I(\mu)$ is maximized by ν_{ONB} :*

$$(1.3) \quad \max_{\mu \in \mathcal{B}(\mathbb{S}^d)} I(\mu) = I(\nu_{ONB}) = \frac{\pi}{2} \cdot \frac{d}{d+1}.$$

Since the maximal value in (1.2) is independent of N , a simple argument based on the weak*-density of the span of Dirac masses in $\mathcal{B}(\mathbb{S}^d)$ shows that the case $N = m(d+1)$ of Conjecture 1.1 implies Conjecture 1.2, and the converse implication is obvious.

The case $d = 1$ of Conjecture 1.1 has been settled in [8, 9]. We shall return to this case in §4 and shall give several alternative proofs.

In dimension $d = 2$, Fejes Tóth confirmed Conjecture 1.1 for $N \leq 6$ and established an asymptotic upper bound $E(Z) \leq \frac{2\pi}{5}$ for large N . In [8] Fodor, Vígh, and Zarnócz proved that for any point distribution $Z \subset \mathbb{S}^2$

$$(1.4) \quad E(Z) \leq \frac{3\pi}{8} \quad \text{when } N \text{ is even,}$$

with a small correction for N odd.

In the present paper, we prove a new upper bound for $I(\mu)$ in all dimensions $d \geq 2$, which is stronger than (1.4) when restricted to \mathbb{S}^2 .

Theorem 1.3. *In all dimensions $d \geq 2$*

$$\max_{\mu \in \mathcal{B}(\mathbb{S}^d)} I(\mu) \leq \frac{\pi}{2} - \frac{69}{50(d+1)}.$$

In particular, for $d = 2$,

$$\max_{\mu \in \mathcal{B}(\mathbb{S}^2)} I(\mu) \leq \frac{\pi}{2} - \frac{69}{150} = 1.110796\dots < \frac{3\pi}{8} = 1.178097\dots$$

We recall that, according to (1.2) and (1.3), the conjectured maximal value in dimension $d = 2$ is $\frac{\pi}{3} = 1.047198\dots$. The proof of Theorem 1.3 is based on bounding the potential $F(t) = \arccos |t|$ above by a quadratic polynomial and employing the estimates for the so-called “frame potential”. It is presented in §2.

It is worth noting that the energy optimization problem considered here has a distinctly different flavor compared to many standard problems of this kind; see, e.g., [5]. In the vast majority of energy minimization problems, the potential $F(t)$ is maximized at $t = 1$ and minimized at $t = -1$. Such is the case, for example, for the classical Riesz potential $F(x \cdot y) = \|x - y\|^{-s}$. If one loosely interprets points as “electrons”, and measures as charge distributions, this results in the strongest repulsion when the electrons are close to each other, and the weakest when they are far away. In this situation one often expects $I_F(\mu)$ to be minimized by σ , the normalized Lebesgue surface measure on \mathbb{S}^d —in other words, minimizing energy induces uniform distribution.

In our setting, however, the smallest value of $-F(t)$ is at $t = 0$, hence the weakest repulsion occurs when two electrons are positioned orthogonally to each other (notice that we are maximizing, not minimizing, the energy). Potentials with such behavior arise in various problems, e.g., $F(t) = t^2$ is known as the “frame potential” (see [1] and Lemma 2.1 below) and $F(t) = |t|^p$ as the p -frame potential [6]. Possible optimizers for such energies naturally include orthonormal bases, other structures which exhibit similar orthogonality properties (e.g., frames), or the uniform distribution σ .

The case of $F(t) = \arccos(t)$, rather than $F(t) = \arccos |t|$ (i.e., the sum of actual angles, rather than the acute line angles), has been studied in [3]: in this situation *every* symmetric measure is a maximizer, with a certain correction in the discrete case for odd N .

In the present paper, in addition to proving Theorem 1.3, we explore some other issues related to this problem. In §3, we observe that the validity of Conjecture 1.2 in some dimension $d \geq 2$ implies its validity in all lower dimensions, and in §4 we revisit the one-dimensional case of Conjectures 1.1 and 1.2 (proved, e.g., in [7], [8]) and provide several alternative proofs: two based on orthogonal expansions, and one based on the Stolarsky principle in discrepancy theory. We hope that these approaches may shed some light on the conjectures in the future.

2. THE FRAME POTENTIAL AND THE PROOF OF THEOREM 1.3

Since our approach is based on bounding the potential $F(t) = \arccos |t|$ by a quadratic function, we first study the behavior of the energy with the potential t^2 . We have the following lemma.

Lemma 2.1. *For any $\mu \in \mathcal{B}(\mathbb{S}^d)$, we have*

$$(2.1) \quad I_{t^2}(\mu) \geq \frac{1}{d+1}.$$

The equality above is achieved precisely for the measures μ whose second moment matrix is a multiple of the identity, i.e., $[\int_{\mathbb{S}^d} x_i x_j d\mu(x)]_{i,j=1}^{d+1} = c \mathbf{I}_{d+1}$.

Proof. We expand the square, throw away off-diagonal terms, and apply the Cauchy–Schwartz inequality:

$$\begin{aligned} I_{t^2}(\mu) &= \int_{\mathbb{S}^d} \int_{\mathbb{S}^d} (x \cdot y)^2 \, d\mu(x) d\mu(y) = \sum_{i,j=1}^{d+1} \left(\int_{\mathbb{S}^d} x_i x_j \, d\mu(x) \right)^2 \\ &\geq \sum_{i=1}^{d+1} \left(\int_{\mathbb{S}^d} x_i^2 \, d\mu(x) \right)^2 \geq \frac{1}{d+1} \left(\sum_{i=1}^{d+1} \int_{\mathbb{S}^d} x_i^2 \, d\mu(x) \right)^2 = \frac{1}{d+1}. \end{aligned}$$

The characterization of minimizers follows immediately from the inequalities above. \square

The energy with the potential $F(t) = t^2$ arises naturally in functional analysis and signal processing. In [1], Benedetto and Fickus defined the discrete “frame potential”, which up to normalization is equal to $E_{t^2}(Z)$, and proved that for $N \geq d+1$

$$(2.2) \quad E_{t^2}(Z) \geq \frac{1}{d+1}$$

with equality achieved if and only if the set $Z = \{z_1, \dots, z_N\} \subset \mathbb{S}^d$ forms a so-called tight frame, i.e., for each $x \in \mathbb{R}^{d+1}$ one has an analogue of Parseval’s identity:

$$c\|x\|^2 = \sum_{k=1}^N |x \cdot z_k|^2. \text{ It is easy to see that Lemma 2.1 implies this discrete result}$$

(although much more is proved in [1], in particular, that every local minimizer is necessarily global). Conversely, since the minimum in (2.2) is independent of N , the weak*-density of probability measures shows that the result of Benedetto and Fickus also implies (2.1). Moreover, inequality (2.1) for arbitrary measures $\mu \in \mathcal{B}(\mathbb{S}^d)$ has been stated in [6]. We have included the nice and short proof above for the sake of completeness.

To prove Theorem 1.3, it suffices to demonstrate that the inequality

$$(2.3) \quad \arccos |t| \leq \frac{\pi}{2} - \frac{69}{50} t^2$$

holds for all $t \in [-1, 1]$, as then (2.1) would imply that, for all $\mu \in \mathcal{B}(\mathbb{S}^d)$,

$$(2.4) \quad I(\mu) \leq \frac{\pi}{2} - \frac{69}{50} I_{t^2}(\mu) \leq \frac{\pi}{2} - \frac{69}{50(d+1)}.$$

A simple calculus exercise shows $\arccos |t| \leq \frac{\pi}{2} - bt^2$ for all $t \in [-1, 1]$ if and only if

$$\frac{\pi}{2} - \frac{b + \sqrt{b^2 - 1}}{2} - \arccos \left(\sqrt{\frac{b + \sqrt{b^2 - 1}}{2b}} \right) \geq 0,$$

and a quick computation shows that for $b \leq \frac{69}{50}$ the inequality above indeed holds, finishing the proof of Theorem 1.3. In Figure 1 we include the graph, illustrating inequality (2.3). \square

It is easy to see that there is very little room for improvement via this method, and it will definitely not yield the Conjecture 1.2.




FIGURE 1. An illustration to inequality (2.3): the graph of the function $f(t) = \frac{\pi}{2} - \frac{69}{50}t^2 - \arccos|t|$ for $0 \leq t \leq 1$.

3. DIMENSION REDUCTION ARGUMENT

Let $F : [-1, 1] \rightarrow \mathbb{R}$ be a bounded, measurable function that achieves its maximum at 0. Let $\nu \in \mathcal{B}(\mathbb{S}^k)$ and $\lambda \in \mathcal{B}(\mathbb{S}^l)$, with $k + l = d - 1$. Construct a measure $\mu \in \mathcal{B}(\mathbb{S}^d)$ as follows:

$$(3.1) \quad \mu = \alpha\tilde{\nu} + \beta\tilde{\lambda},$$

where $0 \leq \alpha, \beta \leq 1$, $\alpha + \beta = 1$, and $\tilde{\nu}$ and $\tilde{\lambda}$ are copies of ν and λ , supported on mutually orthogonal subspheres of \mathbb{S}^d of dimensions k and l , respectively. It is easy to see that

$$(3.2) \quad I_F(\mu) = \alpha^2 I_F(\nu) + \beta^2 I_F(\lambda) + 2\alpha\beta F(0),$$

which easily implies that if μ is a maximizer of I_F over $\mathcal{B}(\mathbb{S}^d)$, then ν and λ maximize I_F , respectively, over $\mathcal{B}(\mathbb{S}^k)$ and $\mathcal{B}(\mathbb{S}^l)$.

The orthonormal basis measure ν_{ONB} is precisely of the form (3.1) with ν and λ also being orthonormal basis measures in lower dimensions. Therefore, in particular, we have just proved that the validity of Conjecture 1.2 on \mathbb{S}^d for some dimension $d \geq 2$ implies its validity on \mathbb{S}^k for $k < d$, i.e., in all lower dimensions.

Moreover, in our case, for $F(t) = \arccos|t|$, we have $F(0) = \frac{\pi}{2}$. Let $l = 0$ and $k = d - 1$. Then $I(\lambda) = F(1) = 0$, and thus (3.2) becomes

$$I(\mu) = \alpha^2 I(\nu) + \pi\alpha(1 - \alpha).$$

Optimizing this quadratic polynomial in α we find that for $\alpha = \frac{\pi}{2(\pi - I_F(\nu))}$

$$I(\nu) = \pi - \frac{\pi^2}{4I(\mu)}.$$

This discussion leads to the following conclusion.

Proposition 3.1 (Dimension reduction).

(1) Denote $M_d = \max\{I(\mu) : \mu \in \mathcal{B}(\mathbb{S}^d)\}$. Then

$$M_{d-1} \leq \pi - \frac{\pi^2}{4M_d}.$$

(2) Assume that Conjecture 1.2 holds on \mathbb{S}^d , i.e., $M_d = \frac{\pi d}{2(d+1)}$. Then it also holds on \mathbb{S}^{d-1} , and consequently in all lower dimensions.

Observe, that part (2) follows both from part (1) and, in a more general setting, from the discussion in the beginning of this section. This proposition implies that, in order to prove Conjecture 1.2 in all dimensions $d \geq 2$, it is enough to establish its validity for infinitely many values of d .

4. THE CASE OF \mathbb{S}^1 REVISITED

We now revisit the case $d = 1$, in which Conjecture 1.2 (and hence also Conjecture 1.1) has been settled. We shall provide several new approaches to this case (two based on orthogonal expansions, and one based on connections to discrepancy theory), which we hope may lead to future progress in higher dimensions.

Before we proceed, we observe that on \mathbb{S}^1

$$(4.1) \quad I(\nu_{ONB}) = I(\sigma) = I(\sigma_{4N}) = \frac{\pi}{4},$$

where σ is the normalized uniform measure $d\sigma = \frac{d\theta}{2\pi}$, and $\sigma_{4N} = \frac{1}{4N} \sum_{k=1}^{4N} \delta_{e^{\pi i k/2N}}$ is the measure with mass equally concentrated at $4N$ equally spaced points. Hence, to prove Conjecture 1.2 it suffices to prove that any of these measures is a maximizer.

4.1. Chebyshev polynomial expansion. It is well known (see, e.g., [2, Proposition 2.1]) that σ is a maximizer of the energy integral I_F on \mathbb{S}^1 if and only if F is a negative definite function on \mathbb{S}^1 (up to the constant term), which is equivalent to the fact that in the orthogonal expansion of F into Chebyshev polynomials, $F(t) \sim \sum_{n=0}^{\infty} \tilde{F}_n T_n(t)$, the coefficients of all nonconstant terms are nonpositive, i.e.,

$$\tilde{F}_n \leq 0 \quad \text{for } n \geq 1.$$

Here T_n is the n th Chebyshev polynomial of the first kind, that is,

$$(4.2) \quad T_n(t) = \cos(n \arccos(t)).$$

A straightforward but technical computation (which we omit) shows that for $F(t) = \arccos |t|$,

$$(4.3) \quad \tilde{F}_n = \frac{1}{\pi} \int_{-1}^1 F(t) T_n(t) (1-t^2)^{-\frac{1}{2}} dt = \begin{cases} \frac{\pi}{4}, & \text{if } n = 0, \\ \frac{-4}{\pi n^2}, & \text{if } n \equiv 2 \pmod{4}, \\ 0, & \text{otherwise.} \end{cases}$$

Therefore, σ maximizes I_F on \mathbb{S}^1 , which together with (4.1) implies Conjecture 1.2.

4.2. Fourier series. Our second orthogonal expansion is a Fourier series. Every point in \mathbb{S}^1 can be defined by its angle, so with a slight abuse of notation we can rewrite our energy integral as

$$I_F(\mu) = \int_{\mathbb{T}} \int_{\mathbb{T}} F(\cos(\theta - \phi)) d\mu(\theta) d\mu(\phi) = \int_{\mathbb{T}} \int_{\mathbb{T}} G(\theta - \phi) d\mu(\theta) d\mu(\phi).$$

Let $G(\theta) = F(\cos(\theta))$ be an even function. Then for $\nu \in \mathcal{B}(\mathbb{S}^d)$ defined by $d\nu(\theta) = \frac{d\mu(\theta) + d\mu(-\theta)}{2}$ for all $\theta \in \mathbb{T}$, we have $I_F(\mu) = I_F(\nu)$. Thus, for the rest of this section, we may assume that μ is an even measure. As long as the Fourier series of G , which

is a cosine series $G(\theta) \sim \sum_{n=0}^{\infty} \widehat{G}(n) \cos n\theta$, converges absolutely, we can use it to compute the energy:

$$\begin{aligned} I_F(\mu) &= \sum_{n=0}^{\infty} \widehat{G}(n) \int_{\mathbb{T}} \int_{\mathbb{T}} \cos(n(\theta - \phi)) \, d\mu(\theta) d\mu(\phi) \\ &= \sum_{n=0}^{\infty} \widehat{G}(n) \int_{\mathbb{T}} \int_{\mathbb{T}} \left(\cos(n\theta) \cos(n\phi) + \sin(n\theta) \sin(n\phi) \right) \, d\mu(\theta) d\mu(\phi). \end{aligned}$$

Since μ is an even measure, the sines do not contribute to the integral, and we have

$$(4.4) \quad I_F(\mu) = \sum_{n=0}^{\infty} \widehat{G}(n) \left(\int_{\mathbb{T}} \cos(n\theta) d\mu(\theta) \right)^2 \leq \widehat{G}(0) + \sum_{n \geq 1: \widehat{G}(n) \geq 0} \widehat{G}(n).$$

The equality above is achieved if and only if $\int_{\mathbb{T}} \cos(n\theta) d\mu(\theta) = 0$ for each value of $n \geq 1$ for which $\widehat{G}(n) < 0$ and $\int_{\mathbb{T}} \cos(n\theta) d\mu(\theta) = 1$ for each value of $n \geq 1$ such that $\widehat{G}(n) > 0$.

Lemma 4.1. *Let G be even and have absolutely convergent Fourier (cosine) series, and let σ_N be the probability measure generated by point masses at N equally spaced points*

$$\sigma_N = \frac{1}{N} \sum_{k=0}^{N-1} \delta_{2\pi k/N}.$$

Assume that for all $n \geq 1$ we have $\widehat{G}(n) \geq 0$ if n is a multiple of N , and $\widehat{G}(n) \leq 0$ otherwise. Then the measure σ_N maximizes I_F over all probability measures on \mathbb{T} .

The proof easily follows from the discussion above and the fact that

$$\int_{\mathbb{T}} \cos(n\theta) d\sigma_N(\theta) = \frac{1}{N} \sum_{k=0}^{N-1} \cos \frac{2\pi nk}{N} = \begin{cases} 1, & \text{if } n \text{ is a multiple of } N, \\ 0, & \text{otherwise.} \end{cases}$$

Remark. The conditions on $\widehat{G}(n)$ in Lemma 4.1 also become necessary if we assume that σ_N maximizes I_F over all *signed* (not just nonnegative) measures of total mass one. This can be proved by considering signed measures of the form $d\mu = d\sigma_N - \cos n\theta \, d\sigma$.

Returning to our specific case, $G(\theta) = F(\cos(\theta)) = \min\{|\theta|, \pi - |\theta|\}$, we observe that according to definition (4.2) and relation (4.3), we have $\widehat{G}(n) = \widetilde{F}_n = -\frac{4}{\pi n^2}$ whenever $n \equiv 2 \pmod{4}$, and $\widehat{G}(n) = 0$ for all other values of $n \geq 1$. Therefore, Lemma 4.1 with $N = 4$ applies and $\sigma_4 = \frac{1}{4}(\delta_0 + \delta_{\frac{\pi}{2}} + \delta_{\pi} + \delta_{\frac{3\pi}{2}})$ maximizes I_F . Symmetry implies that $\nu_{ONB} = \frac{1}{2}(\delta_0 + \delta_{\frac{\pi}{2}})$ is also a maximizer, i.e., Conjecture 1.2 holds on \mathbb{S}^1 .

Notice that alternatively, since $\widehat{G}(n) \leq 0$ for all $n \geq 1$, one could deduce from (4.4) that $I_F(\mu) \leq \widehat{G}(0) = I_F(\sigma)$. Therefore σ is a maximizer of the energy integral I_F , and, due to (4.1), so is ν_{ONB} , leading to another proof of Conjecture 1.2 (almost identical to §4.1). We, however, wanted to introduce Lemma 4.1 into play.

4.3. Stolarsky principle. Finally, our last approach is based on an intriguing connection between energy optimization and discrepancy. The classical *Stolarsky invariance principle* [10] relates the L^2 spherical cap discrepancy to a certain discrete energy (the sum of pairwise Euclidean distances, i.e., $F(x \cdot y) = \|x - y\|$). Some generalizations of this principle were obtained in [2, 3]. Moreover, in [3] a version of this principle has been used to characterize the extremizers of the energy integral and discrete energy with the potential $F(t) = \arccos t$ (i.e., $F(x \cdot y)$ represents the angle or the geodesic distance between x and $y \in \mathbb{S}^d$). We shall apply this idea to our problem—the main task is to find an appropriate version of L^2 discrepancy.

For $x \in \mathbb{S}^1$, define the *antipodal quadrants* in the direction of x as $Q(x) = \{y : |x \cdot y| > \frac{\sqrt{2}}{2}\}$, i.e., $Q(x)$ is a union of two symmetric quartercircle arcs with midpoints at x and $-x$. Observe that the size of the intersection of two such sets depends linearly on the acute angle between their directions:

$$(4.5) \quad \sigma(Q(y) \cap Q(z)) = \frac{1}{2} - \frac{1}{\pi} \arccos |y \cdot z|.$$

We define the L^2 discrepancy of a point set $Z = \{z_1, \dots, z_N\} \subset \mathbb{S}^1$ with respect to these antipodal quadrants as the quadratic average of the difference between the empirical measure $\frac{\#\{Q(x) \cap Z\}}{N}$ and the uniform measure $\sigma(Q(x)) = \frac{1}{2}$:

$$(4.6) \quad D_{L^2, \text{quad}}^2(Z) = \int_{\mathbb{S}^1} \left| \frac{1}{N} \sum_{i=1}^N \mathbb{1}_{Q(x)}(z_i) - \sigma(Q(x)) \right|^2 d\sigma(x).$$

More generally, for a measure $\mu \in \mathcal{B}(\mathbb{S}^1)$ we can define its discrepancy as

$$(4.7) \quad D_{L^2, \text{quad}}^2(\mu) = \int_{\mathbb{S}^1} |\mu(Q(x)) - \sigma(Q(x))|^2 d\sigma(x),$$

i.e., $D_{L^2, \text{quad}}^2(Z) = D_{L^2, \text{quad}}^2(\frac{1}{N} \sum \delta_{z_i})$. The following version of the Stolarsky principle holds.

Lemma 4.2 (Stolarsky principle for quadrants). *For any measure $\mu \in \mathcal{B}(\mathbb{S}^1)$ we have*

$$(4.8) \quad D_{L^2, \text{quad}}^2(\mu) = \frac{1}{\pi} (I(\sigma) - I(\mu)) = \frac{1}{4} - \frac{1}{\pi} I(\mu).$$

In particular, for a discrete point set $Z = \{z_1, \dots, z_N\} \subset \mathbb{S}^1$

$$(4.9) \quad D_{L^2, \text{quad}}^2(Z) = \frac{1}{4} - \frac{1}{\pi} E(Z).$$

Proof. We use relations (4.7) and (4.5) to obtain

$$\begin{aligned} D_{L^2, \text{quad}}^2(\mu) &= \int_{\mathbb{S}^1} \int_{\mathbb{S}^1} \int_{\mathbb{S}^1} \mathbb{1}_{Q(x)}(y) \cdot \mathbb{1}_{Q(x)}(z) d\sigma(x) d\mu(y) d\mu(z) \\ &\quad - \int_{\mathbb{S}^1} \int_{\mathbb{S}^1} \mathbb{1}_{Q(x)}(y) d\sigma(x) d\mu(y) + \frac{1}{4} \\ &= \int_{\mathbb{S}^1} \int_{\mathbb{S}^1} \sigma(Q(y) \cap Q(z)) - \frac{1}{4} = \frac{1}{4} - \frac{1}{\pi} \int_{\mathbb{S}^1} \int_{\mathbb{S}^1} \arccos |y \cdot z| d\mu(y) d\mu(z), \end{aligned}$$

which proves (4.8). □

Remark. Discrepancy with respect to similar sets, but with arbitrary apertures, (“wedges”) on \mathbb{S}^d arose in [4] in connection to sphere tessellations and one-bit sensing, which led to a Stolarsky principle involving energies with the potential $F(t) = (\arcsin t)^2$.

Our Stolarsky principle (4.8) proves that $I(\mu) \leq \frac{\pi}{4}$ and σ maximizes this energy, and by (4.1) so does ν_{ONB} , proving Conjecture 1.2. In the discrete case, (4.9) shows that $E(Z) \leq \frac{\pi}{4}$ for even N and $E(Z) \leq \frac{\pi}{4} \cdot \frac{N^2-1}{N^2}$ for odd N . It also allows one to characterize the maximizers of $E(Z)$ and prove Conjecture 1.1: (4.9) implies that maximizing $E(Z)$ is equivalent to minimizing the discrepancy $D_{L^2, \text{quad}}^2(Z)$. It is easy to see from (4.6) that this happens exactly when the following holds: for σ -almost every $x \in \mathbb{S}^d$ the difference between the number of points of Z in $Q(x)$ and in its complement $(Q(x))^c$ is zero (when N is even) or ± 1 (when N is odd). Equivalently, this should hold for any x such that the boundary of $Q(x)$ doesn’t intersect Z , which recovers the characterization in [8]. The extremizing configurations in Conjecture 1.1 obviously satisfy this condition.

Unfortunately, none of these approaches immediately extends to higher dimensions: for $d \geq 2$ the function $F(t) = \arccos |t|$ is not negative definite, (4.1) fails, and the absence of analogues of (4.5) prevents one from obtaining a Stolarsky principle. However, we hope that some of these ideas with additional ingredients may bring progress on the conjectures in the future.

REFERENCES

- [1] John J. Benedetto and Matthew Fickus, *Finite normalized tight frames*, Adv. Comput. Math. **18** (2003), no. 2-4, 357–385, DOI 10.1023/A:1021323312367. Frames. MR1968126
- [2] D. Bilyk and F. Dai, *Geodesic Riesz Energy on the Sphere* (2017), arXiv:1612.08442.
- [3] Dmitriy Bilyk, Feng Dai, and Ryan Matzke, *The Stolarsky principle and energy optimization on the sphere*, Constr. Approx. **48** (2018), no. 1, 31–60, DOI 10.1007/s00365-017-9412-4. MR3825946
- [4] Dmitriĭ Bilik and Maĭkl T. Lëisi, *One bit sensing, discrepancy, and the Stolarsky principle* (Russian, with Russian summary), Mat. Sb. **208** (2017), no. 6, 4–25, DOI 10.4213/sm8656; English transl., Sb. Math. **208** (2017), no. 5-6, 744–763. MR3659577
- [5] S. Borodachov, D. Hardin, and E. Saff, *Minimal Discrete Energy on Rectifiable Sets*, Springer, Monographs in Math. (to appear).
- [6] M. Ehler and K. A. Okoudjou, *Minimization of the probabilistic p -frame potential*, J. Statist. Plann. Inference **142** (2012), no. 3, 645–659, DOI 10.1016/j.jspi.2011.09.001. MR2853573
- [7] L. Fejes Tóth, *Über eine Punktverteilung auf der Kugel* (German, with Russian summary), Acta Math. Acad. Sci. Hungar **10** (1959), 13–19 (unbound insert), DOI 10.1007/BF02063286. MR0105654
- [8] F. Fodor, V. Vigh, and T. Zarnócz, *On the angle sum of lines*, Arch. Math. (Basel) **106** (2016), no. 1, 91–100, DOI 10.1007/s00013-015-0847-1. MR3451371
- [9] F. Petrov (<https://mathoverflow.net/users/4312/fedor-petrov>), *Maximum sum of angles between n lines*, URL (version: 2014-07-09): <https://mathoverflow.net/q/173712>.
- [10] Kenneth B. Stolarsky, *Sums of distances between points on a sphere. II*, Proc. Amer. Math. Soc. **41** (1973), 575–582, DOI 10.2307/2039137. MR0333995

SCHOOL OF MATHEMATICS, UNIVERSITY OF MINNESOTA, MINNEAPOLIS, MINNESOTA 55408
Email address: dbilyk@math.umn.edu

SCHOOL OF MATHEMATICS, UNIVERSITY OF MINNESOTA, MINNEAPOLIS, MINNESOTA 55408
Email address: matzk053@umn.edu