

ON THE LOCAL WELL-POSEDNESS FOR A FULL-DISPERSION BOUSSINESQ SYSTEM WITH SURFACE TENSION

HENRIK KALISCH AND DIDIER PILOD

(Communicated by Catherine Sulem)

ABSTRACT. In this note, we prove local-in-time well-posedness for a fully dispersive Boussinesq system arising in the context of free surface water waves in two and three spatial dimensions. Those systems can be seen as a weak nonlocal dispersive perturbation of the shallow-water system. Our method of proof relies on energy estimates and a compactness argument. However, due to the lack of symmetry of the nonlinear part, those traditional methods have to be supplemented with the use of a modified energy in order to close the a priori estimates.

1. INTRODUCTION

Consideration is given to the one-dimensional fully dispersive Boussinesq system

$$(1.1) \quad \begin{cases} \partial_t \eta + \mathcal{K}(D)\partial_x u + \partial_x(\eta u) = 0, \\ \partial_t u + \partial_x \eta + u\partial_x u = 0, \end{cases}$$

where $x \in \mathbb{R}$, $t \in \mathbb{R}$, $\eta(x, t) \in \mathbb{R}$, and $u(x, t) \in \mathbb{R}$, and its two-dimensional counterpart

$$(1.2) \quad \begin{cases} \partial_t \eta + \mathcal{K}(D)\nabla \cdot \mathbf{u} + \nabla \cdot (\eta \mathbf{u}) = 0, \\ \partial_t \mathbf{u} + \nabla \eta + \frac{1}{2}\nabla |\mathbf{u}|^2 = 0, \end{cases}$$

for $x \in \mathbb{R}^2$, $t \in \mathbb{R}$, $\eta(x, t) \in \mathbb{R}$, and $\mathbf{u}(x, t) \in \mathbb{R}^2$, where $\mathcal{K}(D)$ is a nonlocal operator related to the dispersion of the linearized water-wave system in finite depth. Namely, $\mathcal{K}(D)$ is defined as a Fourier multiplier associated with the symbol

$$(1.3) \quad K(\xi) = \frac{\tanh(|\xi|)}{|\xi|} (1 + \beta|\xi|^2),$$

where β is a nonnegative dimensionless number related to the surface tension (see [25]).

Those systems were proposed in [1, 19, 21] as approximate models for the study of surface water waves and provide a two-directional alternative to the well-known Whitham equation. We also refer to [6–8, 14] for other versions of full-dispersion Boussinesq type systems. The unknowns η and u in (1.1) represent respectively the deflection of the free surface from its equilibrium position ($\eta = 0$) and the velocity at the free surface, while the bottom is assumed to be at constant depth $h = -1$.

Received by the editors May 22, 2018, and, in revised form, August 30, 2018.

2010 *Mathematics Subject Classification*. Primary 35Q53, 35A01, 76B15; Secondary 35E05, 35E15.

This research was supported by the Bergen Research Foundation (BFS), the Research Council of Norway, and the University of Bergen.

The one-dimensional Whitham equation

$$(1.4) \quad \partial_t u + \mathcal{W}(D)\partial_x u + u\partial_x u = 0,$$

where $x \in \mathbb{R}, t \in \mathbb{R}, u = u(x, t) \in \mathbb{R}$ and $\mathcal{W}(D)$ is the Fourier multiplier associated with the symbol $W(\xi) := \sqrt{K(\xi)}$, was introduced by Whitham in [30] as an alternative to the Korteweg-de Vries (KdV) equation by keeping the exact dispersion of the linearized water waves system in finite depth. This equation has drawn quite a bit of attention lately. In particular, it displays, in the case of pure gravity waves ($\beta = 0$), several interesting phenomena already predicted by Whitham: a solitary wave regime close to KdV [9], the existence of a wave of greatest height (Stokes wave) [11], the existence of shocks [13], and modulational instability of steady periodic waves [15, 26]. Note that when surface tension is taken into account ($\beta > 0$), the dynamics of (1.4) appears to be completely different (see [19] and the references therein). Moreover, it was proved to be a relevant water wave model in the long wave regime on the same time scale as the KdV equation [19, 21].¹ We also refer to [4, 19, 29] for other interesting numerical simulations.

Returning to the full-dispersion systems (1.1) and (1.2), it has been shown in [10] that (1.1) is locally well-posed in the case of pure gravity waves ($\beta = 0$) if one makes the assumption that the initial elevation η is bounded below by a positive constant. No results seem to be known when surface tension is taken into account, i.e., in the case $\beta > 0$.

Our main result in this note is a proof of well-posedness for systems (1.1) and (1.2) in the case of $\beta > 0$ under a noncavitation assumption on the initial surface elevation $\eta(\cdot, 0)$.

Definition 1.1. Let $d = 1$ or 2 and let $s > \frac{d}{2}$. We say that the initial elevation $\eta_0 \in H^s(\mathbb{R}^2)$ satisfies the *noncavitation condition* if

$$(1.5) \quad \exists h_0 \in (0, 1) \quad \text{such that} \quad 1 + \eta_0(x) \geq h_0, \quad \forall x \in \mathbb{R}^d.$$

Remark 1.2. The noncavitation condition (1.5) is a physical condition, meaning that the elevation of the initial wave cannot touch the bottom of the fluid.

Theorem 1.3. Assume that $\beta > 0$.

(i) Let $s > \frac{5}{2}$. Let $(\eta_0, u_0) \in H^s(\mathbb{R}) \times H^{s+\frac{1}{2}}(\mathbb{R})$ satisfying the noncavitation condition (1.5). Then, there exists a positive time $T = T(\|(\eta_0, u_0)\|_{H^s \times H^{s+\frac{1}{2}}})$ (which can be chosen as a nondecreasing function of its argument) and a unique solution (η, u) to (1.1) satisfying

$$(1.6) \quad (\eta, u) \in C([0, T] : H^s(\mathbb{R}) \times H^{s+\frac{1}{2}}(\mathbb{R})) \quad \text{and} \quad (\eta(\cdot, 0), u(\cdot, 0)) = (\eta_0, u_0).$$

In addition, the flow function mapping initial data to solutions is continuous.

(ii) Let $s > 3$. Let $(\eta_0, \mathbf{u}_0) \in H^s(\mathbb{R}^2) \times H^{s+\frac{1}{2}}(\mathbb{R}^2)^2$ satisfy the noncavitation condition (1.5) such that $\text{curl} \mathbf{u}_0 = 0$. Then, there exists a positive time $T = T(\|(\eta_0, \mathbf{u}_0)\|_{H^s \times H^{s+\frac{1}{2}} \times H^{s+\frac{1}{2}}})$ (which can be chosen as a nondecreasing function of its argument) and a unique solution (η, u) to (1.2) satisfying

$$(1.7) \quad (\eta, \mathbf{u}) \in C([0, T] : H^s(\mathbb{R}) \times H^{s+\frac{1}{2}}(\mathbb{R}^2)^2) \quad \text{and} \quad (\eta(\cdot, 0), \mathbf{u}(\cdot, 0)) = (\eta_0, \mathbf{u}_0).$$

In addition, the flow function mapping initial data to solutions is continuous.

¹More precisely, it was proved to be consistent with the KdV equation on those time scales.

Remark 1.4. The same results also hold in the periodic case. The proof is similar up to small changes in the commutator estimates (see for example [17]).

Remark 1.5. The time of existence T in Theorem 1.1 with respect to the parameter β satisfies $T(\beta) \rightarrow 0$ when $\beta \rightarrow 0$. Note that in the case of pure gravity waves ($\beta = 0$), system (1.1) is probably ill-posed² unless one makes the nonphysical assumption that $\eta \geq c_0 > 0$ as in [10]. One interesting observation is that the present situation appears to be similar to the case of the nonlinear Kelvin-Helmoltz problem for two-fluid interfaces, where the criterion established in [22] explains why capillarity is necessary for the well-posedness of the system, but does not affect the long-time dynamics.

For the sake of simplicity, we will renormalize the system and assume that $\beta = 1$ in the following.

Remark 1.6. We do not consider here the system in the long-wave regime as was done in [19], since our method of proof does not seem to provide, at least directly, good lower bounds for the existence time with respect to the small parameter ϵ measuring the size of the dispersive and nonlinear effects, which are of the same order in this regime. It remains nevertheless an interesting issue to prove that systems (1.1) and (1.2) are locally well-posed over large time as was done for some of the (a, b, c, d) -Boussinesq systems [3, 27, 28].

The proof of Theorem 1.3 is based on energy estimates and a standard compactness argument. The main difficulty lies in the lack of symmetry of the nonlinearity in (1.1). Indeed, a direct energy estimate at the $H^s \times H^{s+\frac{1}{2}}$ level³ in the one-dimensional case gives only (for s large enough)

$$\begin{aligned}
 (1.8) \quad & \frac{d}{dt} (\|\eta(t)\|_{H^s}^2 + \|u(t)\|_{H^{s+\frac{1}{2}}}^2) \\
 & \lesssim (1 + \|\eta(t)\|_{H^s} + \|u(t)\|_{H^{s+\frac{1}{2}}}) (\|\eta(t)\|_{H^s}^2 + \|u(t)\|_{H^{s+\frac{1}{2}}}^2) + \left| \int \eta J_x^s \partial_x u J_x^s \eta \right|,
 \end{aligned}$$

where J_x^s denotes the Bessel potential of order $-s$. Note that the last term on the right-hand side of (1.8) cannot be handled directly by integration by parts or commutator estimates.

In the absence of dispersion, it is well known that one can symmetrize the system by using hyperbolic symmetrizers. We refer for example to [21] for the shallow water system. This technique can be adapted in a nontrivial way when one adds a local dispersive perturbation to the system [27, 28]. However, it is not clear whether it still applies for the systems (1.1) and (1.2).⁴

Here, we use instead a modified-energy method. The idea is to add the lower-order cubic term $\int \eta (J_x^s u)^2$ to the energy. The linear contribution of the derivative of this term will cancel out the last term on the right-hand side of (1.8), while the contribution coming from the nonlinear terms can easily be controlled. This approach enables us to close the energy estimate. A similar argument can be used to derive an energy estimate for the difference of two solutions. Once these estimates

²We refer to [19] for a heuristic argument of this fact.

³The scaling $H^s \times H^{s+\frac{1}{2}}$ is needed to cancel out the linear terms.

⁴Note that the technique may work for some other systems with a nonlocal dispersion. We refer for example to [31] for a nonlocal dispersive system in the context of internal wave.

are established, the proof proceeds using bootstrapping and classical compactness arguments. Finally, it is worth noticing that the noncavitation condition on the initial data, which propagates through the flow of (1.1), is needed to ensure the coercivity of the modified energy.

The proof in the two-dimensional case is very similar. This time the energy needs to be modified by the term $\int \eta |J_x^s \mathbf{u}|^2$. Moreover, we also need to assume a curl-free condition on the initial velocity \mathbf{u}_0 . Note that this condition is preserved by the flow of (1.2). When \mathbf{u} is curl-free, the term $\frac{1}{2} \nabla |\mathbf{u}|^2$ can be written as two transport terms, namely $(\mathbf{u} \cdot \nabla u_1, \mathbf{u} \cdot \nabla u_2)^T$, where $\mathbf{u} = (u_1, u_2)^T$. This fact enables us to close the energy estimates for this term by using the Kato–Ponce commutator estimates (see for example Lemma 4.2 in [23]). Note that our proof would also work, without the curl-free assumption on \mathbf{u} , when considering a nonlinearity of the form $(\mathbf{u} \cdot \nabla) \mathbf{u}$ instead of $\nabla |\mathbf{u}|^2$ in the second line of (1.2). For the sake of simplicity, we will focus below on the proof in the one-dimensional case and will indicate in the last section what are the main changes in the two-dimensional case.

The use of a modified energy is well known to be a powerful tool in the study of nonlinear partial differential equations. We refer among others to [17, 20] (well-posedness results for higher-order KdV-type equations), [12] (long-time existence results for small initial data for the Burgers–Hilbert equation), and [24] (growth of Sobolev norm for NLS) for some applications of the modified energy methods in related contexts. The method of proof introduced here seems to be quite general, and we hope that it will have further applications to other weakly dispersive and nonlocal perturbations of nonlinear hyperbolic systems.

The paper is organized as follows: in Section 2, we give the notation and recall some commutator estimates. Sections 3 and 4 are devoted to the proof of the energy estimates respectively for a solution and for the difference of two solutions. Finally, we give the proof of Theorem 1.3(i) in Section 5 and explain the main changes for the two-dimensional case in Section 6.

2. NOTATION AND PRELIMINARY ESTIMATES

2.1. Notation.

- Throughout the text, c will denote a positive constant which may change from line to line. Also, for any positive numbers a and b , the notation $a \lesssim b$ means that $a \leq cb$.
- The operator \mathcal{F} denotes the Fourier transform. We often write $\mathcal{F}(f)(\xi) = \widehat{f}(\xi)$.
- In one dimension, \mathcal{H} will denote the Hilbert transform, i.e., $(\mathcal{H}f)^\wedge(\xi) = -i \operatorname{sgn}(\xi) \widehat{f}(\xi)$.
- In two dimensions, \mathcal{R}_j , $j = 1, 2$, will denote the Riesz transforms, i.e., $(\mathcal{R}_j f)^\wedge(\xi) = -i \frac{\xi_j}{|\xi|} \widehat{f}(\xi)$.
- For any $\alpha \in \mathbb{R}$, D_x^α will denote the Riesz potential of order $-\alpha$, defined via Fourier transform by $(D_x^\alpha f)^\wedge(\xi) = |\xi|^{-\alpha} \widehat{f}(\xi)$. In particular, it follows that $D_x^1 = \mathcal{H} \partial_x$.
- For any $\alpha \in \mathbb{R}$, J_x^α will denote the Bessel potential of order $-\alpha$, defined via Fourier transform by $(J_x^\alpha f)^\wedge(\xi) = (1 + \xi^2)^{\frac{\alpha}{2}} \widehat{f}(\xi)$. In particular, it is well known that the L^2 -based Sobolev space H^s can be defined by the norm $\|f\|_{H^s} = \|J_x^s f\|_{L^2}$.

- If A and B are two operators, then $[A, B]$ denotes the commutator between A and B , i.e., $[A, B]f = ABf - BAf$.

2.2. Fourier multiplier. We reformulate system (1.1) as

$$(2.1) \quad \begin{cases} \partial_t \eta + \mathcal{M}(D)(1 - \partial_x^2)u - \mathcal{H}u + \mathcal{H}\partial_x^2 u + \partial_x(\eta u) = 0, \\ \partial_t u + \partial_x \eta + u\partial_x u = 0, \end{cases}$$

where \mathcal{H} is the Hilbert transform and $\mathcal{M}(D)$ is the Fourier multiplier associated to the symbol

$$(2.2) \quad M(\xi) = i(\tanh(\xi) - \operatorname{sgn}(\xi)).$$

By recalling the pointwise estimate (see for example [14])

$$|\tanh(\xi) - \operatorname{sgn}(\xi)| \leq e^{-|\xi|}, \quad \forall \xi \in \mathbb{R},$$

it follows easily from the Plancherel identity that

$$(2.3) \quad \|J_x^s \mathcal{M}(D)f\|_{L^2} \lesssim \|f\|_{L^2}, \quad \forall s \in \mathbb{R}.$$

Note that the implicit constant in the former inequality depends of course on s . Moreover, we also have from Young’s theorem on convolution

$$\|\mathcal{M}(D)(1 - \partial_x^2)f\|_{L^\infty} = \|(M(\xi)(1 + \xi^2))^\vee * f\|_{L^\infty} \leq \|(M(\xi)(1 + \xi^2))^\vee\|_{L^1} \|f\|_{L^\infty},$$

so that

$$(2.4) \quad \|\mathcal{M}(D)(1 - \partial_x^2)f\|_{L^\infty} \lesssim \|f\|_{L^\infty},$$

since $\xi \mapsto M(\xi)(1 + \xi^2)$ is a Schwartz function.

Finally, we will also need an estimate comparing the Bessel and Riesz potentials. We claim that

$$(2.5) \quad \|(J_x^1 - D_x^1)\partial_x f\|_{L^2} \lesssim \|f\|_{L^2}.$$

Indeed, it follows from Plancherel’s identity that

$$\begin{aligned} \| (J_x^1 - D_x^1)\partial_x f\|_{L^2}^2 &= \int |\xi|^2 |(1 + \xi^2)^{\frac{1}{2}} - |\xi||^2 |\widehat{f}(\xi)|^2 d\xi \\ &= \int |\xi|^4 |(1 + \frac{1}{\xi^2})^{\frac{1}{2}} - 1|^2 |\widehat{f}(\xi)|^2 d\xi, \end{aligned}$$

which implies (2.5), since the function $\xi \mapsto |\xi|^4 |(1 + \frac{1}{\xi^2})^{\frac{1}{2}} - 1|^2$ is bounded on \mathbb{R} .

2.3. Commutator estimates. First, we state the Kato–Ponce commutator estimate [16].

Lemma 2.1 (Kato–Ponce commutator estimates). *Let $s \geq 1$, $p, p_2, p_3 \in (1, \infty)$ and $p_1, p_4 \in (1, \infty]$ be such that $\frac{1}{p} = \frac{1}{p_1} + \frac{1}{p_2} = \frac{1}{p_3} + \frac{1}{p_4}$. Then,*

$$(2.6) \quad \| [J_x^s, f]g\|_{L^p} \lesssim \|\partial_x f\|_{L^{p_1}} \|J_x^{s-1}g\|_{L^{p_2}} + \|J_x^s f\|_{L^{p_3}} \|g\|_{L^{p_4}},$$

for any f, g defined on \mathbb{R} .

We also state the fractional Leibniz rule proved in the appendix of [18].

Lemma 2.2. *Let $\sigma = \sigma_1 + \sigma_2 \in (0, 1)$ with $\sigma_i \in (0, \sigma)$ and $p, p_1, p_2 \in (1, \infty)$ satisfy $\frac{1}{p} = \frac{1}{p_1} + \frac{1}{p_2}$. Then,*

$$(2.7) \quad \|D_x^\sigma(fg) - fD_x^\sigma g - gD_x^\sigma f\|_{L^p} \lesssim \|D_x^{\sigma_1} f\|_{L^{p_1}} \|D_x^{\sigma_2} g\|_{L^{p_2}}.$$

Moreover, the case $\sigma_2 = 0, p_2 = \infty$ is also allowed.

The following commutator estimate was derived in Proposition 3.2 of [5].

Lemma 2.3. *Let $\alpha \in [0, 1)$, $\beta \in (0, 1)$ with $\alpha + \beta \in [0, 1]$. Then, for any $p, q \in (1, \infty)$ and for any $\delta > 1/q$, there exists $c = c(\alpha; \beta; p; q; \delta) > 0$ such that*

$$\|D_x^\alpha [D_x^\beta, a] D_x^{1-(\alpha+\beta)} f\|_{L^p} \leq c \|J_x^\delta \partial_x a\|_{L^q} \|f\|_{L^p}.$$

Corollary 2.4. *Let $s > \frac{3}{2}$. Then,*

$$(2.8) \quad \|[D_x^{\frac{1}{2}}, a] D_x^{\frac{1}{2}} f\|_{L^2} \lesssim \|a\|_{H^s} \|f\|_{L^2}.$$

Proof. The proof of estimate (2.8) follows directly by combining Lemma 2.3 with $p = 2$, $\alpha = 0$, $\beta = \frac{1}{2}$ with the Sobolev embedding by choosing q and δ such that $0 < \delta - \frac{1}{q} < s - \frac{3}{2}$. □

3. ENERGY ESTIMATES

The main goal of this section is to prove the following energy estimate for the solutions of (1.1).

Proposition 3.1. *Let $s > 2$ and $(\eta, u) \in C([0, T] : H^s(\mathbb{R}) \times H^{s+\frac{1}{2}}(\mathbb{R}))$ be a solution to (1.1) on a time interval $[0, T]$ for some $T > 0$. Let us define the modified energy $E^s(\eta, u)$ by*

$$(3.1) \quad E^s(\eta, u)(t) = \frac{1}{2} \|\eta(t)\|_{H^s}^2 + \frac{1}{2} \|u(t)\|_{H^{s+\frac{1}{2}}}^2 + \frac{1}{2} \int \eta(J_x^s u)^2(t),$$

for all $t \in [0, T]$. Assume moreover that η satisfies the condition

$$(3.2) \quad \exists \tilde{h}_0 \in (0, 1), h_1 > 0 \quad \text{such that} \quad \tilde{h}_0 - 1 \leq \eta(x, t) \leq h_1, \quad \forall (x, t) \in \mathbb{R}^2.$$

Then, the following estimates hold true for all $t \in [0, T]$.

(1) Coercivity.

$$(3.3) \quad \frac{1}{2} (\|\eta\|_{H^s}^2 + c_0 \|u\|_{H^{s+\frac{1}{2}}}^2) \leq E^s(\eta, u) \leq \frac{1}{2} (\|\eta\|_{H^s}^2 + (1 + h_1) \|u\|_{H^{s+\frac{1}{2}}}^2),$$

where $c_0 = c_0(\tilde{h}_0)$ is a positive constant.

(2) Energy estimate.

$$(3.4) \quad \frac{d}{dt} E^s(\eta, u) \lesssim E^s(\eta, u) + E^s(\eta, u)^2.$$

Proof. We observe by using condition (3.2) and Plancherel’s identity that

$$\begin{aligned} \int (J_x^{s+\frac{1}{2}} u)^2 + \int \eta(J_x^s u)^2 &\geq \tilde{h}_0 \int (J_x^s u)^2 + \int (1 + \xi^2)^s ((1 + \xi^2)^{\frac{1}{2}} - 1) |\widehat{u}(\xi)|^2 \\ &\geq \tilde{h}_0 \int (J_x^s u)^2 + \tilde{c}_0 \int_{|\xi| \geq 1} (1 + \xi^2)^{s+\frac{1}{2}} |\widehat{u}(\xi)|^2, \end{aligned}$$

where \tilde{c}_0 is a universal constant depending only on s . This implies the first inequality in (3.3) in view of the definition of (3.1) by choosing $c_0 := \min\{\tilde{h}_0, \tilde{c}_0\}$. The proof of the second inequality in (3.3) is a direct consequence of (3.1) and (3.2).

To prove estimate (3.4), we will work on the reformulated version (2.1) of (1.1). We compute the time derivative of each term on the left-hand side of (3.4) separately.

First, we get by using the Cauchy–Schwarz inequality, (2.3), and the identity $D_x^1 = \mathcal{H}\partial_x$ that

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \int (J_x^s \eta)^2 \\ &= - \int J_x^s \mathcal{M}(D)(1 - \partial_x^2) u J_x^s \eta + \int J_x^s \mathcal{H} u J_x^s \eta - \int J_x^s \mathcal{H} \partial_x^2 u J_x^s \eta - \int J_x^s \partial_x(\eta u) J_x^s \eta \\ &\leq c \|u\|_{H^s} \|\eta\|_{H^s} - \int J_x^s D_x^1 \partial_x u J_x^s \eta - \int J_x^s \partial_x(\eta u) J_x^s \eta. \end{aligned}$$

Moreover, it follows after integration by parts that

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \int (J_x^{s+\frac{1}{2}} u)^2 &= - \int J_x^{s+\frac{1}{2}} \partial_x \eta J_x^{s+\frac{1}{2}} u - \int J_x^{s+\frac{1}{2}} (u \partial_x u) J_x^{s+\frac{1}{2}} u \\ &= \int J_x^s \eta J_x^s (J_x^1 - D_x^1) \partial_x u + \int J_x^s \eta J_x^s D_x^1 \partial_x u - \int J_x^{s+\frac{1}{2}} (u \partial_x u) J_x^{s+\frac{1}{2}} u. \end{aligned}$$

Hence, we deduce by using (2.5) that

$$\begin{aligned} (3.5) \quad & \frac{1}{2} \frac{d}{dt} (\|\eta(t)\|_{H^s}^2 + \|u(t)\|_{H^{s+\frac{1}{2}}}^2) \\ & \leq c \|u\|_{H^s} \|\eta\|_{H^s} - \int J_x^s \partial_x(\eta u) J_x^s \eta - \int J_x^{s+\frac{1}{2}} (u \partial_x u) J_x^{s+\frac{1}{2}} u. \end{aligned}$$

Now, we deal with the nonlinear terms appearing on the right-hand side of (3.5). First, we observe that

$$\begin{aligned} \int J_x^s \partial_x(\eta u) J_x^s \eta &= \int J_x^s (\eta \partial_x u) J_x^s \eta + \int J_x^s (\partial_x \eta u) J_x^s \eta \\ &= \int [J_x^s, \eta] \partial_x u J_x^s \eta + \int \eta J_x^s \partial_x u J_x^s \eta + \int [J_x^s, u] \partial_x \eta J_x^s \eta + \int u J_x^s \partial_x \eta J_x^s \eta. \end{aligned}$$

On the one hand, we get by using the commutator estimate (2.6)

$$\left| \int [J_x^s, \eta] \partial_x u J_x^s \eta \right| + \left| \int [J_x^s, u] \partial_x \eta J_x^s \eta \right| \lesssim (\|\partial_x \eta\|_{L^\infty} \|u\|_{H^s} + \|\partial_x u\|_{L^\infty} \|\eta\|_{H^s}) \|\eta\|_{H^s}.$$

On the other hand, integration by parts and Hölder’s inequality yield

$$\left| \int u J_x^s \partial_x \eta J_x^s \eta \right| \lesssim \|\partial_x u\|_{L^\infty} \|\eta\|_{H^s}^2.$$

Then, we deduce by gathering the above estimates that

$$\begin{aligned} (3.6) \quad & \int J_x^s \partial_x(\eta u) J_x^s \eta \\ &= \int \eta J_x^s \partial_x u J_x^s \eta + \mathcal{O}((\|\partial_x \eta\|_{L^\infty} \|u\|_{H^s} + \|\partial_x u\|_{L^\infty} \|\eta\|_{H^s}) \|\eta\|_{H^s}). \end{aligned}$$

To deal with the second one, we get by integrating by parts that

$$\begin{aligned} \int J_x^{s+\frac{1}{2}} (u \partial_x u) J_x^{s+\frac{1}{2}} u &= \int [J_x^{s+\frac{1}{2}}, u] \partial_x u J_x^{s+\frac{1}{2}} u + \int u J_x^{s+\frac{1}{2}} \partial_x u J_x^{s+\frac{1}{2}} u \\ &= \int [J_x^{s+\frac{1}{2}}, u] \partial_x u J_x^{s+\frac{1}{2}} u - \frac{1}{2} \int \partial_x u (J_x^{s+\frac{1}{2}} u)^2. \end{aligned}$$

Then, it follows from the commutator estimate (2.6) and Hölder’s inequality that

$$(3.7) \quad \left| \int J_x^{s+\frac{1}{2}} (u \partial_x u) J_x^{s+\frac{1}{2}} u \right| \lesssim \|\partial_x u\|_{L^\infty} \|u\|_{H^{s+\frac{1}{2}}}^2.$$

Therefore, we conclude by gathering (3.5), (3.6), and (3.7) and using the Sobolev embedding that

$$(3.8) \quad \begin{aligned} & \frac{1}{2} \frac{d}{dt} (\|\eta\|_{H^s}^2 + \|u\|_{H^{s+\frac{1}{2}}}^2) \\ & \leq - \int \eta J_x^s \partial_x u J_x^s \eta + c(1 + \|\eta\|_{H^s}) \|u\|_{H^s} \|\eta\|_{H^s} + c\|u\|_{H^s} \|u\|_{H^{s+\frac{1}{2}}}^2. \end{aligned}$$

Finally, we derive the cubic contribution of the energy with respect to time. By using (2.1), we get

$$(3.9) \quad \frac{1}{2} \frac{d}{dt} \int \eta (J_x^s u)^2 = \frac{1}{2} \int \partial_t \eta (J_x^s u)^2 + \int \eta J_x^s \partial_t u J_x^s u = I_1 + I_2 + I_3,$$

where

$$I_1 := -\frac{1}{2} \int \mathcal{M}(D)(1 - \partial_x^2)u (J_x^s u)^2 + \int \mathcal{H}u (J_x^s u)^2 - \int \mathcal{H} \partial_x^2 u (J_x^s u)^2 - \int \partial_x(\eta u) (J_x^s u)^2,$$

$$I_2 := - \int \eta J_x^s \partial_x \eta J_x^s u = \int \eta J_x^s \eta J_x^s \partial_x u + \int \partial_x \eta J_x^s \eta J_x^s u,$$

after integrating by parts, and

$$I_3 := - \int \eta J_x^s (u \partial_x u) J_x^s u.$$

We have by using Hölder’s inequality, (2.4), and the Sobolev embedding that

$$(3.10) \quad \begin{aligned} |I_1| & \lesssim (\|u\|_{L^\infty} + \|\mathcal{H}u\|_{L^\infty} + \|\mathcal{H} \partial_x^2 u\|_{L^\infty} + \|\partial_x(\eta u)\|_{L^\infty}) \|u\|_{H^s}^2 \\ & \lesssim (1 + \|\eta\|_{H^s}) \|u\|_{H^s}^3 + c\|u\|_{H^{s+\frac{1}{2}}} \|u\|_{H^s}^2, \end{aligned}$$

where we used the restriction $s + \frac{1}{2} > 2 + \frac{1}{2}$, i.e., $s > 2$. Moreover, we observe that I_2 will cancel out the first term on the right-hand side of (3.8). This is why we modify the energy by the cubic term $\frac{1}{2} \int \eta (J_x^s u)^2$. We rewrite I_3 by using the commutator notation and integration by parts as

$$\begin{aligned} I_3 & = - \int \eta [J_x^s, u] \partial_x u J_x^s u - \int \eta u J_x^s \partial_x u J_x^s u \\ & = - \int \eta [J_x^s, u] \partial_x u J_x^s u + \frac{1}{2} \int \partial_x(\eta u) (J_x^s u)^2. \end{aligned}$$

Then, it follows from the Kato–Ponce commutator estimate (2.6) and the Sobolev embedding that

$$(3.11) \quad |I_3| \leq \|\eta\|_{L^\infty} \|[J_x^s, u] \partial_x u\|_{L^2} \|J_x^s u\|_{L^2} + \|\partial_x(\eta u)\|_{L^\infty} \|J_x^s u\|_{L^2}^2 \lesssim \|\eta\|_{H^s} \|u\|_{H^s}^3.$$

Hence, we deduce by gathering (3.9)–(3.11) that

$$(3.12) \quad \frac{1}{2} \frac{d}{dt} \int \eta (J_x^s u)^2 \leq \int \eta J_x^s \eta J_x^s \partial_x u + c(1 + \|\eta\|_{H^s}) \|u\|_{H^s}^3 + c\|u\|_{H^{s+\frac{1}{2}}} \|u\|_{H^s}^2.$$

Therefore, we conclude the proof of estimate (3.4) by combining (3.8) and (3.12) with (3.3). □

4. ESTIMATES FOR THE DIFFERENCES OF TWO SOLUTIONS

In this subsection, we derive energy estimates for the difference of two solutions (η_1, u_1) and (η_2, u_2) of (2.1) in $H^1(\mathbb{R}) \times H^{\frac{3}{2}}(\mathbb{R})$.

Let us define $(\tilde{\eta}, \tilde{u}) = (\eta_1 - \eta_2, u_1 - u_2)$. Then $(\tilde{\eta}, \tilde{u})$ is a solution to

$$(4.1) \quad \begin{cases} \partial_t \tilde{\eta} + \mathcal{M}(D)(1 - \partial_x^2)\tilde{u} - \mathcal{H}\tilde{u} + \mathcal{H}\partial_x^2\tilde{u} + \partial_x(\eta_1\tilde{u} + \tilde{\eta}u_2) = 0, \\ \partial_t \tilde{u} + \partial_x \tilde{\eta} + \frac{1}{2}\partial_x((u_1 + u_2)\tilde{u}) = 0, \end{cases}$$

where the symbol $M(\xi)$ of the Fourier multiplier $\mathcal{M}(D)$ is defined in (2.2).

Proposition 4.1. *Let $s > \frac{5}{2}$ and $(\eta_1, u_1), (\eta_2, u_2) \in C([0, T] : H^s(\mathbb{R}) \times H^{s+\frac{1}{2}}(\mathbb{R}))$ be two solutions to (2.1) on a time interval $[0, T]$ for some $T > 0$.*

Let $(\tilde{\eta}, \tilde{u}) = (\eta_1 - \eta_2, u_1 - u_2)$ denote the difference between the two solutions. We define the modified energy $\tilde{E}(\tilde{\eta}, \tilde{v})$ by

$$(4.2) \quad 2\tilde{E}(\tilde{\eta}, \tilde{v})(t) = \|\tilde{\eta}(t)\|_{L^2}^2 + \|\partial_x \tilde{\eta}(t)\|_{L^2}^2 + \|\tilde{u}(t)\|_{L^2}^2 + \|D_x^{\frac{1}{2}}\partial_x \tilde{u}(t)\|_{L^2}^2 + \int \eta_1(\partial_x \tilde{u})^2,$$

for all $t \in [0, T]$. Assume moreover that η_1 satisfies the condition (3.2). Then, the following estimates hold true on $[0, T]$.

(1) Coercivity. *There exists $\alpha_0 > 0$ such that*

$$(4.3) \quad \frac{1}{2}(\|\tilde{\eta}\|_{H^1}^2 + c_0\|\tilde{u}\|_{H^{\frac{3}{2}}}^2) \leq \tilde{E}(\tilde{\eta}, \tilde{v}) \leq \frac{1}{2}(\|\tilde{\eta}\|_{H^1}^2 + (1 + h_1)\|\tilde{u}\|_{H^{\frac{3}{2}}}^2),$$

where $c_0 = c_0(h_0)$ is a positive constant.

(2) Energy estimate.

$$(4.4) \quad \frac{d}{dt}\tilde{E}(\tilde{\eta}, \tilde{v}) \lesssim \left(1 + \|\eta_1\|_{H^s} + \|\eta_2\|_{H^s} + \|u_1\|_{H^s} + \|u_2\|_{H^s}\right)^2 (\|\tilde{\eta}\|_{H^1}^2 + \|\tilde{u}\|_{H^{\frac{3}{2}}}^2).$$

Proof. The proof of estimate (4.3) is similar to the one of (3.3).

To prove (4.4), we compute separately the time derivative of each term on the right-hand side of (4.2). First, it follows directly by using (4.1) and integrating by parts that

$$(4.5) \quad \begin{aligned} \frac{1}{2} \frac{d}{dt} \int \tilde{u}^2 &= - \int \tilde{u} \partial_x \tilde{\eta} - \frac{1}{2} \int \tilde{u} \partial_x ((u_1 + u_2)\tilde{u}) \\ &\lesssim \|\tilde{u}\|_{L^2} \|\partial_x \tilde{\eta}\|_{L^2} + (\|\partial_x u_1\|_{L^\infty} + \|\partial_x u_2\|_{L^\infty}) \|\tilde{u}\|_{L^2}^2. \end{aligned}$$

By using (2.3), integration by parts, and Hölder’s inequality, we get that

$$(4.6) \quad \begin{aligned} \frac{1}{2} \frac{d}{dt} \int \tilde{\eta}^2 &= - \int \tilde{\eta} \mathcal{M}(D)(1 - \partial_x^2)\tilde{u} + \int \tilde{\eta} \mathcal{H}\tilde{u} - \int \tilde{\eta} \mathcal{H}\partial_x^2\tilde{u} - \int \tilde{\eta} \partial_x(\eta_1\tilde{u} + \tilde{\eta}u_2) \\ &\lesssim \|\tilde{u}\|_{H^1} \|\tilde{\eta}\|_{H^1} + (\|\eta_1\|_{L^\infty} + \|\partial_x \eta_1\|_{L^\infty}) \|\tilde{u}\|_{H^1} \|\tilde{\eta}\|_{L^2} + \|\partial_x u_2\|_{L^\infty} \|\tilde{\eta}\|_{L^2}^2. \end{aligned}$$

Now, we turn to the higher-order part of the $H^1 \times H^{\frac{3}{2}}$ norm of $(\tilde{\eta}, \tilde{u})$. On the one hand, we have that

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \int (\partial_x \tilde{\eta})^2 &= - \int \partial_x \tilde{\eta} \partial_x \mathcal{M}(D)(1 - \partial_x^2)\tilde{u} + \int \partial_x \tilde{\eta} \partial_x \mathcal{H}\tilde{u} - \int \partial_x \tilde{\eta} \mathcal{H}\partial_x^3\tilde{u} \\ &\quad - \int \partial_x \tilde{\eta} \partial_x^2(\eta_1\tilde{u} + \tilde{\eta}u_2). \end{aligned}$$

To deal with the nonlinear term, we integrate by parts and use Hölder’s inequality. It follows that

$$\int \partial_x \tilde{\eta} \partial_x^2 (\eta_1 \tilde{u} + \tilde{\eta} u_2) \leq \int \eta_1 \partial_x \tilde{\eta} \partial_x^2 \tilde{u} + c(\|\partial_x \eta_1\|_{L^\infty} + \|\partial_x^2 \eta_1\|_{L^\infty}) \|\tilde{u}\|_{H^1} \|\tilde{\eta}\|_{H^1} + c(\|\partial_x u_2\|_{L^\infty} + \|\partial_x^2 u_2\|_{L^\infty}) \|\tilde{\eta}\|_{H^1}^2,$$

which implies together with (2.3) that

$$(4.7) \quad \begin{aligned} \frac{1}{2} \frac{d}{dt} \int (\partial_x \tilde{\eta})^2 &\leq c\|\tilde{u}\|_{H^1} \|\tilde{\eta}\|_{H^1} - \int \partial_x \tilde{\eta} \mathcal{H} \partial_x^3 \tilde{u} - \int \eta_1 \partial_x \tilde{\eta} \partial_x^2 \tilde{u} \\ &+ c(\|\partial_x \eta_1\|_{L^\infty} + \|\partial_x^2 \eta_1\|_{L^\infty}) \|\tilde{u}\|_{H^1} \|\tilde{\eta}\|_{H^1} \\ &+ c(\|\partial_x u_2\|_{L^\infty} + \|\partial_x^2 u_2\|_{L^\infty}) \|\tilde{\eta}\|_{H^1}^2. \end{aligned}$$

On the other hand, we compute

$$\frac{1}{2} \frac{d}{dt} \int (D_x^{\frac{1}{2}} \partial_x \tilde{u})^2 = - \int D_x^{\frac{1}{2}} \partial_x \tilde{u} D_x^{\frac{1}{2}} \partial_x^2 \tilde{\eta} - \int D_x^{\frac{1}{2}} \partial_x \tilde{u} D_x^{\frac{1}{2}} \partial_x^2 ((u_1 + u_2) \tilde{u}).$$

By using the identity $D_x^1 = \mathcal{H} \partial_x$ and integration by parts, we have

$$- \int D_x^{\frac{1}{2}} \partial_x \tilde{u} D_x^{\frac{1}{2}} \partial_x^2 \tilde{\eta} = \int \mathcal{H} \partial_x^3 \tilde{u} \partial_x \tilde{\eta},$$

so that this term will cancel out the second one on the right-hand side of (4.7). Now, we deal with the nonlinear term. It follows by using the standard Leibniz rule that

$$\begin{aligned} &\int D_x^{\frac{1}{2}} \partial_x \tilde{u} D_x^{\frac{1}{2}} \partial_x^2 ((u_1 + u_2) \tilde{u}) \\ &= \int D_x^{\frac{1}{2}} \partial_x \tilde{u} D_x^{\frac{1}{2}} (\partial_x^2 (u_1 + u_2) \tilde{u} + 2\partial_x^2 (u_1 + u_2) \partial_x \tilde{u} + (u_1 + u_2) \partial_x^2 \tilde{u}) \\ &=: \mathcal{I}_1 + \mathcal{I}_2 + \mathcal{I}_3. \end{aligned}$$

We deduce from the fractional Leibniz rule (2.7) that

$$\begin{aligned} |\mathcal{I}_1| &\lesssim \|D_x^{\frac{1}{2}} \partial_x \tilde{u}\|_{L^2} (\|\partial_x^2 u_1\|_{L^\infty} + \|\partial_x^2 u_2\|_{L^\infty}) \|D_x^{\frac{1}{2}} \tilde{u}\|_{L^2} \\ &+ \|D_x^{\frac{1}{2}} \partial_x \tilde{u}\|_{L^2} (\|\partial_x^2 D_x^{\frac{1}{2}} u_1\|_{L^2} + \|\partial_x^2 D_x^{\frac{1}{2}} u_2\|_{L^2}) \|\tilde{u}\|_{L^\infty} \end{aligned}$$

and

$$\begin{aligned} |\mathcal{I}_2| &\lesssim \|D_x^{\frac{1}{2}} \partial_x \tilde{u}\|_{L^2} (\|\partial_x u_1\|_{L^\infty} + \|\partial_x u_2\|_{L^\infty}) \|D_x^{\frac{1}{2}} \partial_x \tilde{u}\|_{L^2} \\ &+ \|D_x^{\frac{1}{2}} \partial_x \tilde{u}\|_{L^2} (\|\partial_x D_x^{\frac{1}{2}} u_1\|_{L^4} + \|\partial_x D_x^{\frac{1}{2}} u_2\|_{L^4}) \|\partial_x \tilde{u}\|_{L^4}. \end{aligned}$$

Moreover, by using $\partial_x = -\mathcal{H} D_x^1$, the commutator notation, and integration by parts, we get

$$\begin{aligned} \mathcal{I}_3 &= \int D_x^{\frac{1}{2}} \partial_x \tilde{u} [D_x^{\frac{1}{2}}, u_1 + u_2] \partial_x^2 \tilde{u} + \int D_x^{\frac{1}{2}} \partial_x \tilde{u} (u_1 + u_2) D_x^{\frac{1}{2}} \partial_x^2 \tilde{u} \\ &= - \int D_x^{\frac{1}{2}} \partial_x \tilde{u} [D_x^{\frac{1}{2}}, u_1 + u_2] D_x^{\frac{1}{2}} \mathcal{H} D_x^{\frac{1}{2}} \partial_x \tilde{u} - \frac{1}{2} \int \partial_x (u_1 + u_2) (D_x^{\frac{1}{2}} \partial_x \tilde{u})^2. \end{aligned}$$

Hence, the commutator estimate (2.8) and the Sobolev embedding yield

$$|\mathcal{I}_3| \lesssim (\|u_1\|_{H^s} + \|u_2\|_{H^s}) \|D_x^{\frac{1}{2}} \partial_x \tilde{u}\|_{L^2}^2.$$

Therefore, we deduce by gathering those estimates that

$$(4.8) \quad \frac{1}{2} \frac{d}{dt} \int (D_x^{\frac{1}{2}} \partial_x \tilde{u})^2 \leq \int \mathcal{H} \partial_x^3 \tilde{u} \partial_x \tilde{\eta} + c(\|u_1\|_{H^s} + \|u_2\|_{H^s}) \|\tilde{u}\|_{H^{\frac{3}{2}}}^2.$$

Finally, to deal with the third term on the right-hand side of (4.8), we need to use the cubic part in the modified energy. Observe by using (2.1) and (4.1) that

$$\frac{1}{2} \frac{d}{dt} \int \eta_1 (\partial_x \tilde{u})^2 = \frac{1}{2} \int \partial_t \eta_1 (\partial_x \tilde{u})^2 + \int \eta_1 \partial_x \partial_t \tilde{u} \partial_x \tilde{u} = \mathcal{J}_1 + \mathcal{J}_2 + \mathcal{J}_3,$$

where

$$\begin{aligned} \mathcal{J}_1 &= -\frac{1}{2} \int \mathcal{M}(D)(1 - \partial_x^2) u_1 (\partial_x \tilde{u})^2 + \int \mathcal{H} u_1 (\partial_x \tilde{u})^2 - \int \mathcal{H} \partial_x^2 u_1 (\partial_x \tilde{u})^2 \\ &\quad - \int \partial_x (\eta_1 u_1) (\partial_x \tilde{u})^2, \end{aligned}$$

$$\mathcal{J}_2 = - \int \eta_1 \partial_x^2 \tilde{\eta} \partial_x \tilde{u} = \int \eta_1 \partial_x \tilde{\eta} \partial_x^2 \tilde{u} + \int \partial_x \eta_1 \partial_x \tilde{\eta} \partial_x \tilde{u},$$

after integrating by parts, and

$$\mathcal{J}_3 = - \int \eta_1 \partial_x^2 ((u_1 + u_2) \tilde{u}) \partial_x \tilde{u}.$$

We have by using Hölder’s inequality, (2.4), and the Sobolev embedding that

$$\begin{aligned} |\mathcal{J}_1| &\lesssim (\|u_1\|_{L^\infty} + \|\mathcal{H} u_1\|_{L^\infty} + \|\mathcal{H} \partial_x^2 u_1\|_{L^\infty} + \|\partial_x (\eta_1 u_1)\|_{L^\infty}) \|\partial_x \tilde{u}\|_{L^2}^2 \\ &\lesssim (\|u_1\|_{H^s} + \|\eta_1\|_{H^s} \|u_1\|_{H^s}) \|\partial_x \tilde{u}\|_{L^2}^2, \end{aligned}$$

where we used the restriction $s > \frac{5}{2}$. Moreover, we observe that the first term on the right-hand side of \mathcal{J}_2 will cancel out the third term on the right-hand side of (4.7). To handle \mathcal{J}_3 , we use the standard Leibniz rule and integration by parts to get

$$\mathcal{J}_3 = - \int \eta_1 \partial_x^2 (u_1 + u_2) \tilde{u} \partial_x \tilde{u} + \int \left(-\frac{3}{2} \eta_1 \partial_x (u_1 + u_2) + \frac{1}{2} \partial_x \eta_1 (u_1 + u_2) \right) (\partial_x \tilde{u})^2.$$

It follows from Hölder and Sobolev inequalities that

$$|\mathcal{J}_3| \lesssim \|\eta_1\|_{H^s} (\|u_1\|_{H^s} + \|u_2\|_{H^s}) \|\tilde{u}\|_{H^1}^2.$$

Hence, we deduce by gathering those estimates that

$$(4.9) \quad \frac{1}{2} \frac{d}{dt} \int \eta_1 (\partial_x \tilde{u})^2 \leq \int \eta_1 \partial_x \tilde{\eta} \partial_x^2 \tilde{u} + c(1 + \|\eta_1\|_{H^s}) (\|u_1\|_{H^s} + \|u_2\|_{H^s}) \|\tilde{u}\|_{H^1}^2.$$

Therefore, we conclude the proof of (4.4) by gathering (4.5)–(4.9). □

Remark 4.2. Observe that the restriction $s > \frac{5}{2}$ in Theorem 1.3(i) appears in Proposition 4.1.

5. PROOF OF THEOREM 1.3

We begin this section by proving an a priori estimate on the solutions (η, u) to (1.1).

Lemma 5.1. *Let $s > \frac{5}{2}$. Let $(\eta, u) \in C([0, T^*) : H^s(\mathbb{R}) \times H^{s+\frac{1}{2}}(\mathbb{R}))$ be a solution to (1.1) corresponding to initial data $(\eta_0, u_0) \in H^s(\mathbb{R}) \times H^{s+\frac{1}{2}}(\mathbb{R})$ satisfying the non-cavitation condition (1.5), defined on its maximal time of existence and satisfying the blow-up alternative*

$$(5.1) \quad \text{If } T^* < \infty, \text{ then } \lim_{t \nearrow T^*} \|(\eta(t), u(t))\|_{H^s \times H^{s+\frac{1}{2}}} = +\infty.$$

Then, there exists $T_0 = T_0(\|(\eta_0, u_0)\|_{H^s \times H^{s+\frac{1}{2}}})$ such that $T^* > T_0$ and

$$(5.2) \quad \sup_{t \in [0, T_0]} \|(\eta, u)(t)\|_{H^s \times H^{s+\frac{1}{2}}} \leq c \|(\eta_0, u_0)\|_{H^s \times H^{s+\frac{1}{2}}},$$

for some positive constant c .

Proof. Let us define

$$\tilde{T} := \sup \left\{ T \in (0, T^*) : \sup_{t \in [0, T]} \|(\eta, u)(t)\|_{H^s \times H^{s+1/2}}^2 \leq 8 \|(\eta_0, u_0)\|_{H^s \times H^{s+1/2}}^2 \right\}.$$

Note that $\tilde{T} < T^*$; otherwise it would contradict the blow-up alternative (5.1).

We define $T_0 := \min\{T_1, T_2\}$ where

$$T_1 = \frac{1}{C_1} \log \left(1 + \frac{1}{1 + C_1 \|(\eta_0, u_0)\|_{H^s \times H^{s+1/2}}^2} \right), \quad T_2 = \frac{h_0}{C_2 (1 + \|\eta_0\|_{H^s}) \|u_0\|_{H^{s+1/2}}}$$

and C_1, C_2 are two large positive constants to be fixed in the proof.

Assume by contradiction that $\tilde{T} < T_0$; otherwise we are done. By continuity, we have that

$$(5.3) \quad \sup_{t \in [0, \tilde{T}]} \|(\eta, u)(t)\|_{H^s \times H^{s+1/2}}^2 \leq 8 \|(\eta_0, u_0)\|_{H^s \times H^{s+1/2}}^2.$$

We first verify that the noncavitation condition (3.2) holds on $[0, \tilde{T}]$. On the one hand, since η_0 satisfies (1.5), it follows from the fundamental theorem of calculus that

$$\eta(x, t) + 1 = \eta_0(x) + 1 + \int_0^t \partial_t \eta(x, s) ds \geq h_0 - \tilde{T} \sup_{s \in [0, \tilde{T}]} \|\partial_t \eta(s)\|_{L_x^\infty},$$

for all $t \in [0, \tilde{T}]$. On the other hand, we estimate trivially by using the first equation in (2.1), the Sobolev embedding, and then (5.3) that

$$\|\partial_t \eta\|_{L_x^\infty} \leq c(1 + \|\eta\|_{H^s}) \|u\|_{H^s} \leq c(1 + \|\eta_0\|_{H^s}) \|u_0\|_{H^{s+1/2}},$$

$\forall t \in [0, T_1]$. Thus, by recalling that $\tilde{T} < T_2$ and by choosing C_2 large enough, we deduce from the above analysis that $\eta(x, t) + 1 \geq h_0/2$ on $[0, \tilde{T}]$. This estimate and a similar argument combined with the Sobolev embedding $\eta_0 \in H^s(\mathbb{R}) \hookrightarrow L^\infty(\mathbb{R})$ show that the condition (3.2) holds on the time interval $[0, \tilde{T}]$.

Let $y(t) := E(\eta, u)(t)$ denote the modified energy defined in (3.1). Then (3.4) leads to the inequality $y'(t) \leq c(y(t) + y^2(t))$, which can be integrated to obtain

$$y(t) \left(1 - \frac{y_0}{1 + y_0} e^{ct} \right) \leq \frac{y_0}{1 + y_0} e^{ct} \quad \text{if} \quad \frac{y_0}{1 + y_0} e^{ct} < 1$$

on $[0, \tilde{T}]$. Since $\tilde{T} < T_1$, we get by choosing $C_1 = C_1(h_0)$ large enough and by using (3.3) that

$$\frac{y_0}{1 + y_0} e^{ct} \leq \frac{1 + 2y_0}{2(1 + y_0)} < 1 \quad \text{and} \quad \|(\eta, u)(t)\|_{H^s \times H^{s+1/2}}^2 \leq 4\|(\eta_0, u_0)\|_{H^s \times H^{s+1/2}}^2,$$

for all $t \in [0, \tilde{T}]$. Thus, we deduce by continuity that there exists some $\tilde{T} < T < T^*$ such that $\|(\eta, u)(T)\|_{H^s \times H^{s+1/2}}^2 \leq 6\|(\eta_0, u_0)\|_{H^s \times H^{s+1/2}}^2$. This contradicts the definition of \tilde{T} . Therefore, $\tilde{T} < T_0$, which concludes the proof of Lemma 5.1. \square

With the a priori estimate in hand, the complete proof of the existence would then result from a standard compactness argument implemented on a regularized version of the system. The uniqueness is a consequence of the estimates for the difference of two solutions (4.4). The strong continuity in time and the continuity of the flow would result from an application of the Bona-Smith argument [2] (we refer to [17] for a detailed demonstration of the use of the Bona-Smith argument in the context of the modified energy).

6. THE TWO-DIMENSIONAL CASE

In this section, we comment briefly on the changes to adapt the proof in the two-dimensional setting. By denoting $\mathbf{u} = (u_1, u_2)^T$, we reformulate the system (1.2) as

$$(6.1) \quad \begin{cases} \partial_t \eta - (\widetilde{\mathcal{M}}(D) + 1)(1 - \Delta)(\mathcal{R}_1 u_1 + \mathcal{R}_2 u_2) + \partial_{x_1}(\eta u_1) + \partial_{x_2}(\eta u_2) = 0, \\ \partial_t u_1 + \frac{1}{2} \partial_{x_1}(u_1^2 + u_2^2) = 0, \\ \partial_t u_2 + \frac{1}{2} \partial_{x_2}(u_1^2 + u_2^2) = 0, \end{cases}$$

where \mathcal{R}_j denotes the Riesz transform and $\widetilde{\mathcal{M}}(D)$ is the Fourier multiplier associated to the symbol $\widetilde{\mathcal{M}}(\xi) = \tanh |\xi| - 1$, $\xi = (\xi_1, \xi_2)$, and $|\xi| = \sqrt{\xi_1^2 + \xi_2^2}$. Note that the pointwise estimate $|\tanh |\xi| - 1| \leq e^{-|\xi|}, \forall \xi \in \mathbb{R}^2$, holds true.

We derive an energy estimate (analogous to Proposition 3.1) at the level $(\eta, u_1, u_2) \in H^s(\mathbb{R}^2) \times H^{s+\frac{1}{2}}(\mathbb{R}^2) \times H^{s+\frac{1}{2}}(\mathbb{R}^2)$, $s > \frac{5}{2}$, by using the modified energy

$$E^s(\eta, u_1, u_2)(t) = \frac{1}{2} \|\eta(t)\|_{H^s}^2 + \frac{1}{2} \|u_1(t)\|_{H^{s+\frac{1}{2}}}^2 + \frac{1}{2} \|u_2(t)\|_{H^{s+\frac{1}{2}}}^2 + \frac{1}{2} \int \eta((J_x^s u_1)^2 + (J_x^s u_2)^2)(t).$$

Note that instead of using the identity $D_x^1 = \mathcal{H} \partial_x$, we use in a crucial way the identities $\mathcal{R}_j \Delta = D_x^1 \partial_{x_j}$, $j = 1, 2$, to cancel out the linear terms. The modified energy is then used to handle the nonlinear term in the first equation of (6.1). To deal with the nonlinear terms in the second and third equations of (6.1), we use Lemma 4.2. in [23].⁵

The proof of the uniqueness is very similar to Proposition 4.1.

ACKNOWLEDGMENTS

The authors would like to thank Jean-Claude Saut, Vincent Duchêne, and Mats Ehrnström for helpful comments on a preliminary version of this work. They also thank the anonymous referee for helpful comments and suggestions.

⁵As already observed in the introduction, it is for this reason that we need to make the curl-free assumption on the velocity \mathbf{u} .

REFERENCES

- [1] P. Aceves-Sánchez, A. A. Minzoni, and P. Panayotaros, *Numerical study of a nonlocal model for water-waves with variable depth*, Wave Motion **50** (2013), no. 1, 80–93, DOI 10.1016/j.wavemoti.2012.07.002. MR2991247
- [2] J. L. Bona and R. Smith, *The initial-value problem for the Korteweg-de Vries equation*, Philos. Trans. Roy. Soc. London Ser. A **278** (1975), no. 1287, 555–601, DOI 10.1098/rsta.1975.0035. MR0385355
- [3] Cosmin Burtea, *New long time existence results for a class of Boussinesq-type systems* (English, with English and French summaries), J. Math. Pures Appl. (9) **106** (2016), no. 2, 203–236, DOI 10.1016/j.matpur.2016.02.008. MR3515301
- [4] John D. Carter, *Bidirectional Whitham equations as models of waves on shallow water*, Wave Motion **82** (2018), 51–61, DOI 10.1016/j.wavemoti.2018.07.004. MR3844340
- [5] L. Dawson, H. McGahagan, and G. Ponce, *On the decay properties of solutions to a class of Schrödinger equations*, Proc. Amer. Math. Soc. **136** (2008), no. 6, 2081–2090, DOI 10.1090/S0002-9939-08-09355-6. MR2383514
- [6] Evgueni Dinvay, *On well-posedness of a dispersive system of the Whitham-Boussinesq type*, Appl. Math. Lett. **88** (2019), 13–20, DOI 10.1016/j.aml.2018.08.005. MR3862707
- [7] Evgueni Dinvay, Daulet Moldabayev, Denys Dutykh, and Henrik Kalisch, *The Whitham equation with surface tension*, Nonlinear Dynam. **88** (2017), no. 2, 1125–1138, DOI 10.1007/s11071-016-3299-7. MR3628376
- [8] V. Duchêne, S. Israwi, and R. Talhouk, *A new class of two-layer Green-Naghdi systems with improved frequency dispersion*, Stud. Appl. Math. **137** (2016), no. 3, 356–415, DOI 10.1111/sapm.12125. MR3564304
- [9] Mats Ehrnström, Mark D. Groves, and Erik Wahlén, *On the existence and stability of solitary-wave solutions to a class of evolution equations of Whitham type*, Nonlinearity **25** (2012), no. 10, 2903–2936, DOI 10.1088/0951-7715/25/10/2903. MR2979975
- [10] M. Ehrnström, L. Pei, and Y. Wang, *A conditional well-posedness result for the bidirectional Whitham equation*, preprint, arXiv:1708.0455 (2017).
- [11] M. Ehrnström and E. Wahlén, *On Whitham’s conjecture of a highest cusped wave for a nonlocal shallow water wave equation*, to appear in Ann. Inst. H. Poincaré, Anal. Nonlin., preprint, arXiv:1602.05384 (2016).
- [12] John K. Hunter, Mihaela Ifrim, Daniel Tataru, and Tak Kwong Wong, *Long time solutions for a Burgers–Hilbert equation via a modified energy method*, Proc. Amer. Math. Soc. **143** (2015), no. 8, 3407–3412, DOI 10.1090/proc/12215. MR3348783
- [13] Vera Mikyoung Hur, *Wave breaking in the Whitham equation*, Adv. Math. **317** (2017), 410–437, DOI 10.1016/j.aim.2017.07.006. MR3682673
- [14] V. M. Hur and A. K. Pandey, *Modulational instability in a full-dispersion shallow water model*, preprint, arXiv:1608.04685 (2016).
- [15] Vera Mikyoung Hur and Mathew A. Johnson, *Modulational instability in the Whitham equation with surface tension and vorticity*, Nonlinear Anal. **129** (2015), 104–118, DOI 10.1016/j.na.2015.08.019. MR3414922
- [16] Tosio Kato and Gustavo Ponce, *Commutator estimates and the Euler and Navier–Stokes equations*, Comm. Pure Appl. Math. **41** (1988), no. 7, 891–907, DOI 10.1002/cpa.3160410704. MR951744
- [17] Carlos E. Kenig and Didier Pilod, *Local well-posedness for the KdV hierarchy at high regularity*, Adv. Differential Equations **21** (2016), no. 9–10, 801–836. MR3513119
- [18] Carlos E. Kenig, Gustavo Ponce, and Luis Vega, *Well-posedness and scattering results for the generalized Korteweg-de Vries equation via the contraction principle*, Comm. Pure Appl. Math. **46** (1993), no. 4, 527–620, DOI 10.1002/cpa.3160460405. MR1211741
- [19] Christian Klein, Felipe Linares, Didier Pilod, and Jean-Claude Saut, *On Whitham and related equations*, Stud. Appl. Math. **140** (2018), no. 2, 133–177, DOI 10.1111/sapm.12194. MR3763731
- [20] Soonsik Kwon, *On the fifth-order KdV equation: local well-posedness and lack of uniform continuity of the solution map*, J. Differential Equations **245** (2008), no. 9, 2627–2659, DOI 10.1016/j.jde.2008.03.020. MR2455780

- [21] David Lannes, *The water waves problem: Mathematical analysis and asymptotics*, Mathematical Surveys and Monographs, vol. 188, American Mathematical Society, Providence, RI, 2013. MR3060183
- [22] David Lannes, *A stability criterion for two-fluid interfaces and applications*, Arch. Ration. Mech. Anal. **208** (2013), no. 2, 481–567, DOI 10.1007/s00205-012-0604-6. MR3035985
- [23] Felipe Linares, Didier Pilod, and Jean-Claude Saut, *Well-posedness of strongly dispersive two-dimensional surface wave Boussinesq systems*, SIAM J. Math. Anal. **44** (2012), no. 6, 4195–4221, DOI 10.1137/110828277. MR3023445
- [24] Fabrice Planchon, Nikolay Tzvetkov, and Nicola Visciglia, *On the growth of Sobolev norms for NLS on 2- and 3-dimensional manifolds*, Anal. PDE **10** (2017), no. 5, 1123–1147, DOI 10.2140/apde.2017.10.1123. MR3668586
- [25] Filippo Remonato and Henrik Kalisch, *Numerical bifurcation for the capillary Whitham equation*, Phys. D **343** (2017), 51–62, DOI 10.1016/j.physd.2016.11.003. MR3606200
- [26] Nathan Sanford, Keri Kodama, John D. Carter, and Henrik Kalisch, *Stability of traveling wave solutions to the Whitham equation*, Phys. Lett. A **378** (2014), no. 30-31, 2100–2107, DOI 10.1016/j.physleta.2014.04.067. MR3226084
- [27] Jean-Claude Saut and Li Xu, *The Cauchy problem on large time for surface waves Boussinesq systems*, J. Math. Pures Appl. (9) **97** (2012), no. 6, 635–662, DOI 10.1016/j.matpur.2011.09.012. MR2921604
- [28] Jean-Claude Saut, Chao Wang, and Li Xu, *The Cauchy problem on large time for surface-waves-type Boussinesq systems II*, SIAM J. Math. Anal. **49** (2017), no. 4, 2321–2386, DOI 10.1137/15M1050203. MR3668593
- [29] S. Trillo, M. Klein, G. F. Clauss, and M. Onorato, *Observation of dispersive shock waves developing from initial depressions in shallow water*, Phys. D **333** (2016), 276–284, DOI 10.1016/j.physd.2016.01.007. MR3523508
- [30] Whitham, G. B. *Variational Methods and Applications to Water Waves*, Proceedings of the Royal Society of London. Series A, Mathematical and Physical Sciences **299**, no. 1456 (1967): 6-25. <http://www.jstor.org/stable/2415780>.
- [31] Li Xu, *Intermediate long wave systems for internal waves*, Nonlinearity **25** (2012), no. 3, 597–640, DOI 10.1088/0951-7715/25/3/597. MR2887985

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BERGEN, POSTBOX 7800, 5020 BERGEN, NORWAY

Email address: Henrik.Kalisch@uib.no

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BERGEN, POSTBOX 7800, 5020 BERGEN, NORWAY

Email address: Didier.Pilod@uib.no