

# A FREE BOUNDARY VALUE PROBLEM FOR THE HEAT EQUATION\*

BY

W. L. MIRANKER

*Bell Telephone Laboratories, Inc.*

**1. Introduction.** In this paper we will prove the existence of a solution of a free boundary value problem for the heat equation. We will accomplish this by demonstrating the existence of a solution to a non-linear integro-differential equation.

Let  $D$  be the domain  $0 \leq t, 0 \leq x \leq R(t)$ ,  $R(0) = A$ , indicated in Fig. 1. The boundary value problem is




FIG. 1.

$$\begin{aligned}
 u_t &= \alpha^2 u_{xx}(x, t) \text{ in } D, \\
 u_x(0, t) &= -G(t) \quad G(t) > 0, \\
 u[R(t), t] &= T_c = \text{constant}, \\
 u(x, 0) &= F(x) \geq T_c \quad 0 \leq x \leq A, \\
 u_x[R(t), t] &= B - CR_t(t) \quad B \geq 0, C > 0 \text{ are constants.}
 \end{aligned} \tag{1}$$

Differentiation is denoted by a subscript whether it is a partial derivative of a function of two variables or an ordinary derivative of a function of one variable.

This problem with  $A = 0$  has been discussed by several authors, see for example [1, 2, 6], but thus far no existence proof has been established\*\*. The problem describes the physical phenomena of evaporation, fusion, sublimation, etc. For example with  $B = 0$ , (1) could refer to the following situation. A long metal rod insulated at the sides has begun to melt at one end ( $x = 0$ ). The layer of liquid metal is  $A$  units deep and has some initial temperature distribution,  $F(x)$ . The critical temperature  $T_c$  is the melting point of the metal. At  $x = 0$  heat is applied to the rod at a known rate proportional

\*Received March 26, 1957.

\*\*For our proof it is essential that  $A \neq 0$ . See, however, the remark at the end of the existence proof below.

to  $G(t)$ . As the process continues the interface,  $R(t)$ , between liquid and solid advances down the rod.

This problem is called a free boundary problem since the part  $R(t)$  of the boundary of  $D$  is unknown. The additional boundary condition (5) which would over-determine the problem were  $R(t)$  known compensates for the free boundary.

We set  $B = 0$  to achieve clarity in our discussion, and we introduce the transformation

$$\begin{aligned} t' &= (A^2/\alpha^2)t, \\ x' &= (A/\alpha^2)t, \\ a' &= a - T_c, \\ R' &= (A^2/\alpha^4)R, \\ g &= G, \\ f &= F, \\ D' &= D. \end{aligned} \tag{2}$$

Upon dropping the primes the free boundary value problem then becomes

$$u_t = u_{xx}, \quad (x, t) \in D, \tag{3}$$

$$u_x(0, t) = -g(t) < 0, \tag{4}$$

$$u[R(t), t] = 0, \tag{5}$$

$$u(x, 0) = f(x) \geq 0, \tag{6}$$

$$u_x[R(t), t] = -R_t(t). \tag{7}$$

We require that  $g(t)$  and  $f(x)$  be continuous and have the following additional properties:  $g(t)$  is differentiable;  $f(x)$  is continuously differentiable for  $0 < x < A$ ;  $f(x) = f(-x)$ ;  $f(x) = 0$  for  $x > A$ ;  $f_x(A) < 0$ ; and  $f(0) = -g(0)$ . An  $f(x)$  with these properties is

$$f(x) = \frac{g(0)}{A} x^2 - g(0) |x| + g(0)A - A^2, \quad |x| < A,$$

$f(x) = 0$ , otherwise.

**2. Method of solution.** Our method of solution will be to apply the method of I. Kolodner [4] and derive a functional equation for  $R(t)$ . We will show that the existence of a solution with certain properties of the functional equation implies the existence of a solution to the free-boundary problem. We will solve the functional equation by the method of contracting maps (Picard iterations).

The method of Kolodner: Let  $x = \rho(t)$  be a continuously differentiable function and such that  $\rho(0) = A$ . Consider the function  $u^\rho(x, t)$  defined as

$$u^\rho = v^\rho + w^\rho + f^\rho, \tag{2.1}$$

where

$$v^\rho = -\frac{1}{2}\pi^{-1/2} \int_0^t (t-\tau)^{-1/2} \rho_\tau(\tau) \exp \{-[\frac{1}{2}[x-\rho(\tau)](t-\tau)^{-1/2}]^2\} d\tau, \tag{2.2}$$

$$w^{\rho} = -\frac{1}{2}\pi^{-1/2} \int_0^t (t-\tau)^{-1/2} \rho_{\tau}(\tau) \exp \left\{ -\left[\frac{1}{2}(x+\rho(\tau))(t-\tau)^{-1/2}\right]^2 \right\} d\tau \quad (2.3)$$

$$+ \pi^{-1/2} \int_0^t (t-\tau)^{-1/2} g(\tau) \exp \left\{ -\left[\frac{1}{2}x(t-\tau)^{-1/2}\right]^2 \right\} d\tau,$$

and

$$f^{\rho} = \frac{1}{2}(\pi t)^{-1/2} \int_0^t f(\xi) \exp \left\{ -\left[\frac{1}{2}(x-\xi)t^{-1/2}\right]^2 \right\} d\tau. \quad (2.4)$$

This function is a solution of the heat equation in the domain of Fig. 1 with  $\rho(t)$  replacing  $R(t)$ . We will now calculate the same boundary and initial conditions of  $u^{\rho}(x, t)$  from 2.1 which are prescribed for  $u(x, t)$  in the statement of the problem (3)–(7). That is by computing  $u_x^{\rho}$  from 2.1 and then letting  $(x, t)$  approach the boundaries of the domain of Fig. 1 we shall obtain the boundary conditions in question for  $u^{\rho}$  and  $u_x^{\rho}$ .

The arguments used in doing this are lengthy and are based upon well-known formulas for the integral solutions of the heat equation (see [3]).

The results are

$$u_x^{\rho}[\rho(t) + 0, t] - u_x^{\rho}[\rho(t) - 0, t] = \rho_t(t), \quad (2.5)$$

$$u_x^{\rho}(0+, t) = -g(t), \quad (2.6)$$

$$u^{\rho}(x, 0+) = f^{\rho}(x, 0+) = f(x). \quad (2.7)$$

(2.6) makes use of the evenness of  $f(x)$ .

(2.6) and (2.7) show that  $u^{\rho}(x, t)$  satisfies the boundary and initial condition (4) and (6) which are required of  $u(x, t)$ . (2.5) almost does the same for (7). If in (2.5) we require that

$$u_x^{\rho}[\rho(t) + 0, t] = 0 \quad (2.8)$$

we will have

$$u_x^{\rho}[\rho(t) - 0, t] = -\rho_t(t) \quad (2.9)$$

which is the condition (7) for  $u$ . We will see that (2.8) is an integro-differential equation for  $\rho(t)$ . To show that a solution to this integro-differential equation furnishes a solution to the free boundary problem, we need only apply Green's formula,

$$2 \iint_D (u_x)^2 dx dt = \oint u^2 dx + 2uu_x dt \quad (2.10)$$

to the domain  $D_{\alpha}$  of Fig. 2 and to the function  $u^{\rho}$ .




FIG. 2.

Using (2.7) and the fact that  $f(x) = 0$  for  $x > A$  and also using (2.8) we have

$$2 \iint_{D_\alpha} (u_x^2)^2 dx dt = 2 \int_0^{\alpha} u^2(\alpha, t) u_x^2(\alpha, t) dt - \int_{\rho(t)}^{\alpha} (u^2)^2(x, t_1) dx - \int_0^{\alpha} (u^2)^2[\rho(t), t] \rho_t dt. \quad (2.11)$$

If we now let  $\alpha \rightarrow \infty$  and observe that (2.1)–(2.4) imply that  $u^2(\alpha, t)$  and  $u_x^2(\alpha, t) \rightarrow 0$ , then (2.11) becomes

$$2 \iint_{D_\infty} (u_x^2)^2 dx dt + \int_{\rho(t)}^{\infty} (u^2)^2(x, t_1) dx = - \int_0^t (u^2)^2[\rho(t), t] \rho_t(t) dt. \quad (2.12)$$

If we require that

$$\rho_t \geq 0 \quad (2.13)$$

then the right-hand side of (2.12) is not positive and (2.12) shows that  $u^2 = 0$  for  $x > \rho(t)$  and all  $t$ . This in turn implies that  $u^2(\rho + 0, t) = 0$ , and since  $u^2$  is continuous along  $x = \rho(t)$  that  $u^2(\rho, t) = 0$ .

Thus the requirement (2.13) makes  $\mu^2$  satisfy the condition (5) for  $u$ . We see that if  $\rho(t)$  is a smooth function for which  $\rho(0) = A$  and  $\rho_t \geq 0$  and such that  $u_x^2(\rho(t) + 0, t) = 0$ , then the function  $u^2$  solves the boundary value problem (3)–(7). The condition (2.8) yields from (2.1)–(2.4) the following integro-differential equation for  $\rho$

$$\begin{aligned} \rho_t(t) &= \rho(t) \pi^{-1/2} \int_0^t g(\tau) (t - \tau)^{-3/2} \exp \left\{ -\left[\frac{1}{2} \rho(t)(t - \tau)^{-1/2}\right]^2 \right\} d\tau \\ &- \frac{1}{2} \pi^{-1/2} \int_0^t \rho_t(\tau) (t - \tau)^{-3/2} [\rho(t) - \rho(\tau)] \exp \left\{ -\left[\frac{1}{2} [\rho(t) - \rho(\tau)](t - \tau)^{-1/2}\right]^2 \right\} d\tau \\ &- \frac{1}{2} \pi^{-1/2} \int_0^t \rho_t(\tau) (t - \tau)^{-3/2} [\rho(t) + \rho(\tau)] \exp \left\{ -\left[\frac{1}{2} [\rho(t) + \rho(\tau)](t - \tau)^{-1/2}\right]^2 \right\} d\tau \\ &+ \frac{1}{2} \pi^{-1/2} t^{-3/2} \int_0^t f(\xi) [\rho(t) - \xi] \exp \left\{ -\left[\frac{1}{2} [\rho(t) - \xi] t^{-1/2}\right]^2 \right\} d\tau, \end{aligned} \quad (2.14)$$

$$\rho(0) = A.$$

We introduce the abbreviation

$$\rho_t(t) = F(\rho, \rho_t, g, f, A, t) = F(\rho) \quad (2.15)$$

for (2.14).

If we can solve this equation and if its solution  $\rho(t)$ , has a non-negative derivative then  $\rho(t)$  is  $R(t)$ , the free boundary, and  $u^2$  is a solution of the boundary value problem.

**3. Properties of  $\rho_t(t)$ .** In this section we deduce some properties of the integro-differential equation (2.14) and of the free boundary which will be of use in our existence proof.

**LEMMA 1.** If  $r(t)$  is a continuously differentiable function for  $t \geq 0$  then

$$\rho_t(0) = \lim_{t \rightarrow 0} F(r) = -f_x(A). \quad (3.1)$$

*Proof.* The proof proceeds according to the methods of [3]. The first three integrals on the right in (2.14) tend to zero when  $t$  tends to zero while the fourth tends to  $-f_z(A)$ .

LEMMA 2. If  $R_i(t)$  exists and is continuous then  $R_i(t) > 0$ .

*Proof.* From (6) and (7) we see that  $R_i(0) = -f_z(A) > 0$ . Suppose to the contrary that  $R_i(t) \geq 0$ . Let  $t'$  be the smallest positive value of  $t$  for which  $R_i(t) = 0$ . Let  $D_{t'}$  be that part of  $D$  where  $t \leq t'$ .  $u_z$  is a solution of the heat equation in  $D_{t'}$ . Since  $g(t)$ ,  $f_z(x)$ , and  $R_i(t)$  are continuous and  $f_z(0) = -g(0)$  and  $-f_z(A) = R_i(0)$ ,  $u_z$  is continuous on that part of the boundary of  $D_{t'}$  which is not on the line  $t = t'$ . By the maximum principle [5], the maximum of  $u_z$  occurs on this part of the boundary of  $D_{t'}$ . Since  $-g < 0$ ,  $f'(x) < 0$ , and  $R_i(t) < 0$  for  $0 \leq t < t'$ , this maximum must occur at  $x = R(t')$ ,  $t = t'$ . Thus  $u_z \leq 0$  everywhere in  $D_{t'}$ . Thus since  $u = 0$  for  $x = R(t)$ , we have that  $u \geq 0$  in  $D_{t'}$ . Then in the closure of  $D_{t'}$ , every point on the free boundary is a minimum point of  $u$ . Now at a minimum point of  $u$ , the outward drawn derivative in a characteristic direction must be strictly negative\*. That is,  $u_z$  must be strictly negative along the free boundary. But at  $t = t'$ ,  $u_z = -R_i = 0$ . This contradiction implies the result.

**4. Existence.** To demonstrate the existence of a solution to the free boundary problem (3)–(7), we must show that the integro-differential equation (2.14) or (2.15) possesses a solution  $\rho(t)$  for which  $\rho_i(t) > 0$ . To do this we will use the principle of contracting mappings. We will first obtain the existence of a solution  $\rho(t)$  with  $\rho_i(t) > 0$  in the small and then using Lemma 2 of Sec. 3, show that this solution exists for all  $t > 0$ .

*Existence in the small.* Let  $B$  be the Banach space of continuously differentiable functions  $\{\rho(t)\}$ ,  $0 \leq t \leq T$  for some fixed  $T$  to be specified. Let the norm be

$$\|\rho(t)\| = \|\rho(0)\| + \sup_{0 \leq t \leq T} |\rho_i(t)|. \quad (4.1)$$

Thus convergence in  $B$  is uniform convergence of the function and its continuous derivative.

Let  $G$  be the closed set in  $B$  whose elements satisfy the following properties

$$\begin{aligned} (a) \quad & \rho(0) = A, \\ (b) \quad & \rho_i(0) = -f_z(A), \\ (c) \quad & 0 < l \leq \rho_i(t) \leq K, \end{aligned} \quad (4.2)$$

where  $l < -f_z(A)$  and  $K > -f_z(A)$  are to be specified.

Now consider the map  $\rho'_i = F(\rho)$ . We will show that if  $\rho_1$  and  $\rho_2 \in G$ , then  $\rho'_1$  and  $\rho'_2 \in G$  and  $\|\rho'_1 - \rho'_2\| \leq C(T) \|\rho_1 - \rho_2\|$ , where  $0 < C(T) < 1$  for  $T$  sufficiently small. These statements show that  $\rho'_i = F(\rho)$  is a map of  $G$  into  $G$  which is continuous and moreover contracting. Thus by the principle of contracting mappings  $\rho'_i = F(\rho)$  has a fixed point in  $G$ . This fixed point is our solution and possesses by [4.2(c)] a positive derivative. We now proceed with the proof.

If  $\rho(0) = A$  and  $\rho_i(0) = -f_z(A)$  then the same is true for  $\rho'(0)$  and  $\rho'_i(0)$ . The first since  $\rho'_i = F(\rho)$  is an integro-differential equation and we may arbitrarily require  $\rho'_i(0) = A$  and the second is the assertion of Lemma 1 of Sec. 3.

In Appendix 1, we show that

$$\|\rho'_1 - \rho'_2\| \leq C(T) \|\rho_1 - \rho_2\|, \quad \rho_1, \rho_2 \in G, \quad (4.3)$$

\*This is an unpublished result due to L. Nirenberg.

where  $C(T) = C(T, l, K, A, g, f) < 1$  for  $T$  sufficiently small. In Appendix 2 we show that  $\rho'_i$  is continuous. In Appendix 3 we show that

$$|\rho'_i(t) - [-f_*(A)]| \leq C_1 t^{1/2}, \quad (4.4)$$

uniformly for  $\rho \in G$ .

Thus we fix  $T$  so that  $C(T) < 1$  and

$$C_1 T^{1/2} < \max \{ |l - [-f_*(A)]|, |K - [-f_*(A)]| \}.$$

Our map is then into and contracting and a solution  $\rho(t)$  of (2.14) exists up to time  $T$ . Moreover this solution has a positive derivative.

*Existence in the large.* The proof of existence in the large proceeds as the above proof. The Banach space is now  $B_1$  the set of continuously differentiable functions  $\rho(t)$ ,  $0 \leq t \leq T_1$ , where  $T_1 > T$  is to be specified. We use the same norm as in  $B$ .  $G$  is replaced by  $G_1$ , those functions in  $B_1$  which up to time  $T$  are equal to the solution  $R(t)$  which we have just shown to exist.  $A$  is replaced by  $R(T)$  and  $-f_*(A)$  by  $R_*(T)$ .  $l$  and  $K$  are replaced by two other numbers  $l_1$  and  $K_1$  with  $0 < l_1 < R_*(T)$  and  $K_1 > R_*(T)$ .

With this setup the requirements of the principle of contracting mappings are satisfied here in essentially the same way as above. Thus we produce a  $T_1$  such that for  $T, -T > 0$  and sufficiently small, we have existence of  $R(T)$  up to time  $T_1$  and moreover  $R_i(t) > 0$  for  $0 \leq t \leq T_1$ . We see that we may iterate this procedure and produce a sequence of  $T_i, i = 1, 2, \dots$ , such that  $R(t)$  exists and  $R_i(t) > 0$  for  $t \leq T_i$ . There remains only to show that  $T_i \rightarrow \infty$ . From the form of the estimates in the appendices and the definitions of the sets  $G_i$  we see that this will be the case if  $R_i(t)$  never vanishes for then we may always extend our solution slightly further. But the vanishing of  $R_i(t)$  is ruled out by Lemma 2 of Sec. 3. Thus  $T_i \rightarrow \infty$  and  $R(t)$  exists for all time.

**Remark.** We have mentioned that for our proof it is essential that  $A \neq 0$ . A passage to the limit as  $A \rightarrow 0$  is suggested to obtain the solution with  $A = 0$ . This limit procedure would be legitimate if for some sequence  $R^A(t)$  with  $A$  tending to zero, the slopes,  $R^A_i(t)$ , have a positive lower bound. For in this event the set of functions  $R^A(t)$  are equi-continuously differentiable and a simple compactness argument justifies the passage to the limit. If the condition (7) is changed to  $u_*[R(t), t] = B - R_i(t)$ ,  $B > 0$ , then an obvious extension of Lemma 2 yields a positive lower bound for the slopes  $R^A_i(t)$  and in this case our method yields existence for the case  $A = 0$ .

**Acknowledgment.** The author is grateful to Professor I. Kolodner and to Professor J. Keller for their assistance in preparing this paper.

**Appendix 1.** *Continuity and contracting of the map  $\rho'_i = F(\rho)$ .* In this appendix we show that the map  $\rho'_i = F(\rho)$  is continuous for  $\rho \in G$  is contracting. Let  $u(t)$  and  $v(t)$  be two generic elements in  $G$ . Let  $u'_i = F(u)$  and  $v'_i = F(v)$ . We have

$$\|u' - v'\| = \max_{0 \leq t \leq T} \|F(u) - F(v)\|$$

since  $u'(0) = v'(0) = A$ .

The computation of the right-hand side of (A1) is divided into four parts corresponding to each of the four integrals occurring in  $F$ . The computations for the first three integrals are so alike that we illustrate the computation involving the third and fourth integrals only.

(i) **THIRD INTEGRAL.** We are led to consider

$$I_1 + I_2 = \frac{1}{2}\pi^{-1/2} \left| \int_0^t u_r(\tau)(t-\tau)^{-3/2}[u(t) + u(\tau) - v(t) - v(\tau)] \right. \\ \left. \cdot \exp \left\{ -\left[\frac{1}{2}[u(t) + u(\tau)](t-\tau)^{-1/2}\right] d\tau \right\} \right. \\ \left. + \frac{1}{2}\pi^{-1/2} \left| \int_0^t [v_r(\tau) - u_r(\tau)][v(t) + v(\tau)](t-\tau)^{-3/2} \right. \right. \\ \left. \cdot \exp \left\{ -\left[\frac{1}{2}[v(t) + v(\tau)](t-\tau)^{-1/2}\right]^2 d\tau \right\} d\tau \right|.$$

By the law of the mean, we have

$$I_1 \leq \frac{1}{2}K\pi^{-1/2} \int_0^t |u(t) + u(\tau) - v(t) - v(\tau)| (t-\tau)^{-3/2} |e^{-z^2}(1-2z^2)| d\tau,$$

where  $z$  is  $[\frac{1}{2}[\rho(t) + \rho(\tau)](t-\tau)^{-1/2}]$  for some  $\rho \in B$ . Thus

$$I_1 \leq Kt\pi^{-1/2} \|u - v\| \int_0^t (t-\tau)^{-3/2} |e^{-z^2}(1-2z^2)| d\tau.$$

Now

$$\frac{2A + lt}{2(t-\tau)^{1/2}} \leq \frac{\rho(t) + \rho(\tau)}{2(t-\tau)^{1/2}} \leq \frac{A + Kt}{(t-\tau)^{1/2}} \quad \text{for } \rho \in B.$$

Then

$$I_1 \leq Kt\pi^{-1/2} \|u - v\| \int_0^t (t-\tau)^{-3/2} [1 + 2(A + Kt)^2(t-\tau)^{-1}] \\ \cdot \exp \left\{ -\left[\frac{1}{2}(2A + lt)(t-\tau)^{-1/2}\right]^2 d\tau \right\} d\tau.$$

Let

$$\sigma = \frac{1}{2}(2A + lt)(t-\tau)^{-1/2}, \quad 2 d\sigma = \frac{1}{2}(2A + lt)(t-\tau)^{-3/2} d\tau.$$

Then

$$I_1 \leq 4K\pi^{-1/2} \|u - v\| t(2A + lt)^{-1} \int_{2A+lt/2t^{1/2}}^{\infty} e^{-\sigma^2} [1 + 8\sigma^2(A + Kt)^2(2A + lt)^{-2}] d\sigma.$$

For  $I_2$  we have

$$I_2 \leq \frac{1}{2}\pi^{-1/2} \|u - v\| \int_0^t [v(t) + v(\tau)](t-\tau)^{-3/2} \exp \left\{ -\left[\frac{1}{2}[v(t) + v(\tau)](t-\tau)^{-1/2}\right]^2 d\tau \right\} d\tau \\ \leq \pi^{-1/2} \|u - v\| (A + Kt) \int_0^t (t-\tau)^{-3/2} \exp \left\{ -\left[\frac{1}{2}(2A + lt)(t-\tau)^{-1/2}\right]^2 d\tau \right\} d\tau \\ = 4\pi^{-1/2} \|u - v\| \frac{A + Kt}{2A + lt} \int_{2A+lt/2t^{1/2}}^{\infty} e^{-\sigma^2} d\sigma \\ \leq 4\left(\frac{t}{\pi}\right)^{1/2} \|u - v\| (A + Kt)(2A + lt)^{-2} \exp \left\{ -\frac{1}{4}(2A + lt)^2 t^{-1} \right\}.$$

ii. **FOURTH INTEGRAL.** For this integral we have

$$\begin{aligned}
 I_3 + I_4 &= \frac{1}{2}\pi^{-1/2}t^{-3/2} \left| \int_{-A}^A f(\xi)[u(t) - v(t)] \exp \left\{ -\frac{1}{2}[u(t) - \xi]^2 t^{-1} \right\} d\xi \right| \\
 &\quad + \frac{1}{2}\pi^{-1/2}t^{-3/2} \left| \int_{-A}^A f(\xi)[v(t) - \xi] \left[ \exp \left\{ -\frac{1}{2}[u(t) - \xi]^2 t^{-1} \right\} \right. \right. \\
 &\quad \left. \left. - \exp \left\{ -\frac{1}{2}[u(t) - \xi]^2 t^{-1} \right\} \right] d\xi \right| \\
 I_3 &\leq \frac{1}{2}(\pi t)^{-1/2} \|u - v\| \int_{-A}^A |f(\xi)| \exp \left\{ -\frac{1}{2}[u(t) - \xi]^2 t^{-1} \right\} d\xi \\
 &= \frac{1}{2}(\pi t)^{-1/2} \|u - v\| \int_{-\lambda}^{\lambda} + \int_{\lambda}^A |f(\xi)| \exp \left\{ -\frac{1}{2}[u(t) - \xi]^2 t^{-1} \right\} d\xi \\
 &= I_5 + I_6.
 \end{aligned}$$

where  $\lambda = \lambda(t)$

$$\begin{aligned}
 I_5 &\leq \frac{1}{2}(\pi t)^{-1/2} \|u - v\| (A - \lambda) \max_{A \leq \xi \leq \lambda} |f(\xi)| \\
 I_5 &\leq \frac{1}{2}(\pi t)^{-1/2} (A - \lambda)^2 \max_{A \leq \xi \leq \lambda} |f_2(\xi)| \quad (A1)
 \end{aligned}$$

since  $f(A) = 0$ .

$$I_6 \leq \frac{1}{2}(\pi t)^{-1/2} (\lambda + A) \exp \left\{ -\frac{1}{2}(A + \lambda)^2 t^{-1} \right\} \max_{-A \leq \xi \leq \lambda} |f(\xi)|. \quad (A2)$$

From (A1) and (A2) we see that if as  $t \rightarrow 0$   $\lambda(t) \rightarrow A$  faster than  $t^{1/4}$  but slower than  $t^{1/2}$  then

$$I_5 + I_6 \leq \|u - v\| \text{const } o(1).$$

The computation involving  $I_4$  is similar to the one just conducted and will be omitted.

**Appendix 2.** *Continuous differentiability of an image under  $F$ .* In this appendix we show that if  $\rho'_t = F(\rho)$ ,  $\rho \in G$  then  $\rho'_t$  is a continuous function of  $t$ . The computations are slight variations of the computations in Appendix 1. Therefore we will carry them out only for the first integral in  $F(\rho)$ .

Let  $a \geq 0$  and consider

$$\begin{aligned}
 &\pi^{-1/2} \rho(a + \Delta t) \int_0^{a+\Delta t} g(\tau)(a + \Delta t - \tau)^{-3/2} \exp \left\{ -\left[\frac{1}{2}\rho(a + \Delta t)(a + \Delta t - \tau)^{-1/2}\right]^2 \right\} d\tau \\
 &- \pi^{-1/2} \rho(a) \int_0^a g(\tau)(a - \tau)^{-3/2} \exp \left\{ -\left[\frac{1}{2}\rho(a)(a - \tau)^{-1/2}\right]^2 \right\} d\tau \\
 &= \pi^{-1/2} \rho(a + \Delta t) \int_0^{a+\Delta t} g(\tau)(a + \Delta t - \tau)^{-3/2} \exp \left\{ -\left[\frac{1}{2}\rho(a + \Delta t)(a + \Delta t - \tau)^{-1/2}\right]^2 \right\} d\tau \\
 &+ \pi^{-1/2} \int_0^a g(\tau) [\rho(a + \Delta t)(a + \Delta t - \tau)^{-3/2} \exp \left\{ -\left[\frac{1}{2}\rho(a + \Delta t)(a + \Delta t - \tau)^{-1/2}\right]^2 \right\} \\
 &- \rho(a)(a - \tau)^{-3/2} \exp \left\{ -\left[\frac{1}{2}\rho(a)(a - \tau)^{-1/2}\right]^2 \right\}] d\tau \\
 &= I_1 + I_2.
 \end{aligned}$$



In what follows, we use the fact that  $\rho \in G$  implies that  $\rho(a)$ ,  $\rho_i(a)$ ,  $\rho(a + \Delta t)$ ,  $\rho_i(a + \Delta t) > 0$ .

Consider first  $I_1$ . Let  $\sigma = \rho(a + \Delta t)/2(a + \Delta t - \tau)^{1/2}$ . Then

$$I_1 = 4\pi^{-1/2} \int_x^\infty g e^{-\sigma^2} dx, \quad x = \frac{1}{2}\rho(a + \Delta t)(\Delta t)^{-1/2}$$

$$< 4M\pi^{-1/2} \int_x^\infty e^{-\sigma^2} d\sigma, \quad M = \max_{0 \leq t \leq \tau} |g(t)|.$$

Since

$$\int_x^\infty e^{-\sigma^2} d\sigma < \frac{1}{2x} e^{-x^2}, \quad x \geq 0$$

we have that

$$I_1 < 4M\pi^{-1/2}(\Delta t)^{1/2}/\rho(a + \Delta t).$$

For  $I_2$  we have on application of the law of the mean of the differential calculus:

$$I_2 \leq \pi^{-1/2} \Delta t M \int_0^a \left\{ \left[ \rho_i(t)(t - \tau)^{-3/2} + \frac{3}{2} \rho(t)(t - \tau)^{-3/2} \right. \right. \\ \left. \left. - \rho(t)(t - \tau)^{-1/2} \left\{ \frac{1}{2} \rho_i(t)(t - \tau)^{-1/2} + \frac{1}{4} \rho(t)(t - \tau)^{-3/2} \right\} \right] \right. \\ \left. \cdot \exp \left[ - \left[ \frac{1}{2} \rho(t)(t - \tau)^{-1/2} \right]^2 \right] \right\} d\tau,$$

where the quantity in the curly brackets is to be evaluated at some value of  $t$  in the open interval  $0 \leq a < t < a + \Delta t$ . The integrand is thus finite and the integral exists.

**Appendix 3.** *The initial value of  $\rho'_i$ .* In this appendix we show that  $|\rho'_i(t) - [-f_x(A)]| \leq C_1(t)^{1/2}$ . We sketch the ideas of the proof since the computations are variations of those in Appendix 1.

We note first that the first three integrals vanish as  $t \rightarrow 0$ . For the first integral,

$$\pi^{-1/2} \rho(\tau) \int_0^t g(\tau)(t - \tau)^{-3/2} \exp \{ - [\frac{1}{2} \rho(t)(t - \tau)^{-1/2}]^2 \} d\tau,$$

the singularity at  $t = \tau$  in the exponential [ $\rho(0) = A > 0$ ] causes the integrand to be bounded in the range of integration. This bound depends on the minimum of  $\rho(t) \in G$  and this is bounded below by  $A + lT$ . Thus the entire integral vanishes with its upper limit uniformly in  $G$ .

For the second integral,

$$\frac{1}{2} \pi^{-1/2} \int_0^t \rho_\tau(\tau) [\rho(t) - \rho(\tau)] (t - \tau)^{-3/2} \exp \{ - [\frac{1}{2} [\rho(t) - \rho(\tau)] (t - \tau)^{-1/2}]^2 \} d\tau,$$

the term  $[\rho(t) - \rho(\tau)]/(t - \tau)$  approaches  $\rho_i(0) = -f_x(A)$  uniformly in  $G$ . Thus the integral tends to zero like  $t^{1/2}$  uniformly in  $G$ .

In the third integral,

$$\frac{1}{2} \pi^{-1/2} \int_0^t \rho_\tau(\tau) [\rho(t) + \rho(\tau)] (t - \tau)^{-3/2} \exp \{ - [\frac{1}{2} [\rho(t) - \rho(\tau)] (t - \tau)^{-1/2}]^2 \} d\tau,$$

the singularity in the exponent causes the integral to vanish with its upper limit as in the first integral.

We have, referring to [3], observed that the fourth integral tends to  $-f_x(A)$  as  $t \rightarrow 0$ . By examining the proof of this process one may observe that the limit is approached like  $t^{1/2}$  as  $t \rightarrow 0$ .

#### BIBLIOGRAPHY

1. G. W. Evans, *A note on the existence of a solution to a problem of Stefan*, Quart. Appl. Math. 9, 185-193 (1951)
2. G. W. Evans, E. Isaacson, and J. K. L. MacDonald, *Stefan-like problems*, Quart. Appl. Math., 8, 312-319 (1950)
3. E. Goursat, *Cours d'analyse mathématique*, 3, Chap. XXIX, 5th ed., Gauthier Villars, Paris, 1942
4. I. I. Kolodner, *Free boundary problem for the heat equation with applications to problems of change of phase*, Commun. Pure and Appl. Math., IX, 1-31 (1956)
5. L. Nirenberg, *A strong maximum principle for parabolic equations*, Commun. Pure Appl. Math., VI, 167-177 (1953)
6. L. Rubenstein, *On the solution of Stefan's problem*, Izvest. Akad. Nauk, USSR, 11, 37-53 (1947)