

## MINIMUM $L_1$ -NORM ESTIMATION FOR FRACTIONAL ORNSTEIN–UHLENBECK TYPE PROCESS

UDC 519.21

B. L. S. PRAKASA RAO

ABSTRACT. We investigate the asymptotic properties of the minimum  $L_1$ -norm estimator of the drift parameter for fractional Ornstein–Uhlenbeck type process satisfying a linear stochastic differential equation driven by a fractional Brownian motion.

### 1. INTRODUCTION

Long range dependence phenomenon is said to occur in a stationary time series  $\{X_n, n \geq 0\}$  if the  $\text{Cov}(X_0, X_n)$  of the time series tend to zero as  $n \rightarrow \infty$  and yet the condition

$$\sum_{n=0}^{\infty} |\text{Cov}(X_0, X_n)| = \infty$$

is satisfied. In other words, the  $\text{Cov}(X_0, X_n)$  tend to zero but so slowly that their sum diverges. This phenomenon was first observed by the hydrologist Hurst (1951) on projects involving the design of reservoirs along the Nile river (cf. Montanari (2003)) and by others in hydrological time series. It was recently observed that a similar phenomenon occurs in problems concerning traffic patterns of packet flows in high-speed data networks such as the Internet (cf. Willinger et al. (2003), Norros (2003)). The long range dependence pattern is also observed in macroeconomics and finance (cf. Henry and Zaffaroni (2003)). Long range dependence is also related to the concept of self-similarity for a stochastic process. A stochastic process  $\{X(t), t \in \mathbf{R}\}$  is said to be  $H$ -self-similar with index  $H > 0$  if for every  $a > 0$ , the processes  $\{X(at), t \in \mathbf{R}\}$  and the process  $\{a^H X(t), t \in \mathbf{R}\}$  have the same finite-dimensional distributions. Suppose a self-similar process has stationary increments. Then the increments form a stationary time series which exhibits long range dependence. A Gaussian  $H$ -self-similar process with stationary increments with  $0 < H < 1$  is called a fractional Brownian motion. A recent monograph by Doukhan et al. (2003) discusses the theory and applications of long range dependence and properties of fractional Brownian motion (Taqqu (2003)). If  $H = \frac{1}{2}$ , then the fractional Brownian motion reduces to the standard Brownian motion also called the Wiener process.

Diffusion processes and diffusion type processes satisfying stochastic differential equations driven by Wiener processes are used for stochastic modelling in wide variety of sciences such as population genetics, economic processes, signal processing as well as for modeling sunspot activity and more recently in mathematical finance. Statistical inference for diffusion type processes satisfying stochastic differential equations driven by

---

2000 *Mathematics Subject Classification.* Primary 62M09; Secondary 60G15.

*Key words and phrases.* Minimum  $L_1$ -norm estimation, fractional Ornstein–Uhlenbeck type process, fractional Brownian motion.

Wiener processes have been studied earlier, and a comprehensive survey of various methods is given in Prakasa Rao (1999). There has been a recent interest in studying similar problems for stochastic processes driven by a fractional Brownian motion to model processes involving long range dependence. Le Breton (1998) studied parameter estimation and filtering in a simple linear model driven by a fractional Brownian motion. In a recent paper, Kleptsyna and Le Breton (2002) studied parameter estimation problems for fractional Ornstein–Uhlenbeck process. Such processes play a potentially important role in the modelling of financial time series. The fractional Ornstein–Uhlenbeck process is a fractional analogue of the Ornstein–Uhlenbeck process, that is, a continuous time first order autoregressive process  $X = \{X_t, t \geq 0\}$  which is the solution of a one-dimensional homogeneous linear stochastic differential equation driven by a fractional Brownian motion (fBm)  $W^H = \{W_t^H, t \geq 0\}$  with Hurst parameter  $H \in (\frac{1}{2}, 1)$ . Such a process is the unique Gaussian process satisfying the linear integral equation

$$(1.1) \quad X_t = X_0 + \theta \int_0^t X_s ds + \sigma W_t^H, \quad t \geq 0.$$

The above-mentioned authors investigate the problem of estimation of the parameters  $\theta$  and  $\sigma^2$  based on the observation  $\{X_s, 0 \leq s \leq T\}$  and prove that the maximum likelihood estimator  $\hat{\theta}_T$  is strongly consistent as  $T \rightarrow \infty$ .

Parametric estimation for more general classes of stochastic processes satisfying the linear stochastic differential equations driven by fractional Brownian motion, observed over a fixed period of time  $T$ , is studied in Prakasa Rao (2003a,b). It is well known that the sequential estimation methods might lead to equally efficient estimators from the process observed possibly over a shorter expected period of observation time. We have investigated the conditions for such a phenomenon for estimating the drift parameter of a fractional Ornstein–Uhlenbeck type process in Prakasa Rao (2004). Novikov (1972) investigated the asymptotic properties of a sequential maximum likelihood estimator for the drift parameter in the Ornstein–Uhlenbeck process.

In spite of the fact that maximum likelihood estimators (MLE) are consistent and asymptotically normal and also asymptotically efficient in general, they have some shortcomings at the same time. Their calculation is often cumbersome as the expressions for MLE involve stochastic integrals which need good approximations for computational purposes. Furthermore, MLE are not robust in the sense that a slight perturbation in the noise component will change the properties of MLE substantially. In order to circumvent such problems, the minimum distance approach is proposed. Properties of the minimum distance estimators (MDE) were discussed in Millar (1984) in a general framework.

Our aim in this paper is to obtain the minimum  $L_1$ -norm estimates of the drift parameter of a fractional Ornstein–Uhlenbeck type process and investigate the asymptotic properties of such estimators following the work of Kutoyants and Pilibossian (1994).

## 2. PRELIMINARIES

Let  $(\Omega, \mathcal{F}, (\mathcal{F}_t), P)$  be a stochastic basis satisfying the usual conditions, and suppose that the processes discussed in the following are  $(\mathcal{F}_t)$ -adapted. Further the natural filtration of a process is understood as the  $P$ -completion of the filtration generated by this process.

Let  $W^H = \{W_t^H, t \geq 0\}$  be a normalized fractional Brownian motion with Hurst parameter  $H \in (\frac{1}{2}, 1)$ , that is, a Gaussian process with continuous sample paths such that  $W_0^H = 0$ ,  $E(W_t^H) = 0$  and

$$(2.1) \quad E(W_s^H W_t^H) = \frac{1}{2} [s^{2H} + t^{2H} - |s - t|^{2H}], \quad t \geq 0, s \geq 0.$$

Consider a stochastic process  $\{X_t, t \geq 0\}$  defined by the stochastic integral equation

$$(2.2) \quad X_t = x_0 + \theta \int_0^t X(s) ds + \varepsilon W_t^H, \quad 0 \leq t \leq T,$$

where  $\theta$  is an unknown drift parameter. For convenience, we write the above integral equation in the form of a stochastic differential equation

$$(2.3) \quad dX_t = \theta X(t) dt + \varepsilon dW_t^H, \quad X_0 = x_0, \quad 0 \leq t \leq T,$$

driven by the fractional Brownian motion  $W^H$ . For a discussion of the equivalence of (2.2) and (2.3), see Cheridito et al. (2003). Even though the process  $X$  is not a semimartingale, one can associate with it a semimartingale  $Z = \{Z_t, t \geq 0\}$ , which is called a *fundamental semimartingale*, such that the natural filtration  $(\mathcal{Z}_t)$  of the process  $Z$  coincides with the natural filtration  $(\mathcal{X}_t)$  of the process  $X$  (Kleptsyna et al. (2000)). Define, for  $0 < s < t$ ,

$$(2.4) \quad k_H = 2H\Gamma\left(\frac{3}{2} - H\right)\Gamma\left(H + \frac{1}{2}\right),$$

$$(2.5) \quad \kappa_H(t, s) = k_H^{-1} s^{1/2-H} (t-s)^{1/2-H},$$

$$(2.6) \quad \lambda_H = \frac{2H\Gamma(3-2H)\Gamma(H+\frac{1}{2})}{\Gamma(\frac{3}{2}-H)},$$

$$(2.7) \quad w_t^H = \lambda_H^{-1} t^{2-2H},$$

$$(2.8) \quad M_t^H = \int_0^t \kappa_H(t, s) dW_s^H, \quad t \geq 0.$$

The process  $M^H$  is a Gaussian martingale, called the *fundamental martingale* (cf. Norros et al. (1999)), and its quadratic variance is  $\langle M_t^H \rangle = w_t^H$ . Furthermore, the natural filtration of the martingale  $M^H$  coincides with the natural filtration of the fBM  $W^H$ . Let

$$(2.9) \quad K_H(t, s) = H(2H-1) \frac{d}{ds} \int_s^t r^{H-1/2} (r-s)^{H-3/2} dr, \quad 0 \leq s \leq t.$$

The sample paths of the process  $\{X_t, t \geq 0\}$  are smooth enough so that the process  $Q$  defined by

$$(2.10) \quad Q(t) = \frac{d}{dw_t^H} \int_0^t \kappa_H(t, s) X_s ds, \quad t \in [0, T],$$

is well defined, where  $w^H$  and  $k_H$  are as defined in (2.7) and (2.5) respectively and the derivative is understood in the sense of absolute continuity with respect to the measure generated by  $w^H$ . Moreover, the sample paths of the process  $Q$  belong to  $L^2([0, T], dw^H)$  a.s. [P]. The following theorem due to Kleptsyna et al. (2000) associates with the process  $X$  a *fundamental semimartingale*  $Z$  such that the natural filtration  $(\mathcal{Z}_t)$  coincides with the natural filtration  $(\mathcal{X}_t)$  of  $X$ .

**Theorem 2.1.** *Let the process  $Z = (Z_t, t \in [0, T])$  be defined by*

$$(2.11) \quad Z_t = \int_0^t \kappa_H(t, s) dX_s,$$

where the function  $\kappa_H(t, s)$  is defined in (2.5). Then the following results hold:

- (i) *The process  $Z$  is an  $(\mathcal{F}_t)$ -semimartingale with the decomposition*

$$(2.12) \quad Z_t = \theta \int_0^t Q(s) dw_s^H + \varepsilon M_t^H,$$

where  $M^H$  is the Gaussian martingale defined by (2.8).

(ii) The process  $X$  admits the representation

$$(2.13) \quad X_t = \int_0^t K_H(t, s) dZ_s,$$

where the function  $K_H$  is as defined in (2.9).

(iii) The natural filtrations of  $(Z_t)$  and  $(X_t)$  coincide.

Even though the fBm  $\{W_t^H, t \geq 0\}$  is not a semimartingale, it is still possible to define stochastic integration with respect to the fBm for deterministic integrands. For instance, for  $f \in L_2(\mathbf{R}_+) \cap L_1(\mathbf{R}_+)$ , one can define a stochastic integral of the form

$$\int_0^T f(s) dW_s^H$$

(cf. Gripenberg and Norris (1996), Norros et al. (1999)). Such a stochastic integral can be represented in terms of another stochastic integral with respect to the fundamental Gaussian martingale  $M^H$ . The following result is due to Kleptsyna et al. (2000).

For any measurable function  $f$  on  $[0, T]$  and for  $t \in [0, T]$ , define

$$(2.14) \quad K_H^f(t, s) = -2H \frac{d}{ds} \int_s^t f(r) r^{H-1/2} (r-s)^{H-3/2} dr, \quad 0 \leq s \leq t,$$

where the derivative exists in the sense of absolute continuity with respect to Lebesgue measure (cf. Samko et al. (1993)).

**Lemma 2.2.** *Let  $M^H$  be the fundamental martingale associated with the fBm  $W^H$ . Then the following equality holds a.s. [P]:*

$$(2.15) \quad \int_0^T f(s) dW_s^H = \int_0^T K_H^f(t, s) dM_s^H, \quad t \in [0, T],$$

provided both integrals on both sides are well defined.

Alos et al. (2001) used the stochastic calculus of variations or Malliavin calculus to develop a stochastic calculus with respect to Gaussian processes, in particular, for a fractional Brownian motion. The fractional Brownian motion  $W_t^H$  can be represented in the form

$$W_t^H = \int_0^t K^H(t, s) dW_s,$$

where  $\{W_t, t \geq 0\}$  is a Brownian motion and the kernel  $K^H$  is singular if  $H < \frac{1}{2}$  and regular if  $H > \frac{1}{2}$ . For the definition of singularity and regularity of a kernel, see Alos et al. (2001). They develop an Itô formula in the regular case and in the singular case for  $H > \frac{1}{4}$ . We will not go into more discussion here.

The following lemma due to Gripenberg and Norris (1996) gives the covariance between the two stochastic integrals

$$\int_0^T f(s) dW_s^H \quad \text{and} \quad \int_0^T g(s) dW_s^H.$$

**Lemma 2.3.** *For  $f, g \in L_2(\mathbf{R}_+) \cap L_1(\mathbf{R}_+)$ ,*

$$(2.16) \quad \begin{aligned} & \mathbb{E} \left( \int_0^\infty f(s) dW_s^H \int_0^\infty g(s) dW_s^H \right) \\ &= H(2H-1) \int_0^\infty \int_0^\infty f(s)g(t) |s-t|^{2H-2} dt ds. \end{aligned}$$

The result in Lemma 2.3 can be proved under weaker conditions. Pipiras and Taqqu (2000) showed that for  $H \in (\frac{1}{2}, 1)$  and for any two real-valued measurable functions

$$f, g \in \left\{ f : \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |f(u)| \cdot |f(v)| \cdot |u - v|^{2H-2} du dv < \infty \right\},$$

the result in Lemma 2.3 holds.

### 3. MINIMUM $L_1$ -NORM ESTIMATION

We now consider the problem of estimation of the parameter  $\theta$  based on the observation of a fractional Ornstein–Uhlenbeck type process  $X = \{X_t, 0 \leq t \leq T\}$  satisfying the stochastic differential equation

$$(3.1) \quad dX_t = \theta X(t) dt + \varepsilon dW_t^H, \quad X_0 = x_0, \quad 0 \leq t \leq T,$$

for a fixed time  $T$ , where  $\theta \in \Theta \subset \mathbf{R}$ , and study its asymptotic properties as  $\varepsilon \rightarrow 0$ .

Let  $x_t(\theta)$  be the solution of the above differential equation with  $\varepsilon = 0$ . It is obvious that

$$(3.2) \quad x_t(\theta) = x_0 e^{\theta t}, \quad 0 \leq t \leq T.$$

Let

$$(3.3) \quad S_T(\theta) = \int_0^T |X_t - x_t(\theta)| dt.$$

We define  $\theta_\varepsilon^*$  to be a *minimum  $L_1$ -norm estimator* if there exists a measurable selection  $\theta_\varepsilon^*$  such that

$$(3.4) \quad S_T(\theta_\varepsilon^*) = \inf_{\theta \in \Theta} S_T(\theta).$$

Conditions for the existence of a measurable selection are given in Lemma 3.1.2 in Prakasa Rao (1987). We assume that there exists a measurable selection  $\theta_\varepsilon^*$  satisfying the above equation. An alternative way of defining the estimator  $\theta_\varepsilon^*$  is by the relation

$$(3.5) \quad \theta_\varepsilon^* = \arg \inf_{\theta \in \Theta} \int_0^T |X_t - x_t(\theta)| dt.$$

**Consistency.** Let

$$W_T^{H*} = \sup_{0 \leq t \leq T} |W_t^H|.$$

The self-similarity of the fractional Brownian motion  $W_t^H$  implies that the random variables  $W_{at}^H$  and  $a^H W_t^H$  have the same probability distribution for any  $a > 0$ . Furthermore, it follows from the self-similarity that the supremum process  $W^{H*}$  has the property that the random variables  $W_{at}^{H*}$  and  $a^H W_t^{H*}$  have the same probability distribution for any  $a > 0$ . Hence we have the following observation due to Novikov and Valkeila (1999).

**Lemma 3.1.** *Let  $T > 0$  and  $\{W_t^H, 0 \leq t \leq T\}$  be an fBm with Hurst index  $H$ . Let  $W_T^{H*} = \sup_{0 \leq t \leq T} W_t^H$ . Then*

$$(3.6) \quad \mathbf{E} (W_T^{H*})^p = K(p, H) T^{pH}$$

for every  $p > 0$ , where  $K(p, H) = \mathbf{E}(W_1^{H*})^p$ .

Let  $\theta_0$  denote the true parameter. For any  $\delta > 0$ , define

$$(3.7) \quad g(\delta) = \inf_{|\theta - \theta_0| > \delta} \int_0^T |X_t(\theta) - x_t(\theta_0)| dt.$$

Note that  $g(\delta) > 0$  for any  $\delta > 0$ .

**Theorem 3.2.** For every  $p > 0$ , there exists a constant  $K(p, H)$  such that for every  $\delta > 0$ ,

$$(3.8) \quad \mathbb{P}_{\theta_0}^{(\varepsilon)}\{|\theta_\varepsilon^* - \theta_0| > \delta\} \leq 2^p T^{pH+p} K(p, H) e^{|\theta_0|Tp} (g(\delta))^{-p} \varepsilon^p = O((g(\delta))^{-p} \varepsilon^p).$$

*Proof.* Let  $\|\cdot\|$  denote the  $L_1$ -norm. Then

$$(3.9) \quad \begin{aligned} \mathbb{P}_{\theta_0}^{(\varepsilon)}\{|\theta_\varepsilon^* - \theta_0| > \delta\} &= \mathbb{P}_{\theta_0}^{(\varepsilon)} \left\{ \inf_{|\theta - \theta_0| \leq \delta} \|X - x(\theta)\| > \inf_{|\theta - \theta_0| > \delta} \|X - x(\theta)\| \right\} \\ &\leq \mathbb{P}_{\theta_0}^{(\varepsilon)} \left\{ \inf_{|\theta - \theta_0| \leq \delta} (\|X - x(\theta_0)\| + \|x(\theta) - x(\theta_0)\|) \right\} \\ &< \inf_{|\theta - \theta_0| > \delta} (\|x(\theta) - x(\theta_0)\| - \|X - x(\theta_0)\|) \\ &= \mathbb{P}_{\theta_0}^{(\varepsilon)} \left\{ 2\|X - x(\theta_0)\| > \inf_{|\theta - \theta_0| > \delta} \|x(\theta) - x(\theta_0)\| \right\} \\ &= \mathbb{P}_{\theta_0}^{(\varepsilon)} \left\{ \|X - x(\theta_0)\| > \frac{1}{2}g(\delta) \right\}. \end{aligned}$$

Since the process  $X_t$  satisfies the stochastic differential equation (3.2), it follows that

$$(3.10) \quad \begin{aligned} X_t - x_t(\theta_0) &= x_0 + \theta_0 \int_0^t X_s ds + \varepsilon W_t^H - x_t(\theta_0) \\ &= \theta_0 \int_0^t (X_s - x_s(\theta_0)) ds + \varepsilon W_t^H \end{aligned}$$

since  $x_t(\theta) = x_0 e^{\theta t}$ . Let  $U_t = X_t - x_t(\theta_0)$ . Then it follows from the above equation that

$$(3.11) \quad U_t = \theta_0 \int_0^t U_s ds + \varepsilon W_t^H.$$

Let  $V_t = |U_t| = |X_t - x_t(\theta_0)|$ . The above relation implies that

$$(3.12) \quad V_t = |X_t - x_t(\theta_0)| \leq |\theta_0| \int_0^t V_s ds + \varepsilon |W_t^H|.$$

Applying the Gronwall–Bellman lemma, we obtain that

$$(3.13) \quad \sup_{0 \leq t \leq T} |V_t| \leq \varepsilon e^{|\theta_0|T} \sup_{0 \leq t \leq T} |W_t^H|.$$

Hence

$$(3.14) \quad \begin{aligned} \mathbb{P}_{\theta_0}^{(\varepsilon)} \left\{ \|X - x(\theta_0)\| > \frac{1}{2}g(\delta) \right\} &\leq \mathbb{P} \left\{ \sup_{0 \leq t \leq T} |W_t^H| > \frac{e^{-|\theta_0|T} g(\delta)}{2\varepsilon T} \right\} \\ &= \mathbb{P} \left\{ W_T^{H*} > \frac{e^{-|\theta_0|T} g(\delta)}{2\varepsilon T} \right\}. \end{aligned}$$

Applying Lemma 3.1 to the estimate obtained above, we get that

$$\begin{aligned} \mathbb{P}_{\theta_0}^{(\varepsilon)}\{|\theta_\varepsilon^* - \theta_0| > \delta\} &\leq 2^p T^{pH+p} K(p, H) e^{|\theta_0|Tp} (g(\delta))^{-p} \varepsilon^p \\ &= O((g(\delta))^{-p} \varepsilon^p). \end{aligned} \quad \square$$

*Remarks.* As a consequence of the above theorem, we obtain that  $\theta_\varepsilon^*$  converges in probability to  $\theta_0$  under  $\mathbb{P}_{\theta_0}^{(\varepsilon)}$ -measure as  $\varepsilon \rightarrow 0$ . Furthermore, the rate of convergence is of order  $O(\varepsilon^p)$  for every  $p > 0$ .

**Asymptotic distribution.** We will now study the asymptotic distribution if any of the estimator  $\theta_\varepsilon^*$  after suitable scaling. It can be checked that

$$(3.15) \quad X_t = e^{\theta_0 t} \left\{ x_0 + \int_0^t e^{-\theta_0 s} \varepsilon dW_s^H \right\}$$

or equivalently

$$(3.16) \quad X_t - x_t(\theta_0) = \varepsilon e^{\theta_0 t} \int_0^t e^{-\theta_0 s} dW_s^H.$$

Let

$$(3.17) \quad Y_t = e^{\theta_0 t} \int_0^t e^{-\theta_0 s} dW_s^H.$$

Note that  $\{Y_t, 0 \leq t \leq T\}$  is a Gaussian process and can be interpreted as the “derivative” of the process  $\{X_t, 0 \leq t \leq T\}$  with respect to  $\varepsilon$ . Applying Lemma 2.2, we obtain that, P-a.s.,

$$(3.18) \quad Y_t e^{-\theta_0 t} = \int_0^t e^{-\theta_0 s} dW_s^H = \int_0^t K_H^f(t, s) dM_s^H, \quad t \in [0, T],$$

where  $f(s) = e^{-\theta_0 s}$ ,  $s \in [0, T]$ , and  $M^H$  is the fundamental Gaussian martingale associated with the fBm  $W^H$ . In particular it follows that the random variable  $Y_t e^{-\theta_0 t}$  and hence  $Y_t$  has normal distribution with mean zero, and furthermore, for any  $h \geq 0$ ,

$$(3.19) \quad \begin{aligned} \text{Cov}(Y_t, Y_{t+h}) &= e^{2\theta_0 t + \theta_0 h} \mathbb{E} \left[ \int_0^t e^{-\theta_0 u} dW_u^H \int_0^{t+h} e^{-\theta_0 v} dW_v^H \right] \\ &= e^{2\theta_0 t + \theta_0 h} H(2H - 1) \int_0^t \int_0^t e^{-\theta_0(u+v)} |u - v|^{2H-2} du dv \\ &= e^{2\theta_0 t + \theta_0 h} \gamma_H(t) \quad (\text{say}). \end{aligned}$$

In particular

$$(3.20) \quad \text{Var}(Y_t) = e^{2\theta_0 t} \gamma_H(t).$$

Hence  $\{Y_t, 0 \leq t \leq T\}$  is a zero mean Gaussian process with  $\text{Cov}(Y_t, Y_s) = e^{\theta_0(t+s)} \gamma_H(t)$  for  $s \geq t$ .

Let

$$(3.21) \quad \zeta = \arg \inf_{-\infty < u < \infty} \int_0^T |Y_t - utx_0 e^{\theta_0 t}| dt.$$

**Theorem 3.3.** *As  $\varepsilon \rightarrow 0$ , the random variable  $\varepsilon^{-1}(\theta_\varepsilon^* - \theta_0)$  converges in probability to a random variable whose probability distribution is the same as that of  $\zeta$  under  $P_{\theta_0}$ .*

*Proof.* Let  $x'_t(\theta) = x_0 t e^{\theta t}$  and let

$$(3.22) \quad Z_\varepsilon(u) = \|Y - \varepsilon^{-1}(x(\theta_0 + \varepsilon u) - x(\theta_0))\|$$

and

$$(3.23) \quad Z_0(u) = \|Y - ux'(\theta_0)\|.$$

Furthermore, let

$$(3.24) \quad A_\varepsilon = \{\omega : |\theta_\varepsilon^* - \theta_0| < \delta_\varepsilon\}, \quad \delta_\varepsilon = \varepsilon^\tau, \quad \tau \in \left(\frac{1}{2}, 1\right), \quad L_\varepsilon = \varepsilon^{\tau-1}.$$

Observe that the random variable  $u_\varepsilon^* = \varepsilon^{-1}(\theta_\varepsilon^* - \theta_0)$  satisfies the equation

$$(3.25) \quad Z_\varepsilon(u_\varepsilon^*) = \inf_{|u| < L_\varepsilon} Z_\varepsilon(u), \quad \omega \in A_\varepsilon.$$

Define

$$(3.26) \quad \zeta_\varepsilon = \arg \inf_{|u| < L_\varepsilon} Z_0(u).$$

Observe that, with probability one,

$$(3.27) \quad \sup_{|u| < L_\varepsilon} |Z_\varepsilon(u) - Z_0(u)| = \left\| \left\| Y - ux'(\theta_0) - \frac{1}{2}\varepsilon u^2 x''(\tilde{\theta}) \right\| - \|Y - ux'(\theta_0)\| \right\| \\ \leq \frac{\varepsilon}{2} L_\varepsilon^2 \sup_{|\theta - \theta_0| < \delta_\varepsilon} \int_0^T |x''(\theta)| dt \leq C\varepsilon^{2\tau-1}.$$

Here  $\tilde{\theta} = \theta_0 + \alpha(\theta - \theta_0)$  for some  $\alpha \in (0, 1)$ . Note that the last term in the above inequality tends to zero as  $\varepsilon \rightarrow 0$ . Furthermore, the process  $\{Z_0(u), -\infty < u < \infty\}$  has a unique minimum  $u^*$  with probability one. This follows from the arguments given in Theorem 2 of Kutoyants and Pilibossian (1994). In addition, we can choose the interval  $[-L, L]$  such that

$$(3.28) \quad \mathbf{P}_{\theta_0}^{(\varepsilon)}\{u_\varepsilon^* \in (-L, L)\} \geq 1 - \beta g(L)^{-p}$$

and

$$(3.29) \quad \mathbf{P}\{u^* \in (-L, L)\} \geq 1 - \beta g(L)^{-p},$$

where  $\beta > 0$ . Note that  $g(L)$  increases as  $L$  increases. The processes  $Z_\varepsilon(u)$ ,  $u \in [-L, L]$ , and  $Z_0(u)$ ,  $u \in [-L, L]$ , satisfy the Lipschitz conditions and  $Z_\varepsilon(u)$  converges uniformly to  $Z_0(u)$  over  $u \in [-L, L]$ . Hence the minimizer of  $Z_\varepsilon(\cdot)$  converges to the minimizer of  $Z_0(u)$ . This completes the proof.  $\square$

*Remarks.* We have seen earlier that the process  $\{Y_t, 0 \leq t \leq T\}$  is a zero mean Gaussian process with the covariance function

$$\text{Cov}(Y_t, Y_s) = e^{\theta_0(t+s)} \gamma_H(t)$$

for  $s \geq t$ . Recall that

$$(3.30) \quad \zeta = \arg \inf_{-\infty < u < \infty} \int_0^T |Y_t - utx_0 e^{\theta_0 t}| dt.$$

It is not clear what the distribution of  $\zeta$  is. Observe that for every  $u$ , the integrand in the above integral is the absolute value of a Gaussian process  $\{J_t, 0 \leq t \leq T\}$  with the mean function  $\mathbf{E}(J_t) = -utx_0 e^{\theta_0 t}$  and the covariance function  $\text{Cov}(J_t, J_s) = e^{\theta_0(t+s)} \gamma_H(t)$  for  $s \geq t$ . It would be interesting to say something about the distribution of  $\zeta$  through simulation studies even if an explicit computation of the distribution seems to be difficult.

#### BIBLIOGRAPHY

1. E. Alos, O. Mazet, and D. Nualart, *Stochastic calculus with respect to Gaussian processes*, Ann. Probab. **29** (2001), 766–801. MR1849177 (2002g:60083)
2. P. Cheridito, H. Kawaguchi and M. Maejima, *Fractional Ornstein–Uhlenbeck processes*, Electronic Journal of Probability **8** (2003), 1–14. MR1961165 (2003m:60096)
3. P. Doukhan, G. Oppenheim, and M. S. Taqqu, *Theory of Long-Range Dependence*, Birkhäuser, Boston, 2003. MR1956041 (2003h:60004)
4. G. Gripenberg and I. Norros, *On the prediction of fractional Brownian motion*, J. Appl. Probab. **33** (1996), 400–410. MR1385349 (97a:60058)
5. M. Henry and P. Zaffaroni, *The long-range dependence paradigm for macroeconomics and finance*, Theory of Long-Range Dependence (P. Doukhan, G. Oppenheim, and M. S. Taqqu, eds.), Birkhäuser, Boston, 2003, pp. 417–438. MR1957502
6. H. E. Hurst, *Long term storage capacity of reservoirs (with discussion)*, Trans. Am. Soc. Civ. Eng. **116** (1951), 770–808.



7. M. L. Kleptsyna and A. Le Breton, *Statistical analysis of the fractional Ornstein–Uhlenbeck type process*, *Statist. Infer. Stoch. Proces.* **5** (2002), 229–248. MR1943832 (2003i:62140)
8. M. L. Kleptsyna, A. Le Breton, M.-C. Roubaud, *Parameter estimation and optimal filtering for fractional type stochastic systems*, *Statist. Infer. Stoch. Proces.* **3** (2000), 173–182. MR1819294 (2002c:60089)
9. Yu. Kutoyants and P. Pilibossian, *On minimum  $L_1$ -norm estimate of the parameter of the Ornstein–Uhlenbeck process*, *Statist. Probab. Lett.* **20** (1994), 117–123. MR1293288 (95g:62161)
10. A. Le Breton, *Filtering and parameter estimation in a simple linear model driven by a fractional Brownian motion*, *Statist. Probab. Lett.* **38** (1998), 263–274. MR1629915 (99c:60088)
11. P. W. Millar, *A general approach to the optimality of the minimum distance estimators*, *Trans. Amer. Math. Soc.* **286** (1984), 377–418. MR0756045 (85m:62057)
12. A. Montanari, *Long-range dependence in hydrology*, *Theory of Long-Range Dependence* (P. Doukhan, G. Oppenheim, and M. S. Taqqu, eds.), Birkhäuser, Boston, 2003, pp. 461–472. MR1957504
13. I. Norros, E. Valkeila, and J. Virtamo, *An elementary approach to a Girsanov type formula and other analytical results on fractional Brownian motion*, *Bernoulli* **5** (1999), 571–587. MR1704556 (2000f:60053)
14. I. Norros, *Large deviations of queues with long-range dependent input*, *Theory of Long-Range Dependence* (P. Doukhan, G. Oppenheim, and M. S. Taqqu, eds.), Birkhäuser, Boston, 2003, pp. 409–415. MR1957501
15. A. A. Novikov, *Sequential estimation of the parameters of diffusion processes*, *Mathematical Notes* **12** (1972), 812–818. MR0317493 (47:6040)
16. A. A. Novikov and E. Valkeila, *On some maximal inequalities for fractional Brownian motion*, *Statist. Probab. Lett.* **44** (1999), 47–54. MR1706311 (2000g:60076)
17. V. Papiras and M. Taqqu, *Integration questions related to fractional Brownian motion*, *Prob. Theory and Rel. Fields* **118** (2000), 121–291. MR1790083 (2002c:60091)
18. B. L. S. Prakasa Rao, *Asymptotic Theory of Statistical Inference*, Wiley, New York, 1987. MR0874342 (88b:62001)
19. B. L. S. Prakasa Rao, *Statistical Inference for Diffusion Type Processes*, Arnold, London and Oxford University Press, New York, 1999. MR1717690 (2000k:62149)
20. B. L. S. Prakasa Rao, *Parametric estimation for linear stochastic differential equations driven by fractional Brownian motion*, *Random Oper. Stoch. Eq.* **11** (2003), 229–242. MR2009183 (2005a:62188)
21. B. L. S. Prakasa Rao, *Berry–Esseen bound for MLE for linear stochastic differential equations driven by fractional Brownian motion*, Preprint, Indian Statistical Institute, New Delhi, 2003b.
22. B. L. S. Prakasa Rao, *Sequential estimation for fractional Ornstein–Uhlenbeck type process*, *Sequential Analysis* **23** (2004), 33–44. MR2066971 (2005d:62127)
23. S. G. Samko, A. A. Kilbas, and O. I. Marichev, *Fractional Integrals and Derivatives*, Gordon and Breach Science, 1993. MR1347689 (96d:26012)
24. M. Taqqu, *Fractional Brownian motion and long-range dependence*, *Theory of Long-Range Dependence* (P. Doukhan, G. Oppenheim, and M. S. Taqqu, eds.), Birkhäuser, Boston, 2003, pp. 5–38. MR1956042
25. W. Willinger, V. Paxson, R. H. Riedi, and M. Taqqu, *Long-range dependence and data network traffic*, *Theory of Long-Range Dependence* (P. Doukhan, G. Oppenheim, and M. S. Taqqu, eds.), Birkhäuser, Boston, 2003, pp. 373–407. MR1957500

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF HYDERABAD, HYDERABAD 500 046,  
INDIA

*E-mail address:* blsprsm@uohyd.ernet.in

Received 25/NOV/2003

Originally published in English