

## PROKHOROV–LOÈVE STRONG LAW OF LARGE NUMBERS FOR MARTINGALES NORMALIZED BY OPERATORS

UDC 519.21

V. V. BULDYGIN AND V. O. KOVAL'

ABSTRACT. We study strong laws of large numbers for multivariate martingales normalized by linear operators in a finite-dimensional Euclidean space. Corollaries of the general results are considered for martingales under moment restrictions.

### 1. INTRODUCTION

Let  $(X_n, n \geq 1)$  be a sequence of independent symmetric random vectors in the Euclidean space  $\mathbf{R}^m$  ( $m \geq 1$ );  $(A_n, n \geq 1)$  a sequence of nonrandom linear operators acting from  $\mathbf{R}^m$  to  $\mathbf{R}^d$  ( $d \geq 1$ );  $\|x\|$  the Euclidean norm of a vector  $x$ ; and  $\mathfrak{N}$  the set of all nondecreasing unbounded sequences of positive integers.

Put  $S_n = \sum_{i=1}^n X_i$ ,  $n \geq 1$ . The Prokhorov–Loève strong law of large numbers is an assertion about the equivalence between the almost sure convergence to zero of the sequence  $(A_n S_n, n \geq 1)$ , that is,

$$(a) \quad \|A_n S_n\| \rightarrow 0 \quad \text{a.s.} \quad (n \rightarrow \infty),$$

and the almost sure convergence to zero of the sequence of independent random vectors  $(A_{n_{j+1}}(S_{n_{j+1}} - S_{n_j}), j \geq 1)$ , that is,

$$(b) \quad \|A_{n_{j+1}}(S_{n_{j+1}} - S_{n_j})\| \rightarrow 0 \quad \text{a.s.} \quad (j \rightarrow \infty),$$

where the sequences  $(n_j, j \geq 1)$  belong to a specified subset  $\tilde{\mathfrak{N}}$  of the set  $\mathfrak{N}$ . The principal problem is to determine the set  $\tilde{\mathfrak{N}}$  and find conditions for the equivalence of assertions (a) and (b). In what follows, the set  $\tilde{\mathfrak{N}}$  is denoted by  $\mathfrak{N}((A_n))$  and called the *test class* for the sequence  $(A_n, n \geq 1)$ .

The first results concerning this form of the strong law of large numbers are obtained in Prokhorov [1] and Loève [2] in the case of  $m = d = 1$ . The most advanced results in this direction for  $m = d = 1$  are obtained in [3, 4]. The test class  $\mathfrak{N}((A_n))$  is simple in this case; namely if  $(a_n, n \geq 1)$  is a normalizing sequence of positive numbers such that  $a_n \rightarrow 0$  as  $n \rightarrow \infty$ , then  $\mathfrak{N}((a_n))$  consists of the single sequence  $(n_j, j \geq 1)$  defined as  $n_j = \max\{n: a_n \geq \lambda^{-j}\}$ ,  $j \geq 1$ , where  $\lambda > 1$ .

The multidimensional case is not so easy. The Prokhorov–Loève strong law of large numbers for sums of independent symmetric random vectors normalized by operators is obtained in [5]. The test class  $\mathfrak{N}((A_n))$  for this case depends on the normalizing sequence of operators  $(A_n, n \geq 1)$  and contains a finite number of sequences. The construction of the test class is used in [5] in the proof of the equivalence of assertions (a) and (b)

---

2000 *Mathematics Subject Classification.* Primary 60F15.

*Key words and phrases.* Strong law of large numbers, almost sure convergence, almost sure boundedness, martingales, normalizations by operators.

for sums of independent symmetric random vectors. It is worth mentioning that this construction cannot be applied for more complicated cases.

The procedure for the construction of the test class is given in Section 2 of this paper. General results on the almost sure convergence to zero and on the almost sure boundedness of sequences of random vectors normalized by operators are also given in Section 2. These results are applied to partial sums of orthogonal random vectors in Section 3. Martingales normalized by operators are considered in Sections 4 and 5. Note that the results of Sections 3–5 are discussed in [6, 7]. However, the test class is not well defined there for the general case.

We use the following notation:  $N$  means the set of positive integers;  $\langle x, y \rangle$  is the scalar product of vectors  $x$  and  $y$ ;  $\|A\| = \sup_{\|x\|=1} \|Ax\|$  is the norm of an operator  $A$ ;  $\xrightarrow{P}$  stands for the convergence in probability; and  $\mathbf{E}$  means the mathematical expectation. For two operators,  $A$  and  $B$ , the notation  $A \geq B$  means that the operator  $A - B$  is positive semidefinite. For the sake of definiteness, we put  $\max_{\emptyset}(\cdot) = 0$  and  $\sum_{i=m}^n(\cdot) = 0$  for  $m > n$ .

## 2. GENERAL CONDITIONS FOR THE ALMOST SURE CONVERGENCE TO ZERO AND ALMOST SURE BOUNDEDNESS OF SEQUENCES OF RANDOM VECTORS NORMALIZED BY OPERATORS

Below we obtain some Prokhorov–Loève type sufficient conditions for the almost sure convergence to zero and the almost sure boundedness of arbitrary sequences of random vectors normalized by operators. First we consider the procedure for the construction of the test class  $\mathfrak{N}((A_n))$  for a given sequence  $(A_n, n \geq 1)$ .

**Construction of the test class  $\mathfrak{N}((A_n))$ .** The procedure for the construction of the test class is rather complicated. Thus we split it into several steps. Without loss of generality we assume that  $(A_n, n \geq 1)$  are  $d \times m$  matrices.

*Case 1:*  $d = 1$ ,  $m = 1$ , and  $\|A_n\| \rightarrow 0$  as  $n \rightarrow \infty$ . In this case,

$$(A_n, n \geq 1) = (a_n, n \geq 1)$$

is a sequence of real numbers. Fix an arbitrary number  $\lambda > 1$  and define the sequence of positive integers  $(n_j, j \geq 1)$  as follows:

$$(1) \quad n_j = \max \{n: |a_n| \geq \lambda^{-j}\}, \quad j \geq 1.$$

If there are infinitely many nonzero terms in the sequence  $(a_n, n \geq 1)$ , then we put

$$(2) \quad \mathfrak{N}((A_n)) = \mathfrak{N}((a_n), \lambda) = \{(n_j, j \geq 1)\}.$$

Otherwise, i.e., if all the terms of the sequence starting with some term are zero, we put

$$(3) \quad \mathfrak{N}((A_n)) = \emptyset.$$

*Case 2:*  $d = 1$ ,  $m \geq 1$ , and  $\|A_n\| \rightarrow 0$  as  $n \rightarrow \infty$ . In this case,

$$(A_n, n \geq 1) = (T_n, n \geq 1)$$

is a sequence of row vectors of dimension  $m$  and the class  $\mathfrak{N}((A_n))$  is defined by induction on  $m$ . The class is already defined for  $m = 1$  (see Case 1). Assume that the class is defined for some  $m = s \geq 1$  and define it for  $m = s + 1$ . The last coordinate of the vector  $T_n \in \mathbf{R}^{s+1}$  is denoted by  $t_n$ .

a) First we consider the case of  $t_n \neq 0$ ,  $n \geq 1$ . Fix an arbitrary number  $\lambda > 1$  and put

$$\hat{n}_j = \max \{n: |t_n| \geq \lambda^{-j}\}, \quad j \geq 1.$$

Since  $\|T_n\| \rightarrow 0$  as  $n \rightarrow \infty$ , we have  $|t_n| \rightarrow 0$  as  $n \rightarrow \infty$ . Thus the sequence  $(\hat{n}_j, j \geq 1)$  is well defined. By  $\tilde{T}_n$  we denote the vector formed by the first  $s$  coordinates of the vector  $T_n$ , that is,  $T_n = (\tilde{T}_n, t_n)$ . For each positive integer  $n \geq 1$ , put

$$(4) \quad B_n = \theta_n \left( t_n^{-1} \tilde{T}_n - t_{\hat{n}_{j+1}}^{-1} \tilde{T}_{\hat{n}_{j+1}} \right), \quad \hat{n}_j \leq n < \hat{n}_{j+1},$$

where  $\theta_n = \min\{|t_n|, |t_{\hat{n}_{j+1}}|\}$ . If  $\hat{n}_j \leq n < \hat{n}_{j+1}$ , then  $\|B_n\| \leq \|\tilde{T}_n\| + \|\tilde{T}_{\hat{n}_{j+1}}\|$ . This implies that

$$(5) \quad \|B_n\| \rightarrow 0 \quad (n \rightarrow \infty),$$

since  $\|\tilde{T}_n\| \rightarrow 0$  as  $n \rightarrow \infty$ . The latter relation follows from  $\|T_n\| \rightarrow 0$  as  $n \rightarrow \infty$ . Since  $B_n \in \mathbf{R}^s, n \geq 1$ , and relation (5) holds, the test class  $\mathfrak{N}((B_n))$  is defined for the sequence  $(B_n, n \geq 1)$  by the induction assumption.

Fix an arbitrary sequence  $(n_j, j \geq 1) \in \mathfrak{N}((B_n))$ . Using this sequence and the sequences  $(\hat{n}_j, j \geq 1)$  we define the sequence of intervals  $(I_n, n \geq 1)$  in the set  $N$  of positive integers as follows. Let  $I_1 = [\hat{n}_j, \hat{n}_{j+1})$  if  $n_1 \in [\hat{n}_j, \hat{n}_{j+1})$ . If an interval  $I_i, i \geq 1$ , is already defined, then  $I_{i+1}$  is the interval that is different from  $I_i$ , is the closest to  $I_i$  from the right among the intervals of the sequence  $([\hat{n}_j, \hat{n}_{j+1}), j \geq 1)$ , and that contains the terms of the sequence  $(n_j, j \geq 1)$ . For  $i \geq 1$ , put  $\bar{j}(i) = \min\{p: n_p \in I_i\}$ ,  $\bar{j}(i) = \max\{p: n_p \in I_i\}$ , and  $m_i = \max\{n: n \in I_i\} + 1$ . Consider the two sequences

$$\begin{aligned} & (n_{j_1}, m_2, n_{j(3)}, m_4, \dots, n_{j(2p-1)}, m_{2p}, \dots), \\ & (m_1, n_{j_2}, m_3, n_{j(4)}, \dots, m_{2p-1}, n_{j(2p)}, \dots). \end{aligned}$$

Since

$$n_{j(i)} < m_i \leq n_{\bar{j}(i+1)} \leq n_{j(i+1)} < m_{i+1}, \quad i \geq 1,$$

each of the two sequences belongs to the set  $\mathfrak{N}$ . The set containing these two sequences and the sequence  $(n_j, j \geq 1)$  is denoted by  $\mathfrak{N}_{(n_j)}$ . Then we put

$$(6) \quad \mathfrak{N}((A_n)) = \mathfrak{N}((T_n)) = \bigcup_{(n_j, j \geq 1) \in \mathfrak{N}((B_n))} \mathfrak{N}_{(n_j)} \cup \{(\hat{n}_j, j \geq 1)\}.$$

b) Now consider a general sequence  $(t_n, n \geq 1)$ . We split the set  $N$  of positive integers into two subsets  $N_1$  and  $N_2$ , where  $N_1 = \{n \in N: t_n \neq 0\}$  and  $N_2 = \{n \in N: t_n = 0\}$ . Then the sequence  $(T_n, n \geq 1)$  is split into two subsequences  $T^1 = (T_n^1, n \geq 1)$  and  $T^{(2)} = (T_n^2, n \geq 1)$  corresponding to the subsets  $N_1$  and  $N_2$ . Note that  $T^2$  is a sequence of  $s$ -dimensional vectors; thus the class  $\mathfrak{N}(T^2)$  is defined for it by the induction assumption. If the set  $N_1$  contains only a finite number of elements, then we let  $\mathfrak{N}((T_n)) = \mathfrak{N}(T^2)$ . Otherwise, that is, if the set  $N_1$  is infinite, the class  $\mathfrak{N}(T^1)$  is defined for  $T^1$  according to a). If both sets  $N_1$  and  $N_2$  are infinite, we let

$$\mathfrak{N}((A_n)) = \mathfrak{N}((T_n)) = \mathfrak{N}(T^1) \cup \mathfrak{N}(T^2).$$

*Case 3:*  $d = 1, m \geq 1$ , and the sequence  $(A_n, n \geq 1) = (T_n, n \geq 1)$  converges. Let  $T = \lim_{n \rightarrow \infty} T_n$  and put  $\check{T}_n = T_n - T, n \geq 1$ . Thus  $\|\check{T}_n\| \rightarrow 0$  as  $n \rightarrow \infty$  and the class  $\mathfrak{N}((\check{T}_n))$  is defined according to Case 2. Then we put

$$\mathfrak{N}((A_n)) = \mathfrak{N}((T_n)) = \mathfrak{N}((\check{T}_n)).$$

*Case 4:*  $d = 1, m \geq 1$ , and the sequence  $(A_n, n \geq 1) = (T_n, n \geq 1)$  does not converge. With the sequence  $(T_n, n \geq 1)$  we associate the set  $\mathfrak{A}$  of sequences of pairs of positive integers such that

$$\mathfrak{A} = \left\{ ((n_j, k_j), j \geq 1) \subset \mathfrak{N} \times N^N: \inf_{j \geq 1} \|T_{n_j+k_j} - T_{n_j}\| > 0 \right\}.$$

Now we introduce the set of limit points of normalized differences between the consecutive elements of the sequence  $(T_n, n \geq 1)$ ; namely let  $\Delta_j = (T_{n_j+k_j} - T_{n_j}) / \|T_{n_j+k_j} - T_{n_j}\|$ ,  $j \geq 1$ , and

$$Q = \bigcup_{(n_j, k_j) \in \mathfrak{A}} \left\{ x \in S_1(0) : x = \lim_{j \rightarrow \infty} \Delta_j \right\},$$

where  $S_1(0) = \{x \in \mathbf{R}^m : \|x\| = 1\}$ . We denote by  $W$  the linear closure of the set  $Q$  and let  $W^\perp$  be the orthogonal complement of  $W$ . Denote by  $\text{Pr}_{W^\perp}$  the orthogonal projection to the subspace  $W^\perp$ . Since the sequence  $(T_n, n \geq 1)$  does not converge,

$$\overline{\lim}_{n \rightarrow \infty} \sup_{k \geq 1} \|T_{n+k} - T_n\| > 0.$$

According to [8, Lemma 3.2.2], the sequence  $(T'_n, n \geq 1) = (\text{Pr}_{W^\perp} T_n, n \geq 1)$  converges. Thus we put

$$\mathfrak{N}((A_n)) = \mathfrak{N}((T_n)) = \mathfrak{N}((T'_n)),$$

where  $\mathfrak{N}((T'_n))$  is defined according to Case 3.

*Case 5:*  $d \geq 1$ ,  $m \geq 1$ , and the sequence  $(A_n, n \geq 1)$  is arbitrary. Let  $A_n^{(k)}$  be the row  $k$  in the matrix  $A_n$ ,  $k = 1, 2, \dots, d$ . Consider the sequences of  $m$ -dimensional vectors  $A^{(k)} = (A_n^{(k)}, n \geq 1)$ ,  $k = 1, \dots, d$ . For each of these sequences, the corresponding class  $\mathfrak{N}(A^{(k)})$  is defined according to the cases considered above. Then we put

$$\mathfrak{N}((A_n)) = \bigcup_{k=1}^d \mathfrak{N}(A^{(k)}).$$

Therefore the test class  $\mathfrak{N}((A_n))$  is constructed for an arbitrary sequence of  $d \times m$  matrices  $(A_n, n \geq 1)$ .

**The convergence to zero of sequences of random vectors normalized by operators.** Now we consider the almost sure convergence to zero of arbitrary sequences of random vectors normalized by operators. Let  $(S_n, n \geq 0)$ ,  $S_0 = 0$ , be an arbitrary sequence of random vectors in  $\mathbf{R}^m$  defined on a probability space  $(\Omega, \mathcal{F}, \mathbf{P})$ .

Below we make use of the following condition: for all  $i \geq 1$ ,

$$(7) \quad \|A_n (S_i - S_{i-1})\| \xrightarrow{\mathbf{P}} 0 \quad (n \rightarrow \infty).$$

Condition (7) holds, in particular, if  $\|A_n\| \rightarrow 0$  as  $n \rightarrow \infty$ .

**Theorem 2.1.** *Let  $\mathfrak{N}((A_n))$  be the test class for a sequence  $(A_n, n \geq 1)$ . If condition (7) holds and*

$$(8) \quad \max_{n_j < n \leq n_{j+1}} \|A_{n_{j+1}} (S_n - S_{n_j})\| \rightarrow 0 \quad a.s. \quad (j \rightarrow \infty)$$

for every sequence  $(n_j, j \geq 1) \in \mathfrak{N}((A_n))$ , then

$$(9) \quad \|A_n S_n\| \rightarrow 0 \quad a.s. \quad (n \rightarrow \infty).$$

The proof of Theorem 2.1 is given after two corollaries useful for several applications in this paper.

**Theorem 2.2.** *Let  $\mathfrak{N}((A_n))$  be the test class for a sequence  $(A_n, n \geq 1)$ . If condition (7) holds and*

$$(10) \quad \sum_{j=1}^{\infty} \mathbf{P} \left( \max_{n_j < n \leq n_{j+1}} \|A_{n_{j+1}} (S_n - S_{n_j})\| > \varepsilon \right) < \infty$$

for all sequences  $(n_j, j \geq 1) \in \mathfrak{N}((A_n))$  and all  $\varepsilon > 0$ , then relation (9) holds.

Theorem 2.2 follows from Theorem 2.1 and the Borel–Cantelli lemma.

The construction of the class  $\mathfrak{N}((A_n))$  is complicated and the assumptions of Theorem 2.2 are not easy to check in the general case. Thus we provide another corollary of Theorem 2.2, useful for several applications.

**Theorem 2.3.** *Let condition (7) hold. If condition (10) holds for every sequence*

$$(n_j, j \geq 1) \in \mathfrak{N}$$

and all  $\varepsilon > 0$ , then relation (9) holds.

In proving Theorem 2.1, we follow the idea of [5] and use the induction on the dimension of the space  $\mathbf{R}^m$ . Since the proof is rather complicated, we split it into three steps.

**Lemma 2.1.** *Let a sequence of vectors  $(T_n, n \geq 1) \subset \mathbf{R}^m$ ,  $m \geq 1$ , satisfy*

$$(11) \quad \|T_n\| \rightarrow 0 \quad (n \rightarrow \infty)$$

and let  $\mathfrak{N}((T_n))$  be the test class for this sequence. If

$$(12) \quad \max_{n_j < n \leq n_{j+1}} |\langle T_{n_{j+1}}, S_n - S_{n_j} \rangle| \rightarrow 0 \quad a.s. \quad (j \rightarrow \infty)$$

for all sequences  $(n_j, j \geq 1) \in \mathfrak{N}((T_n))$ , then

$$\langle T_n, S_n \rangle \rightarrow 0 \quad a.s. \quad (n \rightarrow \infty).$$

*Proof.* Our further argument applies to an arbitrary elementary random event  $\omega \in \Omega$ , where condition (12) holds for all  $(n_j, j \geq 1) \in \mathfrak{N}((T_n))$ . Thus we do not write “almost surely”. Now we start the induction on the dimension of the space  $\mathbf{R}^m$ .

First we prove the lemma for  $m = 1$ . Let  $(T_n, n \geq 1) = (a_n, n \geq 1)$  be a sequence of real numbers such that  $a_n \rightarrow 0$  as  $n \rightarrow \infty$ . Then the test class is defined according to (1) and (2), (3). The case of (3) is trivial; thus consider the case of (2). Put

$$Z_j = \max_{n_j < n \leq n_{j+1}} |a_{n_{j+1}} (S_n - S_{n_j})|, \quad j \geq 0,$$

where  $n_0 = 0$  and  $a_0 = 1$ . By the assumption of the lemma

$$(13) \quad Z_j \rightarrow 0 \quad (j \rightarrow \infty).$$

If  $n_j < n \leq n_{j+1}$ , then

$$|a_n S_n| \leq \lambda^{1-j} \sum_{k=0}^j \lambda^k Z_k.$$

Now the Toeplitz lemma and relation (13) imply that  $a_n S_n \rightarrow 0$  as  $n \rightarrow \infty$ . This completes the proof of Lemma 2.1 in the case  $m = 1$ .

Next we assume that the lemma is true for some  $m = s \geq 1$  and prove it for  $m = s + 1$ .

We denote the last coordinate of the vector  $T_n \in \mathbf{R}^{s+1}$  by  $t_n$ . Relation (11) implies that  $t_n \rightarrow 0$  as  $n \rightarrow \infty$ . Without loss of generality we assume that  $t_n \neq 0$ ,  $n \geq 1$ , in Case 2b) of the construction of the test class. Fix an arbitrary number  $\lambda > 1$  and put

$$\hat{n}_j = \max \{n: |t_n| \geq \lambda^{-j}\}, \quad j \geq 1.$$

It is clear that

$$(14) \quad t_{\hat{n}_{j+1}} \langle T_n, S_n \rangle = \langle t_{\hat{n}_{j+1}} T_n - t_n T_{\hat{n}_{j+1}}, S_n \rangle + t_n (\langle T_{\hat{n}_{j+1}}, S_n - S_{\hat{n}_j} \rangle + \langle T_{\hat{n}_{j+1}}, S_{\hat{n}_{j+1}} \rangle - \langle T_{\hat{n}_{j+1}}, S_{\hat{n}_{j+1}} - S_{\hat{n}_j} \rangle).$$

The last coordinate of the vector  $t_{\hat{n}_{j+1}} T_n - t_n T_{\hat{n}_{j+1}}$  is equal to zero. We denote the vector formed by the first  $s$  coordinates of the vector  $T_n$  or  $S_n$  by  $\tilde{T}_n$  or  $\tilde{S}_n$ , respectively. Now

we introduce the sequence of vectors  $(B_n, n \geq 1)$  in  $\mathbf{R}^s$  according to (4). Then the test class  $\mathfrak{N}((T_n))$  for the sequence  $(T_n, n \geq 1)$  is defined by (6).

If  $\hat{n}_j < n \leq \hat{n}_{j+1}$ , then it follows from (14) that

$$(15) \quad \begin{aligned} |\langle T_n, S_n \rangle| &\leq \lambda \max_{\hat{n}_j < n < \hat{n}_{j+1}} \left| \langle B_n, \tilde{S}_n \rangle \right| \\ &\quad + 2\lambda \max_{\hat{n}_j < n \leq \hat{n}_{j+1}} \left| \langle T_{\hat{n}_{j+1}}, S_n - S_{\hat{n}_j} \rangle \right| + \lambda \left| \langle T_{\hat{n}_{j+1}}, S_{\hat{n}_{j+1}} \rangle \right|. \end{aligned}$$

Indeed, inequality (15) is obvious for  $n = \hat{n}_{j+1}$ . Let  $\hat{n}_j < n < \hat{n}_{j+1}$ . If  $n$  is such that  $|t_n| \geq |t_{\hat{n}_{j+1}}|$ , then  $|t_n|/|t_{\hat{n}_{j+1}}| < \lambda$  and  $\theta_n = |t_{\hat{n}_{j+1}}|$ . Thus we obtain (15) from (14). If  $n$  is such that  $|t_n| < |t_{\hat{n}_{j+1}}|$ , then we obtain from (14) that

$$\begin{aligned} |\langle T_n, S_n \rangle| &\leq \max_{\hat{n}_j < n < \hat{n}_{j+1}} \left| \langle B_n, \tilde{S}_n \rangle \right| \\ &\quad + 2 \max_{\hat{n}_j < n \leq \hat{n}_{j+1}} \left| \langle T_{\hat{n}_{j+1}}, S_n - S_{\hat{n}_j} \rangle \right| + \left| \langle T_{\hat{n}_{j+1}}, S_{\hat{n}_{j+1}} \rangle \right|. \end{aligned}$$

This implies inequality (15), since  $\lambda > 1$ .

The second term on the right hand side of (15) tends to zero as  $j \rightarrow \infty$  in view of condition (12). Now we show that also the first term tends to zero. Moreover we show that

$$(16) \quad \left\langle B_n, \tilde{S}_n \right\rangle \rightarrow 0 \quad (n \rightarrow \infty).$$

By the induction assumption, if

$$(17) \quad \max_{n_{j-1} < n \leq n_j} \left| \langle B_{n_j}, \tilde{S}_n - \tilde{S}_{n_{j-1}} \rangle \right| \rightarrow 0 \quad (j \rightarrow \infty)$$

for all sequences  $(n_j, j \geq 1) \in \mathfrak{N}((B_n))$ , then relation (16) holds. Now we prove that condition (12) implies relation (17). Fix an arbitrary sequence  $(n_j, j \geq 1) \in \mathfrak{N}((B_n))$ . Since condition (12) holds for every sequence of the set  $\mathfrak{N}_{(n_j)}$  (see Case 2 in the construction of the test class), we have

$$(18) \quad \begin{aligned} \max_{n_{j(2p-1)} < n \leq m_{2p}} \left| \langle T_{m_{2p}}, S_n - S_{n_{j(2p-1)}} \rangle \right| &\rightarrow 0 \quad (p \rightarrow \infty), \\ \max_{n_{j(2p)} < n \leq m_{2p+1}} \left| \langle T_{m_{2p+1}}, S_n - S_{n_{j(2p)}} \rangle \right| &\rightarrow 0 \quad (p \rightarrow \infty), \\ L_j = \max_{n_{j-1} < n \leq n_j} \left| \langle T_{n_j}, S_n - S_{n_{j-1}} \rangle \right| &\rightarrow 0 \quad (j \rightarrow \infty). \end{aligned}$$

Here and in what follows we use the notation introduced in Case 2 of the construction of the test class. Since  $n_{j(2p)} < m_{2p}$  and  $n_{j(2p+1)} < m_{2p+1}$ ,

$$\begin{aligned} \max_{n_{j(2p-1)} < n \leq n_{j(2p)}} \left| \langle T_{m_{2p}}, S_n - S_{n_{j(2p-1)}} \rangle \right| &\rightarrow 0 \quad (p \rightarrow \infty), \\ \max_{n_{j(2p)} < n \leq n_{j(2p+1)}} \left| \langle T_{m_{2p+1}}, S_n - S_{n_{j(2p)}} \rangle \right| &\rightarrow 0 \quad (p \rightarrow \infty). \end{aligned}$$

Combining these two relations, we obtain

$$(19) \quad \max_{n_{j(i)} < n \leq n_{j(i+1)}} \left| \langle T_{m_{i+1}}, S_n - S_{n_{j(i)}} \rangle \right| \rightarrow 0 \quad (i \rightarrow \infty).$$

Note that  $N = \bigcup_{i=1}^{\infty} [\bar{j}(i), j(i)]$ . Furthermore, let  $j \in [\bar{j}(i+1), j(i+1)]$ . Then

$$n_j \in I_{i+1} = [\hat{n}_p, \hat{n}_{p+1})$$

and

$$\left\langle B_{n_j}, \tilde{S}_n - \tilde{S}_{n_{j-1}} \right\rangle = \theta_{n_j} t_{n_j}^{-1} \langle T_{n_j}, S_n - S_{n_{j-1}} \rangle - \theta_{n_j} t_{\hat{n}_{p+1}}^{-1} \langle T_{\hat{n}_{p+1}}, S_n - S_{n_{j-1}} \rangle$$

for all  $n$  and  $j$ . Indeed, after simple algebra in the right hand side of the latter equality we get

$$\begin{aligned} \theta_{n_j} \left\langle t_{n_j}^{-1} T_{n_j} - t_{\hat{n}_{p+1}}^{-1} T_{\hat{n}_{p+1}}, S_n - S_{n_{j-1}} \right\rangle &= \theta_{n_j} \left\langle t_{n_j}^{-1} \tilde{T}_{n_j} - t_{\hat{n}_{p+1}}^{-1} \tilde{T}_{\hat{n}_{p+1}}, \tilde{S}_n - \tilde{S}_{n_{j-1}} \right\rangle \\ &= \left\langle \theta_{n_j} \left( t_{n_j}^{-1} \tilde{T}_{n_j} - t_{\hat{n}_{p+1}}^{-1} \tilde{T}_{\hat{n}_{p+1}} \right), \tilde{S}_n - \tilde{S}_{n_{j-1}} \right\rangle = \left\langle B_{n_j}, \tilde{S}_n - \tilde{S}_{n_{j-1}} \right\rangle. \end{aligned}$$

Since  $\hat{n}_{p+1} = m_{i+1}$ ,

$$\begin{aligned} \max_{n_{j-1} < n \leq n_j} \left| \left\langle B_{n_j}, \tilde{S}_n - \tilde{S}_{n_{j-1}} \right\rangle \right| &\leq \max_{n_{j-1} < n \leq n_j} \left| \left\langle T_{n_j}, S_n - S_{n_{j-1}} \right\rangle \right| \\ (20) \qquad \qquad \qquad &+ \max_{n_{j-1} < n \leq n_j} \left| \left\langle T_{m_{i+1}}, S_n - S_{n_{j-1}} \right\rangle \right| \\ &= L_j + Q_j, \end{aligned}$$

where

$$Q_j = \max_{n_{j-1} < n \leq n_j} \left| \left\langle T_{m_{i+1}}, S_n - S_{n_{j-1}} \right\rangle \right|.$$

Since  $n_{j(i)} \leq n_{j-1} \leq n_j \leq n_{j(i+1)}$ , we have

$$\begin{aligned} Q_j &\leq \max_{n_{j-1} < n \leq n_j} \left| \left\langle T_{m_{i+1}}, S_n - S_{n_{j(i)}} \right\rangle \right| + \left| \left\langle T_{m_{i+1}}, S_{n_{j-1}} - S_{n_{j(i)}} \right\rangle \right| \\ &\leq 2 \max_{n_{j-1} < n \leq n_j} \left| \left\langle T_{m_{i+1}}, S_n - S_{n_{j(i)}} \right\rangle \right| \\ &\leq 2 \max_{n_{j(i)} < n \leq n_{j(i+1)}} \left| \left\langle T_{m_{i+1}}, S_n - S_{n_{j(i)}} \right\rangle \right|, \quad j \in [\bar{j}(i+1), j(i+1)]. \end{aligned}$$

This together with relation (19) implies that  $\max_{\bar{j}(i+1) \leq j \leq j(i+1)} Q_j \rightarrow 0$  as  $i \rightarrow \infty$ . Hence  $Q_j \rightarrow 0$  as  $j \rightarrow \infty$ . Using the latter result we deduce (17) from (18) and (20).

To complete the proof of the lemma, it remains to show that the third term in (15) approaches zero, that is,

$$(21) \qquad \qquad \qquad V_j = \left| \left\langle T_{\hat{n}_j}, S_{\hat{n}_j} \right\rangle \right| \rightarrow 0 \quad (j \rightarrow \infty).$$

Note that if  $\hat{n}_j = \hat{n}_{j+1} = \dots = \hat{n}_{j+l}$ , then  $V_j = V_{j+1} = \dots = V_{j+l}$  by construction. This allows one to assume (without any loss of generality) that the sequence  $(\hat{n}_j, j \geq 1)$  is increasing.

We have

$$\begin{aligned} t_{\hat{n}_j} \left\langle T_{\hat{n}_{j+1}}, S_{\hat{n}_{j+1}} \right\rangle &= t_{\hat{n}_j} \left\langle T_{\hat{n}_{j+1}}, S_{\hat{n}_{j+1}} - S_{\hat{n}_j} \right\rangle + t_{\hat{n}_{j+1}} \left\langle T_{\hat{n}_j}, S_{\hat{n}_j} \right\rangle \\ &\quad - \left\langle t_{\hat{n}_{j+1}} T_{\hat{n}_j} - t_{\hat{n}_j} T_{\hat{n}_{j+1}}, S_{\hat{n}_j} \right\rangle. \end{aligned}$$

Thus

$$(22) \qquad \qquad \qquad V_{j+1} \leq \left| t_{\hat{n}_{j+1}} t_{\hat{n}_j}^{-1} \right| \cdot V_j + \delta_j, \quad j \geq 1,$$

where

$$\delta_j = \left| \left\langle T_{\hat{n}_{j+1}}, S_{\hat{n}_{j+1}} - S_{\hat{n}_j} \right\rangle \right| + \left| \left\langle B_{\hat{n}_j}, \tilde{S}_{\hat{n}_j} \right\rangle \right|.$$

It follows from (12) and (16) that

$$(23) \qquad \qquad \qquad \delta_j \rightarrow 0 \quad (j \rightarrow \infty).$$

In turn, inequality (22) implies that

$$(24) \qquad \qquad \qquad V_{j+2} \leq \left| t_{\hat{n}_{j+2}} t_{\hat{n}_j}^{-1} \right| \cdot V_{j+1} + \left| t_{\hat{n}_{j+2}} t_{\hat{n}_{j+1}}^{-1} \right| \cdot \delta_j + \delta_{j+1}, \quad j \geq 1.$$

Since  $|t_{\hat{n}_{j+2}} t_{\hat{n}_j}^{-1}| \leq \lambda^{-1} < 1$  and  $|t_{\hat{n}_{j+2}} t_{\hat{n}_{j+1}}^{-1}| \leq 1$ , we obtain relation (21) from (23) and (24) by using the Toeplitz lemma. Lemma 2.1 is proved.  $\square$

Now we replace (11) with a weaker assumption in the following result.

**Lemma 2.2.** *Let a sequence of vectors  $(T_n, n \geq 1)$  in  $\mathbf{R}^m$  be such that*

$$(25) \quad \langle T_n, S_i - S_{i-1} \rangle \xrightarrow{P} 0 \quad (n \rightarrow \infty)$$

for all  $i \geq 1$ . Let  $\mathfrak{N}((T_n))$  be the test class for this sequence. If condition (12) holds for all sequences  $(n_j, j \geq 1) \in \mathfrak{N}((T_n))$ , then

$$\langle T_n, S_n \rangle \rightarrow 0 \quad \text{a.s.} \quad (n \rightarrow \infty).$$

*Proof.* First we assume that the sequence  $(T_n, n \geq 1)$  is such that

$$(26) \quad \langle T_n, S_i - S_{i-1} \rangle \rightarrow 0 \quad \text{a.s.} \quad (n \rightarrow \infty)$$

for all  $i \geq 1$ . We distinguish between the two cases in the proof, namely a) the sequence  $(T_n, n \geq 1)$  converges; and b) the sequence  $(T_n, n \geq 1)$  does not converge.

a) Suppose the sequence  $(T_n, n \geq 1)$  converges. Put

$$T = \lim_{n \rightarrow \infty} T_n, \quad \check{T}_n = T_n - T, \quad n \geq 1.$$

Then  $\mathfrak{N}((T_n))$  is defined according to Case 3 of the construction of the test class. Condition (26) implies that  $\langle T, S_i - S_{i-1} \rangle = 0$  almost surely for all  $i \geq 1$ . This means that

$$\max_{n_j < n \leq n_{j+1}} |\langle T_{n_{j+1}}, S_n - S_{n_j} \rangle| = \max_{n_j < n \leq n_{j+1}} \left| \langle \check{T}_{n_{j+1}}, S_n - S_{n_j} \rangle \right| \quad \text{a.s.}$$

for any sequence  $(n_j, j \geq 1) \in \mathfrak{N}$ , and  $\langle T_n, S_n \rangle = \langle \check{T}_n, S_n \rangle$  almost surely for all  $n \geq 1$ . Now the lemma follows from Lemma 2.1.

b) Suppose the sequence  $(T_n, n \geq 1)$  does not converge. Then  $\mathfrak{N}((T_n))$  is defined according to Case 4 of the construction of the test class. Recall that the terms of the sequence  $(T'_n, n \geq 1)$  satisfy  $T'_n = \text{Pr}_{W^\perp} T_n$ ,  $n \geq 1$ . Condition (26) and [8, Lemma 3.2.3] imply that

$$\langle T_n, S_i - S_{i-1} \rangle = \langle T'_n, S_i - S_{i-1} \rangle \quad \text{a.s.}$$

for all  $n$  and  $i \geq 1$ , and

$$\langle T'_n, S_i - S_{i-1} \rangle \rightarrow 0 \quad \text{a.s.} \quad (n \rightarrow \infty)$$

for all  $i \geq 1$ . Since

$$\max_{n_j < n \leq n_{j+1}} |\langle T_{n_{j+1}}, S_n - S_{n_j} \rangle| = \max_{n_j < n \leq n_{j+1}} \left| \langle T'_{n_{j+1}}, S_n - S_{n_j} \rangle \right| \quad \text{a.s.}$$

for any sequence  $(n_j, j \geq 1) \in \mathfrak{N}$ , and  $\langle T_n, S_n \rangle = \langle T'_n, S_n \rangle$  almost surely for all  $n \geq 1$ , the general case of the lemma follows from case a).

One can reduce condition (26) to condition (25) in view of [8, Lemma 3.2.5]. Lemma 2.2 is proved.  $\square$

*Proof of Theorem 2.1.* Let  $(e_1, \dots, e_d)$  be an orthonormal basis in  $\mathbf{R}^d$ , and let  $A_n^*$  be the conjugate operator to  $A_n$ . Since

$$A_n S_n = \sum_{k=1}^d \langle A_n S_n, e_k \rangle e_k = \sum_{k=1}^d \langle A_n^* e_k, S_n \rangle e_k,$$

relation (9) holds if and only if

$$(27) \quad \left\langle A_n^{(k)}, S_n \right\rangle \rightarrow 0 \quad \text{a.s.} \quad (n \rightarrow \infty)$$

for all  $k = 1, \dots, d$ , where  $A_n^{(k)} = A_n^* e_k$ . The set  $\mathfrak{N}((A_n))$  is defined according to Case 5 of the construction of the test class. It follows from (7) that

$$(28) \quad \left\langle A_n^{(k)}, S_i - S_{i-1} \right\rangle \xrightarrow{P} 0 \quad (n \rightarrow \infty)$$



for all  $k = 1, \dots, d$  and for all  $i \geq 1$ . Furthermore, condition (8) implies that

$$(29) \quad \max_{n_j < n \leq n_{j+1}} \left| \left\langle A_{n_{j+1}}^{(k)}, S_n - S_{n_j} \right\rangle \right| \rightarrow 0 \quad \text{a.s.} \quad (j \rightarrow \infty)$$

for all  $k = 1, \dots, d$  and all sequences  $(n_j, j \geq 1) \in \mathfrak{N}(A^{(k)})$ . In view of (28) and (29) Lemma 2.2 implies relation (27), whence (9) follows. Theorem 2.1 is proved.  $\square$

If  $(S_n, n \geq 1)$  is a sequence of partial sums of independent symmetric random vectors, then conditions (7) and (10) are necessary for relation (9). This follows from Theorem 3.2.3 in [8]. Indeed, condition (7) is obvious in view of Theorem 3.2.3 in [8]. Since the series in (10) does not exceed the series

$$2 \sum_{j=1}^{\infty} \mathbb{P} \left( \|A_{n_{j+1}}(S_{n_{j+1}} - S_{n_j})\| > \varepsilon \right)$$

by the Lévy inequality, condition (10) follows. The convergence of the latter series is implied by the Borel–Cantelli lemma, by the relation  $\|A_{n_{j+1}}(S_{n_{j+1}} - S_{n_j})\| \rightarrow 0$  a.s. as  $j \rightarrow \infty$ , and by the independence of the vectors  $(S_{n_{j+1}} - S_{n_j}, j \geq 1)$ .

**The boundedness of sequences of random vectors normalized by operators.** Consider conditions for the almost sure boundedness of arbitrary sequences of random vectors normalized by operators. The following result is proved similarly to Theorem 2.1.

**Theorem 2.4.** *Let  $\mathfrak{N}((A_n))$  be the test class for a sequence  $(A_n, n \geq 1)$ . If condition (7) holds and*

$$\sup_{j \geq 1} \max_{n_j < n \leq n_{j+1}} \|A_{n_{j+1}}(S_n - S_{n_j})\| < \infty \quad \text{a.s.}$$

for all sequences  $(n_j, j \geq 1) \in \mathfrak{N}((A_n))$ , then

$$(30) \quad \sup_{n \geq 1} \|A_n S_n\| < \infty \quad \text{a.s.}$$

Theorem 2.4 and the Borel–Cantelli lemma imply the following result.

**Theorem 2.5.** *Let  $\mathfrak{N}((A_n))$  be the test class for a sequence  $(A_n, n \geq 1)$ . If condition (7) holds and for any sequence  $(n_j, j \geq 1) \in \mathfrak{N}((A_n))$  there exists a number  $\varepsilon > 0$  (possibly depending on the sequence  $(n_j, j \geq 1)$ ) such that condition (10) holds, that is,*

$$(31) \quad \sum_{j=1}^{\infty} \mathbb{P} \left( \max_{n_j < n \leq n_{j+1}} \|A_{n_{j+1}}(S_n - S_{n_j})\| > \varepsilon \right) < \infty,$$

then inequality (30) holds.

The following result also follows from Theorem 2.5.

**Theorem 2.6.** *Let condition (7) hold. If for any sequence  $(n_j, j \geq 1) \in \mathfrak{N}$  there exists a number  $\varepsilon > 0$  (possibly depending on  $(n_j, j \geq 1)$ ) such that condition (31) holds, then inequality (30) holds.*

The following result shows that condition (7) can be weakened.

**Theorem 2.7.** *Let*

$$\sup_{n \geq 1} \|A_n\| < \infty.$$

for every sequence  $(n_j, j \geq 1) \in \mathfrak{N}$  there exists a number  $\varepsilon > 0$  (possibly depending on the sequence  $(n_j, j \geq 1)$ ) such that condition (31) holds, then inequality (30) holds.

The proof of Theorem 2.7 follows from Theorem 2.1 and Lemma 3 in [9].

If  $(S_n, n \geq 1)$  is the sequence of partial sums of independent symmetric random vectors, then condition (31) for all sequences  $(n_j, j \geq 1) \in \mathfrak{N}$  and some number  $\varepsilon > 0$  (possibly depending on the sequence  $(n_j, j \geq 1)$ ) is necessary for inequality (30). This follows from [10, Theorem 3.4.1] and Lévy's inequality.

### 3. STRONG LAW OF LARGE NUMBERS FOR SUMS OF ORTHOGONAL RANDOM VECTORS NORMALIZED BY OPERATORS

A sequence of random vectors  $(X_i, i \geq 1)$  in  $\mathbf{R}^m$  is called orthogonal if  $\mathbf{E} \|X_i\|^2 < \infty$ ,  $i \geq 1$ , and  $\mathbf{E} (\langle T_1, X_i \rangle \langle T_2, X_j \rangle) = 0$  for all vectors  $T_1, T_2 \in \mathbf{R}^m$  and all  $i \neq j$ . Put  $S_n = \sum_{i=1}^n X_i$ ,  $n \geq 1$ , and  $\log_+ t = \ln \max \{t, e\}$ ,  $t \geq 0$ .

**Theorem 3.1.** *Let  $\mathfrak{N}((A_n))$  be the test class for a sequence  $(A_n, n \geq 1)$  and let*

$$(X_i, i \geq 1)$$

*be an orthogonal sequence in  $\mathbf{R}^m$ . If*

$$(32) \quad \|A_n X_i\| \xrightarrow{P} 0 \quad (n \rightarrow \infty)$$

*for all  $i \geq 1$ , and*

$$(33) \quad \sum_{j=1}^{\infty} \sum_{i=n_j+1}^{n_{j+1}} \mathbf{E} \|A_{n_{j+1}} X_i\|^2 \log_+^2 (n_{j+1} - n_j) < \infty$$

*for all sequences  $(n_j, j \geq 1) \in \mathfrak{N}((A_n))$ , then*

$$(34) \quad \|A_n S_n\| \rightarrow 0 \quad a.s. \quad (n \rightarrow \infty).$$

*Proof.* We check assumptions of Theorem 2.2. Condition (7) holds in view of relation (32). Using Chebyshev's and Rademacher–Men'shov's inequalities we obtain

$$\begin{aligned} \mathbf{P} \left( \max_{n_j < n \leq n_{j+1}} \|A_{n_{j+1}} (S_n - S_{n_j})\| > \varepsilon \right) &\leq \varepsilon^{-2} \mathbf{E} \left( \max_{n_j < n \leq n_{j+1}} \left\| \sum_{i=n_j+1}^n A_{n_{j+1}} X_i \right\|^2 \right) \\ &\leq (\varepsilon \ln 2)^{-2} \ln^2 [4(n_{j+1} - n_j)] \sum_{i=n_j+1}^{n_{j+1}} \mathbf{E} \|A_{n_{j+1}} X_i\|^2 \end{aligned}$$

for all  $\varepsilon > 0$ .

Thus condition (33) implies (10), and this completes the proof of Theorem 3.1.  $\square$

Theorem 3.1 yields the following result.

**Corollary 3.1.** *Let  $(X_i, i \geq 1)$  be an orthogonal sequence in  $\mathbf{R}^m$ . Assume that condition (32) holds. If*

$$(35) \quad \sum_{i=1}^{\infty} \sup_{n \geq i} \left( \mathbf{E} \|A_n X_i\|^2 \ln^2 n \right) < \infty,$$

*then relation (34) holds.*

If  $A_n = a_n I$ ,  $n \geq 1$ , in Corollary 3.1, where  $I$  is the identity operator, and if the sequence of numbers  $(a_n \ln n, n \geq n_0 > 1)$  is decreasing to zero, then condition (35) becomes the classical Rademacher–Men'shov condition

$$\sum_{i=2}^{\infty} \mathbf{E} \|X_i\|^2 a_i^2 \ln^2 i < \infty$$

(see, for example, [2]).

4. CONVERGENCE TO ZERO AND BOUNDEDNESS OF MARTINGALES  
NORMALIZED BY OPERATORS

Given a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$  with filtration  $(\mathcal{F}_n, n \geq 0)$  let  $(S_n, n \geq 0)$  be a martingale with respect to  $(\mathcal{F}_n, n \geq 0)$ . Put  $S_0 = 0$  and denote by  $(X_n, n \geq 1)$  the corresponding martingale difference, that is,

$$X_n = S_n - S_{n-1}, \quad n \geq 1.$$

In what follows we use the following form of condition (7): for all  $i \geq 1$ ,

$$(36) \quad \|A_n X_i\| \xrightarrow{\mathbb{P}} 0 \quad (n \rightarrow \infty).$$

For the sake of brevity we use the notation

$$(37) \quad U_{n_j} = A_{n_{j+1}} (S_{n_{j+1}} - S_{n_j}), \quad j \geq 1, (n_j, j \geq 1) \in \mathfrak{N}.$$

The indicator of a random event is denoted by  $I(\cdot)$ .

**Theorem 4.1.** *Let  $\mathfrak{N}((A_n))$  be the test class for a sequence  $(A_n, n \geq 1)$ . Assume that condition (36) holds. If*

$$(38) \quad \sum_{j=1}^{\infty} \mathbb{E} [\|U_{n_j}\| I(\|U_{n_j}\| > \varepsilon)] < \infty$$

for all sequences  $(n_j, j \geq 1) \in \mathfrak{N}((A_n))$  and all numbers  $\varepsilon > 0$ , or, equivalently, if

$$(39) \quad \sum_{j=1}^{\infty} \int_{\varepsilon}^{\infty} \mathbb{P} (\|U_{n_j}\| > t) dt < \infty$$

for all sequences  $(n_j, j \geq 1) \in \mathfrak{N}((A_n))$ , then

$$(40) \quad \|A_n S_n\| \rightarrow 0 \quad a.s. \quad (n \rightarrow \infty).$$

*Proof.* The proof of the theorem is based on Theorem 2.2. Condition (7) is equivalent to condition (36) in this case. Now we show that condition (10) follows from (38). Note that

$$(A_{n_{j+1}} (S_n - S_{n_j}), \mathcal{F}_n, n_j \leq n \leq n_{j+1})$$

is a martingale with zero mean for any sequence  $(n_j, j \geq 1) \in \mathfrak{N}$ . Using the Brown inequality [11], we get

$$\begin{aligned} & \sum_{j=1}^{\infty} \mathbb{P} \left( \max_{n_j < n \leq n_{j+1}} \|A_{n_{j+1}} (S_n - S_{n_j})\| > \varepsilon \right) \\ & \leq \frac{2d^{3/2}}{\varepsilon} \sum_{j=1}^{\infty} \mathbb{E} \left[ \|U_{n_j}\| I \left( \|U_{n_j}\| > \varepsilon/2d^{1/2} \right) \right] < \infty. \end{aligned}$$

Thus Theorem 2.2 implies relation (40). The equivalence of conditions (38) and (39) is proved by integrating by parts. Theorem 4.1 is proved.  $\square$

*Remark 4.1.* Condition (39) is equivalent to the following two conditions:

- (i) for all sequences  $(n_j, j \geq 1) \in \mathfrak{N}((A_n))$  and all numbers  $\varepsilon > 0$ ,

$$\sum_{j=1}^{\infty} \mathbb{P} (\|U_{n_j}\| > \varepsilon) < \infty;$$

- (ii) for every sequence  $(n_j, j \geq 1) \in \mathfrak{N}((A_n))$  there exists a number  $\varepsilon_0 > 0$  (possibly depending on the sequence  $(n_j, j \geq 1)$ ) such that

$$\sum_{j=1}^{\infty} \int_{\varepsilon_0}^{\infty} \mathbf{P}(\|U_{n_j}\| > t) dt < \infty.$$

Now we discuss whether the assumptions of Theorem 4.1 can be weakened in some sense.

- 1) If there exists a constant  $c > 0$  such that  $\|A_n S_n\| \leq c$  almost surely for all  $n \geq 1$ , then condition (i) is sufficient for relation (40). This obviously follows from (38), since

$$\begin{aligned} \|U_{n_j}\| &\leq \|A_{n_{j+1}} S_{n_{j+1}}\| + \|A_{n_{j+1}} S_{n_j}\| \leq c + \|\mathbf{E}(A_{n_{j+1}} S_{n_{j+1}} \mid \mathcal{F}_{n_j})\| \\ &\leq c + \mathbf{E}(\|A_{n_{j+1}} S_{n_{j+1}}\| \mid \mathcal{F}_{n_j}) \leq 2c \quad \text{a.s.} \end{aligned}$$

in view of the martingale property.

- 2) Let  $(X_i, i \geq 1)$  be a sequence of independent centered ( $\mathbf{E} X_i = 0, i \geq 1$ ) random vectors. Consider the martingale

$$S_n = \sum_{i=1}^n X_i, \quad n \geq 1.$$

It follows from [12] that if (36) holds, then (i) and

$$(41) \quad \|A_n S_n\| \xrightarrow{\mathbf{P}} 0 \quad (n \rightarrow \infty)$$

imply relation (40). This form of the strong law of large numbers originates from the paper [13] for the one-dimensional case (see also [14, p. 160]). Condition (39) implies (i) and (41) in view of Remark 4.1 and Theorem 4.1. An example is constructed in [6] demonstrating that conditions (i) and (41) are not sufficient for (40) in the case of general martingales. This also means that condition (39) cannot be weakened in general.

**The boundedness of martingales normalized by operators.** Consider conditions for the boundedness of martingales normalized by operators.

The next result follows from Theorem 2.5 and Brown's inequality [11].

**Theorem 4.2.** *Let  $\mathfrak{N}((A_n))$  be the test class for a sequence  $(A_n, n \geq 1)$ . Assume that condition (36) holds. If for any sequence  $(n_j, j \geq 1) \in \mathfrak{N}((A_n))$  there exists a number  $\varepsilon > 0$  (possibly depending on the sequence  $(n_j, j \geq 1)$ ) such that*

$$\sum_{j=1}^{\infty} \mathbf{E} [\|U_{n_j}\| I(\|U_{n_j}\| > \varepsilon)] < \infty,$$

then

$$(42) \quad \sup_{n \geq 1} \|A_n S_n\| < \infty \quad \text{a.s.}$$

The following result demonstrates that condition (36) can be weakened.

**Theorem 4.3.** *Let*

$$\sup_{n \geq 1} \|A_n\| < \infty.$$

*If for any sequence  $(n_j, j \geq 1) \in \mathfrak{N}$  there exists a number  $\varepsilon > 0$  (possibly depending on the sequence  $(n_j, j \geq 1)$ ) such that*

$$\sum_{j=1}^{\infty} \int_{\varepsilon}^{\infty} \mathbf{P}(\|U_{n_j}\| > t) dt < \infty,$$

then relation (42) holds.

The proof of Theorem 4.3 follows from Theorem 4.1 and Lemma 3 in [9].

Comparing Theorem 4.3 with Remark 4.1, we see that condition (ii) (which is true for the set  $\mathfrak{N}$ ) implies relation (42), that is, (ii) implies the almost sure boundedness.

5. STRONG LAW OF LARGE NUMBERS FOR MARTINGALES  
THAT ARE NORMALIZED BY OPERATORS AND HAVE HIGHER MOMENTS

In this section, we consider a number of corollaries of Theorem 4.1 for martingales that have a finite absolute moment of order  $\nu \geq 1$ . A comprehensive study of the strong law of large numbers for martingales normalized by sequences of real numbers can be found in [14]–[17].

**Theorem 5.1.** *Let  $\mathfrak{N}((A_n))$  be the test class for a sequence  $(A_n, n \geq 1)$ . Assume that condition (36) holds. If*

$$(43) \quad \sum_{j=1}^{\infty} \mathbf{E} \|U_{n_j}\|^\nu < \infty$$

for some  $\nu \geq 1$  and all sequences  $(n_j, j \geq 1) \in \mathfrak{N}((A_n))$ , then

$$(44) \quad \|A_n S_n\| \rightarrow 0 \quad a.s. \quad (n \rightarrow \infty).$$

*Proof.* The case of  $\nu = 1$  is obvious. Consider the case of  $\nu > 1$ . Using the Chebyshev inequality we get

$$\sum_{j=1}^{\infty} \int_{\varepsilon}^{\infty} \mathbf{P} (\|U_{n_j}\| > t) dt \leq \sum_{j=1}^{\infty} \mathbf{E} \|U_{n_j}\|^\nu \int_{\varepsilon}^{\infty} \frac{dt}{t^\nu} = \frac{\varepsilon^{1-\nu}}{\nu-1} \sum_{j=1}^{\infty} \mathbf{E} \|U_{n_j}\|^\nu < \infty.$$

Thus Theorem 4.1 implies relation (44). The theorem is proved. □

**Theorem 5.2.** *Let  $\mathfrak{N}((A_n))$  be the test class for a sequence  $(A_n, n \geq 1)$ . Assume that condition (36) holds. If*

$$\sum_{j=1}^{\infty} \sum_{i=n_j+1}^{n_{j+1}} \mathbf{E} \|A_{n_{j+1}} X_i\|^\nu < \infty$$

for some number  $\nu \in [1, 2]$  and all sequences  $(n_j, j \geq 1) \in \mathfrak{N}((A_n))$ , then relation (44) holds.

The proof of Theorem 5.2 follows from Theorem 5.1 and the inequality (see [18])

$$\mathbf{E} \|U_{n_j}\|^\nu = \mathbf{E} \left\| \sum_{i=n_j+1}^{n_{j+1}} A_{n_{j+1}} X_i \right\|^\nu \leq 2^{2-\nu} \sum_{i=n_j+1}^{n_{j+1}} \mathbf{E} \|A_{n_{j+1}} X_i\|^\nu$$

(we use notation (37)).

*Remark 5.1.* An equivalent form of Theorem 5.2 is obtained in [4] for sequences of independent centered random variables  $(X_n, n \geq 1)$ .

**Corollary 5.1.** *If condition (36) holds and*

$$\sum_{i=1}^{\infty} \sup_{n \geq i} \mathbf{E} \|A_n X_i\|^\nu < \infty$$

for some number  $\nu \in [1, 2]$ , then relation (44) holds.

**Definition 5.1.** A sequence of operators  $(A_n, n \geq 1)$  is called *monotone* if

$$(45) \quad \|A_n x\| \geq \|A_{n+1} x\|$$

for all  $x \in \mathbf{R}^m$  and  $n \geq 1$ .

Note that condition (45) is equivalent to  $A_n^* A_n \geq A_{n+1}^* A_{n+1}$ ,  $n \geq 1$ , where  $A_n^*$  is the conjugate operator to  $A_n$ .

Corollary 5.1 and condition (45) imply the following result originally proved in [18, Theorem 2].

**Corollary 5.2.** *Let  $(A_n, n \geq 1)$  be a monotone sequence of operators and let*

$$\|A_n\| \rightarrow 0, \quad n \rightarrow \infty.$$

If

$$\sum_{i=1}^{\infty} \mathbb{E} \|A_i X_i\|^\nu < \infty$$

for some number  $\nu \in [1, 2]$ , then relation (44) holds.

Now we consider the case where a martingale has a finite absolute moment of order  $\nu \geq 2$ .

**Theorem 5.3.** *Let  $\mathfrak{N}((A_n))$  be the test class of a sequence  $(A_n, n \geq 1)$ . Assume that condition (36) holds. If*

$$\sum_{j=1}^{\infty} (n_{j+1} - n_j)^{(\nu/2)-1} \sum_{i=n_j+1}^{n_{j+1}} \mathbb{E} \|A_{n_{j+1}} X_i\|^\nu < \infty$$

for some number  $\nu \in [2, +\infty)$  and all sequences  $(n_j, j \geq 1) \in \mathfrak{N}((A_n))$ , then relation (44) holds.

The proof of Theorem 5.3 follows from Theorem 5.1 and the Dharmadhikari–Fabian–Jogdeo inequality [19].

Theorem 5.3 implies the following result.

**Corollary 5.3.** *Suppose condition (36) holds. If*

$$\sum_{i=1}^n \sup_{n \geq i} \left( n^{\nu/2-1} \mathbb{E} \|A_n X_i\|^\nu \right) < \infty$$

for some number  $\nu \in [2, +\infty)$ , then relation (44) holds.

Corollary 5.3 contains the following result obtained in [20].

**Corollary 5.4.** *Let  $A_n = a_n I$ ,  $n \geq 1$ , where  $I$  is the identity operator and  $(a_n, n \geq 1)$  is a sequence of positive numbers such that  $a_n \rightarrow 0$  as  $n \rightarrow \infty$  and the sequence*

$$\left( a_n n^{(\nu-2)/(2\nu)}, n \geq n_0 \geq 1 \right)$$

decreases for some  $\nu \in [2, +\infty)$ . If

$$\sum_{i=1}^{\infty} i^{\nu/2-1} a_i^\nu \mathbb{E} \|X_i\|^\nu < \infty,$$

then

$$a_n \|S_n\| \rightarrow 0 \quad \text{a.s.} \quad (n \rightarrow \infty).$$

The following result is useful for some cases.

**Corollary 5.5.** *Suppose condition (36) holds. Assume that*

$$\mathbb{E} \|A_n (S_n - S_k)\|^\nu \leq c (f_k - f_n)$$

for some number  $\nu \in [1, +\infty)$  and for all positive integers  $k \leq n$ , where  $c$  is a positive constant (that does not depend on  $n$  and  $k$ ). If  $(f_n, n \geq 1)$  is a sequence of real numbers such that  $f_n \downarrow 0$  as  $n \rightarrow \infty$ , then relation (44) holds.

The proof of Corollary 5.5 follows from Theorem 5.1, since relation (42) holds for all sequences

$$(n_j, j \geq 1) \in \mathfrak{N}.$$

Indeed,

$$\sum_{j=1}^{\infty} \mathbf{E} \|U_{n_j}\|^\nu = \sum_{j=1}^{\infty} \mathbf{E} \|A_{n_{j+1}} (S_{n_{j+1}} - S_{n_j})\|^\nu \leq c \sum_{j=1}^{\infty} (f_{n_j} - f_{n_{j+1}}) = cf_{n_1} < \infty.$$

Consider an application of Corollary 5.5 to the proof of the strong consistency of the least squares estimator of an unknown parameter for vector linear regression. This problem is studied in [21]. The most general result for this problem is obtained in [22].

Assume that elements of  $\mathbf{R}^m$  are represented as vector columns and denote by  $\|\cdot\|$  the Euclidean norm of a matrix (or a vector). Consider the model of vector linear regression

$$Y_i = D_i^T V + Z_i, \quad i \geq 1,$$

where  $(Z_i, i \geq 1)$  is a sequence of random vectors in  $\mathbf{R}^m$ ;  $(D_i, i \geq 1)$  is a sequence of nonrandom  $d \times m$  matrices;  $V$  is an unknown parameter; and  $^T$  means transposition. Assume that the matrices

$$B_n = \sum_{i=1}^n D_i D_i^T, \quad n \geq 1,$$

are nonsingular. Then the least squares estimator of the unknown parameter  $V$  can be written as

$$\hat{V}_n = B_n^{-1} \sum_{i=1}^n D_i Y_i, \quad n \geq 1,$$

where  $B_n^{-1}$  is the inverse function to  $B_n$ .

It is shown in [7] that Corollary 5.5 implies the following result on the strong consistency of the estimator  $(\hat{V}_n, n \geq 1)$  (this result is proved earlier in [22]).

**Theorem 5.4.** *Let  $(Z_i, i \geq 1)$  be a martingale difference and let  $\sup_{i \geq 1} \mathbf{E} \|Z_i\|^2 < \infty$ . If*

$$(46) \quad \|B_n^{-1}\| \rightarrow 0 \quad (n \rightarrow \infty),$$

then

$$\|\hat{V}_n - V\| \rightarrow 0 \quad a.s. \quad (n \rightarrow \infty).$$

Note that condition (46) is equivalent to

$$\lambda_{\min}(B_n) \rightarrow \infty \quad (n \rightarrow \infty),$$

where  $\lambda_{\min}(B_n)$  is the minimal eigenvalue of the matrix  $B_n$ .

#### BIBLIOGRAPHY

1. Yu. V. Prokhorov, *On the strong law of large numbers*, Izv. AN SSSR Ser. Matem. **14** (1950), no. 6, 523–536. MR0038592 (12:425c)
2. M. Loève, *Probability Theory*, Second edition, D. van Nostrand Company Inc., Princeton–Toronto–New York–London, 1960. MR0123342 (23:A670)
3. V. V. Buldygin, *The strong laws of large numbers and convergence to zero of Gaussian sequences*, Teor. Veroyatn. Mat. Stat. **19** (1978), 33–41; English transl. in Theory Probab. Math. Statist. **19** (1979), 35–44. MR0494429 (58:13294)
4. A. I. Martikainen, *On necessary and sufficient conditions for the strong law of large numbers*, Teor. Veroyatnost. i Primenen. **24** (1979), no. 4, 814–820; English transl. in Theory Probab. Appl. **24** (1980), 813–819. MR550536 (81g:60030)
5. V. V. Buldygin and S. A. Solntsev, *The strong law of large numbers for sums of independent random vectors with operator normalizations and null convergence of Gaussian sequences*, Teor. Veroyatnost. i Primenen. **32** (1987), no. 2, 266–281; English transl. in Theory Probab. Appl. **32** (1988), 243–256. MR902755 (88i:60013)

6. V. O. Koval', *Limit theorems for random vectors normalized by operators. I*, Teor. ĭmovir. Mat. Stat. **61** (1999), 47–58; English transl. in Theory Probab. Math. Statist. **61** (2000), 49–60. MR1866970 (2002j:60050)
7. V. V. Buldygin and V. A. Koval', *The strong law of large numbers for martingales and sums of orthogonal random vectors with operator normalizations*, Ukrain. Mat. Zh. **52** (2000), no. 8, 1045–1061; English transl. in Ukrainian Math. J. (2001). MR1819715 (2001m:60064)
8. V. V. Buldygin and S. A. Solntsev, *Functional Methods in Problems of Summation of Random Variables*, “Naukova Dumka”, Kiev, 1989. (Russian) MR1007589 (91i:60013)
9. V. A. Egorov, *On the law of the iterated logarithm*, Teor. Veroyatnost. i Primenen. **14** (1969), no. 4, 722–729; English transl. in Theory Probab. Appl. **14** (1970), 693–700. MR0266286 (42:1193)
10. V. Buldygin and S. Solntsev, *Asymptotic Behaviour of Linearly Transformed Sums of Random Variables*, Kluwer, Dordrecht, 1997. MR1471203 (98m:60002)
11. B. M. Brown, *Martingale central limit theorems*, Ann. Math. Statist. **42** (1971), no. 1, 59–66. MR0290428 (44:7609)
12. S. A. Solntsev, *Limit Theorems for Linear Transformations of Sums of Independent Random Vectors*, Doctoral dissertation, Kiev, 1994. (Russian)
13. K. L. Chung, *The strong law of large numbers*, Proc. Second Berkeley Symp. on Math. Statist. and Probability (July 31–August 12, 1950), University of California Press, Berkeley, 1951, pp. 341–352. MR0045328 (13:567c)
14. W. Stout, *Almost Sure Convergence*, Academic Press, New York, 1974. MR0455094 (56:13334)
15. P. Hall and C. C. Heyde, *Martingale Limit Theory and its Applications*, Academic Press, New York, 1980. MR624435 (83a:60001)
16. V. A. Egorov, *On the strong law of large numbers and law of the iterated logarithm for martingales and sums of independent random variables*, Teor. Veroyatnost. i Primenen. **35** (1990), no. 4, 691–703; English transl. in Theory Probab. Appl. **35** (1991), 653–666. MR1090495 (92g:60040)
17. H. Teicher, *Strong laws for martingale differences and independent random variables*, J. Theor. Probab. **11** (1998), no. 4, 979–996. MR1660911 (99i:60074)
18. H. Kaufmann, *On the strong law of large numbers for multivariate martingales*, Stochastic Process. Appl. **26** (1987), no. 1, 73–85. MR917247 (88m:60134)
19. S. W. Dharmadhikari, V. Fabian, and K. Jogdeo, *Bounds on the moments of martingales*, Ann. Math. Statist. **39** (1968), no. 5, 1719–1723. MR0230363 (37:5925)
20. I. Fazekas and O. Klesov, *A general approach to the strong laws of large numbers*, Teor. Veroyatnost. i Primenen. **45** (2000), no. 3, 568–583. MR1967791 (2003k:60066)
21. T. W. Anderson and J. Taylor, *Strong consistency of least squares estimates in normal linear regression*, Ann. Statist. **4** (1976), no. 4, 788–790. MR0415899 (54:3977)
22. T. L. Lai, *Some almost sure convergence properties of weighted sums of martingale difference sequences*, Almost Everywhere Convergence II, Academic Press, Boston, 1991, pp. 179–190. MR1131791 (92k:60065)

NATIONAL TECHNICAL UNIVERSITY OF UKRAINE “KPI”, DEPARTMENT OF MATHEMATICAL ANALYSIS  
AND PROBABILITY THEORY, PR. PEREMOGY 37, 03056 KYIV–56, UKRAINE  
*E-mail address:* valbuld@comsys.ntu-kpi.kiev.ua

ZHYTOMYR STATE UNIVERSITY FOR TECHNOLOGY, DEPARTMENT OF MATHEMATICS, CHERNYAKHOVSKIĬ  
STREET 103, 10005 ZHYTOMYR, UKRAINE  
*E-mail address:* vkoval@com.zt.ua

Received 23/FEB/2005

Translated by OLEG KLESOV