

ON THE REDUCIBILITY OF LINEAR GROUPS*

BY

LEONARD EUGENE DICKSON

The object of this note is a two-fold generalization of LOEWY'S theorem proved in these Transactions, vol. 4, pp. 171-177. His theorem may be conveniently stated as follows: If R is the domain of all real numbers and C the domain of all complex numbers, any group of linear homogeneous transformations with coefficients in R which is irreducible in R , but reducible in C , can be transformed linearly into a decomposable group $\begin{pmatrix} G & 0 \\ 0 & \bar{G} \end{pmatrix}$, where G and \bar{G} are two groups irreducible in C , with coefficients not all in R , such that the coefficients in every transformation of \bar{G} are the conjugate imaginaries of the corresponding coefficients for G .

In seeking a generalization, we note that the domain C may be considered as derived from R by the adjunction of a root i of the quadratic equation $x^2 + 1 = 0$ belonging to and irreducible in R . For the generalization, R is replaced by a general domain F (or field not having a modulus) and $R(i)$ is replaced by the domain $F(\rho_0)$ given by the extension of F by the adjunction of a root ρ_0 of an equation $f(x) = 0$ of degree r belonging to and irreducible in F . The generalization will therefore be two-fold. Let the roots of $f(x) = 0$ be $\rho_0, \rho_1, \dots, \rho_{r-1}$. If G_{11} is a group of transformations with coefficients $C_{ij}(\rho_0)$ in the domain $F(\rho_0)$, let $G_{11}^{(s)}$ denote the group of transformations with the coefficients $C_{ij}(\rho_s)$; in particular, $G_{11}^{(0)} = G_{11}$. The coefficients of $G_{11}, G_{11}^{(1)}, \dots, G_{11}^{(r-1)}$ are thus conjugate with respect to F . The generalized theorem is as follows:

*Let G be a group of linear homogeneous transformations with coefficients in a domain F , such that G is irreducible in F but is reducible in the domain $F(\rho_0)$ given by the extension of F by the adjunction of a root ρ_0 of an equation belonging to and irreducible in F and having as its roots $\rho_0, \rho_1, \dots, \rho_{r-1}$. Then G can be transformed linearly into a decomposable group**

* Presented to the Society at the Boston summer meeting, August 31-September 1, 1903. Received for publication April 27, 1903.

† When the irreducible equation is a normal equation, the groups $G_{11}^{(s)}$ ($s = 0, 1, \dots, r-1$) are all irreducible in the same (normal) domain. LOEWY'S case furnishes an example.

$$\begin{matrix}
 G_{11} & 0 & 0 & \dots & 0 \\
 0 & G'_{11} & 0 & \dots & 0 \\
 0 & 0 & G''_{11} & \dots & 0 \\
 \dots & \dots & \dots & \dots & \dots \\
 0 & 0 & 0 & \dots & G_{11}^{(r-1)}
 \end{matrix}$$

where $G_{11}^{(s)}$ is a group irreducible in $F(\rho_s)$ with coefficients not all in F , and $G_{11}, G'_{11}, \dots, G_{11}^{(r-1)}$ are conjugate with respect to F .

The proof starts as in LOEWY, §1. The first variation * occurs at the bottom of p. 173; we now take r -fold decomposable matrices

$$H = \begin{bmatrix} G & 0 & \dots & 0 \\ 0 & G & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & G \end{bmatrix}, \quad Q = \begin{bmatrix} P & 0 & \dots & 0 \\ 0 & P' & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & P^{(r-1)} \end{bmatrix}.$$

Corresponding changes are to be made in the first two statements on p. 174. Thus, the diagonal groups in (6) are to be replaced by

$$G_{11}, G_{22}, G'_{11}, G'_{22}, G''_{11}, G''_{22}, \dots, G_{11}^{(r-1)}, G_{22}^{(r-1)}.$$

In place of the transformation † (7), we have

$$(7') \quad y_{jk} = \sum_{i=1}^n C_{ki}^{(j)} y_{ji}^* \quad (k=1, \dots, n; j=0, \dots, r-1),$$

where $C_{ki}^{(j)}$ is a rational function of ρ_j with coefficients in F , and

$$(7'') \quad C_{ki}^{(j)} = 0 \quad (k=1, \dots, m; i=m+1, \dots, n; j=0, \dots, r-1).$$

Introduce two pairs each of rn new variables defined by

$$\begin{aligned}
 (8') \quad y_{sk} &= Y_{0k} + \rho_s Y_{1k} + \rho_s^2 Y_{2k} + \dots + \rho_s^{r-1} Y_{r-1k} \\
 (8'') \quad y_{sk}^* &= Y_{0k}^* + \rho_s Y_{1k}^* + \rho_s^2 Y_{2k}^* + \dots + \rho_s^{r-1} Y_{r-1k}^*
 \end{aligned} \quad \left(\begin{matrix} s=0, \dots, r-1 \\ k=1, \dots, n \end{matrix} \right).$$

This may be done since the determinant

$$\Delta \equiv \begin{vmatrix} 1 & \rho_0 & \rho_0^2 & \dots & \rho_0^{n-1} \\ 1 & \rho_1 & \rho_1^2 & \dots & \rho_1^{n-1} \\ \dots & \dots & \dots & \dots & \dots \\ 1 & \rho_{r-1} & \rho_{r-1}^2 & \dots & \rho_{r-1}^{n-1} \end{vmatrix} = \Pi (\rho_i - \rho_j) \neq 0.$$

* The statement on p. 173, lines 7-8, is apparently not used later; a proof follows readily from the main theorem under consideration.

† LOEWY's notation is unwieldy even in his simple case. I write y_{0k}, y_{1k} for his y_k, z_k . The transformed variables are marked * instead of being primed.

Solving (8') for fixed k , while $s = 0, \dots, r - 1$, we get

$$(e) \quad \Delta Y_{tk} = \sum_{s=0}^{r-1} (-1)^s D_{ts} y_{sk} \quad (t=0, 1, \dots, r-1),$$

where

$$D_{ts} \equiv \begin{pmatrix} 1 & \rho_0 & \dots & \rho_0^{t-1} & \rho_0^{t+1} & \dots & \rho_0^{r-1} \\ 1 & \rho_1 & \dots & \rho_1^{t-1} & \rho_1^{t+1} & \dots & \rho_1^{r-1} \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 1 & \rho_{s-1} & \dots & \rho_{s-1}^{t-1} & \rho_{s-1}^{t+1} & \dots & \rho_{s-1}^{r-1} \\ 1 & \rho_{s+1} & \dots & \rho_{s+1}^{t-1} & \rho_{s+1}^{t+1} & \dots & \rho_{s+1}^{r-1} \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 1 & \rho_{r-1} & \dots & \rho_{r-1}^{t-1} & \rho_{r-1}^{t+1} & \dots & \rho_{r-1}^{r-1} \end{pmatrix}.$$

Substituting for y_{sk} in (e) its value from (7') and then eliminating y_{si}^* by (8'), we obtain

$$(9) \quad Y_{tk} = \sum_{\substack{i=1, \dots, n \\ t=0, \dots, r-1}} \alpha_{ii}^{tk} Y_{ii}^* \quad (k=1, \dots, n; t=0, \dots, r-1),$$

where

$$\alpha_{ii}^{tk} \equiv \frac{(-1)^t}{\Delta} \sum_{s=0}^{r-1} (-1)^s D_{ts} C_{ki}^{(s)} \rho_s^t.$$

The coefficients of transformation (9) belong to the domain F' . It suffices to show that each α_{ii}^{tk} is unaltered by the interchange of ρ_0 with ρ_j (j being any one of the series $1, 2, \dots, r - 1$), since it is then a symmetric function of $\rho_0, \rho_1, \dots, \rho_{r-1}$ with coefficients in F' . To show that, for example, it is unaltered by the interchange of ρ_0 with ρ_1 , we note that under this interchange, D_{t0} and D_{t1} are interchanged, D_{ts} ($s > 1$) is changed into $-D_{ts}$ while $C_{ki}^{(0)} \equiv C_{ki}(\rho_0)$ and $C_{ki}^{(1)} \equiv C_{ki}(\rho_1)$ are interchanged, and $C_{ki}^{(s)}$ ($s > 1$) is unaltered. Hence the factor of α given by the sum is changed in sign; likewise the factor $1/\Delta$.

Moreover, from (7_a) follows at once

$$\alpha_{ii}^{tk} = 0 \quad (i = m + 1, \dots, n; k = 1, \dots, m; t, l = 0, 1, \dots, r - 1).$$

The group of transformations (9) is therefore of LOEWY'S form (10), \bar{H}_{11} being always a matrix of rm rows and rm columns. The proof is then readily completed as in LOEWY'S case (bottom of p. 175 and 176).

THE UNIVERSITY OF CHICAGO,
April 27, 1903.