

NOTES AND ERRATA, VOLUMES 10, 11.

VOLUME 10.

Page 315. W. B. FITE: *Irreducible homogeneous linear groups in an arbitrary domain.*

The results of this article are embodied in the theorem, *A necessary and sufficient condition that any group of finite order be simply isomorphic with an irreducible group in any domain is that its central be cyclic.* But that this condition is not sufficient is shown by the following example to which Professor BURNSIDE has kindly called my attention, namely the group defined by the operations  $A$ ,  $P$ ,  $Q$ , and  $R$  with the following relations:

$P^7 = 1$ ,  $Q^7 = 1$ ,  $R^n = 1$  ( $n$  any positive integer),  $A^3 = 1$ ,  $A^{-1}PA = P^2$ ,  $A^{-1}QA = Q^2$ ,  $A^{-1}RA = R$ , and with  $P$ ,  $Q$ , and  $R$  commutative among themselves.

The argument rests upon the supposed fact that  $r_1 > 1$ , and I can now see no reason why it might not equal unity.

I have been unable to determine what is both a sufficient and a necessary condition that a group of finite order may be simply isomorphic with an irreducible group. It is easy however to give sufficient conditions that are not necessary ones; for example, in order that a group of finite order be simply isomorphic with an irreducible group it is *sufficient* that its direct product with some group of finite order be simply isomorphic with an irreducible group.

VOLUME 11.

Page 489. G. E. WAHLIN: *On the base of a relative number field, with an application to the composition of fields.*

The assertion that  $O_{\tau}^{(1)}$  is prime to  $I_{\tau}$  is not true for every  $\vartheta$ . I wish therefore to replace lines 7-19 of p. 489 by the following.

Since  $\omega_{\tau}^{(1)} = \bar{\omega}_{\tau} / D_k(\vartheta)$  is an integer in  $K$ ,  $\bar{\omega}_{\tau}$  is divisible by  $D_k(\vartheta)$ , and consequently also by  $O_{\tau}$ . That is,

$$\bar{\omega}_{\tau} = O_{1\tau}^{(1)} + O_{2\tau}^{(1)}\vartheta + \dots + O_{\tau\tau}^{(1)}\vartheta^{\tau-1} \equiv 0(O_{\tau}).$$

From the fact,\* that if  $J$  and  $J_1$  are two ideals in a number field, then there exists an ideal  $J'$  such that  $JJ'$  is a principal ideal and  $J'$  is prime to  $J_1$ , we know that there exists in  $k$  an ideal  $J_{\tau}$ , prime to  $[D_k(\vartheta)]$  and such that

\* DIRICHLET: *Vorlesungen über Zahlentheorie*, vierte Auflage, p. 559, Theorem XI.

$O_\tau \cdot J_\tau = (\alpha)$ , a principal ideal. In the same way there exists an ideal  $J'_\tau$ , prime to  $[D_k(\vartheta)]$ , such that  $J_\tau \cdot J'_\tau = (\beta)$ , a principal ideal, and since  $J_\tau$  and  $J'_\tau$  are both prime to  $D_k(\vartheta)$ ,  $\beta$  is prime to  $D_k(\vartheta)$ .

Since  $(\beta) = J_\tau \cdot J'_\tau$  and  $(\alpha) = O_\tau \cdot J_\tau$  and since  $\bar{\omega}_\tau$  is divisible by  $O_\tau$ , it follows that  $\beta \bar{\omega}_\tau$  is divisible by  $(\alpha)$ , and must therefore be a multiple of  $\alpha$ , and

$$\frac{\beta \bar{\omega}_\tau}{\alpha} = \frac{\beta O_{1\tau}^{(1)} + \beta O_{2\tau}^{(1)} \vartheta + \dots + \beta O_{\tau\tau}^{(1)} \vartheta^{\tau-1}}{\alpha}$$

is an integer in  $K$ .

But since  $\bar{\omega}_\tau$  and  $O_{\tau\tau}^{(1)}$  are both divisible by  $O_\tau$  it follows that

$$X = O_{1\tau}^{(1)} + O_{2\tau}^{(1)} \vartheta + \dots + O_{\tau-1,\tau}^{(1)} \vartheta^{\tau-2} = \bar{\omega}_\tau - O_{\tau\tau}^{(1)} \vartheta^{\tau-1}$$

must be divisible by  $O_\tau$  and therefore, in the same way as above, we see that  $\beta X/\alpha$  is an integer in  $K$ , and  $R = \beta O_{\tau\tau}^{(1)}/\alpha$  an integer in  $k$ . We can now write

$$\frac{\beta \bar{\omega}_\tau}{\alpha} = \frac{\beta X + \beta O_{\tau\tau}^{(1)} \vartheta^{\tau-1}}{\alpha} = \frac{\beta X}{\alpha} + R \vartheta^{\tau-1}$$

and since  $O_\tau$  is the highest common factor of  $O_{\tau\tau}^{(1)}$  and  $D_k(\vartheta)$ , the ideals  $O_{\tau\tau}^{(1)}/O_\tau$  and  $I_\tau = D_k(\vartheta)/O_\tau$  are relatively prime, and since  $\beta$  is prime to  $I_\tau$ , it follows that  $R$  is prime to  $I_\tau$  and there exists in  $k$  an integer  $R'$  such that

$$R'R \equiv 1 (I_\tau).$$

We have therefore

$$\frac{R'\beta \bar{\omega}_\tau}{\alpha} = \frac{R'\beta X}{\alpha} + R'R \vartheta^{\tau-1} \equiv \bar{\Omega}_\tau (I_\tau),$$

where

$$\bar{\Omega}_\tau = \frac{R'\beta O_{1\tau}^{(1)} + R'\beta O_{2\tau}^{(1)} \vartheta + \dots + R'\beta O_{\tau-1\tau}^{(1)} \vartheta^{\tau-2}}{\alpha} + \vartheta^{\tau-1}.$$

Since  $\bar{\omega}_\tau$  is divisible by  $D_k(\vartheta)$ , and  $\alpha$  contains no other factor of  $D_k(\vartheta)$  than  $O_\tau$ , we see that the integer  $\beta \bar{\omega}_\tau/\alpha$  must be divisible by  $I_\tau$  and from the last congruence we have

$$\bar{\Omega}_\tau \equiv O (I_\tau).$$

While  $\Omega_\tau = \bar{\Omega}_\tau/D_k(\vartheta)$  is not necessarily an integer in  $K$ , the products  $\Omega_\tau \cdot O_{\tau\tau}^{(1)}$  and  $\Omega_\tau \cdot D_k(\vartheta)$  are both integers in  $K$ . Moreover, since  $\beta$  is divisible by  $J_\tau$  and since  $D_k(\vartheta)$  and  $O_{\tau\tau}^{(1)}$  are both divisible by  $O_\tau$ , the coefficients  $R'\beta O_{\tau\tau}^{(1)}$  when multiplied by  $O_{\tau\tau}^{(1)}$  or  $D_k(\vartheta)$  will be divisible by  $\alpha$  and the integers  $\Omega_\tau \cdot D_k(\vartheta)$  and  $\Omega_\tau \cdot O_{\tau\tau}^{(1)}$  are therefore of the form  $\omega_\tau^{(i)}$ . The number  $\Omega_\tau$  thus found can therefore take the place of the  $\Omega_\tau$  in my original paper.