

A NEW METHOD FOR WARING THEOREMS WITH POLYNOMIAL SUMMANDS, II*

BY

L. E. DICKSON

1. In a paper† with the same title, I showed how to deduce instantaneously a Waring theorem for an even polynomial $f(x)$ of degree $2n$ from a known Waring theorem for a polynomial $g(x)$ of degree n . Here I extend the method to the new case in which $f(x)$ contains also a term in x .

2. First, let $n = 2$ and

$$(1) \quad f(x) = ux^4 + vx^2 - wx + k, \quad g(x) = ux^2 + vx + 2k.$$

We have the identity in a, b, c, d, u, v, w, k

$$(2) \quad 6q(s) \equiv \sum f(z), \quad s = a^2 + b^2 + c^2 + d^2,$$

in which z takes the following twelve values:

$$(3) \quad b \pm a, \quad c \pm a, \quad d \pm a, \quad \pm b - c, \quad \pm d - b, \quad \pm c - d,$$

whose sum is zero. Since some of the numbers (3) are negative, we impose the condition

$$(4) \quad f(x) \text{ is an integer } \geq 0 \text{ for all integers } x.$$

But when x ranges over all integers (positive, negative, or zero), evidently $f(-x)$ takes the same values as $f(x)$. Without loss of generality we may therefore take $w \geq 0$. Since $f(-x) = f(x) + 2wx$, $f(-x)$ will be ≥ 0 for all integers $x \geq 0$ if the same is true for $f(x)$. Hence (4) follows from

$$(5) \quad f(x) \text{ is an integer } \geq 0 \text{ for every integer } x \geq 0.$$

Since $u > 0$, only a limited number of integers x yield negative values of $f(x) - k$. If one of these values is $-P$, while all the remaining are $\geq -P$, then (5) holds if and only if $k \geq P$. In brief, we need only take k sufficiently large in (1).

Consider triangular, pyramidal, and figurate numbers

$$(6) \quad \begin{aligned} T(x) &= (x^2 - x)/2, & P(x) &= (x^3 - x)/6, \\ F(x) &= (x + 2)(x + 1)x(x - 1)/24 = (x^4 + 2x^3 - x^2 - 2x)/24. \end{aligned}$$

* Presented to the Society, November 29, 1935; received by the editors June 14, 1935.

† These Transactions, vol. 36 (1934), pp. 731-748. In (36) read $\pm d$ for $+d$. In (38) delete exponent 6.

Any quartic function with rational coefficients can evidently be expressed in the form

$$(7) \quad f = AF + gP + hT + tx + k,$$

where A, \dots, k are rational numbers. Let (5) hold. Taking $x=0, \dots, 4$, we see that

$$k, \quad t + k, \quad A + g + h + 2t + k, \quad 5A + 4g + 3h + 3t + k, \\ 15A + 10g + 6h + 4t + k$$

must be integers. Hence k, t, A, g, h are integers. The coefficients of x^3 and x in (7) are

$$A/12 + g/6, \quad - (A/12 + g/6) - h/2 + t.$$

These must be 0 and ≤ 0 respectively if (7) shall be of the form (1) with $w \geq 0$. Hence

$$(8) \quad A + 2g = 0, \quad h \geq 2t.$$

3. Of special interest are functions f for which

$$(9) \quad \text{Every integer } \geq 0 \text{ is a sum of } V \text{ values of } f(x)$$

for integers x . The smaller A is, the more slowly will f increase with x , and the smaller V will be in general. Hence we give to A its minimum (even) value 2. Evidently (9) requires that

$$(10) \quad f(y) = 0, \quad f(z) = 1 \text{ for certain integers } y, z.$$

The functions (7) satisfying (4), (8), and (10) and having $A = 2, t \geq -5$, are found to be those with

t	1	0	0	-1	-1	-2	-2	-2	-2	-3	-3	-3
h	≥ 2	0	1	-1	≥ 1	-2	-1	0	2	-4	1	3
k	0	0	0	2	1	6	4	3	2	16	4	3

t	-4	-4	-4	-4	-4	-5	-5	-5
h	-3	-1	0	2	4	-6	3	5
k	15	9	7	5	4	36	6	5

Hence each of these functions represents 0 and 1, and is an integer ≥ 0 for every integer x . The general theory therefore yields a value of V in (9) and hence a universal Waring theorem for summands $f(x)$.

4. Consider, for example, the seventh set $t = -2$, $h = -1$, $k = 4$. Then

$$(11) \quad \begin{aligned} f(x) &= 2F - P - T - 2x + 4 = (x^4 - 7x^2)/12 - \frac{2}{3}x + 4, \\ Q &= 6q = H + 42, \quad H = \frac{1}{2}(x-3)(x-4). \end{aligned}$$

Every integer ≥ 0 is a sum of three values of the triangular number H for integers $x \geq 4$. Thus every integer ≥ 126 is a sum of three values of Q . Hence by (2), every integer ≥ 126 is a sum of 36 values of $f(x)$. We next verify this fact also for positive integers < 126 and indicate a probable reduction from 36 to 5. By a table to 1000 of sums of three values of $f(x)$, we find that 415, 734, and 749 are the only positive integers < 1000 which are not sums of four values. We find at once that all integers ≤ 5114 are sums of five values of $f(x)$ for integers x (positive, negative, or zero).

5. Waring theorem for sextic polynomials. Let

$$(12) \quad \begin{aligned} f(x) &= ux^6 + vx^4 + wx^2 - hx + k, \\ q(x) &= 120ux^3 + 72vx^2 + 60wx + 108k. \end{aligned}$$

Then a like generalization of (38) of the former paper gives

$$(13) \quad q(s) \equiv \sum f(y) + 8 \sum f(z) + f(2a) + f(2b) + f(2c) + f(2d),$$

where z ranges over the twelve values (3), and y ranges over the eight values

$$(14) \quad \begin{aligned} -a - b - c \pm d, \quad \pm a - b + c - d, \quad \mp a + b - c - d, \\ \pm (a - b - c) + d, \end{aligned}$$

whose sum is $-2a - 2b - 2c - 2d$. Hence the sum of the 108 arguments of f in (13) is zero. In case $f(x)$ is an integer ≥ 0 for every integer x (which is true when k is sufficiently large), a Waring theorem for q leads instantly to one for f . This condition (4) holds if $f(x)$ is

$$(15) \quad x^6 + x^2 - x, \quad \frac{1}{2}(x^6 + 3x^2) - x,$$

each of which represents 0 and 1. Hence each yields at once a universal Waring theorem.

6. Quartics with property (4). Replacing x by $-x-1$ in (7), we get

$$(16) \quad AF(x) - gP(x) + (h-g)T(x) + (2h-g-t)x + h-t+k.$$

Hence (7) remains unaltered if and only if

$$(17) \quad g = 0, \quad h = t.$$

In this case the values of $f(x)$ for negative integers coincide with its values for integers $x \geq 0$. Such unfavorable cases are

$$(18) \quad f = F(x), \quad F - T - x + 2, \quad F - 6T - 6x + 56,$$

whose values are ≥ 0 for an integer $x \geq 0$ and hence for all integers. By tables to 1000, all integers from 0 to 3366 inclusive are sums of 7 values of $F(x)$ except only 64, 99, 119, 189, 314, 774. Hence all ≤ 23841 are sums of 8.

Since $T(-x) = T(x+1)$, $F+T \geq 0$ for all integers. By a table to 1000 of sums by three, it was verified that all integers from 0 to 3900 are sums of four values of $F+T$.

Miss H. Rees found that (7) has property (4) if

$$A = 1, \quad g = -p - 2, \quad h = p + 1 + \frac{1}{6}p(2p + 1) + m, \quad t = 1, \quad k = 0, \\ p \geq 0, \quad m \geq 0.$$

To obtain an integer h take m to be the sum of an integer ≥ 0 and $\frac{1}{2}$ if $p \equiv 1$ or $3 \pmod{6}$; 0 if $p \equiv 0$ or 4 ; $\frac{1}{3}$ if $p \equiv 2$; $\frac{5}{6}$ if $p \equiv 5 \pmod{6}$. We may remove the term involving P by the transformation $x = y + p + 2$. We get

$$(19) \quad F + (m - r)T + \{1 - r + m(p + 2)\}y + (p + 2)\{1 + (p + 1)J\}, \\ r = \frac{1}{6}p(p + 2), \quad J = (p + 1)(p - 12)/24 + \frac{1}{2}(m + p + 1).$$

When $p = 1$, $m = \frac{1}{2}$, (19) is $f = F + 2y + 5$. By a table of sums of three values from 0 to 3000 and from 9000 to 11000, it was found that every integer ≤ 16151 , except only 11784, is a sum of four (positive) values of (16) for positive and negative integers x . It follows that all ≤ 210739 are sums of five such values.

Another favorable function $f = F + y + 2$ is the case $p = m = 0$ of (19); it represents 0, 1, 2, 3, 5, 10, 12, 21, etc.

UNIVERSITY OF CHICAGO,
CHICAGO, ILL.