

# DIFFERENTIABLE FUNCTIONS DEFINED IN ARBITRARY SUBSETS OF EUCLIDEAN SPACE\*

BY  
HASSLER WHITNEY

1. **Introduction.** In a former paper† we studied the differentiability of a function defined in closed subsets of Euclidean  $n$ -space  $E$ . We consider here the differentiability “about” an arbitrary point of a function defined in an arbitrary subset of  $E$ . We show in Theorem 1 that any function defined in a subset  $A$  of  $E$  which is differentiable about a subset  $B$  of  $E$  may be extended over  $E$  so that it remains differentiable about  $B$ . This theorem is a generalization of AE Lemma 2. We show further that any function of class  $C^m$  about a set  $B$  is of class  $C^{m-1}$  about an open set  $B'$  containing  $B$ . In the second part of the paper we consider some elementary properties of differentiable functions, such as: the sum or product of two such functions is such a function.‡ We end with the theorem that differentiability is a local property.§

2. **Definitions and elementary properties.** We use a one-dimensional notation as in AE. Thus  $f_k(x) = f_{k_1 \dots k_n}(x_1, \dots, x_n)$ ,  $x^l = x_1^{l_1} \dots x_n^{l_n}$ ,  $l! = l_1! \dots l_n!$ ,  $D_k f(x) = \partial^{k_1 + \dots + k_n} f(x) / \partial x_1^{k_1} \dots \partial x_n^{k_n}$ , etc.; we set  $\sigma_k = k_1 + \dots + k_n$ ,  $r_{xy}$  = distance from  $x$  to  $y$ . We always set  $f(x) = f_0(x)$ . Suppose the functions  $f_k(x)$  for  $\sigma_k \leq m$  are defined in the subset  $A$  of Euclidean  $n$ -space  $E$ . Define  $R_k(x'; x)$  for  $x, x'$  in  $A$  by

---

\* Presented to the Society, January 2, 1936; received by the editors October 26, 1935.

† *Analytic extensions of differentiable functions defined in closed sets*, these Transactions, vol. 36 (1934), pp. 63–89. We refer to this paper as AE. See also *Functions differentiable on the boundaries of regions*, Annals of Mathematics, vol. 35 (1934), pp. 482–485, and *Differentiable functions defined in closed sets*, I, these Transactions, vol. 36 (1934), pp. 369–387, which we refer to as F and D respectively.

P. Franklin in Theorem 1 of a paper *Derivatives of higher order as single limits*, Bulletin of the American Mathematical Society, vol. 41 (1935), pp. 573–582, has given a necessary and sufficient condition for the existence of a continuous  $m$ th derivative. We remark that this theorem is exactly the special case of Theorem I of D obtained by letting  $f(x)$  be defined in an interval. It is also a special case of Theorem 2 of the author's *Derivatives, difference quotients, and Taylor's formula*, Bulletin of the American Mathematical Society, vol. 40 (1934), pp. 89–94 (see also Errata, p. 894). For his assumption is easily seen to imply the needed uniformity condition; it also implies at once that  $f(x)$  is continuous, so that no considerations of measurability are necessary. His Theorem 2 should be compared with Theorems II and III of D.

‡ If the set is closed, these theorems may be proved by first extending the functions throughout  $E$ .

§ For the case of one variable this follows from D, Theorem I.

$$(1) \quad f_k(x') = \sum_{\sigma_l \leq m - \sigma_k} \frac{f_{k+l}(x)}{l!} (x' - x)^l + R_k(x'; x).$$

Let  $x^0$  be an arbitrary point of  $E$ . If for each  $k$  ( $\sigma_k \leq m$ ) and every  $\epsilon > 0$  there is a  $\delta > 0$  such that

$$(2) \quad |R_k(x'; x)| \leq r_{xx'}^{m-\sigma_k} \epsilon \text{ if } x, x' \text{ in } A, r_{xx^0} < \delta, r_{x'x^0} < \delta,$$

we shall say that  $f(x)$  is of class  $C^m$  in  $A$  about  $x^0$  in terms of the  $f_k(x)$ , or,  $f(x)$  is  $(C^m, A, x^0, f_k(x))$ . If this is true for each  $x^0$  in  $B$ , we say  $f(x)$  is  $(C^m, A, B, f_k(x))$ , and replace "about  $x^0$ " by "about  $B$ ." We say  $f(x)$  (defined in  $A$ ) is of class  $C^m$  in  $A$  about  $B$ , or,  $f(x)$  is  $(C^m, A, B)$ , if there exist functions  $f_k(x)$  ( $\sigma_k \leq m$ ) defined in  $A$  such that  $f(x)$  is  $(C^m, A, B, f_k(x))$ . If  $B=A$  in the last two definitions, we leave out the words "about  $B$ "; this is in agreement with the previous definitions. We say  $f(x)$  is  $(C^\infty, A, B, f_k(x))$  if  $f(x)$  is  $(C^m, A, B, f_k(x))$  for each  $m$ . Any function defined in  $A$  is  $(C^{-1}, A, E)$ .

**Remark.** We might define in an obvious manner such relations as  $(C^m, A, x^0)$ ,  $(C^\infty, A, B)$ . To study them would require a study of the different possible definitions of the  $f_k(x)$  if  $f(x)$  is  $(C^m, A, B)$ . The  $f_k(x)$  are not in general determined by  $f(x)$ . Thus if  $A=B$  is the  $x_1$ -axis, only the  $f_k(x)$  with  $k_2 = \dots = k_n = 0$  are determined by  $f(x)$ . It is not obvious for what point sets  $A$  the  $f_k(x)$  are all determined by  $f(x)$ .

If  $f(x)$  is  $(C^m, A, B, f_k(x))$  ( $m \geq 0$ ), then the  $f_k(x)$  are continuous at each point of  $B$ ;<sup>\*</sup> that is, the  $f_k(x)$  may be defined in  $B - B \cdot A$  so that this will be true. To show this, take  $x^0$  in  $B$ , set  $\epsilon = 1$ , and choose  $\delta$  so that (2) holds for any  $k$  ( $\sigma_k \leq m$ ). Take  $x$  in  $A$  within  $\delta$  of  $x^0$  (if there is such a point); then (1) and (2) show that  $f_k(x')$  is bounded for  $x'$  in  $A$  within  $\delta$  of  $x^0$  ( $\sigma_k \leq m$ ). Now let  $\{x^i\}$  be any sequence of points of  $A$ ,  $x^i \rightarrow x^0$ ; (1) and (2) show that  $\{f_k(x^i)\}$  is a regular sequence.

If  $A$  is open and  $f(x)$  is  $(C^m, A, A, f_k(x))$ , then  $D_k f(x)$  exists and equals  $f_k(x)$  in  $A$  ( $\sigma_k \leq m$ ). (See AE.) If  $x^0$  is an isolated point of  $A$  or is at a positive distance from  $A$ , then  $f(x)$  is  $(C^m, A, x^0, f_k(x))$  for any  $f_k(x)$ . If  $f(x)$  is  $(C^m, A, B, f_k(x))$  [or  $(C^m, A, B)$ ], and  $A'$  is in  $A$ ,  $B'$  is in  $B$ , then  $f(x)$  is  $(C^m, A', B', f_k(x))$  [or  $(C^m, A', B')$ ]. Also  $f(x)$  is  $(C^0, A, B)$  if and only if it is continuous at each point of  $B$ . If  $f(x)$  is  $(C^m, A, B, f_k(x))$ , then it is  $(C^{m'}, A, B, f_k(x))$  for all  $m' < m$ ; a stronger theorem is proved in Theorem 2. If  $f(x)$  is  $(C^m, A, B, f_k(x))$ , then  $f_k(x)$  is  $(C^{m-\sigma_k}, A, B, f_l(x))$ .

**3. Extension theorems.** We prove here a theorem which gives the maximum range of differentiability of a function, and a theorem about the still larger range of differentiability of a function to an order one less.

<sup>\*</sup> Or better, "continuous in  $A$  about  $B$ ."

**THEOREM 1.** *If  $f(x)$  is  $(C^m, A, B, f_k(x))^*$  ( $m$  finite or infinite), then the  $f_k(x)$  may be extended throughout  $E$  so that  $f(x)$  is  $(C^m, E, B, f_k(x))$ .†*

We note, conversely, that if  $f(x)$  is not  $(C^m, A, x^0, f_k(x))$ , then no extension of  $f(x)$  will be so. We remark also that  $f(x)$  may be made analytic in  $E - \bar{A}$  ( $\bar{A} = A$  plus limit points).

To prove the theorem, we first extend the  $f_k(x)$  through  $\bar{A} - A$  as follows: Take any  $x^0$  in  $\bar{A} - A$ . Let  $f_k(x^0)$  be the upper limit of  $f_k(x^i)$  for sequences  $\{x^i\}$ ,  $x^i \rightarrow x^0$ ,  $x^i$  in  $A$ , if this is finite; otherwise, set  $f_k(x^0) = 0$ . Next we extend the  $f_k(x)$  throughout  $E - \bar{A}$  by the method of AE Lemma 2. We shall assume in the proof that  $m$  is finite. If  $m = \infty$ , we prove  $C^{m'}$  for every integer  $m'$ . The only alteration needed in the proof is that AE §12 should be used; but this makes no essential change.

As  $E - \bar{A}$  is open,  $f(x)$  is  $(C^m, E, E - \bar{A}, f_k(x))$ ; we must show that  $f(x)$  is  $(C^m, E, B \cdot \bar{A}, f_k(x))$ . Take a fixed point  $x^0$  in  $B \cdot \bar{A}$ . Let us say  $(k, \epsilon, A_1, A_2)$  holds if there is a  $\delta > 0$  such that (2) holds whenever  $x$  is in  $A_1$ ,  $x'$  is in  $A_2$ , and  $r_{xx^0} < \delta$ ,  $r_{x'x^0} < \delta$ . We must prove  $(k, \epsilon, E, E)$  for each  $k$  ( $\sigma_k \leq m$ ) and each  $\epsilon > 0$ .

First we prove  $(k, \epsilon, \bar{A}, \bar{A})$ . Set  $\epsilon' = \epsilon / [2(m+1)^n]$ , and let  $\delta$  be the smallest of the  $\delta$ 's given by  $(l, \epsilon', A, A)$  for  $\sigma_l \leq m$ . Let  $U$  be the spherical neighborhood of  $x^0$  of radius  $\delta$ ; then  $f_l(x)$  is bounded in  $U \cdot A$  ( $\sigma_l \leq m$ ). Given  $x, x'$  in  $U \cdot \bar{A}$ , choose sequences  $\{x^i\}$ ,  $\{x'^i\}$  of points of  $U \cdot A$ , with  $x^i \rightarrow x$ ,  $x'^i \rightarrow x'$ . Suppose first  $\sigma_k = m$ . Then we may take these sequences so that  $f_k(x^i) \rightarrow f_k(x)$ ,  $f_k(x'^i) \rightarrow f_k(x')$ , and the desired inequality for  $R_k(x'; x)$  follows from that for  $R_k(x'^i; x^i)$ . Suppose now that  $\sigma_k < m$ . Relations (1) and (2) with  $k, x'$ ,  $x$  replaced by  $l, x^i, x^j$  show that for any such  $\{x^i\}$ ,  $\{f_l(x^i)\}$  is a regular sequence ( $\sigma_l < m$ ); hence  $f_l(x^i) \rightarrow f_l(x)$ , and similarly  $f_l(x'^i) \rightarrow f_l(x')$  ( $\sigma_l < m$ ). Relation (1) now shows that for  $i$  large enough,  $\Delta = R_k(x'; x) - R_k(x'^i; x^i)$  differs as little as we please from

$$-\sum_{\sigma_l = m - \sigma_k} \frac{f_{k+l}(x) - f_{k+l}(x^i)}{l!} (x' - x)^l.$$

As  $|f_j(x') - f_j(x^i)| \leq \epsilon'$  ( $\sigma_j = m$ ) and  $|(x' - x)^l| \leq r_{xx^0}^{\sigma_l}$ ,  $|\Delta| \leq (m+1)^n \epsilon' r_{xx^0}^{m - \sigma_k}$  for  $i$  large enough; the inequality again follows.

Next we prove  $(k, \epsilon, \bar{A}, E - \bar{A})$ . Set  $\epsilon' = \epsilon / [2 \cdot 4^m (m+1)^n]$ , and define  $\eta$  in terms of  $\epsilon'$  and then  $\delta$  as in AE §11, using  $(k, \eta, \bar{A}, \bar{A})$ . Take  $x$  in  $\bar{A}$  and  $x'$  in  $E - \bar{A}$ , each within  $\delta/4$  of  $x^0$ . By AE (6.3) and the equation following (11.6),

\* Or merely locally  $(C^m, A, B)$ ; see Theorem 6.

† If  $A = B$  is closed, then  $B$  may be replaced by  $E$ ; the present proof then gives a proof of AE Lemma 2 which makes no use of AE Lemma 1.

$$\begin{aligned}
 R_k(x'; x) &= D_k f(x') - \psi_k(x'; x) \\
 &= \sum_l \frac{R_{k+l}(x^*; x)}{l!} (x' - x^*)^l + \sum_{s=1}^l \sum_l \binom{k}{l} D_l \phi_{\lambda_s}(x') \zeta_{\lambda_s; k-l}(x'),
 \end{aligned}$$

where  $x^*$  is a point of  $\bar{A}$  distant  $\delta_*/4$  from  $x'$ ,  $\delta_*/4$  being the distance from  $x'$  to  $\bar{A}$ . As  $r_{x^*x} \leq 2r_{xx'}$ ,  $r_{x'x^*} \leq 2r_{xx'}$ , and  $\delta_* \leq 4r_{xx'}$ , we find with the help of AE (11.8)

$$|R_k(x'; x)| \leq (m+1)^n (2r_{xx'})^{m-\sigma_k} \eta + (4r_{xx'})^{m-\sigma_k} \epsilon'/2 < r_{xx'}^{m-\sigma_k} \epsilon.$$

Next we prove  $(k, \epsilon, E-\bar{A}, \bar{A})$ . As is easily seen from AE (6.3) or by F (6) with  $x^{i-1}, x^i$  replaced by  $x, x'$ ,

$$R_k(x'; x) = \sum_l \frac{R_{k+l}(x; x')}{l!} (x' - x)^l.$$

Set  $\epsilon' = \epsilon/(m+1)^n$ , and take the smallest  $\delta$  given by  $(k+l, \epsilon', \bar{A}, E-\bar{A})$  for  $\sigma_l \leq m - \sigma_k$ . The required inequality now follows at once.

Finally we must show  $(k, \epsilon, E-\bar{A}, E-\bar{A})$ . Set  $\epsilon' = \epsilon/[2n(m+1)^n]$ , and take  $\delta$  smaller than the  $\delta/4$  given by AE §11 with  $\epsilon$  replaced by  $\epsilon'$  and smaller than the  $\delta$ 's given by  $(k+l, \epsilon', \bar{A}, E-\bar{A})$  and  $(k+l, \epsilon', E-\bar{A}, \bar{A})$  for  $\sigma_l \leq m - \sigma_k$ . Now take  $x$  and  $x'$  in  $E-\bar{A}$  within  $\delta$  of  $x^0$ ; we must consider two cases. Case I: The line segment  $S = xx'$  lies wholly in  $E-\bar{A}$ . By AE (11.2),  $|f_i(y) - f_i(x')| < 2\epsilon'$  for  $y$  on  $S$  ( $\sigma_l \leq m$ ); the desired inequality now follows from F, Lemma 3. Case II: There is a point  $x^*$  of  $\bar{A}$  on  $S$ . From AE (6.3), or F (6) with  $x^{i-1}, x^i$  replaced by  $x, x^*$ , we find

$$R_k(x'; x) = R_k(x'; x^*) + \sum_l \frac{R_{k+l}(x^*; x)}{l!} (x' - x^*)^l,$$

and the inequality again follows.

**THEOREM 2.** *If  $f(x)$  is  $(C^m, A, B, f_k(x))$  ( $m$  finite), then there is an open set  $B'$  containing  $B$  such that  $f(x)$  is  $(C^{m-1}, A, B', f_k(x))$ .*

For each  $x$  in  $B$ , let  $\delta(x)$  be the largest of the numbers  $\delta$  for which (2) holds for all  $k$  ( $\sigma_k \leq m$ ) with  $\epsilon$  replaced by 1. Let  $U(x)$  be the set of all points  $x'$  within  $\delta(x)$  of  $x$ ; then  $B'$  is the sum of all  $U(x)$ . The set  $B'$  is open. To prove  $(C^{m-1}, A, B', f_k(x))$ , take any  $x^0$  in  $B'$  and any  $\epsilon > 0$ . For some  $x^*$  in  $B$ ,  $r_{x^*x^0} < \delta(x^*)$ . There is an  $M$  such that  $|f_k(y)| < M$  for  $y$  in  $A \cdot U(x^*)$  ( $\sigma_k \leq m$ ). † Let  $\delta$  be the smaller of  $\delta(x^*) - r_{x^*x^0}$  and  $\epsilon/[2(m+1)^n M + 2]$ . Now take any  $x$  and  $x'$  in  $A$  within  $\delta$  of  $x^0$ . We are interested in the remainders

† For the proof, see the paragraph following the remark.

$$R'_k(x'; x) = \sum_{\sigma_l = m - \sigma_k} \frac{f_{k+l}(x)}{l!} (x' - x)^l + R_k(x'; x)$$

with  $\sigma_k < m$ . As  $r_{xx'} < 2\delta$ ,

$$|R'_k(x'; x)| \leq (m+1)^n M r_{xx'}^{m-\sigma_k} + r_{xx'}^{m-\sigma_k} < r_{xx'}^{m-1-\sigma_k} \epsilon.$$

**COROLLARY.** *If  $f(x)$  is of class  $C^m$  in any given point set about  $B$ , then it may be extended through an open set  $B'$  containing  $B$  so that it is of class  $C^{m-1}$  in  $B'$  and of class  $C^m$  in  $B'$  about  $B$ .*

**4. Composite functions, etc.** We prove here three theorems.

**THEOREM 3.** *If  $f$  and  $g$  are of class  $C^m$  in  $A$  about  $B$ , then so are  $f+g$  and  $f-g$ , with*

$$(3) \quad (f \pm g)_k = f_k \pm g_k.$$

This is obvious.

**THEOREM 4.** *If  $f$  and  $g$  are of class  $C^m$  in  $A$  about  $B$ , then so is  $fg$ , and  $f/g$  if  $g \neq 0$ . The derivatives are given by the ordinary formulas. Thus*

$$(4) \quad (fg)_k = \sum_l \binom{k}{l} f_l g_{k-l}.$$

We might prove this theorem directly, but it follows from Theorem 5:  $fg$  and  $f/g$  are functions (of two variables) of class  $C^\infty$  of the functions  $f$  and  $g$ . (The condition  $B$  in  $A$  is obtained by using Theorem 1.)

**THEOREM 5.** *Let  $A$  and  $B$  be subsets of  $n$ -space  $E_n$ , and let  $A'$  and  $B'$  be subsets of  $\nu$ -space  $E_\nu$ . Let  $f^i(x)$  be  $(C^m, A, B, f^i(x))$  ( $i=1, \dots, \nu$ ), and let  $g(y)$  be  $(C^m, A', B', g_k(y))$  ( $m$  finite or infinite). Suppose  $B$  is in  $A$ ,  $x$  in  $A$  implies*

$$y = (y_1, \dots, y_\nu) = (f^1(x), \dots, f^\nu(x)) = f(x)$$

in  $A'$ , and  $x$  in  $B$  implies  $f(x)$  in  $B'$ . Then the function

$$h(x) = g(f^1(x), \dots, f^\nu(x)) = g(f(x))$$

is  $(C^m, A, B, h_k(x))$ ; the  $h_k(x)$  are given by the ordinary formulas (9) for derivatives.

As a consequence of this theorem, the definition of being of class  $C^m$  is independent of the coordinate system chosen. If the condition  $x$  in  $A$  [or  $B$ ] does not imply  $f(x)$  in  $A'$  [or  $B'$ ], we may apply the theorem to any subset  $A_1$  [or  $B_1$ ] of  $A$  [or  $B$ ] for which it does. We shall suppose  $m$  is finite; if  $m = \infty$ , we merely apply the reasoning below for each positive integer.

Suppose first  $u^1(x), \dots, u^r(x)$  are functions of class  $C^m$  in an open set  $\Gamma$  of  $E_n$ , suppose  $v(y)$  is of class  $C^m$  in an open set  $\Gamma'$  of  $E_r$ , and suppose  $x$  in  $\Gamma$  implies  $u(x)$  in  $\Gamma'$ . Letting  $R^i, S'$  denote remainders for  $u^i, v$ , Taylor's formula gives

$$(5) \quad u_k^i(x') = D_k u^i(x') = \sum_{\sigma_l \leq m - \sigma_k} \frac{u_{k+l}^i(x)}{l!} (x' - x)^l + R_k^i(x'; x),$$

$$(6) \quad v_k(y') = D_k v(y') = \sum_{\sigma'_l \leq m - \sigma'_k} \frac{v_{k+l}(y)}{l!} (y' - y)^l + S'_k(y'; y),$$

certain inequalities on the  $R_k^i$  and  $S'_k$  being satisfied. We have set  $\sigma'_k = k_1 + \dots + k_r$ . Set  $w(x) = v(u(x))$ ; then (5) and (6) with  $k=0$  give

$$(7) \quad w(x') = \sum_t \frac{v_t(u(x))}{t!} \left\{ \sum_{\sigma_j \geq 1} \frac{u_j(x)}{j!} (x' - x)^j + R'(x'; x) \right\}^t + S'(u(x'); u(x)),$$

where  $S' = S'_0$ . Also, by Taylor's formula,

$$(8) \quad w_k(x') = \sum_l \frac{w_{k+l}(x)}{l!} (x' - x)^l + T'_k(x'; x).$$

Subtract (8) with  $k=0$  from (7); then as  $R^i, S',$  and  $T'$  all approach 0 to the  $m$ th order as  $x' \rightarrow x$ , † we may equate coefficients of  $(x' - x)^k$  for  $\sigma_k \leq m$ . ‡ Thus we find polynomials

$$P_k(u_p^i, v_q) \quad (\sigma_p \leq \sigma_k, \sigma'_q \leq \sigma_k; \sigma_k \leq m)$$

such that, for any  $x$  in  $\Gamma$ ,

$$(9) \quad w_k(x) = P_k(u_p^i(x), v_q(u(x))).$$

Using (8) gives for  $w_k(x')$

$$(10) \quad w_k(x') = \sum_l \frac{P_{k+l}(u_p^i(x), v_q(u(x)))}{l!} (x' - x)^l + T'_k(x'; x).$$

We may also evaluate it by replacing  $x$  by  $x'$  in (9) and using (5) and (6). (In (6) we replace  $y'$  by  $u(x')$  and use (5) again.) Each variable in the resulting polynomial  $P_k$  consists of a polynomial in quantities  $R', S',$  and other quantities; if we multiply out and collect all terms with an  $R'$  or an  $S'$  as a factor, we obtain

† This is clear for  $S'$  if  $m=0$ ; if  $m>0$ , then  $S'/r_{xx'}^m = [S'/|u(x') - u(x)|^m] \cdot [|u(x') - u(x)|/r_{xx'}]^m$ ,

where  $|y' - y| = r_{yy'}$ , and the last factor is bounded in  $U \cdot A$ .

‡ This is easily proved in succession for  $\sigma_k=0, 1, \dots$  on letting  $x' \rightarrow x$ .

$$(11) \quad w_k(x') = P_k \left[ \sum_s \frac{u_{p+s}^i(x)}{s!} (x' - x)^s, \right. \\ \left. \sum_t \frac{v_{q+t}(u(x))}{t!} \left\{ \sum_{j \geq 1} \frac{u_j(x)}{j!} (x' - x)^j \right\}^t \right] + Q_k,$$

where  $Q_k$  is a polynomial containing an  $R'$  or an  $S'$  as a factor in each term. It must be understood that  $\sum u_{p+s}^i(x)(x' - x)^s/s!$  appears as the variable in the position of  $u_p^i$ , etc., in  $P_k(u_p^i, v_q)$ .

We now prove: *If  $u_k^i$  ( $\sigma_k \leq m; i = 1, \dots, \nu$ ),  $v_k$  ( $\sigma'_k \leq m$ ) are any numbers, then*

$$(12) \quad P_k^*(x; u_p^i, v_q) = P_k \left[ \sum_s \frac{u_{p+s}^i}{s!} x^s, \sum_t \frac{v_{q+t}}{t!} \left\{ \sum_{j \geq 1} \frac{u_j}{j!} x^j \right\}^t \right] \\ - \sum_l \frac{P_{k+l}(u_p^i, v_q)}{l!} x^l,$$

considered as a polynomial in  $x$ , contains no terms of degree  $\leq m - \sigma_k$ . To prove this, define the polynomials

$$(13) \quad u^i(x) = \sum_{\sigma_i \leq m} \frac{u_i^i}{l!} x^l, \quad v(y) = \sum_{\sigma'_i \leq m} \frac{v_i}{l!} (y - u_0)^l;$$

then  $u_k^i(0) = D_k u^i(0) = u_k^i$ ,  $v_k(u_0) = D_k v(u_0) = v_k$ . Set  $w(x) = v(u(x))$ . Replacing  $x'$ ,  $x$  by  $x$ , 0 in (10) and (11) and putting in (12) gives, as  $Q_k = 0$  in this case,

$$(14) \quad P_k^*(x; u_p^i, v_q) = T_k'(x; 0).$$

As  $T_k' \rightarrow 0$  to the  $(m - \sigma_k)$ th order as  $x \rightarrow 0$ ,  $P_k^*$  cannot contain any terms of degree  $\leq m - \sigma_k$ .

We return now to the functions  $f^i(x)$ ,  $g(y)$ ,  $h(x)$ . Set  $h_k(x) = P_k(f_p^i(x), g_q(f(x)))$ . The formulas (10) and (11) hold equally well for the  $f^i$ ,  $g$ ,  $h$ . Hence using (10), (11), and (12), we find for the remainder for  $h_k(x)$

$$(15) \quad T_k(x'; x) = P_k^*(x' - x; f_p^i(x), g_q(f(x))) + Q_k.$$

To show that  $h(x)$  is  $(C^m, A, B, h_k(x))$ , take any  $x^0$  in  $B$ , and set  $y^0 = f(x^0)$ . As  $f(x)$  is continuous in  $A$  about  $B$ , for each neighborhood  $V$  of  $y^0$  there is a neighborhood  $U(V)$  of  $x^0$  such that  $x$  in  $U(V) \cdot A$  implies  $f(x)$  in  $V \cdot A'$ . As  $y^0$  is in  $B'$ , we may take  $V$  so that the  $g_k(y)$  are bounded in  $V \cdot A'$ . We may take  $U$  in  $U(V)$  so small that the  $f_k(x)$  are bounded in  $U \cdot A$ . Because of the property of  $P_k^*$ , we may obviously take  $\delta$  small enough so that  $P_k^*$  satisfies an inequality of the nature of (2). Moreover each term in  $Q_k$  contains an  $R_r(x'; x)$  or an  $S_q(u(x'); u(x))$  with  $\sigma_p \leq \sigma_k$  or  $\sigma'_q \leq \sigma_k$ ; as each such remainder satisfies

an inequality (2) (see a recent footnote) and all other quantities entering into  $Q_k$  are bounded, we may take  $\delta$  small enough so that  $Q_k$  also satisfies an inequality (2). Hence the same is true of  $T_k$ , and the theorem is proved.

5. Differentiability a local property. Our object is to prove

**THEOREM 6.** *Let  $f(x)$  be locally  $(C^m, A, B)$  ( $m$  finite or infinite). For each point  $x^0$  of  $B$  there is a neighborhood  $U$  of  $x^0$  and functions  $f_k^{(x^0)}(x)$  defined in  $U \cdot A$  such that  $f(x)$  is  $(C^m, U \cdot A, U \cdot B, f_k^{(x^0)}(x))$ .<sup>†</sup> Then  $f(x)$  is  $(C^m, A, B)$ . If the  $f_k^{(x^0)}(x)$  for  $\sigma_k \leq p$  are independent (at any  $x$  for which they are defined) of  $x^0$ , then these functions may be included among the  $f_k(x)$  ( $\sigma_k \leq m$ ).*

We may take each neighborhood  $U$  as an open  $n$ -cube, so small that the  $f_k^{(x^0)}(x)$  are bounded in  $U$ . A finite or denumerable number of them,  $C_1, C_2, \dots$ , cover  $B$ ; we may take them so that any one touches at most a finite number of the others, and so that any boundary point of any  $C_i$  is interior to some  $C_j$ .<sup>‡</sup> By hypothesis, to each  $i$  there correspond functions  $f_k^i(x)$ ,  $\sigma_k \leq m$ , such that  $f(x)$  is  $(C^m, C_i \cdot A, C_i \cdot B, f_k^i(x))$ . In each  $C_i$  we define the function  $\phi_i(x)$  as it was defined in  $I_i$  in AE §9; set

$$(16) \quad \phi_i(x) = \pi_i(x) / \sum_j \pi_j(x)$$

in  $C_1 + C_2 + \dots$ . Set  $g^i(x) = \phi_i(x)f(x)$  in  $C_i \cdot A$ . By Theorem 4,  $g^i(x)$  is  $(C^m, C_i \cdot A, C_i \cdot B)$ , and

$$(17) \quad g_k^i(x) = \sum_l \binom{k}{l} D_l \phi_i(x) f_{k-l}^i(x).$$

As the  $f_k^i(x)$  are bounded in  $C_i \cdot A$  and the  $D_l \phi_i(x) \rightarrow 0$  to infinite order as  $x$  approaches the boundary of  $C_i$  (see AE §9), the latter statement is true also of the  $g_k^i(x)$ . Hence, evidently, if we set  $g_k^i(x) = 0$  in  $A - C_i \cdot A$ ,  $g^i(x)$  is  $(C^m, A, B, g_k^i(x))$ . Set

$$(18) \quad f_k(x) = g_k^1(x) + g_k^2(x) + \dots,$$

which in any  $C_i \cdot A$  is a finite sum; this reduces to  $f(x)$  for  $k=0$ . Theorem 3 shows at once that  $f(x)$  is  $(C^m, A, B, f_k(x))$ . (Given  $x^0$  in  $B$ , to apply Theorem

<sup>†</sup> Note that " $f(x)$  is locally  $(C^\infty, \dots)$ " is not the same statement as " $f(x)$  is locally  $(C^m, \dots)$  for each  $m$ ."

<sup>‡</sup> Let  $C^1, C^2, \dots$  be a denumerable set of the cubes which cover  $B$ . Express each  $C^i$  as the sum of a denumerable number of cubes  $C_j^i$  with the following properties: Each  $C_j^i$  is, with its boundary, interior to  $C^i$ ; the diameter of  $C_j^i$ ,  $\delta(C_j^i)$ , is  $< 1/i$ ;  $\delta(C_j^i) \rightarrow 0$  as  $j \rightarrow \infty$ ; the cubes  $C_j^i$  approach the boundary of  $C^i$  as  $j \rightarrow \infty$ . Now drop out all cubes  $C_j^i$  which are interior to larger cubes  $C_k^i$ ; the remaining cubes  $C_1, C_2, \dots$  still cover  $B$ . To each cube  $C_j^i$  corresponds a number  $\eta > 0$  such that any point set of diameter  $< \eta$  having points in common with  $C_j^i$  lies interior to some  $C_k^i$ ; using this fact, it is easily seen that any  $C_i$  has points in common with but a finite number of the  $C_j$ .

3, we choose  $\delta$  so small that the points within  $\delta$  of  $x^0$  lie in but a finite number of the  $C_i$ .)

To prove the second statement, let  $f'_l(x)$  denote the common value of  $f'_i(x)$  for  $\sigma_i \leq p$ . Differentiating  $\sum \phi_i = 1$  gives

$$(19) \quad \sum_i D_l \phi_i(x) = \begin{cases} 1 & \text{if } l = 0, \\ 0 & \text{if } \sigma_l > 0. \end{cases}$$

Define the  $f_k(x)$  as before. Take any  $k$  with  $\sigma_k \leq p$ ; then (17) and (18) give

$$f_k(x) = \sum_l \binom{k}{l} f'_{k-l}(x) \sum_i D_l \phi_i(x) = f'_k(x)$$

in  $C_1 + C_2 + \dots$ . It does not matter how  $f_k(x)$  is defined outside this set.

The second statement in the theorem does not hold for an arbitrary set of  $f_k(x)$ , at least using the above method. To see this, take  $n = m = 2$ ,  $A = B$  = the interval  $(-1, 1)$  of the  $x_1$ -axis,  $C_1 = C_2$  = the square with corners  $(\pm 1, \pm 1)$ ; set  $f = 0$ ,

$$f_{10}^1 = f_{20}^1 = f_{11}^1 = f_{02}^1 = 0, \quad f_{01}^1 = 1,$$

and  $f_{ij}^2 = -f_{ij}^1$  on  $A$ . Also set

$$\phi_1(x, y) = \frac{1}{2} + \frac{3}{4}x - \frac{1}{4}x^3, \quad \phi_2(x, y) = \frac{1}{2} - \frac{3}{4}x + \frac{1}{4}x^3.$$

(Though  $\phi_1$  and  $\phi_2$  are not the functions defined above, they have the necessary properties.) We find on  $A$

$$g_{11}^1(x, y) = g_{11}^2(x, y) = \frac{3}{4} - \frac{3}{4}x^2, \quad f_{11}(x, y) = \frac{3}{2} - \frac{3}{2}x^2 \neq 0.$$

HARVARD UNIVERSITY,  
CAMBRIDGE, MASS.