

ON COMPLEMENTARY MANIFOLDS AND PROJECTIONS IN SPACES L_p AND l_p †

BY
F. J. MURRAY

Introduction. If Λ is a Banach space, \mathfrak{M} a closed linear subset of Λ , then a closed linear subset \mathfrak{N} such that every $f \in \Lambda$ is uniquely expressible as $g + h$, $g \in \mathfrak{M}$, $h \in \mathfrak{N}$, is called a complementary manifold to \mathfrak{M} .

In his treatise on linear operations, Banach‡ presents the following two problems ((B), pp. 244–245).

(a) *To every closed linear subset \mathfrak{M} in L_p , $1 < p \neq 2$, does there exist a complementary manifold?*

(b) *To every closed linear subset \mathfrak{M} in l_p , $1 < p \neq 2$, does there exist a complementary manifold?*

We show in this paper that the answer to both questions is “no.”

In Chapter 1 of this paper we show that if a certain limit has the value ∞ , then the answer is negative.§ In Chapters 2 and 3, it is proved that this is indeed the case. In the concluding section we discuss the relation of various other problems to (a) and (b).

CHAPTER 1. $C(\mathfrak{M})$ AND $\overline{C}(\Lambda)$

1.1. Let Λ denote a separable space with a p -norm, i.e., Λ is either L_p or l_p or the set of ordered n -tuples of real numbers $\{(a_1, \dots, a_n)\}$, $l_{p,n}$, with the norm $\|(a_1, \dots, a_n)\| = (\sum_{i=1}^n |a_i|^p)^{1/p}$. We also let $l_{p,\infty} = l_p$. The notation $p' = 1/(p-1)$, $1/p + 1/p' = 1$ will be used throughout.

Let \mathfrak{M} be a closed linear manifold in Λ . Let R denote the set of real numbers $0 \leq a \leq \infty$, and let $r(a, b) = a/(1+a) - b/(1+b)$, ($\infty/(1+\infty) \equiv 1$). It is easy to see that R with the metric $|r(a, b)|$ is a complete metric space and is homeomorphic to the closed interval $(0, 1)$.

If \mathfrak{M} is a closed linear manifold in Λ , a limited transformation E such that $E\Lambda = \mathfrak{M}$, $E^2 = E$, is said to project Λ on \mathfrak{M} .

If E is a limited transformation, we denote by $|E|$ the bound of E .

LEMMA 1.1.1. *Let \mathfrak{M} be a closed linear manifold in Λ . The existence of a*

† Presented to the Society, September 5, 1936; received by the editors August 27, 1936.

‡ *Théorie des Opérations Linéaires*, Warsaw, 1932. We shall refer to this work as (B). The quantities considered in the sequel are assumed to be real valued unless explicitly stated to the contrary.

§ This result was obtained while the author was a National Research Fellow at Brown University, Providence, R. I.

complementary manifold \mathfrak{N} to \mathfrak{M} is equivalent to the existence of a projection E of Λ on \mathfrak{M} .

Suppose \mathfrak{N} exists. Let E be the transformation which is such that $f = g + h$, $g \in \mathfrak{M}$, $h \in \mathfrak{N}$, then $Ef = g$. Owing to the properties of \mathfrak{N} , E is single-valued, additive, homogeneous and defined everywhere. Now let $\{f_i\}$ be a sequence, which approaches f and such that if $f_i = g_i + h_i$, $g_i \in \mathfrak{M}$, $h_i \in \mathfrak{N}$ the g_i form a convergent sequence with limit g' . Then g' is $\in \mathfrak{M}$ and the sequence $h_i = f_i - g_i$ also converges to a $h' \in \mathfrak{N}$. By continuity we have $f = g' + h'$. The uniqueness of the resolution of f now implies that $Ef = g'$ or that E is closed. Theorem 7 of (B), Chapter III, p. 41, now implies that E is bounded. Since the range of E is included in \mathfrak{M} and for every $f \in \mathfrak{M}$, $Ef = f$, we see that the range of E is \mathfrak{M} and $E^2 = E$ or E is a projection of Λ on \mathfrak{M} .

Now suppose E exists. Let \mathfrak{N} be the set of g 's in Λ for which $Eg = 0$. Since E is limited and linear, \mathfrak{N} is a closed linear manifold. If $f \in \Lambda$, $f = Ef + (1 - E)f$ where $Ef \in \mathfrak{M}$ and $(1 - E)f \in \mathfrak{N}$, since $E(1 - E)f = (E - E^2)f = 0$. On the other hand if $h \in \mathfrak{N}$, $h = Ef$ for some $f \in \Lambda$, and hence $Eh = E^2f = Ef = h$. Thus if $h \in \mathfrak{N} \cdot \mathfrak{M}$, $0 = Eh = h$, or $\mathfrak{M} \cdot \mathfrak{N} = \{0\}$. Now let f again be $\in \Lambda$, $f = g + h = g' + h'$, $g, g' \in \mathfrak{M}$, $h, h' \in \mathfrak{N}$. Then $g - g' = h' - h$, and since $h - h' \in \mathfrak{N}$, $g' - g \in \mathfrak{M}$, and $\mathfrak{M} \cdot \mathfrak{N} = \{0\}$, this implies $g - g' = h' - h = 0$. This shows that $f \in \Lambda$ can only be expressed in one way as $h + g$, $h \in \mathfrak{N}$, $g \in \mathfrak{M}$.

We may therefore consider problems (a) and (b) in the following equivalent form.

(A) To every closed linear manifold \mathfrak{M} of L_p , $1 < p \neq 2$, is there a projection of L_p on \mathfrak{M} ?

(B) To every closed linear manifold \mathfrak{M} of l_p , $1 < p \neq 2$, is there a projection of l_p on \mathfrak{M} ?

Let $\Lambda_1, \dots, \Lambda_n, n = 1, 2, \dots, \infty, \Lambda_\infty \equiv \Lambda$ be a set of spaces. Let $\sum_{\alpha=1}^n \oplus \Lambda_\alpha = \Lambda_1 \oplus \dots \oplus \Lambda_n$ denote the space of ordered sets of elements $\{f_1, f_2, \dots, f_n\}$ ($f_\alpha \in \Lambda_\alpha$), such that $\sum_{\alpha=1}^n \|f_\alpha\|^p < \infty$, with a norm defined by the equation

$$\|\{f_1, f_2, \dots, f_n\}\| = \left(\sum_{\alpha=1}^n \|f_\alpha\|^p \right)^{1/p}.$$

$\Lambda_\alpha \cong \Lambda_\beta$ is to mean that there exists a one-to-one isometric mapping of Λ_α on Λ_β .

LEMMA 1.1.2. (a) $\sum_{\alpha=1}^n \oplus l_{p, n_\alpha} = l_{p, m}, n_\alpha = 1, 2, \dots, \infty$, if $\sum_{\alpha=1}^n n_\alpha = m$.

(b) $\sum_{\alpha=1}^n \oplus \Lambda_\alpha \cong L_p$ if $\Lambda_\alpha = L_p$, for each α .

The proof of this lemma may be left to the reader.

1.2. Let \mathfrak{M} be a closed linear manifold in Λ . We define a function $C(\mathfrak{M})$,

which takes on values in R as follows. If there exists no projection of Λ on \mathfrak{M} , then $C(\mathfrak{M}) = \infty$. Otherwise $C(\mathfrak{M}) = \text{g.l.b.} (|E|; E\Lambda = \mathfrak{M}, E^2 = E)$. Similarly we define the function $\bar{C}(\Lambda)$ as l.u.b. $(C(\mathfrak{M}), \mathfrak{M} \subseteq \Lambda)$.

LEMMA 1.2.1. *Let Λ_1 and Λ_2 be such that Λ_1 is equivalent [(B), p. 180] to a closed linear manifold \mathfrak{M} of Λ_2 . Let \mathfrak{M} be such that there exists a projection E of Λ_2 on \mathfrak{M} , with $|E| = 1$, \mathfrak{N} the set of f 's $\epsilon \Lambda$, for which $Ef = 0$. Let \mathfrak{P} be any closed linear manifold of Λ_2 , such that if $f \epsilon \mathfrak{P}$, then $f = g + h$, $g \epsilon \mathfrak{P} \cdot \mathfrak{M}$, $h \epsilon \mathfrak{P} \cdot \mathfrak{N}$. Let \mathfrak{P}_1 in Λ_1 be the manifold which corresponds to $\mathfrak{P} \cdot \mathfrak{M}$. Then $C(\mathfrak{P}_1) \leq C(\mathfrak{P})$.*

If $C(\mathfrak{P}) = \infty$, our statement is true. Suppose $C(\mathfrak{P})$ is $< \infty$. Let F be any projection of Λ_2 on \mathfrak{P} . Then EF is a projection on $\mathfrak{P} \cdot \mathfrak{M}$. For if $f_1 \epsilon \Lambda_2$, $f = Ff_1 = g + h$, $g \epsilon \mathfrak{P} \cdot \mathfrak{M}$, $h \epsilon \mathfrak{P} \cdot \mathfrak{N}$, and $EFf = g$ or the range of EF is included in $\mathfrak{P} \cdot \mathfrak{M}$. Also for every $h \epsilon \mathfrak{P} \cdot \mathfrak{N}$, we have $EFh = Eh = h$. This with our previous statement shows that $(EF)^2 = EF$ and that the range of EF is exactly $\mathfrak{P} \cdot \mathfrak{M}$.

Let $(EF)'$ be EF considered only on \mathfrak{M} . Obviously $(EF)'$ projects \mathfrak{M} on $\mathfrak{P} \cdot \mathfrak{M}$. Let G be the corresponding transformation on Λ_1 . Then $C(\mathfrak{P}_1) \leq |G| = |(EF)'| \leq |EF| \leq |E| \cdot |F| = |F|$ or $C(\mathfrak{P}_1) \leq |F|$. Since F was any projection on \mathfrak{P} , $C(\mathfrak{P}_1) \leq C(\mathfrak{P})$.

LEMMA 1.2.2. *If Λ_1 and Λ_2 are as in Lemma 1.2.1, $\bar{C}(\Lambda_1) \leq \bar{C}(\Lambda_2)$. In particular if $\Lambda_2 = \Lambda_0 \oplus \Lambda_1$, $\bar{C}(\Lambda_1) \leq \bar{C}(\Lambda_2)$.*

Let \mathfrak{P}_1 be any closed linear manifold of Λ_1 , \mathfrak{P} the corresponding set of elements in \mathfrak{M} . \mathfrak{P} is a closed linear manifold satisfying the conditions given in Lemma 1.2.1, since $\mathfrak{P} \cdot \mathfrak{M} = \mathfrak{P}$, $\mathfrak{P} \cdot \mathfrak{N} = \{0\}$. Lemma 1.2.1 now implies that $C(\mathfrak{P}_1) \leq C(\mathfrak{P}) \leq \bar{C}(\Lambda_2)$. But \mathfrak{P}_1 was any closed linear manifold in Λ_1 , hence $\bar{C}(\Lambda_1) \leq \bar{C}(\Lambda_2)$.

To show the second statement, we take $\mathfrak{M} \subseteq \Lambda_0 \oplus \Lambda_1$ as the set of elements $\{0, f\}$ of $\Lambda_0 \oplus \Lambda_1$, E as the transformation of $\Lambda_0 \oplus \Lambda_1$, such that $E\{f, g\} = \{0, g\}$. One readily sees that \mathfrak{M} is equivalent to Λ_1 and that E projects $\Lambda_0 \oplus \Lambda_1$ on \mathfrak{M} and $|E| = 1$. We may now apply the first part of this lemma to obtain the desired result.

LEMMA 1.2.3. *If $\Lambda \cong \sum_{\alpha=1}^{\infty} \Lambda_{\alpha}$ and k is $\lim \sup_{\alpha \rightarrow \infty} \bar{C}(\Lambda_{\alpha})$, then there exists a manifold $\mathfrak{P} \subseteq \Lambda$, such that $C(\mathfrak{P}) \geq k$.*

It follows from the definition of k , that if ϵ is > 0 , then there exists an infinite number of the α 's for which $r(\bar{C}(\Lambda_{\alpha}), k) \geq -\epsilon$. Thus we can find a sequence of integers $\{\alpha_i\}$ such that $\alpha_i < \alpha_{i+1}$, for which $r(\bar{C}(\Lambda_{\alpha_i}), k) \geq -2^{-i-1}$.

Now since $r(\bar{C}(\Lambda_{\alpha_i}), k) \geq -2^{-i-1}$, we can find a \mathfrak{P}_{α_i} , in Λ_{α_i} , such that $r(C(\mathfrak{P}_{\alpha_i}), k) > -2^{-i}$. Let \mathfrak{P} be the closed linear manifold consisting of those elements $\{f_1, f_2, f_3, \dots\} \epsilon \Lambda$, such that $f_{\beta} = 0$ if β is not $\epsilon \{\alpha_i\}$ and $f_{\alpha_i} \epsilon \mathfrak{P}_{\alpha_i}$. As we saw in the proof of Lemma 1.2.2, Λ_{α_i} and Λ are as Λ_1 and Λ_2 in Lemma 1.2.1

and it is easily seen that \mathfrak{B} satisfies the conditions given in Lemma 1.2.1 also. Thus Lemma 1.2.1 now implies that $C(\mathfrak{B})$ is $\geq C(\mathfrak{B}_{\alpha_i})$. Hence $r(C(\mathfrak{B}), k) \geq -2^{-i}$ for every i . This implies that $r(C(\mathfrak{B}), k) \geq 0$, $C(\mathfrak{B}) \geq k$.

1.3. We now prove the following lemma.

LEMMA 1.3.1. $\overline{C}(L_p) \geq \overline{C}(l_{p,\infty})$.

In (B), Theorem 9, Chapter XII, p. 206, it is shown that the manifold $\mathfrak{M} \subseteq L_p$, determined by the functions y_i , is equivalent to l_p when

$$y_i(t) = 2^{i/p} \text{ for } 1/2^i \leq t \leq 1/2^{i-1}, \quad y_i(t) = 0, \text{ otherwise.}$$

Now for any $z(t) \in L_p$, let

$$E(z(t)) = \sum_{i=1}^{\infty} \int_0^1 z(s) y_i^{p-1}(s) ds \cdot y_i(t).$$

Then by a direct calculation one can verify that $|E| = 1$ and that if $z \in \mathfrak{M}$ (i.e., if $z = \sum_i \alpha_i y_i$, $\sum |\alpha_i|^p < \infty$), then $Ez = z$. Hence E projects L_p on \mathfrak{M} and we may apply Lemma 1.2.2 so that it yields $\overline{C}(L_p) \geq \overline{C}(l_{p,\infty})$.

LEMMA 1.3.2. *There exists a linear manifold $\mathfrak{M} \subseteq L_p$, such that $C(\mathfrak{M}) = \overline{C}(L_p)$.*

This follows from Lemma 1.1.2, (b) (with $n = \infty$) and Lemma 1.2.3 for k is in this case $\overline{C}(L_p)$.

LEMMA 1.3.3. *There exists a linear manifold $\mathfrak{M} \subseteq l_{p,\infty}$, such that $C(\mathfrak{M}) = \overline{C}(l_{p,\infty})$.*

In Lemma 1.1.2, (a), let $n_\alpha = \infty$ for every α . Then apply Lemma 1.2.3.

LEMMA 1.3.4. $\overline{C}(l_{p,n}) \geq \overline{C}(l_{p,m})$ if $n \geq m$.

This follows from Lemma 1.1.2, (a) and Lemma 1.2.2.

THEOREM I. *$C(\mathfrak{M})$ and $\overline{C}(\Lambda)$ are to be as in §1.2. There exists an \mathfrak{M} in L_p , and an \mathfrak{N} in l_p , such that $C(\mathfrak{M}) = \overline{C}(L_p)$ and $C(\mathfrak{N}) = \overline{C}(l_p)$. Furthermore*

$$1 = \overline{C}(l_{p,1}) \leq \overline{C}(l_{p,2}) \leq \dots \leq \overline{C}(l_p) \leq \overline{C}(L_p).$$

The lemmas of this section imply this theorem.

Now if we are able to show that $\lim_{n \rightarrow \infty} \overline{C}(l_{p,n}) = \infty$, it follows from this theorem that $\overline{C}(L_p) = \overline{C}(l_p) = \infty$ and then in each of them we have a manifold \mathfrak{M} for which $C(\mathfrak{M}) = \infty$. Hence from the definition of $C(\mathfrak{M})$, we can answer problems (a) and (b) negatively. The next two chapters of this paper contain the proof of the fact that $\lim_{n \rightarrow \infty} \overline{C}(l_{p,n}) = \infty$.

CHAPTER 2. \mathfrak{M} IN SITUATION A

2.1. Let $f = \{a_1, \dots, a_n\}$ be an n -dimensional vector, which may be regarded as $\epsilon l_{p,n}$. We define for $k > 0$

$$\begin{aligned} \{f\}^k &= \{ |a_1|^k \text{ sign } a_1, \dots, |a_n|^k \text{ sign } a_n \} \\ [f]^k &= \{ |a_1|^k, \dots, |a_n|^k \}, \end{aligned}$$

which may be regarded as elements of $l_{p/k,n}$. If $g = \{b_1, \dots, b_n\}$, we define $(f, g) = \sum_{i=1}^n a_i b_i$. The linearity and homogeneity of this expression will be used without comment.

The following two lemmas can be easily shown.

LEMMA 2.1.1. *If $p > 1$,*

$$\frac{d}{dt} \|f + tg\|^p \Big|_{t=0} = p(\{f\}^{p-1}, g).$$

LEMMA 2.1.2. *If $p > 2$,*

$$\frac{d^2}{dt^2} \|f + tg\|^p = p(p - 1)([f + tg]^{p-2}, [g]^2).$$

We now prove

LEMMA 2.1.3. *If $p > 2$, and $(\{f\}^{p-1}, g) \geq 0$, then $\|f + g\|^p \geq \|f\|^p$.*

By Lemma 2.1.2, $H(t) = \|f + tg\|^p$ is convex in t and hence increasing for $t \geq 0$ since $dH/dt|_{t=0} \geq 0$ by Lemma 2.1.1.

2.2. If \mathfrak{M} is a linear manifold in $l_{p,n}$, let \mathfrak{M}^\perp consist of those elements $g \in l_{p',n}$, $1/p + 1/p' = 1$, for which $(f, g) = 0$ for all $f \in \mathfrak{M}$. If \mathfrak{M} is k -dimensional, it is well known that \mathfrak{M}^\perp is $(n - k)$ -dimensional and also that $(\mathfrak{M}^\perp)^\perp = \mathfrak{M}$. The following lemma is of a standard type in the theory of linear manifolds of a finite number of dimensions and the proof of it may be omitted.

LEMMA 2.2.1. *Let \mathfrak{M} be a k -dimensional linear manifold in $l_{p,n}$. Let ϕ_1, \dots, ϕ_k be k linearly independent elements of \mathfrak{M} . If E is a projection of $l_{p,n}$ on \mathfrak{M} , there exist k elements ψ_1, \dots, ψ_k , of $l_{p',n}$ such that for every $f \in l_{p,n}$,*

$$(\alpha) \quad Ef = \sum_{i=1}^k (\psi_i, f)\phi_i$$

and $(\psi_i, \phi_j) = \delta_{i,j}$. *If ψ_i 's with this last property are given, the E defined by (α) is a projection. If E' is any other projection of $l_{p,n}$ on \mathfrak{M} , then $\psi_i' = \psi_i + g_i$, $i = 1, \dots, k$, where g_i is $\in \mathfrak{M}^\perp$.*

2.3. If E is a linear transformation in $l_{p,n}$, we denote by E^* (the adjoint of E) the transformation in $l_{p',n}$, such that if g and $g^* \in l_{p',n}$, are related so that for every $f \in l_{p,n}$, $(Ef, g) = (f, g^*)$, then $E^*g = g^*$. It is well known that E^* is linear, $|E| = |E^*|$ and $(FE)^* = E^*F^*$.

LEMMA 2.3.1. *If \mathfrak{M} and E are as in Lemma 2.2.1, then*

$$E^*g = \sum_{i=1}^k (\phi_i, g)\psi_i$$

for all $g \in l_{p',n}$.

For every $f \in l_{p,n}$, we have

$$(Ef, g) = \left(\sum_{i=1}^k (\psi_i, f)\phi_i, g \right) = \sum_{i=1}^k (\psi_i, f)(\phi_i, g) = \left(f, \sum_{i=1}^k (\phi_i, g)\psi_i \right).$$

LEMMA 2.3.2. *If \mathfrak{M} and E are as in Lemma 2.2.1, then $1 - E^*$ is a projection on \mathfrak{M}^\perp .*

The range of $1 - E^*$ is \mathfrak{M}^\perp . For if $f = (1 - E^*)g$ and h is $\epsilon\mathfrak{M}$, then $Eh = h$ and

$$(h, f) = (h, (1 - E^*)g) = (h, g) - (h, E^*g) = (Eh, g) - (h, E^*g) = 0.$$

Hence f is $\epsilon\mathfrak{M}^\perp$ or the range of $1 - E^*$ is included in \mathfrak{M}^\perp . Furthermore by Lemma 2.3.1, if f is $\epsilon\mathfrak{M}^\perp$,

$$E^*f = \sum_{i=1}^k (\phi_i, f)\psi_i = 0$$

and $(1 - E^*)f = f$. This with our previous result shows that the range $1 - E^*$ is exactly \mathfrak{M}^\perp and $(1 - E^*)^2 = 1 - E^*$.

LEMMA 2.3.3. $\overline{C}(l_{p',n}) \leq \overline{C}(l_{p,n}) + 1$.

Let \mathfrak{M}' be any linear manifold of $l_{p',n}$. Let $\mathfrak{M} = \mathfrak{M}'^\perp$. Then if ϵ is > 0 , there exists a projection E of $l_{p,n}$ on \mathfrak{M} with $|E| \leq C(\mathfrak{M}) + \epsilon \leq \overline{C}(l_{p,n}) + \epsilon$. By Lemma 2.3.2, $1 - E^*$ is a projection on $\mathfrak{M}^\perp = (\mathfrak{M}'^\perp)^\perp = \mathfrak{M}'$. Thus $C(\mathfrak{M}') \leq |1 - E^*| \leq 1 + |E^*| = 1 + |E| \leq 1 + \overline{C}(l_{p,n}) + \epsilon$, which implies our lemma.

Of course p and p' are interchangeable and so we see that the answer to our question is the same for both p and p' . Thus we may confine ourselves to the case $p < 2$. This is not an essential step in our proof but merely a convenient one. We suppose from now on that p is < 2 .

2.4. We say that Situation A holds in a k -dimensional manifold \mathfrak{M} of $l_{p,n}$, if

(a) we have k linearly independent elements, $\phi_1, \dots, \phi_k, \epsilon\mathfrak{M}$, and k elements of $l_{p',n}$, ψ_1, \dots, ψ_k such that $(\phi_i, \psi_j) = \delta_{i,j}$ (the transformation E given by the equation $Ef = \sum_{i=1}^k (\psi_i, f)\phi_i$ is a projection of $l_{p,n}$ on \mathfrak{M});

(b) we have r elements h_1, \dots, h_r of \mathfrak{M} , with $\|h_i\| = 1$;

(c) there exists a constant $C > 1$, such that $\|E^*\{h_i\}^{p-1}\| = C$ for every i ;

(d) there exist r constants $c_1, \dots, c_r, c_i > 0$, such that for every $f \in \mathfrak{M}$ and $g \in \mathfrak{M}^\perp$

$$\sum_{i=1}^r c_i(\{h_i\}^{p-1}, f)(\{E^*\{h_i\}^{p-1}\}^{p'-1}, g) = 0.$$

LEMMA 2.4.1. *If \mathfrak{M} is in Situation A, then $C(\mathfrak{M}) \geq C$ (cf. (c) above).*

Since $|E| = |E^*|$, we must show that for every projection E' of $l_{p,n}$ on \mathfrak{M} , $|E'^*| \geq C$. Since $\|h_i\| = 1$ and hence $\|\{h_i\}^{p-1}\| = 1$, it will be sufficient to show that $\|E'^*\{h_i\}^{p-1}\| \geq C$ for at least one i .

Now

$$E'^*\{h_i\}^{p-1} = \sum_{j=1}^k (\{h_i\}^{p-1}, \phi_j)\psi'_j,$$

where $\psi'_j = \psi_j + g_j$, where g_j is an element of \mathfrak{M}^\perp (Lemmas 2.2.1 and 2.3.1). Let E_i be the projection of $l_{p,n}$ on \mathfrak{M} given by

$$E_i f = \sum_{i=1}^k (\psi_i + t g_i, f)\phi_i.$$

By Lemma 2.3.1,

$$E_i^* g = \sum_{i=1}^k (\phi_i, g)(\psi_i + t g_i)$$

and

$$\begin{aligned} E_i^*\{h_i\}^{p-1} &= \sum_{j=1}^k (\phi_j, \{h_i\}^{p-1})(\psi_j + t g_j) \\ &= \sum_{j=1}^k (\phi_j, \{h_i\}^{p-1})\psi_j + t \sum_{j=1}^k (\phi_j, \{h_i\}^{p-1})g_j \\ &= E^*\{h_i\}^{p-1} + t \sum_{j=1}^k (\phi_j, \{h_i\}^{p-1})g_j. \end{aligned}$$

Now by Lemma 2.1.1

$$\begin{aligned} \frac{d}{dt} \|E_i^*\{h_i\}^{p-1}\|_{t=0}^{p'} &= \left(\{E^*\{h_i\}^{p-1}\}^{p'-1}, \sum_{j=1}^k (\phi_j, \{h_i\}^{p-1})g_j \right) \\ &= \sum_{j=1}^k (\{E^*\{h_i\}^{p-1}\}^{p'-1}, g_j)(\phi_j, \{h_i\}^{p-1}). \end{aligned}$$

Since $g_j \in \mathfrak{M}^\perp$, $\phi_j \in \mathfrak{M}$, (d) implies that

$$\sum_{i=1}^r c_i \frac{d}{dt} \|E_i^*\{h_i\}^{p-1}\|_{t=0}^{p'} = \sum_{i=1}^r c_i \sum_{j=1}^k (\{E^*\{h_i\}^{p-1}\}^{p'-1}, g_j)(\phi_j, \{h_i\}^{p-1}) = 0.$$

Since $c_i > 0$, for $i = 1, \dots, r$ this implies that there must be an i' such that $d\|E_{i'}^*\{h_{i'}\}^{p-1}\|^{p'}/dt]_{t=0}$ is ≥ 0 . Hence by the above $d\|E^*\{h_{i'}\}^{p-1} + tg'_{i'}\|^{p'}/dt]_{t=0}$ is ≥ 0 , when $g'_{i'} = \sum_{i=1}^k (\phi_i, \{h_{i'}\}^{p-1})g_i$. Lemma 2.1.3 now yields

$$\|E^*\{h_{i'}\}^{p-1} + g'_{i'}\|^{p'} \geq \|E^*\{h_{i'}\}^{p-1}\|^{p'}$$

since $p' > 2$. But

$$E'^*\{h_{i'}\}^{p-1} = E_1^*\{h_{i'}\}^{p-1} = E^*\{h_{i'}\}^{p-1} + g'_{i'}$$

and $\|E^*\{h_{i'}\}^{p-1}\|^{p'} = C$. Substituting these values on both sides of our inequality, we get $\|E'^*\{h_{i'}\}^{p-1}\|^{p'} \geq Cp'$ or $\|E'^*\{h_{i'}\}^{p-1}\| \geq C$. As we remarked at the beginning of the proof this is sufficient.

CHAPTER 3. THE PRODUCT OF $l_{p,n}$ AND $l_{p,m}$

3.1. We define $l_{p,n} \otimes l_{p,m}$ as $l_{p,nm}$. If $f = \{a_1, \dots, a_n\} \in l_{p,n}$, and $g = \{b_1, \dots, b_m\} \in l_{p,m}$, we define $f \otimes g$ as $\{a_1b_1, a_1b_2, \dots, a_1b_m, a_2b_1, a_2b_2, \dots, a_2b_m, \dots, a_nb_1, a_nb_2, \dots, a_nb_m\}$ or if $f \otimes g = \{c_1, \dots, c_{nm}\}$, then $c_{(s-1)m+t} = a_s b_t$. The proofs of the following Lemmas 3.1.1–3.1.4 do not present any difficulty and may be left to the reader.

LEMMA 3.1.1. (i) $\|f \otimes g\| = \|f\| \cdot \|g\|$,

(ii) $(\phi_1 \otimes \phi_2, f \otimes g) = (\phi_1, f)(\phi_2, g)$,

(iii) $\{f \otimes g\}^k = \{f\}^k \otimes \{g\}^k$,

(iv) $\alpha(f^{(1)} \otimes g) + \beta(f^{(2)} \otimes g) = (\alpha f^{(1)} + \beta f^{(2)}) \otimes g$.

LEMMA 3.1.2. Let $f_r = \{a_{1,r}, \dots, a_{n,r}\}$, $r = 1, \dots, k$, $k \leq n$, be k linearly independent elements of $l_{p,n}$, and $g = \{b_{1,r}, \dots, b_{m,r}\}$ be k elements of $l_{p,m}$, such that $\sum_{r=1}^k f_r \otimes g_r = 0$. Then $g_r = 0$, $r = 1, \dots, k$.

LEMMA 3.1.3. Let f_r , $r = 1, \dots, k$, $k \leq n$, be k linearly independent elements of $l_{p,n}$ and for every $r = 1, \dots, k$ let $g_{r,s}$, $s = 1, \dots, k_r$, $k_r \leq m$, be k_r linearly independent elements of $l_{p,m}$. Then the set of elements $f_r \otimes g_{r,s}$, $r = 1, \dots, k$, $s = 1, \dots, k_r$ are linearly independent.

LEMMA 3.1.4. Let f_1, \dots, f_n be n linearly independent elements of $l_{p,n}$, g_1, \dots, g_m , m linearly independent elements $l_{p,m}$. Then the set of elements $f_i \otimes g_j$, $i = 1, \dots, n$, $j = 1, \dots, m$, determine $l_{p,mn}$.

3.2. Let $\mathfrak{M}^{(1)} \subseteq l_{p,n}$ and $\mathfrak{M}^{(2)} \subseteq l_{p,m}$ be linear manifolds. We define $\mathfrak{M}^{(1)} \otimes \mathfrak{M}^{(2)}$ as the linear manifold in $l_{p,mn}$ determined by the elements $f \otimes g$, $f \in \mathfrak{M}^{(1)}$, $g \in \mathfrak{M}^{(2)}$.

LEMMA 3.2.1. If $\phi_1^{(1)}, \dots, \phi_{k^{(1)}}^{(1)}$ is a set of linearly independent elements which determine $\mathfrak{M}^{(1)}$ and $\phi_1^{(2)}, \dots, \phi_{k^{(2)}}^{(2)}$ is a set of linearly independent elements which determine $\mathfrak{M}^{(2)}$, then $\phi_i^{(1)} \otimes \phi_j^{(2)}$, $i = 1, \dots, k^{(1)}$, $j = 1, \dots, k^{(2)}$, determine the manifold $\mathfrak{M}^{(1)} \otimes \mathfrak{M}^{(2)}$ which is $k^{(1)}k^{(2)}$ -dimensional.

The proof is easily derived from Lemma 3.1.3.

LEMMA 3.2.2. *Let $\mathfrak{M}^{(1)}$ and $\mathfrak{M}^{(2)}$ be as above. Then $(\mathfrak{M}^{(1)} \otimes \mathfrak{M}^{(2)})^\perp$ is determined by the elements of the form $f \otimes g$, where either $f \in \mathfrak{M}^{(1)\perp}$ or $g \in \mathfrak{M}^{(2)\perp}$.*

We first show that if $f \otimes g$ is such that either $f \in \mathfrak{M}^{(1)\perp}$ or $g \in \mathfrak{M}^{(2)\perp}$, then $f \otimes g \in (\mathfrak{M}^{(1)} \otimes \mathfrak{M}^{(2)})^\perp$. Indeed by the definition of $\mathfrak{M}^{(1)} \otimes \mathfrak{M}^{(2)}$ and the linearity of the operation $(\ , \)$, if $(\phi^{(1)} \otimes \phi^{(2)}, f \otimes g) = 0$ for all $\phi^{(1)} \in \mathfrak{M}^{(1)}$ and $\phi^{(2)} \in \mathfrak{M}^{(2)}$, then $f \otimes g \in (\mathfrak{M}^{(1)} \otimes \mathfrak{M}^{(2)})^\perp$. But Lemma 3.1.1, (ii) implies that $(\phi^{(1)} \otimes \phi^{(2)}, f \otimes g) = (\phi^{(1)}, f)(\phi^{(2)}, g)$. Since either $(\phi^{(1)}, f) = 0$ or $(\phi^{(2)}, g) = 0$, we obtain that $f \otimes g \in (\mathfrak{M}^{(1)} \otimes \mathfrak{M}^{(2)})^\perp$.

Now let $\tilde{\phi}_1^{(1)}, \dots, \tilde{\phi}_{n-k^{(1)}}^{(1)}$ and $\tilde{\phi}_1^{(2)}, \dots, \tilde{\phi}_{n-k^{(2)}}^{(2)}$ be sets of $n-k^{(1)}$ and $m-k^{(2)}$ linearly independent elements of $\mathfrak{M}^{(1)\perp}$ and $\mathfrak{M}^{(2)\perp}$ respectively. Let $\psi_1^{(1)}, \dots, \psi_{k^{(1)}}^{(1)}$ ($\psi_1^{(2)}, \dots, \psi_{k^{(2)}}^{(2)}$) be elements of $l_{p',n}$ ($l_{p',m}$) such that $\tilde{\phi}_i^{(1)}$'s and $\psi_j^{(1)}$'s together determine $l_{p',n}$ ($\tilde{\phi}_i^{(2)}$'s and $\psi_j^{(2)}$'s determine $l_{p',m}$). In Lemma 3.1.3, let $f_r = \psi_r^{(1)}$ for $r = 1, \dots, k^{(1)}$, $f_{k^{(1)}+t} = \tilde{\phi}_t^{(1)}$ for $t = 1, \dots, n-k^{(1)}$; $k = n$. For $r = 1, \dots, k^{(1)}$, let $k_r = m-k^{(2)}$, $g_{r,s} = \tilde{\phi}_s^{(2)}$, $s = 1, \dots, m-k^{(2)}$, and for $r = k+1, \dots, n$; $k = m$, $g_{r,s} = \psi_s^{(2)}$, $s = 1, \dots, k^{(2)}$, $g_{r,k^{(2)}+t} = \phi_t^{(2)}$, $t = 1, \dots, m-k^{(2)}$. Thus the $f_r \otimes g_{r,s}$ are linearly independent and such that either $f_r \in \mathfrak{M}^{(1)\perp}$, or $g_{r,s} \in \mathfrak{M}^{(2)\perp}$. Since there are $k^{(1)}(m-k^{(2)}) + (n-k^{(1)})m = mn - k^{(1)}k^{(2)}$ of them and the dimensionality of $\mathfrak{M}^{(1)} \otimes \mathfrak{M}^{(2)}$ is $k^{(1)}k^{(2)}$ by Lemma 3.2.1, the $f_r \otimes g_{r,s}$ determine $(\mathfrak{M}^{(1)} \otimes \mathfrak{M}^{(2)})^\perp$.

3.3. Let $T^{(1)}$ be a linear transformation in $l_{p,n}$ and $T^{(2)}$ a linear transformation in $l_{p,m}$. Let $\phi_1^{(1)}, \dots, \phi_n^{(1)}$ be n linearly independent elements of $l_{p,n}$ and $\phi_1^{(2)}, \dots, \phi_m^{(2)}$, m linearly independent elements of $l_{p,m}$. Then the elements $\phi_i^{(1)} \otimes \phi_j^{(2)}$, $i = 1, \dots, n, j = 1, \dots, m$ determine $l_{p,mn}$ by Lemma 3.1.4 and are linearly independent. Hence given any set of mn elements $f_{i,j} \in l_{p,mn}$, there exists a unique linear transformation T' such that

$$T'(\phi_i^{(1)} \otimes \phi_j^{(2)}) = f_{i,j} \quad (i = 1, \dots, n, j = 1, \dots, m).$$

Now let $f_{i,j} = T^{(1)}\phi_i^{(1)} \otimes T^{(2)}\phi_j^{(2)}$ and denote the corresponding T' by $T^{(1)} \otimes T^{(2)}$. Apparently this definition of $T^{(1)} \otimes T^{(2)}$ depends on the choice of the $\phi_i^{(1)}$'s and $\phi_j^{(2)}$'s, but this is not the case as is shown by the following

LEMMA 3.3.1. *If $f \in l_{p,n}$, $g \in l_{p,m}$, then $T^{(1)} \otimes T^{(2)} f \otimes g = T^{(1)} f \otimes T^{(2)} g$. Thus $T^{(1)} \otimes T^{(2)}$ does not depend on the choice of the $\{\phi_i^{(1)}\}$ or the $\{\phi_j^{(2)}\}$.*

The proof follows immediately from the definition of $T^{(1)} \otimes T^{(2)}$ and Lemma 3.1.1, (iv).

We also have

LEMMA 3.3.2. *A linear transformation T' of $l_{p,mn}$, equals $T^{(1)} \otimes T^{(2)}$ if and only if $T' f \otimes g = T^{(1)} f \otimes T^{(2)} g$ for every $f \in l_{p,n}$, and $g \in l_{p,m}$.*

The sufficiency of the condition follows from the definition. Lemma 3.3.1 implies its necessity.

LEMMA 3.3.3. $(T^{(1)} \otimes T^{(2)})^* = (T^{(1)})^* \otimes (T^{(2)})^*$

By Lemma 3.3.2, it suffices to show that if $f \in l_{p',n}$, $g \in l_{p',m}$, then $(T^{(1)} \otimes T^{(2)})^* f \otimes g = T^{(1)*} f \otimes T^{(2)*} g$. Now if $h \in l_{p,mn}$, it follows from Lemma 3.1.4, that $h = \sum_{i=1}^n \sum_{j=1}^m a_{i,j} \phi_i^{(1)} \otimes \phi_j^{(2)}$ and by the definition of $T^{(1)} \otimes T^{(2)}$, $T^{(1)} \otimes T^{(2)} h = \sum_{i=1}^n \sum_{j=1}^m a_{i,j} T^{(1)} \phi_i^{(1)} \otimes T^{(2)} \phi_j^{(2)}$.

By the definition of T^* (cf. §2.3) and Lemma 3.1.1, (ii), we have

$$\begin{aligned} (T^{(1)} \otimes T^{(2)} h, f \otimes g) &= \sum_{i=1}^n \sum_{j=1}^m a_{i,j} (T^{(1)} \phi_i^{(1)} \otimes T^{(2)} \phi_j^{(2)}, f \otimes g) \\ &= \sum_{i=1}^n \sum_{j=1}^m a_{i,j} (T^{(1)} \phi_i^{(1)}, f) (T^{(2)} \phi_j^{(2)}, g) \\ &= \sum_{i=1}^n \sum_{j=1}^m a_{i,j} (\phi_i^{(1)}, T^{(1)*} f) (\phi_j^{(2)}, T^{(2)*} g) \\ &= \sum_{i=1}^n \sum_{j=1}^m a_{i,j} (\phi_i^{(1)} \otimes \phi_j^{(2)}, T^{(1)*} f \otimes T^{(2)*} g) \\ &= \left(\sum_{i=1}^n \sum_{j=1}^m a_{i,j} \phi_i^{(1)} \otimes \phi_j^{(2)}, T^{(1)*} f \otimes T^{(2)*} g \right) \\ &= (h, T^{(1)*} f \otimes T^{(2)*} g). \end{aligned}$$

Or for every h of $l_{p,mn}$,

$$(T^{(1)} \otimes T^{(2)} h, f \otimes g) = (h, T^{(1)*} f \otimes T^{(2)*} g).$$

The definition of T^* , then implies

$$(T^{(1)} \otimes T^{(2)})^* f \otimes g = T^{(1)*} f \otimes T^{(2)*} g$$

which is the desired result.

3.4. We have the following lemma.

LEMMA 3.4.1. (i) If $E^{(1)}$ is a projection on $\mathfrak{M}^{(1)} \subseteq l_{p,n}$, and $E^{(2)}$ a projection on $\mathfrak{M}^{(2)} \subseteq l_{p,m}$, then $E^{(1)} \otimes E^{(2)}$ is a projection of $l_{p,mn}$ on $\mathfrak{M}^{(1)} \otimes \mathfrak{M}^{(2)}$.

(ii) Let $\phi_i^{(1)}$ and $\psi_i^{(1)}$ ($\phi_i^{(2)}$ and $\psi_i^{(2)}$) be in the same relation to $E^{(1)}$ ($E^{(2)}$) as ϕ_i and ψ_i are to E in Lemma 2.2.1, i.e.,

$$E^{(1)} f = \sum_{i=1}^{k^{(1)}} (\psi_i^{(1)}, f) \phi_i^{(1)}; \quad E^{(2)} f = \sum_{i=1}^{k^{(2)}} (\psi_i^{(2)}, f) \phi_i^{(2)}.$$

If E is the transformation on $l_{p,mn}$ defined by the equation

$$Eh = \sum_{i=1}^{k^{(1)}} \sum_{j=1}^{k^{(2)}} (\psi_i^{(1)} \otimes \psi_j^{(2)}, h) \phi_i^{(1)} \otimes \phi_j^{(2)},$$

then E is a projection of $l_{p,mn}$ on $\mathfrak{M}^{(1)} \otimes \mathfrak{M}^{(2)}$ and $E = E^{(1)} \otimes E^{(2)}$.

(iii) $E^* = E^{(1)*} E^{(2)*}$.

Since (ii) implies (i) and (ii) and Lemma 3.3.3 implies (iii), we need only prove (ii).

We have $(\psi_i^{(1)} \otimes \psi_j^{(2)}, \phi_l^{(1)} \otimes \phi_k^{(2)}) = (\psi_i^{(1)}, \phi_l^{(1)}) (\psi_j^{(2)}, \phi_k^{(2)}) = \delta_{i,l} \delta_{j,k}$, by Lemma 3.1.1 and Lemma 2.2.1. By Lemma 3.2.1 the $\phi_i^{(1)} \otimes \phi_k^{(2)}$ determine $\mathfrak{M}^{(1)} \otimes \mathfrak{M}^{(2)}$. Lemma 2.2.1 now implies that E is a projection on $\mathfrak{M}^{(1)} \otimes \mathfrak{M}^{(2)}$.

It remains to show that $E = E^{(1)} \otimes E^{(2)}$. If $f \in l_{p,n}$, $g \in l_{p,m}$, then by Lemma 3.1.1, (ii) and (iv), and Lemma 2.2.1,

$$\begin{aligned} Ef \otimes g &= \sum_{i=1}^{k^{(1)}} \sum_{j=1}^{k^{(2)}} (\psi_i^{(1)} \otimes \psi_j^{(2)}, f \otimes g) \phi_i^{(1)} \otimes \phi_j^{(2)} \\ &= \sum_{i=1}^{k^{(1)}} \sum_{j=1}^{k^{(2)}} (\psi_i^{(1)}, f) (\psi_j^{(2)}, g) \phi_i^{(1)} \otimes \phi_j^{(2)} \\ &= \left(\sum_{i=1}^{k^{(1)}} (\psi_i^{(1)}, f) \phi_i^{(1)} \right) \otimes \left(\sum_{j=1}^{k^{(2)}} (\psi_j^{(2)}, g) \phi_j^{(2)} \right) = E^{(1)} f \otimes E^{(2)} g. \end{aligned}$$

Lemma 3.3.2 now implies that $E = E^{(1)} \otimes E^{(2)}$.

3.5. Next we prove

LEMMA 3.5.1. *If $\mathfrak{M}^{(1)}$ in $l_{p,n}$ and $\mathfrak{M}^{(2)}$ in $l_{p,m}$ are in Situation A (cf. §2.4), then $\mathfrak{M}^{(1)} \otimes \mathfrak{M}^{(2)}$ is in Situation A with*

- (a) $\phi_{(s-1)k^{(1)}+t} = \phi_t^{(1)} \otimes \phi_j^{(2)}, \quad t = 1, \dots, k^{(1)}, \quad s = 1, \dots, k^{(2)},$
 $\psi_{(s-1)k^{(1)}+t} = \psi_t^{(1)} \otimes \psi_s^{(2)}, \quad t = 1, \dots, k^{(1)}, \quad s = 1, \dots, k^{(2)};$
- (b) $h_{(s-1)r^{(1)}+t} = h_t^{(1)} \otimes h_s^{(2)}, \quad t = 1, \dots, r^{(1)}, \quad s = 1, \dots, r^{(2)};$
- (c) $C = C^{(1)} C^{(2)};$
- (d) $c_{(s-1)r^{(1)}+t} = c_t^{(1)} c_s^{(2)}, \quad t = 1, \dots, r^{(1)}; \quad s = 1, \dots, r^{(2)}.$

That the $\phi_i^{(1)} \otimes \phi_k^{(2)}, t=1, \dots, k^{(1)}, s=1, \dots, k^{(2)}$ are linearly independent and determine $\mathfrak{M}^{(1)} \otimes \mathfrak{M}^{(2)}$ has been shown in Lemma 3.1.3 and Lemma 3.2.1. The remaining statements of (a) were shown in the proof of Lemma 3.4.1.

To prove (b) we have $h_i^{(1)} \otimes h_s^{(2)} \in \mathfrak{M}^{(1)} \otimes \mathfrak{M}^{(2)}$ by definition (cf., §3.2). Also by Lemma 3.1.1 $\|h_i^{(1)} \otimes h_s^{(2)}\| = \|h_i^{(1)}\| \cdot \|h_s^{(2)}\| = 1$.

Now consider (c). If $i = (s-1)r^{(1)} + t$, then by Lemma 3.1.1, (ii); Lemma 3.4.1, (iii); Lemma 3.3.2; Lemma 3.1.1, (i); and §2.4, (c),

$$\begin{aligned} \|E^* \{h_i\}^{p-1}\| &= \|E^* \{h_i^{(1)} \otimes h_s^{(2)}\}^{p-1}\| = \|E^* \{h_i^{(1)}\}^{p-1} \otimes \{h_s^{(2)}\}^{p-1}\| \\ &= \|E^{(1)*} \otimes E^{(2)*} \{h_i^{(1)}\}^{p-1} \otimes \{h_s^{(2)}\}^{p-1}\| \\ &= \|E^{(1)*} \{h_i^{(1)}\}^{p-1} \otimes E^{(2)*} \{h_s^{(2)}\}^{p-1}\| \\ &= \|E^{(1)*} \{h_i^{(1)}\}^{p-1}\| \cdot \|E^{(2)*} \{h_s^{(2)}\}^{p-1}\|. \end{aligned}$$

We now prove (d). If $h \in \mathfrak{M}^{(1)} \otimes \mathfrak{M}^{(2)\perp}$, then by Lemma 3.2.2, h is a linear combination of elements in the form $f^{(1)} \otimes g^{(2)}$, where either $f^{(1)}$ is in $\mathfrak{M}^{(1)\perp}$ or $g^{(2)}$ is in $\mathfrak{M}^{(2)\perp}$. Hence since

$$\sum_{i=1}^{r^{(1)r^{(2)}}} c_i(\{h_i\}^{p-1}, f)(h, \{E^* \{h_i\}^{p-1}\}^{p'-1}),$$

$f \in \mathfrak{M}^{(1)} \otimes \mathfrak{M}^{(2)}$, $h \in (\mathfrak{M}^{(1)} \otimes \mathfrak{M}^{(2)\perp})^\perp$, is linear in h , it is enough to show (d) for h in the form $f^{(1)} \otimes g^{(2)}$, where either $f^{(1)}$ is in $\mathfrak{M}^{(1)\perp}$ or $g^{(2)}$ is in $\mathfrak{M}^{(2)\perp}$. It is also linear in f ; hence by §3.2 it suffices to show (d) for f in the form $\phi^{(1)} \otimes \phi^{(2)}$, $\phi_1 \in \mathfrak{M}^{(1)}$, $\phi_2 \in \mathfrak{M}^{(2)}$.

Now it was shown in the proof of (c) above that $E^* \{h_i\}^{p-1} = E^{(1)*} \{h_i^{(1)}\}^{p-1} \otimes E^{(2)*} \{h_s^{(2)}\}^{p-1}$. Then by Lemma 3.1.1, (iii),

$$\{E^* \{h_i\}^{p-1}\}^{p'-1} = \{E^{(1)*} \{h_i^{(1)}\}^{p-1}\}^{p'-1} \otimes \{E^{(2)*} \{h_s^{(2)}\}^{p-1}\}^{p'-1}$$

and $\{h_i\}^{p-1} = \{h_i^{(1)}\}^{p-1} \otimes \{h_s^{(2)}\}^{p-1}$. Hence by Lemma 3.1.1, (ii), we see that

$$\begin{aligned} &\sum_{i=1}^{r^{(1)r^{(2)}}} c_i(\{h_i\}^{p-1}, \phi^{(1)} \otimes \phi^{(2)})(f^{(1)} \otimes g^{(2)}, \{E^* \{h_i\}^{p-1}\}^{p'-1}) \\ &= \left(\sum_{i=1}^{r^{(1)}} c_i^{(1)}(\{h_i^{(1)}\}^{p-1}, \phi^{(1)})(f^{(1)}, \{E^{(1)*} \{h_i^{(1)}\}^{p-1}\}^{p'-1}) \right) \\ &\quad \times \left(\sum_{s=1}^{r^{(2)}} c_s^{(2)}(\{h_s^{(2)}\}^{p-1}, \phi^{(2)})(g^{(2)}, \{E^{(2)*} \{h_s^{(2)}\}^{p-1}\}^{p'-1}) \right). \end{aligned}$$

Since either $f^{(1)} \in \mathfrak{M}^{(1)\perp}$, or $g^{(2)} \in \mathfrak{M}^{(2)\perp}$, this is zero for $\mathfrak{M}^{(1)}$ and $\mathfrak{M}^{(2)}$ are in Situation A.

3.6. Now it follows from Lemma 2.4.1 and Lemma 3.5.1, that we can show that $\lim_{n \rightarrow \infty} (\overline{C}(l_{p,n})) = \infty$ if we can find a \mathfrak{M} in $l_{p,n}$ in Situation A (cf. §2.4, (c)). For let N be any integer > 0 . Then using Lemma 3.5.1 we can find a manifold \mathfrak{M}_N in l_{p,n^N} in Situation A with $C_{\mathfrak{M}_N} = C_{\mathfrak{M}}^N$. Lemma 2.4.1 now implies that $C(\mathfrak{M}_N) \geq C_{\mathfrak{M}}^N$ and since

$$\overline{C}(l_{p,n^N}) \geq C(\mathfrak{M}_N), \quad \lim_{n \rightarrow \infty} (\overline{C}(l_{p,n})) = \infty.$$

As we remarked in §1.3, this implies that the answer to both (a) and (b) is negative.

Let \mathfrak{M} be the manifold in $l_{p,3}$, determined by the vectors $(1, 1, 0)$ and $(0, 1, 1)$. Let

$$\begin{aligned} \phi_1 &= (2^{-1/p}, 2^{-1/p}, 0), & \phi_2 &= (0, 2^{-1/p}, 2^{-1/p}), \\ \psi_1 &= (2^{1/p+1}/3, 2^{1/p}/3, -2^{1/p}/3), & \psi_2 &= (-2^{1/p}/3, 2^{1/p}/3, 2^{1/p+1}/3); \end{aligned}$$

if $\alpha = 1/(2 + 2^p)^{1/p}$, $h_1 = (\alpha, -\alpha, -2\alpha)$, $h_2 = (\alpha, 2\alpha, \alpha)$, $h_3 = (2\alpha, \alpha, -\alpha)$. Also

$$\begin{aligned} C &= ((2^{p'-1} + 1)/3)^{1/p'}((2^{p-1} + 1)/3)^{1/p}, \\ c_1 &= c_2 = c_3 = 1. \end{aligned}$$

We show that \mathfrak{M} is in Situation A (§2.4) and thus complete the proof.

We have (a) $(\phi_i, \psi_j) = \delta_{i,j}$.

(b) $h_1 = \alpha 2^{1/p}(\phi_1 - 2\phi_2)$, $h_2 = \alpha 2^{1/p}(\phi_1 + \phi_2)$, $h_3 = \alpha 2^{1/p}(2\phi_1 - \phi_2)$, and thus h_i is $\epsilon\mathfrak{M}$, $i = 1, 2, 3$. We also have $\|h_i\| = 1$, $i = 1, 2, 3$.

Before showing (c) and (d) we make certain calculations. From the definitions in §2.1, we get $\{h_1\}^{p-1} = \alpha^{p-1}(1, -1, -2^{p-1})$, $\{h_2\}^{p-1} = \alpha^{p-1}(1, 2^{p-1}, 1)$, $\{h_3\}^{p-1} = (2^{p-1}, 1, 1)$, $(\{h_1\}^{p-1}, \phi_1) = (\{h_3\}^{p-1}, \phi_2) = 0$, $-(\{h_1\}^{p-1}, \phi_2) = (\{h_2\}^{p-1}, \phi_1) = (\{h_2\}^{p-1}, \phi_2) = (\{h_3\}^{p-1}, \phi_1) = 2^{-1}(1 + 2^{p-1})^{1/p}$.

By Lemma 2.3.1

$$\begin{aligned} E^*\{h_1\}^{p-1} &= (\{h_1\}^{p-1}, \phi_1)\psi_1 + (\{h_1\}^{p-1}, \phi_2)\psi_2 = (\{h_1\}^{p-1}, \phi_2)\psi_2 \\ &= -3^{-1}2^{-1/p'}(2^{p-1} + 1)^{1/p}(-1, 1, 2), \\ E^*\{h_2\}^{p-1} &= (\{h_2\}^{p-1}, \phi_1)\psi_1 + (\{h_2\}^{p-1}, \phi_2)\psi_2 = 2^{-1}(2^{p-1} + 1)^{1/p}(\psi_1 + \psi_2) \\ &= 3^{-1}2^{-1/p'}(2^{p-1} + 1)^{1/p}(1, 2, 1). \end{aligned}$$

Similarly

$$E^*\{h_3\}^{p-1} = 3^{-1}2^{-1/p'}(2^{p-1} + 1)^{1/p}(2, 1, -1).$$

Finally (cf. §2.1)

$$\begin{aligned} \{E^*\{h_1\}^{p-1}\}^{p'-1} &= -3^{-(p'-1)}2^{-(p'-1)/p'}(2^{p-1} + 1)^{(p'-1)/p}(-1, 1, 2^{p'-1}) \\ &= -K(-1, 1, 2^{p'-1}) \\ \{E^*\{h_2\}^{p-1}\}^{p'-1} &= K(1, 2^{p'-1}, 1) \\ \{E^*\{h_3\}^{p-1}\}^{p'-1} &= K(2^{p'-1}, 1, -1). \end{aligned}$$

(c) By direct calculation, we obtain $\|E^*\{h_1\}^{p-1}\| = \|E^*\{h_2\}^{p-1}\| = \|E^*\{h_3\}^{p-1}\| = C$ using the above. For $p \neq 2$, C is > 1 since by the Hölder inequality,

$$6C = (2^p + 2)^{1/p}(2^{p'} + 2)^{1/p'} \geq 2 \cdot 2 + 2^{1/p}2^{1/p'} = 6,$$

where the equality sign holds only for $p = 2$.

(d) Now if $f \in \mathfrak{M}^\perp$, then $f = k\tilde{\phi}$, where $\tilde{\phi} = (1, -1, 1)$. Thus

$$\begin{aligned} (\{E^*\{h_1\}^{p-1}\}^{p'-1}, f) &= Kk(2 - 2^{p'-1}), \\ (\{E^*\{h_2\}^{p-1}\}^{p'-1}, f) &= Kk(2 - 2^{p'-1}), \\ (\{E^*\{h_3\}^{p-1}\}^{p'-1}, f) &= Kk(2^{p'-1} - 2). \end{aligned}$$

We can now verify by a direct calculation that (d) holds.

Conclusion. Our results permit us to conclude that

There exists a manifold \mathfrak{M}_0 in l_p and L_p such that there exists no biorthogonal set $\{\phi_i, \psi_i\}$ where $\{\phi_i\}$ is a basis for \mathfrak{M}_0 (cf. (B), Chapter VII, p. 110, §3), while the expansion

$$(*) \quad \sum_{i=1}^{\infty} a_i \phi_i, \quad a_i = (f, \psi_i),$$

converges for each $f \in l_p$ or L_p .

Let us suppose that (*) converges for every f . Let \mathfrak{M} be the manifold determined by the ϕ_i 's. The ϕ_i 's are a basis for \mathfrak{M} (cf. (B), loc. cit.) for if $f \in \mathfrak{M}$, then

$$f = \sum_{i=1}^{\infty} a_i \phi_i, \quad a_i = (f, \psi_i)$$

by (B), Chapter VII, Theorem 2, p. 107.

We will show that under these circumstances $C(\mathfrak{M})$ is $< \infty$. For let E be the transformation defined by the equation

$$Ef = \sum_{i=1}^{\infty} (f, \psi_i) \phi_i.$$

Ef is defined for every f since we assume that the series is convergent for every f . The same assumption implies that E is limited for the partial sums are uniformly limited (cf. (B), Chapter VII, Theorem 2 and Theorem 5). E is obviously additive and homogeneous. If $f \in \mathfrak{M}$, $Ef = f$ by the above and the range of E is included in \mathfrak{M} , $E\Lambda = \mathfrak{M}$ and $E^2 = E$. Thus E is a projection of Λ on \mathfrak{M} . Hence $C(\mathfrak{M})$ is $< \infty$.

Our construction also permits us to show that no statement concerning the norms of the ϕ_i and ψ_i will insure convergence by itself. We can assume that $\|\phi_i\| = 1$ for every i . The least possible value for $\|\psi_i\|$ is then 1 since $(\phi_i, \psi_i) = 1$ and $|\langle \phi_i, \psi_i \rangle| \leq \|\phi_i\| \cdot \|\psi_i\|$. We will show that *there exists in both l_p and L_p a biorthogonal set $\{\phi_i, \psi_i\}$ for which $\|\phi_i\| = \|\psi_i\| = 1$ and for which the associated expansion (*) does not always converge.*

It is a consequence of the proof of Lemma 1.3.1 that if such a set exists in l_p , there must be a similar one in L_p . So we need only consider l_p . Owing to the nature of biorthogonal sets in l_p , we need only consider the case $1 < p < 2$.

Let \mathfrak{M} be the manifold of §3.6, and let $f_1 = (2^{-1/p}, 2^{-1/p}, 0) = \phi_1, f_2 = (\alpha, -\alpha, -2\alpha) = h_1$. We have $(f_1, \{f_2\}^{p-1}) = (\{f_1\}^{p-1}, f_2) = 0; (f_1, \{f_1\}^{p-1}) = (f_2, \{f_2\}^{p-1}) = 1, \|f_1\| = \|f_2\| = \|\{f_1\}^{p-1}\| = \|\{f_2\}^{p-1}\| = 1$. Of course f_1 and $f_2 \in l_{p,3}, \{f_1\}^{p-1}$ and $\{f_2\}^{p-1} \in l_{p',3}$.

We define $f_{i_1} \otimes f_{i_2} \otimes \dots \otimes f_{i_n}, i_j = 1, 2; j = 1, \dots, n$, as an element of $l_{p,3^n}$ as follows: $f_{i_1} \otimes f_{i_2}$ in $l_{p,3^2}$ has already been defined (cf. §3.1). Let us suppose that $f_{i_1} \otimes \dots \otimes f_{i_{n-1}}$ in $l_{p,3^{n-1}}$ has been defined. We define $f_{i_1} \otimes f_{i_2} \otimes \dots \otimes f_{i_{n-1}} \otimes f_{i_n}$ as $(f_{i_1} \otimes \dots \otimes f_{i_{n-1}}) \otimes f_{i_n}$ in $l_{p,3^n}$ using §3.1. Let $\mathfrak{M}_1 = \mathfrak{M}, \mathfrak{M}_n$ in $l_{p,3^n}$ be $\mathfrak{M}_{n-1} \otimes \mathfrak{M}$. Then by successive applications of Lemma 3.2.1, we see that the set of elements $f_{i_1} \otimes \dots \otimes f_{i_n}$ determine \mathfrak{M}_n .

By Lemma 1.1.2, $l_p = \sum_{-1}^{\infty} l_{p,3^\alpha}$. Let \mathfrak{B} be the closed linear manifold in $\sum_{-1}^{\infty} l_{p,3^\alpha}$ consisting of those elements $\{g_1, g_2, \dots\}$ for which $g_\alpha \in \epsilon \mathfrak{M}_\alpha$ for every α . Let S consist of those elements which are such that every $g_\alpha = 0$ except for one g_n and $g_n = f_{i_1} \otimes \dots \otimes f_{i_n}$. Let S' consist of elements of $l_{p'}$ in the form $\{g\}^{p-1}, g \in S$. Since as we have seen above the $f_{i_1} \otimes \dots \otimes f_{i_n}$ determine \mathfrak{M}_n, S determines \mathfrak{B} .

Now the sets S and S' are denumerable and it is easily seen by using Lemma 3.1.1 that with a suitable enumeration they form a biorthogonal set with $\|\phi_i\| = \|\psi_i\| = 1$. Since S determines \mathfrak{B} , we see from the above that this series (*) cannot converge always if $C(\mathfrak{B}) = \infty$.

Let C be as in §3.6, (c). By repeated applications of Lemma 3.5.1 and then using Lemma 2.4.1, one may prove that $C(\mathfrak{M}_n) \geq C^n$. It follows from the proof of Lemma 1.2.3 that $C(\mathfrak{B}) \geq C(\mathfrak{M}_n) \geq C^n$ for every n and since $C > 1, p \neq 2$ this implies that $C(\mathfrak{B}) = \infty$. As we have remarked above, this proves our statement.

Incidentally we have explicitly constructed a manifold \mathfrak{B} in l_p , for which there exists no projection. Lemma 1.3.1 indicates how we can find a \mathfrak{B} in L_p with the corresponding property.

In L_p , the space of complex-valued functions whose p th power is summable, the situation is the same. As pointed out in a previous paper by the writer,† the theorems given in (B) and used here can be generalized to the complex case. Chapter 1 of this paper also falls into this category. Some variations are necessary in Chapters 2 and 3 but they are not basic.

Finally it should be pointed out that the negative answer to (a) and (b) precludes the possibility of a spectral theory in l_p and L_p similar to the theory of self-adjoint operators in Hilbert space.

† These Transactions, vol. 39 (1936), pp. 83-100.