

ON n -ARC CONNECTEDNESS

BY

G. T. WHYBURN

1. Introduction. In this paper a simple elementary proof will be given for the well known theorem that any two points a and b of a locally connected continuum M which are not separated in M by any set of less than n points can be joined in M by a set of n independent simple arcs, that is, arcs intersecting by pairs in just $a+b$. The theorem in this form was first proven by Rutt [1]⁽¹⁾ for plane continua M and for general M by Nöbeling [2] and Menger [3] and has been extended by Zippin [4]. Our proof represents an extension to arbitrary n of the inductive type of reasoning used by the author [5] in simplifying the proof of the cyclic connectedness theorem (which is the case $n=2$ of the above theorem). It holds for locally compact M , as in Zippin's generalized form

Our terminology in the main will be that of Menger-Nöbeling and of the author's book [5]. We recall that a *continuum* is a connected compact metric space and a *generalized continuum* is the same except that compactness is replaced by local compactness. A metric set W is *regular* at a point $x \in W$ if x has arbitrarily small neighborhoods in W whose boundaries are finite sets. A subset X of W *separates* two disjoint sets P and Q in W provided $W-X$ is the sum of two separated sets one containing P and the other Q . Such a set X *separates P and Q in the broad sense* in W provided $W-X$ is the sum of two separated sets (possibly empty), one containing $P-X \cdot P$ and the other $Q-X \cdot Q$. The set W is *n -point connected* between P and Q provided no set of less than n points separates P and Q in W , and W is *n -point strongly connected* between P and Q provided no set of less than n points separates P and Q in the broad sense in W . An arc *joins* two sets if it has an end point in each of them.

2. Theorem⁽²⁾. *If the locally connected separable complete metric space M is n -point strongly connected between two disjoint closed sets P and Q , then M contains n disjoint arcs joining P and Q .*

Presented to the Society, April 26, 1947; received by the editors March 5, 1947.

⁽¹⁾ Numbers in brackets refer to the bibliography at the end of the paper.

⁽²⁾ This is called the *second n -arc theorem* by Menger [3]. It seems to have been proved heretofore only for locally compact spaces M , except in the cases $n=1$, where it becomes the arcwise connectedness theorem established for complete spaces by Moore [6] and Menger [7], and $n=2$, where it was proved for complete spaces by the author [8]. It should be noted that the main theorem, §3, *does not hold* for complete spaces even in case $n=2$, as was shown by the author in [8].

The proof will be by induction on n . The case $n = 1$ is an immediate consequence of the arcwise connectedness theorem, since then some component of M must intersect both P and Q . Suppose the theorem holds for all positive integers less than n .

Let S denote the set of all $x \in M$ such that there exists a sum S_x of n disjoint arcs, $n - 1$ of which join P and Q and the remaining one px joins P and x . Then S is open in M and nonempty. For if $x \in S$ and R is any region in M with $x \in R \subset M - S_x + px$, the arc px can be extended in R to any $y \in R$ thus giving a system S_y for y so that $S \supset R$. Also if $y \in M$, $M - y$ is $(n - 1)$ -point strongly connected between $(P + y) - y$ and $(Q + y) - y$, because if a set X of less than $n - 1$ points separated these two sets in $M - y$ in the broad sense, $X + y$ would similarly separate P and Q in M . Hence $P \subset S$ since for any y , $M - y$ contains a set of $n - 1$ disjoint arcs joining P and Q .

The set S is also closed in M . For let y be any limit point of S . Then as just shown, $M - y$ contains a sum A of $n - 1$ disjoint arcs $\alpha_1, \alpha_2, \dots, \alpha_{n-1}$ each joining P and Q . Let R be a region about y with $\bar{R} \subset M - (P + A)$. Let $x \in R \cdot S$ and let S_x be a sum of n arcs determined by x as above. Then $S_x - R \cdot S_x$ contains a sum B of n disjoint arcs $\beta_1, \beta_2, \dots, \beta_n$ each joining P and $Q + \bar{R}$. Let $\alpha_i = s_i q_i$, $\beta_i = p_i r_i$ where $p_i, s_i \in P$, $q_i \in Q$, $r_i \in (Q + \bar{R})$. On α_i , in the order q_i, s_i let q_i^1 be the first point of $B + P$, let $Q_1 = \sum_1^{n-1} q_i^1$ and $A_1 = \sum_1^{n-1} q_i q_i^1$. On β_i , in the order p_i, r_i let p_i^1 be the first point of $Q + \bar{R} + Q_1$ and let $B_1 = \sum_1^n p_i p_i^1$. Repeating: on α_i , in the order q_i, s_i let q_i^2 be the first point of $B_1 + P$, let $Q_2 = \sum_1^{n-1} q_i^2$ and $A_2 = \sum_1^{n-1} q_i q_i^2$; on β_i , in the order p_i, r_i let p_i^2 be the first point of $Q + \bar{R} + Q_2$ and $B_2 = \sum_1^n p_i p_i^2$; and so on. Continue this process. For some integer m we must have $A_j = A_m$ and $B_j = B_m$ for all $j > m$. For otherwise $B_j \neq B_{j+1}$ for all j ; and hence for some $i \leq n$ and infinitely many j 's: j_1, j_2, \dots , we have $p_i^{j_1} \neq p_i^{j_1+1}$. But this is impossible since the open subarcs $[p_i^{j_k} p_i^{j_k+1}]$ of β_i are disjoint and $p_i^{j_k}$ and $p_i^{j_k+1}$ lie on different arcs α_i .

Now A_m and B_m are sums of $n - 1$ and n disjoint arcs respectively; and $A_m \cdot B_m$ is contained in the set Q_m of end points q_i^m of the arcs in A_m . Thus if we keep intact each arc in A_m having its end point q_i^m in P and to each remaining arc of A_m , say $q_k q_k^m$, we add the unique arc $p_i p_i^m$ which it intersects so that $p_i^m = q_k^m$, we obtain a set of $n - 1$ disjoint arcs e_1, e_2, \dots, e_{n-1} each joining P and Q .

Now since Q_m contains exactly $n - 1$ points, there is at least one arc β_i , say β_k , with $Q_m \cdot \beta_k = 0$. But this gives $Q_i \cdot \beta_k = 0$ for all $i \leq m$, since by construction $\beta_k \cdot Q_i \neq 0$ gives $\beta_k \cdot Q_j \neq 0$ for all $j > i$ and hence $\beta_k \cdot Q_m \neq 0$. Accordingly, $p_k r_k = \beta_k = p_k p_k^m \subset B_m$ so that $\beta_k \cdot E = 0$ where $E = \sum_1^{n-1} e_i$.

Now if $r_k \in \bar{R}$, $\beta_k + R + W$ (W a region about r_k not intersecting E) contains an arc $p_k y$ which, together with the arcs in E , forms a system S_y so that $y \in S$. On the other hand, if $r_k \in Q$, $E + \beta_k$ consists of n disjoint arcs joining P and Q ; and even in this case, $R +$ the arc of S_x containing $x +$ one of the

arcs of $E + \beta_k$ contains an arc py which together with the remaining $n - 1$ arcs in $E + \beta_k$ forms a system S_y so that $y \in S$.

Thus S is both open and closed in M ; and since M contains arcs joining P and Q and $P \subset S$, we have $S \cdot Q \neq \emptyset$. Hence if $x \in S \cdot Q$, the system S_x determined by x satisfies our theorem.

3. Theorem. *If the locally connected generalized continuum M is n -point connected between two of its points a and b , it contains a set of n independent arcs joining a and b .*

Again the proof is by induction on n . For $n = 1$ the conclusion follows by arcwise connectedness. We suppose it holds for all $k < n$. If $M - a - b$ contains more than one component Q with $\bar{Q} \supset a + b$, $M_1 = \bar{Q}$ and $M_2 = M - Q$ are locally connected generalized continua; and since if a set K_i separates a and b in M_i ($i = 1, 2$), $K_1 + K_2$ separates a and b in M , it follows that M_i is n_i -point connected between a and b ($i = 1, 2$), where $0 < n_i \leq n - 1$ and $n_1 + n_2 = n$; and our induction hypothesis gives a set of n_i independent arcs in M_i joining a and b whose sum is a set of n such arcs as required by the theorem. Thus we may suppose there is only one such component Q and we set $\bar{Q} = E$. Since any set separating a and b in E also separates them in M , E is n -point connected between a and b .

LEMMA. *E contains a locally connected generalized continuum N which is n -point connected between a and b and has a as a regular point.*

Proof of the lemma. Let U be a neighborhood of a such that \bar{U} is compact and does not contain b . Since each point of E is interior to an arbitrarily small locally connected continuum in E and the boundary $F(U)$ of U is compact, $F(U)$ is contained in the interior of the sum L of a finite number of locally connected continua in $E - a - b$; and if we add to L an arc in $E - a - b$ joining each pair of components of L , we obtain a locally connected continuum X_1 in $E - a - b$ containing $F(U)$ in its interior. Thus if we set $H_1 = G + X_1$, where G is the component of $E - F(U)$ containing b , H_1 is a locally connected generalized continuum because every $x \in H_1$ is interior either to G or to X_1 in H_1 .

Now let (V_i) be a sequence of neighborhoods of a such that $(U - U \cdot X_1) \supset \bar{V}_1 \supset V_1 \supset \bar{V}_2 \supset V_2 \supset \dots$ and $\delta(V_i) < 1/i$. Let us set $Y_1 = 0$, and for each $i > 1$ let X_i be the sum of a finite number of locally connected continua such that $F(V_i) \subset X_i \subset V_{i-1} - V_{i-1} \cdot X_{i-1} - \bar{V}_{i+1}$, and let Y_i be the sum of a finite number of arcs in $E - a$ each joining X_i and H_{i-1} and having just one point in H_{i-1} and such that Y_i intersects every component of X_i . Let

$$H_i = H_{i-1} + Y_i + X_i \qquad \text{for } i > 1, \text{ and } H = a + \sum_1^\infty H_i.$$

Then since H_1 is a locally connected generalized continuum and, for each

$i > 1$, $X_i + Y_i \subset V_{i-1}$, it follows that H is a locally connected generalized continuum.

Since for any $x \in (E - a - b)$, $E - x$ is $(n - 1)$ -point connected between a and b , it contains a sum S_x of $n - 1$ independent arcs joining a and b . Since the open sets $E - S_x$ cover the compact sets $X_i + Y_i$, for each i there exists a finite collection $S_1^i, S_2^i, \dots, S_{m_i}^i$ of the sets S_x such that $X_i + Y_i \subset \sum_{j=1}^{m_i} (E - S_j^i)$. Set $S_j^i = T_j^i$, $j \leq m_i$, and for each $i > 1$, and $j \leq m_i$, let T_j^i be the closure of the component of $S_j^i - S_j^i \cdot H_{i-1}$ containing a . Finally, set

$$N = H + \sum_{i=1}^{\infty} \sum_{j=1}^{m_i} T_j^i.$$

Then since for each $i > 1$ and all $j \leq m_i$, $T_j^i \subset V_i$, it follows that N is a locally connected generalized continuum. Further, since for any $i > 1$, $W_i = N \cdot (V_{i-1} - \bar{V}_i) \subset X_{i-1} + X_i$ plus a finite number of the arcs of Y_s and of T_j^s for $s \leq i$, since W_i contains a compact set not intersecting $X_{i-1} + X_i$ and separating a and $F(V_{i-1})$ in N , and since any finite arc sum is everywhere regular as it can contain no continuum of condensation, it follows that N is regular at a .

It remains to show that N is n -point connected between a and b . Suppose, on the contrary, that some set K of less than n points separates a and b irreducibly in N . We shall prove this impossible by showing that $K \cdot H = 0$. In the first place, $K \cdot X_1 = 0$ because for any $x \in X_1$ some T_j^1 does not contain x whereas every T_j^1 must contain K , as otherwise one of the arcs of T_j^1 from a to b fails to intersect K . Also, $K \cdot G = 0$. For if not, then if y is the last point of X_1 on an arc ab in $E - K$, $yb \subset G + X_1 \subset N$. Then since $K - K \cdot G$ would not separate a and b in N , $N - (K - K \cdot G)$ would contain an arc aub ; and if u is the first point of X_1 on this arc, $au \cdot K = 0$ and we have a connected subset $au + X_1 + yb$ of $N - K$ containing $a + b$. Thus $K \cdot H_1 = 0$. Hence if $K \cdot H \neq 0$ and k is the least integer such that $K \cdot H_k \neq 0$, we have $k > 1$. But $H_k = H_{k-1} + Y_k + X_k$; and if $Y_k + X_k$ contains a point x of K , some set T_j^k fails to contain x and hence contains an arc az with $az \cdot K = 0$, $z \in H_{k-1}$, since T_j^k consists of $n - 1$ arcs joining a and H_{k-1} and intersecting by pairs in just a . This is impossible since then $H_{k-1} + az$ is connected and contains $a + b$ but does not intersect K . Thus $K \cdot H = 0$, which also is absurd because $N \supset H \supset a + b$. Thus the lemma is proven.

Returning to the proof of the theorem, we note that since the lemma can be applied in N at b , we may suppose N regular at both a and b . Thus there exist infinite sequences (P_i) and (Q_i) of regions in N such that $\bar{P}_1 \cdot \bar{Q}_1 = 0$ and, for each $i > 1$,

$$a \subset P_i \subset \bar{P}_i \subset P_{i-1} \subset V_{1/i}(a), \quad b \subset Q_i \subset \bar{Q}_i \subset Q_{i-1} \subset V_{1/i}(b),$$

the boundaries $F(P_i)$ and $F(Q_i)$ contain finite numbers p_i and q_i of points, and the boundary of any open set lying in P_i (or Q_i) and containing a (b) contains not less than p_i (q_i) distinct points. The set $N_0 = N - P_1 - Q_1$ is

n -point strongly connected between $F(P_1)$ and $F(Q_1)$, because any set separating N_0 between these two sets in the broad sense also separates N between a and b . Hence, by §2, N_0 contains a sum T_0 of n disjoint arcs joining $F(P_1)$ and $F(Q_1)$ and we suppose these arcs reduced so that each has only its end points in the sum of these two sets.

Let $P'_1 = T_0 \cdot F(P_1)$, $Q'_1 = T_0 \cdot F(Q_1)$. Then each of these sets contains exactly n points, namely, one end point from each of the n arcs in T_0 . Now if $N_1 = \bar{P}_1 - P_2$, N_1 must be n -point strongly connected between P'_1 and $F(P_2)$. For if some set H of less than n points separated these two sets in N_1 in the broad sense, the set $K = H + F(P_1) - P'_1$ contains less than p_1 points; and the component of $N - K$ containing a is in P_1 and has less than p_1 boundary points, contrary to the choice of P_1 . Hence N_1 contains a sum T_1 of n disjoint arcs joining P'_1 and $F(P_2)$. Similarly $M_1 = \bar{Q}_1 - Q_2$ contains a sum S_1 of n disjoint arcs joining Q'_1 and $F(Q_2)$. We suppose the arcs in T_1 and S_1 reduced so that the set $P'_2 = T_1 \cdot F(P_2)$ and $Q'_2 = S_1 \cdot F(Q_2)$ consist of exactly n end points as before. Continuing, $N_2 = \bar{P}_2 - P_3$ and $M_2 = \bar{Q}_2 - Q_3$ contain sums T_2 and S_2 respectively of n disjoint arcs joining P'_2 to $F(P_3)$ and Q'_2 to $F(Q_3)$, and so on indefinitely. Clearly the set

$$S = a + b + T_0 + \sum_1^{\infty} (T_i + S_i)$$

consists of exactly n independent arcs in N (and thus in M) from a to b as required by the theorem.

BIBLIOGRAPHY

1. N. E. Rutt, *Concerning the cut points of a continuous curve when the arc-curve AB contains exactly N independent arcs*, Amer. J. Math. vol. 51 (1929) pp. 217-246.
2. G. Nöbeling, *Eine verschärfung des n -Beinsatzes*, Fund. Math. vol. 18 (1931) pp. 23-38.
3. K. Menger, *Kurventheorie*, Teubner, Berlin-Leipzig, 1932, chap. VI.
4. L. Zippin, *Independent arcs of a continuous curve*, Ann. of Math. vol. 34 (1933) pp. 95-113.
5. G. T. Whyburn, *On the cyclic connectivity theorem*, Bull. Amer. Math. Soc. vol. 37 (1931) pp. 429-433; *Analytic topology*, Amer. Math. Soc. Colloquium Publications, vol. 28, 1942.
6. R. L. Moore, *Foundations of point set theory*, Amer. Math. Soc. Colloquium Publications, vol. 13, 1932.
7. K. Menger, *Zur Begründung einer axiomatischen Theorie der Dimension*, Monatshefte für Mathematik und Physik vol. 36 (1929) pp. 193-218.
8. G. T. Whyburn, *The cyclic and higher connectivity of locally connected spaces*, Amer. J. Math. vol. 53 (1931) pp. 427-442.

UNIVERSITY OF VIRGINIA,
CHARLOTTESVILLE, VA.