

A THEORY OF ANALYTIC FUNCTIONS IN BANACH ALGEBRAS

BY

E. K. BLUM

Introduction. The present paper is concerned with the general problem of extending the classical theory of analytic functions of a complex variable. This question received the attention of Hilbert and F. Riesz, and probably goes back to Volterra. More recently N. Dunford, L. Fantappi , I. Gelfand, E. R. Lorch, A. D. Michal, and A. E. Taylor have contributed to the subject (see bibliography).

Our approach differs from most of the others in two main respects, namely, in the type of domain and range of the functions and in the definition of analyticity. We consider functions which have for their domains and ranges subsets of an abstract commutative Banach algebra with unit and we use a definition of analyticity introduced by E. R. Lorch [1]. It is known [4] that a function analytic by this definition is differentiable in the Fr chet sense but not every Fr chet-differentiable function on a commutative Banach algebra is analytic in the Lorch sense. Accordingly, the Lorch theory is the richer.

For the most part, the development of the primary aspects of the Lorch theory parallels that of the classical theory. Interesting departures occur in the more advanced stages. As one would expect, the Cauchy integral theorem and formula occupy a central position and yield the Taylor expansion in the usual way. With Lorch's work as a foundation, we have extended the theory to include a study of Laurent expansions and analytic continuation. There are also some results on the zeros of polynomials over the algebra, on rational functions and their integrals, and on the singularities of analytic functions. Although the objective of this investigation was essentially analytical, we have also obtained results of an algebraic-topological character (e.g. distribution of singular elements). This was a natural outcome of the algebraic character of the techniques used:

1. Basic concepts. A set, B , of elements (denoted by Latin letters a, b, c, x, y, z, \dots) is a "Banach algebra" if (1) B is an algebra over the complex numbers (denoted by greek letters), (2) B is a complex Banach space, and (3) the norm satisfies the inequality $\|ab\| \leq \|a\| \|b\|$. In addition, we assume that multiplication is commutative and that B contains a unit element, e , with $\|e\| = 1$.

We recall without proof some of the salient facts about Banach algebras. Multiplication is continuous in both factors together in the metric topology

Received by the editors March 9, 1953 and, in revised form, January 14, 1954.

defined by the norm. If a^{-1} exists ($aa^{-1} = a^{-1}a = e$), then a is a "regular" element. The set of regular elements will be denoted by G . Since z^{-1} is a continuous function of z , G is a topological group relative to multiplication.

G is an open set and is therefore the union of disjoint maximal open connected sets, the components of G . The component G_1 is the maximal connected subgroup of G [4]. If G has a component other than G_1 , then G has infinitely many components [1]. We note that G_1 contains the complex plane with the origin removed, i.e. the set of elements of the form λe , $\lambda \neq 0$.

An element which is not regular is "singular." The set of singular elements is closed and contains at least the zero element, 0. If it contains only 0, then B is isomorphic to the field of complex numbers [8].

If $I \subset B$ is a closed proper ideal ($I \neq B$), then the quotient algebra B/I is also a commutative Banach algebra with unit [3]. If M is a maximal ideal, then B/M is isomorphic to the field of complex numbers. The natural homomorphism $B \rightarrow B/M$ induces a homomorphism of the algebra onto the complex numbers. If $x \in B$, we let $x(M)$ denote the complex number onto which x is mapped [3]. If x is fixed and M varies over the set of maximal ideals, \mathfrak{M} , then $x(M)$ may be regarded as a complex-valued function on \mathfrak{M} . Gelfand shows that this function has the following properties: (1) If $x = x_1 + x_2$, then $x(M) = x_1(M) + x_2(M)$. (2) If $x = x_1 \cdot x_2$, then $x(M) = x_1(M) \cdot x_2(M)$. If λ is complex, then $\lambda \cdot x(M) = (\lambda x)(M)$. (3) $e(M) = 1$. (4) $|x(M)| \leq \|x\|$. (5) If $M \neq M'$, there is an x in B such that $x(M) \neq x(M')$. (6) If $x(M)$ is not zero for any M in \mathfrak{M} , then there is a y in B such that $y(M) = 1/x(M)$.

An element b is "nilpotent" if $b^n = 0$ for some positive integer n . It is "quasi-nilpotent" if for all complex λ , $\lim_{n \rightarrow \infty} (\lambda b)^n = 0$ [1]. The set of quasi-nilpotents is called the "radical" of B . An element b is quasi-nilpotent if and only if $\lim_{n \rightarrow \infty} (\|b^n\|)^{1/n} = 0$. For every x in B , $\lim_{n \rightarrow \infty} (\|x^n\|)^{1/n} = \max \{ |x(M)| \}$, M in \mathfrak{M} . Thus the radical is the intersection of all the maximal ideals of B . \mathfrak{M} may be topologized by defining a neighborhood of M_0 in \mathfrak{M} as the set of all M in \mathfrak{M} such that $|x_i(M) - x_i(M_0)| < \epsilon$ where the x_i , $i = 1, 2, \dots, n$, are any finite number of arbitrary elements in B and $\epsilon > 0$ is any real number. With this topology, \mathfrak{M} is a compact Hausdorff space and the functions $x(M)$ are continuous on \mathfrak{M} .

The algebra B is "reducible" if it is the direct sum of proper ideals which are also B -algebras with units, that is, $B = B_1 + B_2$, where $B_i \neq \{0\}$. Reducibility is equivalent to the existence of an idempotent element e_j ($e_j^2 = e_j$) such that $e_j \neq e$, $e_j \neq 0$. A notion closely related to reducibility is that of the spectrum of an element. The "spectrum" of b is the set of complex numbers λ such that $(b - \lambda e)$ is singular. The number λ is in the spectrum of b if and only if there is an M in \mathfrak{M} such that $b(M) = \lambda$. The algebra B is irreducible if and only if the spectrum of every element is a connected set in the complex plane. An element b is regular if and only if $b(M) \neq 0$ for every M in \mathfrak{M} .

2. Analytic functions. All functions in §§2-4 are assumed to be single-

valued. The concepts "curve," "rectifiable," "length," "integration on a curve" are defined in the natural way [1]. The classical theorems on Riemann integration are valid since the proofs require only those properties of the complex numbers which are also possessed by the algebra B , the norm in B playing the role of the modulus.

We now state the definition of analyticity, given by Lorch [1], which shall be used herein.

DEFINITION 2.1. Let z_0 be a point in B and let $N(z_0)$ be a neighborhood of z_0 . Let $f(z)$ be a function defined on $N(z_0)$ with range in B . $f(z)$ is "differentiable" at z_0 if there is an element, $f'(z_0)$, in B with the property that for any real $\epsilon > 0$ there is a $\delta > 0$ such that for all z in B with $\|z - z_0\| < \delta$

$$\|f(z) - f'(z_0) - (z - z_0)f'(z_0)\| \leq \epsilon \|z - z_0\|.$$

The element $f'(z_0)$ is the "derivative" of $f(z)$ at z_0 . The function $f(z)$ is "analytic at z_0 " if it has a derivative at all points in some neighborhood of z_0 .

3. The Cauchy theorem and formula. The following theorem was stated by Lorch [1]:

THEOREM 3.1. *Let E be a convex region of B . If $f(z)$ is analytic in E and K is an arbitrary rectifiable closed curve in E , then $\int_K f(z) dz = 0$.*

The convexity of E permits the construction of a proof along classical lines in which the theorem is established first for a triangle, then a polygon, and finally for the curve K . The proof for the triangle is based on its compactness.

We now prove the Cauchy integral theorem for a Banach algebra.

THEOREM 3.2. *Let $f(z)$ be analytic in a region E . Let K be an arbitrary rectifiable closed curve which is homotopic to a point in E . Then $\int_K f(z) dz = 0$.*

(REMARK. When dealing with homotopy in an open set, there is no loss of generality in assuming that all the curves are rectifiable. For example, in the proof which follows we can assume that the curves $K_i, i = 1, \dots, m$, are rectifiable since each K_i can be replaced by the homotopic polygonal curve consisting of the line segments joining the points $z_0, z_{i1}, z_{i2}, \dots, z_{in-1}, z_n$ on K_i .)

Proof. Let K be parametrized by $z = \phi(t)$, where $\phi(0) = \phi(1) = z_0$. Since K is homotopic to z_0 , there is a function of two real variables, $F(s, t)$, continuous on the unit square $0 \leq s \leq 1, 0 \leq t \leq 1$, in both variables together and such that $F(0, t) = \phi(t), F(1, t) \equiv z_0, F(s, 0) = F(s, 1) = z_0$, and $F(s, t)$ is in E for every (s, t) in the unit square.

Since the mapping F is continuous on the compact unit square, its image, $\{F(s, t)\}$, is a compact set in E . Hence, there is a minimum distance, $\rho > 0$, from this image set to the boundary of E .

Now, F is uniformly continuous. Hence, there is a $\delta > 0$ such that $\|F(s', t') - F(s, t)\| < \rho$ whenever $\|s' - s\| + \|t' - t\| < \delta$.

$-F(s, t) \parallel < \rho/2$ for all pairs $(s, t), (s', t')$ with $|s' - s| < \delta$ and $|t' - t| < \delta$. We choose s_1 such that $|s_1| < \delta$ and $0 = t_0 < t_1 < \dots < t_n = 1$ such that $|t_i - t_{i-1}| < \delta$. Let $S(z_0)$ be the sphere $\|z - z_0\| < \rho$. Let $z_{0i} = F(0, t_i), i > 0$, and $z_{1i} = F(s_1, t_i)$ and let K_1 be the curve defined by $F(s_1, t), 0 \leq t \leq 1$. Now, $\|z_{11} - z_0\| < \rho/2, \|z_{12} - z_{11}\| < \rho/2, \|z_{02} - z_{01}\| < \rho/2$, and $\|z_{01} - z_0\| < \rho/2$. Therefore, for $|t - t_1| < \delta$ we have $\|F(s_1, t) - z_0\| \leq \|F(s_1, t) - z_{11}\| + \|z_{11} - z_0\| < \rho, \|F(0, t) - z_0\| \leq \|F(0, t) - z_{01}\| + \|z_{01} - z_0\| < \rho$. Thus, these points are in $S(z_0)$, which is a convex set contained in E . Hence, the path consisting of the arc of K joining z_0 to z_{01} , the arc of K_1 joining z_0 to z_{11} , and the line segment joining z_{01} to z_{11} (denoted by L_1) is contained in $S(z_0)$. By Theorem 3.1 we have

$$\int_{z_0, K}^{z_{01}} f(x) dx + \int_{z_{01}, L_1}^{z_{11}} f(x) dx = \int_{z_0, K_1}^{z_{11}} f(x) dx.$$

Next we let $S(z_{01})$ be the sphere $\|z - z_{01}\| < \rho$. As above, the arc of K from z_{01} to z_{02} , the line segment L_2 joining z_{02} to z_{12} , the arc of K_1 from z_{12} to z_{11} , and the line segment L_1 are all in $S(z_{01})$. By Theorem 3.1 again,

$$-\int_{z_{01}, L_1}^{z_{11}} f(x) dx + \int_{z_{01}, K}^{z_{02}} f(x) dx + \int_{z_{02}, L_2}^{z_{12}} f(x) dx = \int_{z_{11}, K_1}^{z_{12}} f(x) dx.$$

Proceeding in this way at each of the points $z_{0i}, i = 0, 1, \dots, n - 1$, we obtain a set of n equations of the above type. Adding these equations, we see that all integrals over the line segments cancel, leaving

$$\int_K f(x) dx = \int_{K_1} f(x) dx.$$

Now consider the curve $K_2: F(s_2, t)$ where $|s_2 - s_1| < \delta$. Applying the above procedure, we obtain $\int_{K_2} f(x) dx = \int_{K_1} f(x) dx$. After a finite number of steps, we obtain $\int_{K_m} f(x) dx = \int_K f(x) dx$, where K_m is the "null" curve, $F(1, t) \equiv z_0$. Thus $\int_{K_m} f(x) dx = 0$ and the theorem is proven.

DEFINITION 3.1. The set of points of the form $z + \lambda e$ where z is a fixed element in B and λ varies over the complex numbers is the " z -plane⁽¹⁾."

DEFINITION 3.2. Let $K: \{z + \lambda(t)e\}$ be a simple closed curve in the z -plane (i.e. $\lambda(t)$ is continuous and complex, $0 \leq t \leq 1$). A point $z + \lambda e$ in the z -plane is "interior to K " if the complex number λ is interior to the simple closed curve, $\zeta = \lambda(t)$, in the complex plane.

An analogue of the Cauchy formula was stated for a neighborhood by Lorch. The following theorem is a somewhat sharper formulation but is not quite the complete analogue (see §4).

THEOREM 3.3. *Let $f(z)$ be analytic in a region $E \subset B$. Let z_0 be a point in E*

(¹) More generally we may consider the set $\{z + \lambda g\}$ where g is a regular element. We shall call this set the $(z + \lambda g)$ -plane.

and $S(z_0)$ the sphere of maximum radius, ρ , with center z_0 which is contained in E . If K is the curve $\{z_0 + (\mu \exp i\theta)g \mid 0 \leq \theta \leq 2\pi, g \in G, \mu \|g\| = \rho/2\}$, and $S(z_0, \delta)$ is the sphere with center z_0 and radius $\delta = \mu/(4\|g^{-1}\|)$, then for all z in $S(z_0, \delta)$

$$(1) \quad f(z) = \frac{1}{2\pi i} \int_K (x - z)^{-1} f(x) dx.$$

Proof. For all points x on K we have $\|x - z_0\| = |\mu| \|g\|$. If z is in $S(z_0, \delta)$, then $z = z_0 + b$ where $\|b\| < \mu/(4\|g^{-1}\|) \leq \mu \|g\|/4$. Thus, $|(x - z)(M)| = |(x - z_0 - b)(M)| \geq |(x - z_0)(M)| - |b(M)| \geq |\mu| \|g(M)\| - \mu/4 \|g^{-1}\| > 0$, for every maximal ideal M . This implies that $(x - z)^{-1}$ exists for all x in K . Since $K \subset S(z_0)$, the integrand is defined.

Now let K' be the curve $\{z + (\mu \exp i\theta)g\}$ and let \bar{K}' be the set $\{z + \lambda g \mid |\lambda| \leq \mu\}$. Clearly $\bar{K}' \subset S(z_0)$. Thus $f(z)$ is analytic in \bar{K}' and since the Cauchy formula holds in the $(z + \lambda g)$ plane,

$$f(z) = \frac{1}{2\pi i} \int_{K'} (x - z)^{-1} f(x) dx.$$

If x is in K , the sphere $S(x, \mu/2\|g^{-1}\|)$ is contained in $S(z_0)$. Let y be in $S(x, \mu/2\|g^{-1}\|)$. Then $y = x + c$ where $\|c\| < \mu/2\|g^{-1}\|$. Hence, for every maximal ideal M ,

$$|(y - z)(M)| \geq |(x - z)(M)| - |c(M)| \geq 3\mu/4\|g^{-1}\| - \mu/2\|g^{-1}\| > 0$$

so that $(y - z)^{-1} f(y)$ is defined and analytic in y in the sphere $S(x, \mu/2\|g^{-1}\|)$.

Now choose $0 = \theta_1 < \theta_2 < \dots < \theta_n = 2\pi$ such that for the corresponding points $x_1, x_2, \dots, x_n = x_1$ on K the arc length from x_j to x_{j+1} is less than $\mu/4\|g^{-1}\|, j = 1, \dots, n - 1$. For $x_j = z_0 + (\mu \exp i\theta_j)g$, choose $x'_j = z + (\mu \exp i\theta_j)g$ on K' . Let L_j denote the line segment joining x_j to x'_j . Clearly, $L_j \subset S(x_j, \mu/2\|g^{-1}\|)$ so that $L_j \subset S(z_0)$ and $(x - z)^{-1}$ exists for all x on L_j .

Finally, let C_j be the closed curve consisting of the arc of K from x_j to x_{j+1} , followed by the line segment L_{j+1} , then by the arc of K' from x'_{j+1} to x'_j and lastly the line segment L_j . The preceding remarks show that $C_j \subset S(x_j, \mu/2\|g^{-1}\|)$. Since the function $g(x) \equiv (1/2\pi i)(x - z)^{-1} f(x)$ is analytic for x in $S(x_j, \mu/2\|g^{-1}\|)$, the Cauchy integral theorem yields the result

$$\begin{aligned} \int_{C_j} g(x) dx &= \int_{x_j, K}^{x_{j+1}} g(x) dx + \int_{x_{j+1}, L_{j+1}}^{x'_{j+1}} g(x) dx + \int_{x'_{j+1}, K'}^{x'_j} g(x) dx \\ &\quad + \int_{x'_j, L_j}^{x_j} g(x) dx = 0. \end{aligned}$$

Adding these equations for $j = 1, \dots, n - 1$, we obtain

$$\frac{1}{2\pi i} \int_K (x - z)^{-1} f(x) dx = \frac{1}{2\pi i} \int_{K'} (x - z)^{-1} f(x) dx.$$

But the right-hand member equals $f(z)$. This establishes the theorem.

The formula for the derivative is obtained in similar fashion and for the higher derivatives by induction as in the classical theory.

4. Taylor expansions.

DEFINITION 4.1. A series $P(z|z_0) = \sum_{n=0}^{\infty} b_n(z-z_0)^n$ where the b_n are elements of B is a "Taylor series" or "power series" around z_0 . It is "super-absolutely" convergent (s.-a. convergent) if the series $\sum_0^{\infty} \|b_n\| \|z-z_0\|^n$ converges. It is "absolutely" convergent if $\sum_0^{\infty} \|b_n(z-z_0)^n\|$ converges.

DEFINITION 4.2. The "radius of convergence" of the series $P(z|z_0)$ is defined as $1/\limsup_{n \rightarrow \infty} (\|b_n\|)^{1/n}$.

The classical theorems on power series are easily established. However, Taylor's theorem must be proven in two steps. First, it is shown that the Taylor series of $f(z)$ equals $f(z)$ in a neighborhood of the center of expansion, z_0 . Then the identity theorem is proven. (Here, it is important to note that we require two analytic functions to be equal throughout a neighborhood of z_0 in order to prove them identical. The classical condition that they be equal at a sequence of points approaching z_0 is known to be insufficient.) Combining these two results, we obtain equality of $f(z)$ and its power series in the largest sphere contained in the region of analyticity.

This, in turn, yields the complete analogue of the Cauchy formula. With reference to Theorem 3.3, we state it as follows.

THEOREM 4.1. Let K be the curve $\{z_0 + (\mu \exp i\theta)e | 0 \leq \theta \leq 2\pi, \mu < \rho\}$. Then formula (1) of Theorem 3.3 holds for all z such that $\|z - z_0\| < \mu$.

5. Analytic continuation. The concepts of analytic continuation, functional element, and complete analytic function are taken over directly from the classical theory. We shall be especially concerned with the singularities of an analytic function. For the precise definitions, the reader is referred to Courant's *Funktionentheorie*, p. 376 ff. The following example will illustrate how the definitions are used here.

Consider the function defined by the formula z^{-1} . In each component of G , z^{-1} is a distinct single-valued analytic function. Consider the principal component, G_1 . G_1 is the domain of a complete analytic function defined by z^{-1} , for suppose d is a boundary point of G_1 . Let K be a path joining e to d such that K is in G_1 except for the end point d . Suppose further that it is possible to continue a functional element $P_1(z|e)$, at e , along K so as to obtain a functional element $P_n(z|d)$ at d . We assume, as we may, that the functional elements $P_1(z|e)$, $P_2(z|z_2)$, \dots , $P_{n-1}(z, z_{n-1})$, $P_n(z, d)$ are power series, the center of each being in the sphere of s.-a. convergence of the preceding. Now, there is a sequence of regular elements $\{g_j\}$ on K such that $\lim_j g_j = d$ and g_j is in the sphere of s.-a. convergence of $P_{n-1}(z|z_{n-1})$. Then we have $P_{n-1}(g_j|z_{n-1}) = g_j^{-1}$ and $e = \lim_j [g_j \cdot P_{n-1}(g_j|z_{n-1})] = d \cdot P_{n-1}(d|z_{n-1})$. This implies d is in G , which is a contradiction. Thus d is a "K-singularity" of z^{-1} and z^{-1} cannot be continued outside of G .

6. The Laurent expansion. A Laurent expansion about a point z_0 is a series of the form $\sum_{-\infty}^{\infty} b_n(z-z_0)^n$, b_n in B , defined for those z such that $(z-z_0)^{-1}$ exists and the two series $\sum_0^{\infty} b_n(z-z_0)^n$ and $\sum_{-1}^{-\infty} b_n(z-z_0)^n$ are convergent. A Laurent expansion may be defined in several disjoint regions in each of which it may define a different analytic function; e.g. z^{-1} . With this possibility in mind, we prove the main theorem.

THEOREM 6.1. *Let g be an arbitrary element in G , and let E denote the set $\{z_0 + \lambda g \mid 0 \leq r_1 < |\lambda| < r_2\}$. Setting $\mu_\lambda = \min \{r_2 - |\lambda|, |\lambda| - r_1\}$, we write $S(\lambda)$ for the sphere $\|z - (z_0 + \lambda g)\| < \mu_\lambda / \|g^{-1}\|$ and $U = \bigcup_{r_1 < |\lambda| < r_2} S(\lambda)$. If D is a region such that $E \subset D \subset U$ and $f(z)$ is a function which is single-valued and analytic in D , then for each z in U there are two curves $K_j(z): \{z_0 + (\rho_j \exp it)g, 0 \leq t \leq 2\pi\}$, $j=1, 2$, with $r_1 < \rho_1 < \rho_2 < r_2$ such that:*

$$(1) \quad \phi(z) = \frac{1}{2\pi i} \int_{K_2(z)} (x - z)^{-1} f(x) dx - \frac{1}{2\pi i} \int_{K_1(z)} (x - z)^{-1} f(x) dx$$

is a well-defined single-valued analytic function of z ;

$$(2) \quad \text{for all } z \text{ in } U, \phi(z) = \sum_{-\infty}^{\infty} a_n(z-z_0)^n \text{ where}$$

$$a_n = \frac{1}{2\pi i} \int_K (x - z_0)^{-n-1} f(x) dx$$

and K is any curve of the form $\{z_0 + (r \exp it)g \mid r_1 < r < r_2, 0 \leq t \leq 2\pi\}$;

$$(3) \quad \text{for all } z \text{ in } D, \phi(z) = f(z).$$

Proof. Let z be in U . $z = z_0 + \lambda g + h$ where $\|h\| = \mu_\lambda - \epsilon / \|g^{-1}\|$, $\epsilon > 0$. For all M in \mathfrak{M} , $|g(M)| = 1 / |g^{-1}(M)| \geq 1 / \|g^{-1}\|$. Hence, $|h(M)| \leq \|h\| < \mu_\lambda / \|g^{-1}\| \leq |g(M)| \cdot \min \{r_2 - |\lambda|, |\lambda| - r_1\}$, or $|h(M)| \leq |g(M)| \cdot (r_2 - \epsilon - |\lambda|)$ and $|h(M)| \leq |g(M)| (|\lambda| - r_1 - \epsilon)$. Then we have $|(z - z_0)(M)| = |\lambda g(M) + h(M)| \leq |\lambda| \cdot |g(M)| + |h(M)| \leq |\lambda| \cdot |g(M)| + |g(M)| (r_2 - \epsilon - |\lambda|) = (r_2 - \epsilon) |g(M)|$. Also $|(z - z_0)(M)| \geq |\lambda| \cdot |g(M)| - |h(M)| \geq |\lambda| |g(M)| - |\lambda| |g(M)| + (r_1 + \epsilon) |g(M)| = (r_1 + \epsilon) |g(M)|$.

Now let $\rho_1 = r_1 + \delta$, $\rho_2 = r_2 - \delta$ where $0 < \delta < \epsilon$, and consider the curves $K_j(z) = \{z_0 + (\rho_j \exp it)g\}$, $j=1, 2$. For all x on $K_2(z)$ we have $|(x - z)(M)| \geq \|g(M)\| \rho_2 - |(z_0 - z)(M)| \geq |g(M)| (r_2 - \delta) - (r_2 - \epsilon) |g(M)| = (\epsilon - \delta) |g(M)| > 0$. Likewise for x on $K_1(z)$, $|(x - z)(M)| \geq |g(M)| (r_1 + \epsilon) - |g(M)| (r_1 + \delta) = |g(M)| (\epsilon - \delta) > 0$. Thus the integrand in both of the integrals of (1) is defined. Furthermore, although the choice of the curves $K_j(z)$ is not uniquely determined by z , any pair of curves which satisfy the specified conditions will yield the same value for each of the integrals in (1). This is an immediate consequence of the Cauchy theorem. Hence, the value of $\phi(z)$ is uniquely determined by z .

Now, for x on $K_2(z)$ we have $(x - z_0)^{-1} = (1/\rho_2)g^{-1} \exp(-it)$. Hence, $(x - z_0)^{-1}(z - z_0) = (1/\rho_2)g^{-1}(\lambda g + h) \exp(-it) = (1/\rho_2)(\lambda + hg^{-1}) \exp(-it)$ and $\|(x - z_0)^{-1}(z - z_0)\| \leq (1/\rho_2) [|\lambda| + \|h\| \|g^{-1}\|] \leq (1/\rho_2) [|\lambda| + (r_2 - |\lambda|) - \epsilon]$

$\cdot (1/||g^{-1}||) ||g^{-1}|| = (r_2 - \epsilon) / \rho_2 = (r_2 - \epsilon) / (r_2 - \delta) < 1$. The last fraction is less than 1. Therefore, we may write $(x - z)^{-1} = (x - z_0)^{-1} \{ e + \sum_{n=1}^{\infty} [(z - z_0)(x - z_0)^{-1}]^n \}$ since the series converges. The convergence is absolute and uniform in x for x on $K_2(z)$.

For x on $K_1(z)$, $(x - z_0) = (\rho_1 \exp it)g$. Since $(z - z_0)^{-1} = (\lambda g + h)^{-1} = (1/\lambda)g^{-1}(e + hg^{-1}/\lambda)^{-1}$, we have

$$(x - z_0)(z - z_0)^{-1} = (\rho_1 \exp it/\lambda) [e - hg^{-1}/\lambda + (hg^{-1}/\lambda)^2 - \dots],$$

the series being super-absolutely convergent because $||hg^{-1}||/|\lambda| \leq ||h|| \cdot ||g^{-1}||/|\lambda| = 1/|\lambda|(\mu_\lambda - \epsilon/||g^{-1}||) ||g^{-1}|| = (\mu_\lambda - \epsilon)/|\lambda| < 1$. In fact, $|(x - z_0)(z - z_0)^{-1}| \leq (\rho_1/|\lambda|) [1 - (\mu_\lambda - \epsilon)/|\lambda|]^{-1} = \rho_1/(|\lambda| - \mu_\lambda + \epsilon) \leq \rho_1/(r_1 + \epsilon) < 1$. Thus we may write $(x - z)^{-1} = -(z - z_0)^{-1} \{ e + \sum_{n=1}^{\infty} [(z - z_0)^{-1}(x - z_0)]^n \}$. Again the series converges uniformly in x for x on $K_1(z)$. Term-by-term integration yields the result (2) of the theorem if we observe that $a_n = (1/2\pi i) \cdot \int_{K_j(z)} (x - z_0)^{-n-1} f(x) dx$ where $j=2$ for $n \geq 0$ and $j=1$ for $n < 0$, and by the Cauchy theorem $\int_{K_j(z)} (x - z_0)^{-n-1} f(x) dx = \int_K (x - z_0)^{-n-1} f(x) dx$ where K is any curve of the form $\{z_0 + (r \exp it)g | r_1 < r < r_2, 0 \leq t \leq 2\pi\}$.

For $n \geq 0$,

$$\begin{aligned} a_n &= \frac{1}{2\pi i} \int_K [f(x)/(r \exp it)^{n+1}] (g^{-1})^{n+1} dx \\ &= \frac{g^{-n-1}}{2\pi} \int_0^{2\pi} [f(x)/(r \exp it)^{n+1}] g \cdot r \exp(it) dt \end{aligned}$$

so that $||a_n|| \leq (1/2\pi) (||g^{-1}||^n ||g^{-1}g|| \cdot m/r^n) 2\pi = ||g^{-1}||^n \cdot m/r^n$, where $m = \max \{ ||f(x)||, x \text{ on } K \}$. Thus the series of positive powers of the Laurent series in (2) must be s.-a. convergent for $||z - z_0|| < r/||g^{-1}|| < r_2/||g^{-1}||$, or since r may be taken arbitrarily close to r_2 , for $||z - z_0|| < r_2/||g^{-1}|| \leq r_2 ||g||$.

For $n < 0$, $||a_n|| \leq (1/2\pi) m r^{-n} ||g||^{-n} \cdot 2\pi$, where $m = \max \{ ||f(x)||, x \text{ on } K \}$. Thus the series of negative powers converges super-absolutely for $|(z - z_0)^{-1}| < 1/(r_1 ||g||)$. For z in U , $|(z - z_0)^{-1}| \leq ||g^{-1}||/r_1$ (by fourth paragraph of this proof). Since $||g^{-1}||/r_1 \geq 1/(r_1 ||g||)$, the series of negative powers may not be s.-a. convergent for all z in U , although it converges for these z . If $g = e$, then $||g^{-1}|| = ||g|| = 1$ and this difficulty does not arise. Likewise, if $r_1 = 0$, the series of negative powers is absolutely convergent and s.-a. convergent.

Now suppose z is in D . As above, $z = z_0 + \lambda g + h$ where $||h|| = (\mu_\lambda - \epsilon)/||g^{-1}||$. Choose $K_j(z)$, $j = 1, 2$, as described in the proof of (1) and choose K_λ as the curve $\{z_0 + \lambda g + (r \exp it)g, 0 \leq t \leq 2\pi\}$ where $r = \mu_\lambda - \delta$, $0 < \delta < \epsilon$. Consider the set $F = \{z_0 + \zeta g | r_1 < |\zeta| < r_2, |\zeta - \lambda| > \mu_\lambda - \epsilon\}$, $F \subset E$ and $K_\lambda \subset F$ and $K_j(z) \subset F$. We have for all M in \mathfrak{M} and x in F $|(x - z)(M)| = |(z_0 + \zeta g - z)(M)| = |(\zeta - \lambda)g(M) - h(M)| \geq |\zeta - \lambda| |g(M)| - ||h|| = |\zeta - \lambda|/||g^{-1}(M)|| - ||h||$ since $1 = |g^{-1}g(M)| = |g^{-1}(M)| \cdot |g(M)|$. Since $-||h|| = (-\mu_\lambda + \epsilon)/||g^{-1}||$,

$$\begin{aligned} |(x-z)(M)| &\geq [(|\zeta - \lambda| \|g^{-1}\| / |g^{-1}(M)|) - \mu_\lambda + \epsilon] / \|g^{-1}\| \\ &\geq [|\zeta - \lambda| - \mu_\lambda + \epsilon] / \|g^{-1}\| > 0. \end{aligned}$$

Hence, $(x-z)^{-1}f(x)$ is defined and analytic for all x in F and

$$\begin{aligned} \frac{1}{2\pi i} \int_{K_\lambda} (x-z)^{-1}f(x)dx &= \frac{1}{2\pi i} \int_{K_2(z)} (x-z)^{-1}f(x)dx \\ &\quad - \frac{1}{2\pi i} \int_{K_1(z)} (x-z)^{-1}f(x)dx \end{aligned}$$

by the Cauchy theorem. The integral on the left is equal to $f(z)$ by the Cauchy formula and the right side equals $\phi(z)$. This proves (3), that is, $\phi(z)$ is an analytic continuation of $f(z)$ from the set D to the set U . We shall show that it is possible to continue beyond U .

If M is in \mathfrak{M} , let $E(M)$ be the open annular ring in the complex plane bounded by the circles $K_1(M): \{z_0(M) + r_1 \exp it, 0 \leq t \leq 2\pi\}$ and $K_2(M): \{z_0(M) + r_2 \exp it, 0 \leq t \leq 2\pi\}$. Let $Q_M = \{z | z(M) \in E(M)\}$ and let $Q = \bigcap_{M \in \mathfrak{M}} Q_M$. Then we prove

LEMMA 6.1. *If z is in Q , there is a $\delta > 0$ such that for all x on the curves $K_1: \{z_0 + (r_1 \exp it)e\}$ and $K_2: \{z_0 + (r_2 \exp it)e\}$ and for all M in \mathfrak{M} ,*

$$|(x-z)(M)| > \delta.$$

Proof. First we prove that the function $|x(M)|$ is continuous in the pair (x, M) for all x in B and all M in \mathfrak{M} . Let x_0 be in B , M_0 in \mathfrak{M} , $\epsilon > 0$. Let $\mathfrak{M}_0 = \{M | |x_0(M) - x_0(M_0)| < \epsilon/2\}$ and $\mathfrak{N}_0 = \{x | \|x - x_0\| < \epsilon/2\}$. The product $\mathfrak{N}_0 \times \mathfrak{M}_0$ is a neighborhood of (x_0, M_0) in $B \times \mathfrak{M}$. If (x, M) is in $\mathfrak{N}_0 \times \mathfrak{M}_0$, then $|x(M)| - |x_0(M_0)| \leq |x(M) - x_0(M_0)| \leq |x(M) - x_0(M)| + |x_0(M) - x_0(M_0)| \leq \|x - x_0\| + \epsilon/2 < \epsilon/2 + \epsilon/2 = \epsilon$. Similarly, $|x_0(M_0)| - |x(M)| < \epsilon$, which establishes the continuity at (x_0, M_0) .

Now, since K_1 and K_2 are compact sets, their union $K_1 \cup K_2$ is also compact. \mathfrak{M} is a compact space ((3)). Therefore, the product $(K_1 \cup K_2) \times \mathfrak{M}$ is compact. Since the function $|(x-z)(M)|$ is continuous in the pair (x, M) , it assumes a minimum value $\delta \geq 0$ on $(K_1 \cup K_2) \times \mathfrak{M}$. If $\delta = 0$, there is a pair (x', M') such that $|(x'-z)(M')| = 0$. Thus $|z(M')| = |x'(M')|$, contradicting the assumption that z is in Q .

The above lemma shows that Q is an open set, for if z is in Q , consider the sphere $\|z' - z\| < \delta/2$. Then $|z'(M) - z(M)| \leq \|z' - z\| < \delta/2$ so that $z'(M)$ is in $E(M)$ for all M ; i.e. z' is in Q .

Obviously, $U \subset Q$ since z in U implies that $r_1 < |z(M) - z_0(M)| < r_2$ for all M . To give some idea of the structure of Q , we point out that the sphere

$\|z - z_0\| \leq r_1$ does not intersect Q since $|z(M) - z_0(M)| \leq r_1$ for all such z . On the other hand, the set of z such that $\|z - z_0\| \geq r_2$ may intersect Q , although Q contains no element of the form $z_0 + \lambda e$, where $|\lambda| > r_2$ since $|z_0(M) + \lambda - z_0(M)| > r_2$. In fact, if $z = z_0 + \lambda e + h$ where $|\lambda| > r_2$ and $\|h\| < |\lambda| - r_2$, then z is not in U , for $|(z - z_0)(M)| = |\lambda + h(M)| \geq |\lambda| - \|h\| > r_2$. Examples may be adduced to show that Q may contain U properly.

Consider the Banach space (m) of bounded sequences $z = \{\zeta_n\}$, ζ_n a complex number, $\|z\| = \sup_n \{|\zeta_n|\}$. If multiplication of sequences is defined by $\{\xi_n\} \cdot \{\zeta_n\} = \{\xi_n \zeta_n\}$ it is easy to see that (m) becomes a B -algebra. Let $z_0 = \{0, 0, \dots\}$. In this case, the set E is a subset of the cartesian product of \aleph_0 annular rings of the complex plane, each of inner radius r_1 , outer radius r_2 , and center at the origin. Let b be the element of (m) defined by the sequence $\{(-1)^{n+1}(r_1 + r_2)/2\}$. We shall prove that b is in Q but not in U .

Note first that if $z = \{\zeta_n\}$ and $z(M) = \lambda$ for some maximal ideal, then there is a sub-sequence of $\{\zeta_n\}$ which converges to λ . (The sub-sequence may of course consist of only one distinct number λ .) Otherwise $(z - \lambda e)^{-1} = \{1/(\zeta_n - \lambda)\}$ and λ is not in the spectrum of z . This sub-sequence determines a sequence of maximal ideals $\{M_k\}$ in (m) such that $\lim_k z(M_k) = \lambda$. Note also that the set M_p of all elements $z = \{\zeta_n\}$ such that $\zeta_p = 0$ is a maximal ideal. If $a = \{\alpha_n\}$ and $c = \{\gamma_n\}$ are in M_p , and if $x = \{\xi_n\}$, $y = \{\eta_n\}$ are arbitrary elements of (m) , then $xa + yc = \{\xi_n \alpha_n + \eta_n \gamma_n\}$ is in M_p since $\alpha_p = \gamma_p = 0$. This makes M_p an ideal. Clearly, $M_p \neq (m)$. Suppose $M_p \subset I$ where I is a proper ideal of (m) , and let x be in I but not in M_p . Then $x = \{\xi_n\}$ where $\xi_p \neq 0$ and $|\xi_n| \leq \|x\|$ for all n . Consider the element $y = \{\eta_n\}$ where $\eta_p = 0$ and $\eta_n = \|x\| + 1$ for $n \neq p$. Since y is in M_p , $x + y = \{\xi_n + \eta_n\}$ is in I . But $|\xi_n + \eta_n| \geq |\eta_n| - |\xi_n| \geq \|x\| + 1 - \|x\| = 1$ for $n \neq p$ and $|\xi_p + \eta_p| = |\xi_p| \neq 0$. Thus $(x + y)^{-1}$ exists and $I = (m)$. This contradiction proves M_p is a maximal ideal. For any element $x = \{\xi_n\}$, $x(M_p) = \xi_p$ since $x - \xi_p e = \{\xi_n - \xi_p\}$ is in M_p .

These remarks show that $b(M) = \pm(r_1 + r_2)/2$ for any maximal ideal M in (m) . Therefore, b is in Q . Now suppose b is in U . Since $z_0 = \{0, 0, \dots\}$, $b = \lambda e + h$ where $r_1 < |\lambda| < r_2$ and $\|h\| < \mu_\lambda$. This implies that $|(b - \lambda e)(M)| \leq \|h\| < \mu_\lambda$ for all M , or $\max_n \{ | [(-1)^{n+1}(r_1 + r_2)/2] - \lambda | \} < \mu_\lambda$. For odd n this means that λ lies within the circle of radius $(r_2 - r_1)/2$ and center at $(r_1 + r_2)/2$, whereas for even n , λ must lie within the circle of radius $(r_2 - r_1)/2$ and center at $-(r_1 + r_2)/2$. This is manifestly impossible so that b is not in U .

Returning to the abstract algebra B , we observe that Q , as an open set, is the union of maximal open connected sets, its components. Let Q_1 be the component containing U . We shall prove that it is possible to continue analytically from U to Q_1 .

THEOREM 6.2. *Let $f(z)$ be the function of Theorem 6.1. If z is in Q_1 , then there are curves $K_1(z): \{z_0 + (\rho_1 \exp it)e\}$ and $K_2(z): \{z_0 + \rho_2 \exp it)e\}$ such that*

$$(1) \quad \psi(z) = (1/2\pi i) \int_{K_2(z)} (x - z)^{-1} f(x) dx - (1/2\pi i) \int_{K_1(z)} (x - z)^{-1} f(x) dx$$

is a well-defined single-valued analytic function in Q_1 ,

$$(2) \quad \psi(z) \text{ is an analytic continuation of } \phi(z) \text{ from } U \text{ to } Q_1.$$

Proof. Lemma 6.1 shows that $z(M)$ is contained in the open annulus bounded by the curves $\{z_0(M) + (r_2 - \delta) \exp it\}$ and $\{z_0(M) + (r_1 + \delta) \exp it\}$ for all M . Thus if ρ_1 and ρ_2 are chosen so that $r_1 < \rho_1 < r_1 + \delta$ and $r_2 - \delta < \rho_2 < r_2$, then the curves $K_1(z)$ and $K_2(z)$ are such that $(x - z)(M) \neq 0$ for all x in $K_1(z) \cup K_2(z)$ and all M ; i.e. $(x - z)^{-1}$ exists and the integrands in (1) are defined. For any other curves chosen in the prescribed way, the integrals have the same value by the Cauchy theorem. Thus $\psi(z)$ is uniquely determined by z .

$\psi(z)$ obviously coincides with $\phi(z)$ for z in U . To show that it is analytic in Q , we let $\|z' - z\| < \delta/2$. Then $|(z' - x)(M)| = |(z' - z)(M) + (z - x)(M)| \geq |(z - x)(M)| - |(z' - z)(M)| > \delta - \delta/2 > 0$ for all x in $K_1(z) \cup K_2(z)$ and all M in \mathfrak{M} . If we choose ρ'_1 and ρ'_2 such that $r_1 < \rho'_1 < r_1 + \delta/2$, $r_2 - \delta/2 < \rho'_2 < r_2$, then the curves $K_j(z')$: $\{z_0 + (\rho'_j \exp it)e\}$, $j = 1, 2$, will not only serve to define $\psi(z')$ but $\psi(z)$ as well. Thus we may write

$$\begin{aligned} \psi(z') - \psi(z) &= \frac{1}{2\pi i} \int_{K_2(z)} [(x - z')^{-1} f(x) - (x - z)^{-1} f(x)] dx \\ &\quad - \frac{1}{2\pi i} \int_{K_1(z')} f(x) [(x - z')^{-1} - (x - z)^{-1}] dx \\ &= \frac{1}{2\pi i} \int_{K_2(z')} f(x) [(z' - z)(x - z')^{-1} (x - z)^{-1}] dx \\ &\quad - \frac{1}{2\pi i} \int_{K_1(z')} f(x) [(z' - z)(x - z')^{-1} (x - z)^{-1}] dx. \end{aligned}$$

A classical argument then shows that $\psi'(z) = (1/2\pi i) \int_{K_2(z')} (x - z)^{-2} f(x) dx - (1/2\pi i) \int_{K_1(z')} (x - z)^{-2} f(x) dx$. Thus $\psi(z)$ is an analytic continuation of $\phi(z)$ to Q_1 . The question of whether Q_1 is the maximal region to which $\phi(z)$ may be continued analytically will be answered negatively by means of an example to be cited in §7.

7. Singularities of analytic functions. Let $f(z)$ be a complete analytic function and let $P_1(z|z_1)$ be a power series functional element of $f(z)$ at the point $z_1 = z_0 + \lambda g$, where g is in G . Let K be a path in the $(z_0 + \lambda g)$ -plane joining z_1 to z_0 and such that $P_1(z|z_1)$ may be continued analytically along K up to but not including z_0 . Thus, z_0 is a K -singularity of $f(z)$.

DEFINITION 7.1. Let z_0 be as above. z_0 is a "g-plane singularity" of $f(z)$ if

there is an $r_2 > 0$ and a connected arc K' of K beginning at z_0 and contained in the set $E = \{z_0 + \lambda g, 0 < |\lambda| < r_2\}$ with the following property: If x is in K' and $P(z|x)$ is the power series functional element at x , then $P(z|x)$ may be continued along every path in E which does not contain z_0 . If $g = e$, we say that z_0 is a "plane singularity" of $f(z)$. In the continuation process within E , it may occur that for each point, $z_0 + \lambda g$, of E all functional elements whose regions of analyticity contain $z_0 + \lambda g$ assume the same value at this point. In this case, we say that z_0 is of "locally single-valued type."

If z_0 is of locally single-valued type, then $f(z)$ has a functional element whose region of analyticity contains E . By Theorem 6.1 this functional element has a Laurent expansion at z_0 which represents the functional element at least in E .

DEFINITION 7.2. The series of negative powers of the Laurent expansion is its "principal part." If the Laurent expansion at a g -plane singularity z_0 has only a finite number of negative powers, then z_0 is called a "pole." If there are an infinite number of negative powers, z_0 is an "essential singularity."

For the remainder of this paper, we assume that all complete analytic functions are single-valued unless the contrary is specifically stated. Thus all g -plane and plane singularities are of single-valued type and there are Laurent expansions at all such points.

If z_0 is a plane singularity of $f(z)$, then it is also a g -plane singularity for all g in G_1 and conversely. In proof, the series of negative powers of the Laurent expansion, $f(z) = \sum_{-\infty}^{\infty} b_n(z-z_0)^n$, converges for $\|(z-z_0)^{-1}\| < \infty$ (see Theorem 6.1), and the series of positive powers for $\|z-z_0\| < r_2$ if z_0 is a plane singularity. Hence the series converges for all $z = z_0 + \lambda g$ where $0 < |\lambda| < r_2/\|g\|$ and g is in G_1 . Thus z_0 is also a g -plane singularity. The converse is proven similarly. (If g is in G but not in G_1 , the Laurent expansion is defined at $z_0 + \lambda g$, $0 < |\lambda| < r_2/\|g\|$ but may represent a different complete analytic function if the region of definition of the Laurent series consists of disjoint components.)

A plane singularity need not be isolated.

THEOREM 7.1. *Let B be a Banach algebra such that (i) the set of regular elements G is dense in B and (ii) the radical of B is $\{0\}$. If z_0 is a plane singularity of the single-valued complete analytic function $f(z)$, then every sphere $\|z-z_0\| < \rho$ contains a point z' which is not contained in the domain of $f(z)$.*

Proof. Let $\sum_{-\infty}^{\infty} b_n(z-z_0)^n$ be the Laurent expansion of $f(z)$ at z_0 . By Theorem 6.1, the Laurent expansion is defined for all z such that $(z-z_0)^{-1}$ exists and $\|z-z_0\| < r_2$. Since $f(z)$ is single-valued, $f(z) = \sum_{-\infty}^{\infty} b_n(z-z_0)^n$ for all such z . Since z_0 is a plane singularity, for some negative integer $-m$, $b_{-m} \neq 0$. Since the radical of B is $\{0\}$, there is an M^* in \mathfrak{M} such that $b_{-m}(M^*) \neq 0$.

Let z_1 be an element of M^* such that $\|z_1\| < \rho$. There are such elements

since M^* contains λz whenever it contains z . Letting $z' = z_0 + z_1$, we have $z'(M^*) = z_0(M^*)$ and $\|z' - z_0\| < \rho$. Suppose z' is in the domain of $f(z)$. Then there is a Taylor expansion $\sum_0^\infty a_n(z - z')^n$ which converges in a sphere $\|z - z'\| < \delta$, and equals $f(z)$ therein.

Since G is dense in B , there is an element g in G such that $\|g - z_1\| < \epsilon$, where $0 < \epsilon < \delta$, but ϵ is otherwise arbitrary. Let $h = z_0 + g$. Since $\|h - z'\| = \|g - z_1\| < \epsilon < \delta$, we have $f(h) = \sum_0^\infty a_n(h - z')^n$. Thus $[f(h)](M^*) = \sum_0^\infty a_n(M^*) [h(M^*) - z_0(M^*)]^n$. Now, $(h - z_0)^{-1}$ exists. Furthermore, $\|h - z_0\| \leq \|h - z'\| + \|z' - z_0\| < \epsilon + \rho$. Since ρ and ϵ may be chosen arbitrarily small, we may write $\|h - z_0\| < r_2$. Therefore, the Laurent expansion is defined for h and $f(h) = \sum_{-\infty}^\infty b_n(h - z_0)^n$. It follows that $[f(h)](M^*) = \sum_{-\infty}^\infty b_n(M^*) [h(M^*) - z_0(M^*)]^n = \sum_0^\infty a_n(M^*) [h(M^*) - z_0(M^*)]^n$ where $b_{-m}(M^*) \neq 0$. The equality of the two series holds for a set of complex numbers approaching $h(M^*)$ as a limit. To show this, we set $h_t = z_0 + tg$, t real. Then $\|h_t - z'\| = \|tg - (z' - z_0)\| \leq \|tg - g\| + \|g - (z' - z_0)\| < |t - 1| \cdot \|g\| + \epsilon < \delta$ for t sufficiently close to 1. It follows that $f(h_t) = \sum_0^\infty a_n(h_t - z')^n$. Further, $h_t(M) - z_0(M) = tg(M) \neq 0$ for all M and $\|h_t - z_0\| \leq \|h_t - z'\| + \|z' - z_0\| < |t - 1| \|g\| + \epsilon + \rho < r_2$ for t close to 1. Thus, $f(h_t) = \sum_{-\infty}^\infty b_n(h_t - z_0)^n$. Finally, $[f(h_t)](M^*) = \sum_0^\infty a_n(M^*) [h_t(M^*) - z_0(M^*)]^n = \sum_{-\infty}^\infty b_n(M^*) [h_t(M^*) - z_0(M^*)]^n$ for $|t - 1| < \epsilon_1$, say. As t approaches 1, $h_t(M^*)$ approaches $h(M^*)$. Applying the classical identity theorem for complex analytic functions, we conclude that $a_n(M^*) = b_n(M^*)$ for all n , $-\infty < n < \infty$ (uniqueness of Laurent expansion). This implies that $b_{-m}(M^*) = 0$ which is a contradiction. Therefore, z' is not in the domain of $f(z)$.

The method of proof of the preceding theorem shows that a weaker hypothesis than G being dense in B would suffice. It is sufficient that B be such that every maximal ideal M contains an element b on the boundary of G . Then we may set $z_1 = \lambda b$ if λ is chosen in the right way, and the existence of a regular element in any neighborhood of z_1 is assured.

We shall give examples of algebras in which G is a dense set and examples of algebras in which G is nondense. In fact, there is an algebra having a maximal ideal consisting entirely of elements which are in the interior of the set of singular elements.

The algebra of bounded sequences (m) is a B -algebra in which G is a dense set. The set of singular elements consists of all sequences $\{\sigma_n\}$ such that $\inf_n \{|\sigma_n|\} = 0$. If a $*$ -operation is introduced into (m) by defining $\{\sigma_n\}^* = \{\bar{\sigma}_n\}$, where $\bar{\sigma}_n$ is the complex conjugate of σ_n , then (m) becomes a B^* -algebra with real bounded $*$ -operation [10] since $\{\sigma_n\}^* = \{\sigma_n\}$ if and only if σ_n is real for all n , i.e. the spectrum of $\{\sigma_n\}$ is real, and $\sup_{x \in (m)} \{\|x^*\|/\|x\|\} = 1$ where $\|x^*\| = \sup_n \{|\bar{\sigma}_n|\}$. It follows from a theorem of Rickart [11] that (m) is a proper algebra; i.e. every singular element is a generalized zero-divisor. (z is a generalized zero-divisor if there is a sequence of elements z_n with $\|z_n\| = 1$ and such that $\lim_n z_n z = 0$.) Furthermore, the set of singular ele-

ments is contained in the closure of G , for if $\{\sigma_n\}$ is singular and $\epsilon > 0$, there is a subsequence $\{\sigma_{n_j}\}$ such that $|\sigma_{n_j}| < \epsilon/2$ and for $n \neq n_j$, $|\sigma_n| \geq \epsilon/2$. Let $\gamma_n = \sigma_n$ for $n \neq n_j$ and otherwise let $\gamma_{n_j} = \epsilon/4$. The element defined by the sequence $\{\gamma_n\}$ is regular since $|\gamma_n| \geq \epsilon/4$ for all n . Also $\|\{\gamma_n\} - \{\sigma_n\}\| = \sup_n \{|\gamma_n - \sigma_n|\} < \epsilon$. Thus every neighborhood of $\{\sigma_n\}$ intersects G . Hence, every singular element is in the closure of G so that (m) is in the closure of G . Thus G is dense in (m) .

In the same way it is shown that the sub-algebra of convergent sequences is also such that the regular elements in it form a dense set.

The algebra $C[0, 1]$ of continuous functions (complex-valued) on the unit interval $[0, 1]$ with the norm defined by $\|f(t)\| = \max_{0 \leq t \leq 1} \{|f(t)|\}$ is another example of an algebra in which the regular elements form a dense set. In proof, let $f(t)$ be a singular element. Then $\min_t \{|f(t)|\} = 0$. Let $T = \{t \mid |f(t)| \leq \epsilon/3\}$ where $\epsilon < \max_t \{|f(t)|\}$. We define the function $g(t)$ as follows: for t not in T , $g(t) = f(t)$. If $t \in T$ and $|f(t)| = \epsilon/3$ then $g(t) = f(t)$. If t is in T and $|f(t)| < \epsilon/3$, then there is an open interval containing t and such that $|f(t)| < \epsilon/3$ for all points in this interval. Let (t_1, t_2) be the largest such interval. By continuity, $|f(t_1)| = |f(t_2)| = \epsilon/3$. Let $f(t_1) = (\epsilon/3) \exp i\theta_1$ and $f(t_2) = (\epsilon/3) \exp i\theta_2$. We define $g(t) = (\epsilon/3) \exp [iu\theta_2 + i(1-u)\theta_1]$ where $u = (t - t_1)/(t_2 - t_1)$ for all t in (t_1, t_2) . This describes $g(t)$ completely. Clearly, it is a continuous function. It is a regular element of $C[0, 1]$ since $\inf_t \{|g(t)|\} = \epsilon/3 > 0$. Since $\max_t \{|g(t) - f(t)|\} < \epsilon$, $g(t)$ is in an ϵ -neighborhood of $f(t)$. Thus every singular element is in the closure of the set of regular elements which therefore form a dense set.

We now consider another example. Let E denote the sub-algebra of $C[0, 1]$ generated by the element $b = \exp 2\pi it$ and the complex numbers. E consists of all polynomials in b (with complex coefficients) and their limits (uniform convergence). The spectrum of b consists of all the complex numbers λ with $|\lambda| \leq 1$. In proof, we note that for $\lambda < 1$, we have

$$\frac{1}{\lambda + \exp(2\pi it)} = \left(\frac{1}{\exp(2\pi it)} \right) [1 - \lambda/\exp(2\pi it) + \lambda^2/(\exp 2\pi it)^2 - \dots]$$

and this element is not in E for if we consider the function $g(t) = [1/(\lambda + \exp 2\pi it) - \sum_0^n \lambda_n \exp 2\pi int]$ we find that $\|g(t)\| \geq 1$. We have $1 = |\int_0^1 (\exp 2\pi it) g(t) dt| \leq \max_t \{|g(t)|\} = \|g(t)\|$. Finally, for $|\lambda| = 1$, there exists t_0 such that

$$\exp 2\pi it_0 = \lambda$$

so that $(\exp 2\pi it - \lambda)^{-1}$ does not exist even in $C[0, 1]$. Since 0 is in the spectrum of b , b is a singular element.

Now we show that the principal ideals of the form $(b - \lambda)$, where $|\lambda| < 1$, are maximal. If $|\lambda| < 1$, the element $b - \lambda$ is singular and the principal ideal $(b - \lambda)$ is contained in a maximal ideal, M . Then $b(M) = \lambda$. Now suppose z is in M and let $P_n(b) = \sum_{k=0}^n \alpha_{kn} b^k$, $n = 1, 2, \dots$, be a sequence of polynomials

such that $\lim_n P_n(b) = z$. Let q_m be an integer such that $\|P_{q_m}(b) - z\| < 1/m$. Writing $\lambda_m = \sum_{k=0}^{q_m} \alpha_k \lambda^k$, we have $p_{q_m}(\lambda) - \lambda_m = 0$ so that $p_{q_m}(x) - \lambda_m = (x - \lambda)g_{q_m}(x)$, where $g_{q_m}(x)$ is a polynomial in x . $|\lambda_m| = |\sum_{k=0}^{q_m} \alpha_k \lambda^k| = |P_{q_m}(b)(M) - z(M)| \leq \|P_{q_m}(b) - z\| < 1/m$. Thus $\lim_{m \rightarrow \infty} [P_{q_m}(b) - \lambda_m] = \lim_{m \rightarrow \infty} P_{q_m}(b) - \lim_m \lambda_m = \lim_m P_{q_m}(b) = z$. But $\lim_m [P_{q_m}(b) - \lambda_m] = \lim_m (b - \lambda) \cdot g_{q_m}(b) = (b - \lambda) \lim_m g_{q_m}(b)$ and we have $z = (b - \lambda)c$, where $c = \lim_m g_{q_m}(b)$ is in E . Thus, M is contained in the principal ideal $(b - \lambda)$, which means that $M = (b - \lambda)$.

Now let us consider the maximal ideal $(b) = (\exp 2\pi it)$ in E . If z_0 is in (b) , then $z_0 = b \lim_n p_n(b)$ where $p_n(b)$ is a polynomial in b for $n = 1, 2, \dots$. The element z_0 is obviously singular in E . If $z_0 \neq 0$, we shall show that there is a spherical neighborhood of z_0 contained in the set of singular elements.

Since z_0 is singular and not zero, there is an $\epsilon, 0 < \epsilon < 1$, such that the sphere $\|z - z_0\| < \epsilon$ contains no scalars. We shall show that all z in this sphere are singular, i.e. all the nonzero elements in (b) are interior points of the set of singular elements. Thus, the regular elements are certainly not dense in this algebra.

Choose δ such that $0 < \epsilon + \delta < 1$. Since z is not a scalar, there is a polynomial $\sum_0^r \lambda_k b^k$ with $\lambda_r \neq 0, r \geq 1$, such that $\|z - \sum_0^r \lambda_k b^k\| < \delta$. Further, since $z_0 = b \lim_n p_n(b)$, given $\gamma, 0 < \epsilon + \delta + \gamma < 1$, we can find n such that $\|z_0 - b p_n(b)\| < \gamma$. Writing $p_n(b) = \sum_0^s \alpha_k b^k$, we have

$$\left\| \sum_0^s \lambda_k b^k - \sum_0^r \alpha_k b^{k+1} \right\| < \epsilon + \delta + \gamma < \mu < 1.$$

Hence, for any M ,

$$|\lambda_0 + (\lambda_1 - \alpha_0)b(M) + (\lambda_2 - \alpha_1)b^2(M) + \dots| < \mu.$$

Since 0 is in the spectrum of b , there is a maximal ideal such that $b(M) = 0$ (i.e. $M = (b)$). Hence, $|\lambda_0| < \mu$. This implies that $\sum_0^s \lambda_k b^k$ is singular, for the complex polynomial $\sum_0^s \lambda_k x^k$ has a complex root $|\zeta_1| < 1$. But ζ_1 is in the spectrum of b and there is a maximal ideal M_1 such that $b(M_1) = \zeta_1$ and we have $(\sum_0^s \lambda_k b^k)(M_1) = \sum_0^s \lambda_k \zeta_1^k = 0$. Since δ may be chosen arbitrarily small, it is seen that every neighborhood of z contains singular elements. Hence z cannot be regular and the sphere $\|z - z_0\| < \epsilon < 1$ is contained in the set of singular elements.

Incidentally, this also affords an example of a commutative B -algebra in which the set of generalized zero-divisors is not contained in the intersection of the closure of G with the set of singular elements [11]. The element $b - e = (\exp 2\pi it - 1)$ is in this intersection since all elements of the form $(\exp 2\pi it - 1 - \epsilon), \epsilon > 0$, in a neighborhood of $(b - e)$ are regular. Rickart [11] shows that every singular element on the boundary of G is a generalized zero-divisor. Hence, the element $(\exp 2\pi it - 1)$ is a generalized zero-divisor. There-

fore the element $(\exp 2\pi it)(\exp 2\pi it - 1)$ is also a generalized zero-divisor. However, this element is also in the maximal ideal $(\exp 2\pi it)$ which means that it is an interior point of the set of singular elements. Hence it is not in the closure of G .

In connection with Theorem 7.1, we remark that the point z' may be a planar singularity or not, that is, both situations do arise. As a simple illustration of the first situation, we consider the cartesian product of the complex plane with itself, i.e. the set of pairs (λ_1, λ_2) with λ_1 and λ_2 complex. With the usual definitions of addition and multiplication and now with norm given by $\max \{ |\lambda_1|, |\lambda_2| \}$, this set becomes a B -algebra. If we let $a_1 = (1, 0)$ and $a_0 = (0, 1)$, then the function $f(z) = a_1 z^{-1} + a_0$, where $z = (\zeta_1, \zeta_2)$, may be written in the form $f(z) = (1/\zeta_1, 0/\zeta_2) + (0, 1) = (1/\zeta_1, 1)$. The formula $a_1 z^{-1} + a_0$ defines $f(z)$ for all z in G and only such z . Thus the set Q of Lemma 6.1 is equal to G , which consists of all pairs (λ_1, λ_2) such that $\lambda_1 \neq 0, \lambda_2 \neq 0$. $f(z)$ may obviously be continued analytically outside of Q by means of the formula $(1/\zeta_1, 1)$ which is defined for all $z = (\zeta_1, \zeta_2)$ with $\zeta_1 \neq 0$ and coincides with $f(z)$ in G . This also is the maximum analytic extension of $f(z)$ and illustrates the possibility of continuing beyond the set Q . The point $(0, 0)$ is clearly a pole of $f(z)$. It is not isolated since all points $z' = (0, \zeta_2)$ with $\zeta_2 \neq 0$ in a neighborhood of $(0, 0)$ are points which cannot be in the domain of the complete analytic function determined by $f(z)$. In fact, these points are also poles.

An example of the second situation is afforded by the algebra (m) . Let $f(z) = z^{-1}$ and $z' = \{1/n\}$. The point 0 is obviously a pole and z' is not in the domain of the complete analytic function z^{-1} . Further, every annulus of the form $\{z' + \lambda e, 0 < |\lambda| < r\}$ must contain elements which are singular and therefore not in the domain of z^{-1} . Thus z' is not a plane singularity. On the other hand, $z'' = \{0, 1, 1, \dots, 1, \dots\}$ is a pole, so that both kinds of points exist in a neighborhood of the pole 0.

The behavior of analytic functions in a neighborhood of a singularity is further illuminated by the following analogue of the classical theorem on poles.

THEOREM 7.2. *Let B be a Banach algebra such that (i) the set of regular elements is dense in B and (ii) the radical of B is $\{0\}$. If the point b in B is a pole of the single-valued complete analytic function $f(z)$, then $\lim_{z \rightarrow b} \|f(z)\| = \infty$ regardless of the manner in which z approaches b .*

Proof. Let $\sum_{n=-m}^{\infty} a_n(z-b)^n$ be the Laurent expansion of $f(z)$ at b , assuming that b is a pole of order m . By Theorem 6.1, the Laurent expansion is defined for all z such that $(z-b)^{-1}$ exists and $\|z-b\| < r_2$. Since $f(z)$ is single-valued, $f(z) = \sum_{n=-m}^{\infty} a_n(z-b)^n$ for all such z , and $(z-b)^m f(z) = [a_{-m} + a_{-m+1}(z-b) + \dots]$. We denote the quantity in brackets by $g(z)$. For a constant p there is a $\delta > 0$ such that $\|g(z)\| \geq p > 0$, whenever $\|z-b\| < \delta$. This follows from $\|g(z)\| \geq \|a_{-m}\| - \|a_{-m+1}\| \|z-b\| - \|a_{-m+2}\| \|z-b\|^2 - \dots$, where $a_{-m} \neq 0$. Hence $\|z-b\|^m \|f(z)\| \geq \|(z-b)^m f(z)\| = \|g(z)\| \geq p$ and $\|f(z)\| \geq p/\|z-b\|^m$.

Thus if z approaches b in the region for which the Laurent expansion is valid, then $\|f(z)\| \rightarrow \infty$.

Now suppose $\{z_k\}$ is a sequence of elements in the domain of $f(z)$ for which the Laurent series is not defined and such that $\lim_k z_k = b$. There is an integer N such that $\|z_k - b\| < r_2/3$ for all $k > N$. For each k there is an element g_k in G such that $\|g_k - (z_k - b)\| < \epsilon_k$, where the $\{\epsilon_k\}$ is a sequence of decreasing positive numbers approaching 0. If we let $c_k = b + g_k$, then $c_k - b$ is regular and $f(c_k) = \sum_{n=-\infty}^{\infty} a_n (c_k - b)^n$ for $k > N$. Since $\|c_k - b\| = \|g_k\| \leq \|z_k - b\| + \epsilon_k$, we see that $\lim_{k \rightarrow \infty} \|g_k\| = 0$. From $\|g_k(M)\| \leq \|g_k\|$ it follows that $\lim_k |(c_k - b)(M)| = 0$. Since the radical of B is $\{0\}$, there is a maximal ideal M' for which $a_{-m}(M') \neq 0$. Thus $[f(c_k)](M') = \sum_{n=-\infty}^{\infty} a_n(M') [(c_k - b)(M')]^n$ approaches infinity as k approaches infinity. Therefore, $\|f(c_k)\|$ becomes infinite also. If the ϵ_k are chosen such that $\|f(c_k) - f(z_k)\| < 1$, which is possible by the continuity of $f(z)$ at the points z_k , it follows that $\|f(z_k)\|$ approaches infinity also.

The following remark shows that the behavior of functions in a neighborhood of a singularity is not always analogous to the classical behavior. For example, the Weierstrass theorem for essential singularities does not always hold. Again we assume that G is dense in the algebra. If the point 0 is an essential singularity of the single-valued complete analytic function $f(z)$, and $\sum_{n=-\infty}^{\infty} a_n z^n$ is its Laurent expansion around 0, where $a_n \neq 0$ for an infinite number of negative n , then it may occur that $a_n(M) = 0$ for all $n < 0$ and some maximal ideal M . Then $[f(z)](M) = \sum_0^n a_n(M) [z(M)]^n$ for all z for which the Laurent series is defined. By an argument like the one in the preceding theorem, it can be shown that $[f(z)](M)$ approaches $a_0(M)$ as z approaches 0. Hence, $f(z)$ cannot approach any element of the form λe , where $\lambda \neq a_0(M)$, in every neighborhood of 0, as the Weierstrass theorem requires.

To obtain more information on the behavior of functions in a neighborhood of a singularity, we turn to a consideration of rational functions. In preparation, we shall discuss some of the properties of polynomials over a B -algebra.

8. Polynomials over a B -algebra. Let $p(x) = a_0 + a_1x + \dots + a_nx_n$ be a polynomial with coefficients a_j in B , and leading coefficient $a_n \neq 0$. Let $D(p)$ denote the set of elements z in B such that $p(z)$ is in G . We note that $D(p)$ may be empty. For example, if there is a maximal ideal M such that $a_j(M) = 0$ for $j = 0, 1, \dots, n$, then $[p(z)](M) = \sum_{j=0}^n a_j(M) [z(M)]^j = 0$ for all z in B . Hence, $p(z)$ is singular for all z in B . Whether this is a necessary condition for $D(p)$ to be empty is not known. In the algebra (m) it is the case, but the proof employed for (m) does not generalize. However, if $D(p)$ is empty, we may state the following.

LEMMA 8.1. *If $p(x)$ is a polynomial such that $D(p)$ is empty, then the set of complex numbers λ such that λ is a zero of one of the polynomials $[p(x)](M) = \sum_{j=0}^n a_j(M)x^j$, where M is a maximal ideal, must contain the entire complex plane.*

Proof. Suppose there is a complex number λ such that λ is not a zero of any polynomial $[p(x)](M)$ for any M in \mathfrak{M} . Then $[p(\lambda e)](M) = \sum_{j=0}^n a_j(M)\lambda^j \neq 0$ for all M , which means that λe is in $D(p)$, contrary to hypothesis.

The converse of the lemma is not true, as an example shows. Again we consider the algebra E generated by $b = \exp 2\pi i t$ and the complex numbers. We have shown that the spectrum of b in E consists of all λ such that $|\lambda| \leq 1$. Hence there is a maximal ideal M_λ such that $b(M_\lambda) = \lambda$ for each λ with $|\lambda| \leq 1$. Let $p(x)$ be the polynomial $bx^2 + (b^2 - 1)x - b$. Thus $p(x) = (x + b) \cdot (bx - 1)$. If ζ is any complex number, there is a maximal ideal M_ζ such that $[p(x)](M_\zeta)$ is a polynomial having ζ as one of its zeros. In proof, if $|\zeta| \leq 1$, then there is a maximal ideal M_ζ such that $b(M_\zeta) = -\zeta$ so that $(x + b)(M_\zeta) = (x - \zeta)$ has ζ as a zero. If $|\zeta| > 1$, then $1/|\zeta| < 1$ and there is a maximal ideal M' such that $b(M') = 1/\zeta$ so that $[b(M')x - 1]$ has ζ as a zero. Therefore, the set of zeros of all the polynomials $[p(x)](M)$, M in \mathfrak{M} , contains every complex number; i.e. $D(p)$ contains no complex number. Yet, $D(p)$ is not empty, for if we let $z = 1 - b/2$, then $(bz - 1) = (-1/2)(b^2 - 2b + 2) = (-1/2) \cdot (b - 1 + i)(b - 1 - i)$, so that $(bz - 1)(M) \neq 0$ for all M . Likewise, $z + b = (1/2)(b + 2)$ so that $(z + b)(M) \neq 0$ for all M . Hence, $(1 - b/2)$ is in $D(p)$. Since $D(p)$ is evidently an open set, it contains infinitely many elements whenever it contains one.

We now state a sufficient condition for $D(p)$ to be nonempty.

LEMMA 8.2. *If either a_n or a_0 is a regular element, then $D(p)$ is not empty.*

Proof. If a_0 is regular, then since $p(0) = a_0$, it follows that $p(z)$ is regular for z near 0.

If a_n is regular, then for any complex λ we have $p(\lambda e) = [a_0 a_n^{-1} / \lambda^n + \dots + a_{n-1} a_n^{-1} / \lambda + 1] \lambda^n a_n$. For $|\lambda|$ sufficiently large, $p(\lambda e)$ is close to the regular element $\lambda^n a_n$ and is therefore regular. Thus λe is in $D(p)$.

The preceding example with $p(x) = bx^2 + (b^2 - 1)x - b$ shows that the above sufficient condition for $D(p)$ to be nonempty is not a necessary condition, since b is not a regular element of the algebra E .

Let $S(p)$ be the complement of $D(p)$, that is, the set of all z such that $p(z)$ is singular.

LEMMA 8.3. *If a_0 is singular or if one of the a_i is not in the radical for $i \neq 0$, then $S(p)$ is not empty.*

Proof. There is a maximal ideal M such that $a_n(M) \neq 0$. For this M , the polynomial $[p(x)](M) \equiv a_n(M)x^n + \dots + a_0(M)$ has at least one complex zero, ζ . Then $[p(\zeta e)](M) = 0$ and ζe is in $S(p)$.

Combining the two preceding lemmas, we may state

THEOREM 8.1. *If $p(x)$ is a polynomial with leading coefficient a_n in G , then $p(x)$ separates the algebra B into an open nonempty set $D(p)$ and a nonempty closed set $S(p)$ such that $D(p)$ is the set of all z in B for which $p(z)$ is in G , $S(p)$*

is its complement, and if at least one a_i is not in the radical for $i < n$, both $D(p)$ and $S(p)$ intersect G .

Concerning the second lemma, we remark that if a_n is in the radical of B and $a_n \neq 0$, then $S(p)$ may or may not be empty according as a_0 is or is not regular. For example, if $p(x) = a_n x^n + a_0$ and a_0 is in the radical while a_0 is in G , then $[p(z)](M) \neq 0$ for all z in B and all M in \mathfrak{M} . Thus $S(p)$ is empty. On the other hand, if a_0 is not in G , then $D(p)$ is empty.

Theorem 8.1 may be regarded as an attempt to obtain something like a fundamental theorem of algebra. However, it affords meager information about the zeros of $p(x)$. Yet, in a sense to be explained, it is a best possible result for abstract algebras.

Let $Z(p)$ denote the set of zeros of $p(x)$. Obviously, $Z(p) \subset S(p)$ and therefore may be empty. However, for $Z(p)$ to be empty it is not necessary that a_n be in the radical. In fact, as the following example shows, a_n may be regular.

Let C' be the sub-algebra of $C[0, 1]$ consisting of those functions $f(t)$ for which $f(0) = f(1)$. Let $p(x)$ be the polynomial $a_0 + a_2 x^2$ where $a_0 = -e$ and $a_2 = \exp 2\pi i i$. Suppose $p(z) = 0$. Then $a_2 z^2 = -a_0$ or $(\exp 2\pi i i) z^2 = 1$. Thus $z^2 = \exp(-2\pi i i)$ and $z = \pm \exp(-\pi i i)$. Since $\exp(-\pi i i)$ is not in C' , $p(x)$ has no zeros in C' . Yet, $a_2 = \exp 2\pi i i$ is regular in C' .

Therefore, for an arbitrary algebra, there can be no fundamental theorem of algebra. At the same time, for some polynomials the set $Z(p)$ may be infinite. In the algebra (m) , if $p(x) = a_n x^n + a_0$ where a_n is the sequence $\{\alpha_{nk}\}$, $k = 1, 2, \dots$, and $\inf_k \{|\alpha_{nk}|\} > 0$ and $a_0 = \{\alpha_{0k}\}$ where $\inf_k \{|\alpha_{0k}|\} > 0$, then $p(x)$ has infinitely many zeros. In (m) , if a_n is regular, then $p(x)$ has at least one zero.

The foregoing indicates to some extent how the algebraic properties of polynomials over a B -algebra differ from those of complex polynomials. We now show that the metric properties differ also.

We recall that for complex polynomials the modulus $|p(x)| > N$ for all x such that $|x| > r$, $N > 0$ being an arbitrary real number. A very simple example shows that this is not so for arbitrary B -algebras. In the algebra of pairs (λ_1, λ_2) , let $p(x) = a_1 x + a_0$ where $a_1 = (1, 0)$ and $a_0 = (1, 1)$. For every $r > 0$ there is a z in B with $\|z\| > r$ and $\|p(z)\| = 1$; e.g. $z = (0, r+1)$.

If the polynomial is restricted by the requirement that a_n be in G , one might expect the classical result to apply. For the algebras (m) and $C[0, 1]$, or any algebra in which the norm has the property $\|z^2\| = \|z\|^2$, this is true. If $\|z^2\| = \|z\|^2$, then $\|z\| = \limsup_n (\|z^n\|)^{1/n} = \max \{ \|z(M)\|, M \text{ in } \mathfrak{M} \}$. Hence, $\|p(z)\| \geq \| [p(z)](M) \| \geq \|z(M)\|^n [\mu - \|a_{n-1}\| / \|z(M)\| - \dots - \|a_0\| / \|z(M)\|^n]$ where $\mu = \min \{ \|a_n(M)\|, M \text{ in } \mathfrak{M} \} > 0$. If $\|z\| > r$, then $\max_M \{ \|z(M)\| \} > r$ so that we need only choose r large enough to obtain $\|p(z)\| > N$.

Of course, if $\max_M \{ \|z(M)\| \} < \|z\|$, then the proof is invalid, as it is in an algebra with radical, or in the algebra (1) of convergent (absolutely) power

series with complex coefficients. (1) is an algebra with multiplication and addition defined as ordinary multiplication and addition of series and the norm of a power series defined by the usual formula $\|\sum_0^\infty \alpha_n x^n\| = \sum_0^\infty |\alpha_n|$. In (1), the element $p(x) = x^2 - x - 2$ has norm $1 + 1 + 2 = 4$ whereas $\max \{ |p(x)|, |x| \leq 1 \} = \max \{ |[p(x)](M)|, M \text{ in } \mathfrak{M} \} < 4$ as an elementary calculation shows. The behavior of the norm in algebras like (1) is not clear. We point out that for $N > 0$ there is an r such that for all λe with $|\lambda| > r$, $\|p(\lambda e)\| > N$ if a_n is in G . If F is the set of elements z such that $\|p(z)\| > N$, then F contains all z such that $\max_M \{ |z(M)| \} > r$. F is a subset of the set of z such that $\|z\| > r$. If the radical of B is not $\{0\}$, F is a proper subset.

We turn now to a study of rational functions.

9. Rational functions. Let $q(x) = ax + b$, where a is in G . If $q(z) = 0$, then $az = -b$, and $z = a^{-1}b$. Thus $q(x)$ has precisely one zero. The rational function $g(x) = [q(x)]^{-1} = a^{-1}(x + a^{-1}b)^{-1} = a^{-1}(x - c)^{-1}$ is defined for all elements of the form $c + g$ where $-c = a^{-1}b$ and g is in G . We denote this set by $D(q)$. It is easily seen that $D(q)$ is homeomorphic to G with components mapping onto components under the translation $g \rightarrow c + g$. Unlike G , however, $D(q)$ need not be a group under multiplication.

Let $D(q)'$ be the subset of $D(q)$ consisting of elements $c + g$ such that g is in the principal component G_1 . The fundamental group of $D(q)'$ is isomorphic to $\pi_1(G_1)$, the fundamental group of G_1 . It has been shown [12] that the curves of the form $\{c + \exp ty \mid \exp y = e, 0 \leq t \leq 1\}$ form a set of generators with base point $c + e$. Any closed curve in $D(q)'$ with initial point $c + e$ is homotopic to one of these generators. Let the curve K in $D(q)'$ be parametrized by $z(t)$ where $z(0) = z(1) = c + e$. Let $K(c)$ be the curve in G parametrized by $x(t) = z(t) - c$ so that $x(0) = x(1) = e$. Then $\int_K (z - c)^{-1} dz = \int_{K(c)} x^{-1} dx = 2\pi iy$, where $\exp 2\pi iy = e$. Thus the residues of $(x - c)^{-1}$ are the same as those of x^{-1} .

The formula $(x - c)^{-1}$ defines a single-valued complete analytic function in each of the components of $D(q)$. The point c is obviously a pole of order one of each of these functions. However, there may be other simple poles. Suppose z in B is such that the complement of the spectrum of $z - c$ contains the interior of a circle of radius r , centered at 0, with the exception of the point 0 itself. Then $(\lambda e + z - c)^{-1}$ is defined for all $|\lambda| < r$ except $\lambda = 0$, so that z is a pole. This is also a necessary condition for an element z to be a pole. There may be an infinitude of poles, possibly nondenumerable. For example, if B is a reducible algebra and e_j is an idempotent, then $z_j = c + e_j$ has the above property since the spectrum of $z_j - c$ consists of the points 0 and 1. There may be a nondenumerable number of idempotents in the algebra. If the radical of B is $\{0\}$, and B is irreducible, then since the spectrum of every element is connected, we may conclude that the only plane singularity of $(x - c)^{-1}$ is the pole c .

We observe that all the residues of x^{-1} may be obtained by integrating along special curves which contain the various singularities of x^{-1} as "center"

in a sense. For example, if e_j is an idempotent, then e_j is a pole of x^{-1} . Let $K(e_j)$ be the curve $\{e_j + (r_j \exp 2\pi it)e, 0 < r_j < 1\}$. Then x^{-1} is defined on $K(e_j)$ and for all points of the e_j -plane interior to $K(e_j)$, and we have

$$\int_K x^{-1} dx = \int_0^1 [(\exp 2\pi it)/(e_j + r_j \exp 2\pi it)] 2\pi i r_j dt = z', \text{ say.}$$

If M is a maximal ideal, then $z'(M) = \int_0^1 (2\pi i r_j \exp 2\pi it)/(e_j(M) + r_j \exp 2\pi it) dt$. If $e_j(M) = 1$, then $z'(M) = 0$. If $e_j(M) = 0$, then $z'(M) = 2\pi i$. Thus $z'(M) = 2\pi i(e - e_j)(M)$ for all M . If the radical of B is $\{0\}$, then $z' = 2\pi i(e - e_j)$.

Now let us consider the polynomial $p(x) = a_2 x^2 + a_1 x + a_0$ where a_2 is in G . We may assume $a_2 = e$. If $p(x)$ has a zero z_1 , then we may write $p(x) = (x - z_1) \cdot (x - z_2)$. The substitution $y = x - z_1$ yields $p(x) = y[y - (z_2 - z_1)] = y^2 - cy$ where $c = z_2 - z_1$. Thus we may confine our attention to polynomials of the form $x^2 - cx$ when studying polynomials which are of degree 2 and have a zero. We assume $p(x) = x^2 - cx$ and point out some facts about its zeros.

If B is reducible and e_j is an idempotent, then clearly ce_j is also a zero. Thus $p(x)$ may have infinitely many zeros. Reducibility is not a necessary condition for $p(x)$ to have more than two zeros. In the algebra $C[0, 1]$, if $c = \sin 2\pi t$, then the polynomial $x^2 - cx$ has the following zeros besides 0 and c :

the function $r_1(t)$ defined by: $\sin 2\pi t$ for $0 \leq t \leq 1/2$, and 0 for $1/2 \leq t \leq 1$;

the function $r_2(t)$ defined by: $\sin 2\pi t$ for $1/2 \leq t \leq 1$ and 0 for $0 \leq t \leq 1/2$.

Thus the polynomial has four zeros in $C[0, 1]$ although this algebra is irreducible. Further, if we let $c = t \sin (1/t)$, then $x^2 - cx$ has infinitely many zeros in $C[0, 1]$.

If B is an irreducible algebra with radical $\{0\}$ and c is a regular element in B , then 0 and c are the only zeros of $x^2 - cx$. For if z is such that $z(z - c) = 0$, then for every maximal ideal M we have $z(M)[z(M) - c(M)] = 0$. Thus $z(M) = 0$ or $z(M) = c(M)$. If $z(M) \neq 0$ for any M , then $z = c$. If $z(M') = 0$ for some M' in \mathfrak{M} , then $z = 0$, for if not, by virtue of the connectedness of the spectrum of z there is a maximal ideal M'' such that $0 < z(M'') < \mu$, where $\mu = \min \{|c(M)| M \text{ in } \mathfrak{M}\} > 0$. Then we have $z(M'')[z(M'') - c(M'')] \neq 0$, which contradicts the assumption that z is a zero of $p(x)$.

Having observed the various modes of behavior of $p(x)$, we proceed to study the rational function $[p(x)]^{-1}$. As above, let $D(q)$ be the set of elements of the form $c + g$ where g varies over G . $[p(x)]^{-1}$ is defined for all x in $G \cap D(q)$. We denote this set by $D(p)$. As the intersection of two open sets, $D(p)$ is also open. We shall show that it is not empty. (Remark: It is clear that $D(p)$ is independent of its representation as the intersection of the domains of the factors of $p(x)$; i.e. if $p(x) = (x - z)(x - c + z)$ is any other factorization, then $D(p) = \{z + g | g \text{ in } G\} \cap \{(c - z) + g | g \text{ in } G\}$.)

We recall that G_1 is a subgroup of G and that gG_1 denotes the coset of

G/G_1 which contains g and that this is also the component of G which contains g . Let $(c+gG_1)$ denote the set $\{c+gg_1 \mid g_1 \in G\}$. We call this set the “ c -translate” of the component gG_1 . Observe that λg is in gG_1 for all complex λ . If x is such that $\|x-\lambda g\| < \|\lambda\|/\|g^{-1}\|$ for a given g , then x is in gG_1 . Thus, if we choose λ so that $\|c\| < \|\lambda\|/\|g^{-1}\|$, then $c+\lambda g$ is in $gG_1 \cap (c+gG_1)$; i.e. for all g in G and all c in B the set $gG_1 \cap (c+gG_1)$ is not empty, or each component of G intersects its c -translate.

A component of G may also intersect the c -translate of another component. For example, if g is in G but not in G_1 , and g_1 is in G_1 , then $(c+G_1) \cap gG_1$ contains the element g if we let $c=g-g_1$. Thus we have $G \cap (c+G) = \bigcup_{g,h \in G} [gG_1 \cap (c+hG_1)]$.

Now, $G \cap (c+G)$ is a union of maximal open connected sets, its components. For arbitrary elements c in an arbitrary algebra B , not much is known about the structure of these components. We have seen that each gG_1 intersects its c -translate. It is not known whether this intersection is connected. If it is connected, then clearly it is a component of $D(p)$. In each component of $D(p)$, the formula $[p(x)]^{-1}$ defines a single-valued complete analytic function. We shall investigate its singularities. We write $f(x) = [p(x)]^{-1}$.

If z is a plane singularity of $f(x)$, then z is in $\sim G \cup \sim (c+G)$, where “ \sim ” denotes the complement of a set. If z is in $\sim G$, then its spectrum contains 0 as an isolated point, and all complex numbers in a neighborhood of 0 are not in the spectrum of $z-c$, with the possible exception of 0. This condition is sufficient for z to be a plane singularity of $f(x)$. If z is not in G and the algebra is irreducible and has radical $\{0\}$, then $z=0$ since the only element which has a spectrum containing 0 as an isolated point is the zero element. In this case, 0 is not in the spectrum of $z-c=-c$ unless $c=0$. If $c \neq 0$, then c is in G . Similarly, if z is not in $(c+G)$, then $z-c=0$. If c is not 0, then the spectrum of z cannot contain 0 since it is connected. Thus $z=c$ is in G again. This shows that 0 and c are the only possible plane singularities of $f(x)$ when the algebra is irreducible and has radical 0. They are plane singularities only if c is in G . This last condition is also sufficient as the next lemma shows.

LEMMA 9.1. *If c is in G , then c and 0 are simple poles of $f(x)$.*

Proof. $f(x)$ is clearly defined for all elements $c+\lambda e$ and λe such that $0 < \|\lambda\| < r$, for some real r . The Laurent expansions are

$$\begin{aligned} f(x) &= [x(x-c)]^{-1} = (x-c)^{-1}[(x-c)+c]^{-1} \\ &= (x-c)^{-1}[e+(x-c)c^{-1}]^{-1}c^{-1} = (x-c)^{-1}c^{-1}[e-(x-c)c^{-1}+\dots], \\ f(x) &= c^{-1}(x-c)^{-1} - c^{-2} + c^{-3}(x-c) - c^{-4}(x-c)^2 + \dots \end{aligned}$$

where the series converges super-absolutely at least for $\|x-c\| < 1/\|c^{-1}\|$; similarly, $f(x) = x^{-1}(-c^{-1})(e-xc^{-1})^{-1} = -c^{-1}x^{-1}[e+xc^{-1}+x^2c^{-2}+\dots]$, where this series converges super-absolutely at least for $0 < \|x\| < 1/\|c^{-1}\|$.

For reducible algebras we have the following

LEMMA 9.2. *If B is a reducible algebra with radical $\{0\}$, any plane singularity is a pole of order at most 2.*

Proof. Let z be a plane singularity of $f(x)$. The coefficients of the Laurent expansion around z are given by $a_n = (1/2\pi i) \int_K (x-z)^{-n-1} f(x) dx$ where K is the curve $\{z + (r \exp 2\pi i t)e\}$ and $r > 0$ is such that both z and $z-c$ have their spectra outside the circle $C = \{r \exp 2\pi i t\}$ except for the point 0. Then we have

$$a_n = (1/2\pi i) \int_C (z + \zeta e)^{-1} (z - c + \zeta e)^{-1} (\zeta e)^{-n-1} d\zeta \quad \text{where } \zeta \text{ is on } C,$$

$a_n(M) = (1/2\pi i) \int_C (z(M) + \zeta)^{-1} [\zeta + z(M) - c(M)]^{-1} \zeta^{-n-1} d\zeta$ for all n and all M in \mathfrak{M} . In particular, for $n = -3$, $a_{-3}(M) = (1/2\pi i) \int_C (z(M) + \zeta)^{-1} [z(M) - c(M) + \zeta]^{-1} \zeta^2 d\zeta$. There are four cases:

(i) $c(M) = 0$ and $z(M) \neq 0$. In this case, $a_{-3}(M) = (1/2\pi i) \int_C \zeta^2 (\zeta + z(M))^{-2} d\zeta$ and since $z(M)$ is exterior to C , $a_{-3}(M) = 0$ for all M .

(ii) $c(M) \neq 0$ and $z(M) = 0$. Then $a_{-3}(M) = (1/2\pi i) \int_C \zeta^{-1} (\zeta - c(M))^{-1} \zeta^2 d\zeta = (1/2\pi i) \int_C (\zeta^2/c(M)) [1/(\zeta - c(M)) - 1/\zeta] d\zeta = 0$ since $c(M)$ is outside C .

(iii) $c(M) = 0$ and $z(M) = 0$. Then $a_{-3}(M) = (1/2\pi i) \int_C c d\zeta = 0$.

(iv) $c(M) \neq 0$ and $z(M) \neq 0$. Then

$$\begin{aligned} a_{-3}(M) &= \frac{1}{2\pi i} \int_C \frac{\zeta^2}{-c(M)} \left[\frac{1}{\zeta + z(M)} - \frac{1}{\zeta + z(M) - c(M)} \right] d\zeta \\ &= -\frac{1}{2\pi i} \int_C \frac{1}{c(M)} \left[\frac{\zeta^2}{\zeta + z(M)} \right] d\zeta \\ &\quad + (1/2\pi i c(M)) \int_C \frac{\zeta^2 d\zeta}{\zeta + z(M) - c(M)} = 0 \end{aligned}$$

since $z(M)$ and $(z-c)(M)$ lie outside C .

Similar treatment may be accorded the coefficients a_n for $n < -3$. Since the radical is $\{0\}$, $a_n = 0$ for $n \leq -3$ so that z is a pole of order at most 2.

In reducible algebras, if c is a pole and e_j is an idempotent, then $e_j c$ is also a pole since $(e_j c)(M) = e_j(M) c(M)$ and $e_j(M) = 0$ or 1 ; i.e. $(e_j c)(M) = 0$ or $c(M)$, and $(e_j c - c)(M) = 0$ or $-c(M)$. If c is a pole, then $c(M) = 0$ or $|c(M)| > \epsilon > 0$. Hence $e_j c$ has the requisite spectral properties.

Of course, c may not be a singularity⁽²⁾ at all; i.e. if c is not in G . In reducible algebras, this does not preclude the existence of other singularities as it does in the irreducible case. For example, in the algebra (m) , let c be a sequence $\{1/n\}$, which is not in G . Then $z = (1, 0, 1, 1, \dots, 1, \dots)$ is a pole of $f(x)$, for the spectrum of z consists of 0 and 1, while the spectrum of $z-c = \{0, -1/2, 1-1/3, \dots, 1-1/n, \dots\}$ consists of all the numbers in

(²) Hereinafter singularity will mean plane-singularity.

that sequence and the number 1. Hence 0 is an isolated point. This is sufficient to make z a plane singularity. The lemma shows that it is a pole.

The above example reveals that a zero of $p(x)$ need not be a singularity of $f(x)$. Conversely, the pole z need not be a zero as the example shows. Thus any attempt to locate zeros by contour integration appears doomed to failure.

We have shown that the residues of x^{-1} can be obtained by integrating on curves of the form $\{z_0 + \phi(t) \cdot e\}$ where z_0 is a pole of x^{-1} . This is a type of residue theorem. We should like to prove a similar result for $f(x) = x^{-1}(x - c)^{-1}$.

Let e_j be an idempotent and let $\mathfrak{M}_{0j} = \{M | e_j(M) = 0\}$ and $\mathfrak{M}_{1j} = \{M | e_j(M) = 1\}$. Then $\mathfrak{M} = \mathfrak{M}_{0j} \cup \mathfrak{M}_{1j}$ and both sets \mathfrak{M}_{0j} and \mathfrak{M}_{1j} are closed in \mathfrak{M} , and therefore compact. We have the following

LEMMA 9.3. *A necessary and sufficient condition for $xc = e_j$ to have solutions is that $c(M) \neq 0$ for all M in \mathfrak{M}_{1j} .*

Proof. Necessity. If $c_jc = e_j$, then $c_j(M)c(M) = e_j(M) = 1$ for all M in \mathfrak{M}_{1j} . Hence $c(M) \neq 0$ for all such M . If $e_j = 0$, then \mathfrak{M}_{1j} is empty and the lemma is trivial. If $e_j = e$, then $c_j = c^{-1}$.

Sufficiency. We shall prove $c_j = [e_jc + (e - e_j)]^{-1}e_j$ is a solution of $xc = e_j$. First, $[e_jc + (e - e_j)](M) = e_j(M) \cdot c(M) + 1 - e_j(M)$. For M in \mathfrak{M}_{1j} , this equals $c(M) \neq 0$. For M in \mathfrak{M}_{0j} , it equals 1. Thus $[e_jc + (e - e_j)]^{-1}$ exists.

Let $d = c_jc$. Then $d[e_jc + (e - e_j)] = e_jc$. But $e_j[e_jc + (e - e_j)] = e_jc$. Therefore, $(d - e_j)[e_jc + (e - e_j)] = 0$. Since $[e_jc + (e - e_j)]^{-1}$ exists, $d = e_j$, which proves the lemma.

We observe that there may be more than one solution of the above equation. In the algebra (m) , for example, if $c = (2, 0, 0, \dots)$ and $e_j = (1, 0, 0, \dots)$, then $c = (1/2, \delta_2, \delta_3, \dots)$ where δ_n may be chosen arbitrarily for $n \geq 2$, except that the $\{|\delta_n|\}$ must be bounded.

We now prove the main theorem on the residues of $f(x) = x^{-1}(x - c)^{-1}$.

THEOREM 9.1. *If the radical of B is $\{0\}$, and d is a residue of $f(x)$, then d is of the form $2\pi i \sum_{j=1}^p n_j c_j e_j$ where $e_j^2 = e_j$, $e_j e_k = 0$ for $j \neq k$, and $c_j c = e_j$ with $|c(M)| > 0$ for M in $\cup_{j=1}^p \mathfrak{M}_{1j}$.*

Conversely, if $|c(M)| > 0$ for all M in $\cup_{j=1}^p \mathfrak{M}_{1j}$, then there is a curve K in $G \cap (c + G)$ such that $\int_K f(x) dx = d$ where $cd = 2\pi i \sum_{j=1}^p n_j e_j$.

Proof. Suppose $\int_K f(x) dx = d$. For all M , we have $d(M) = \int_{K(M)} d\xi / \xi [\xi - c(M)]$. Thus $d(M) = (1/c(M)) 2\pi i n$ where $n = 0, \pm 1, \pm 2, \dots$ if $c(M) \neq 0$. If $c(M) = 0$, then $d(M) = 0$. Hence, $(cd)(M) = 2\pi i n$ where n is a rational integer or zero. This implies $[\exp(cd)](M) = 1$ for all M . Since the radical is $\{0\}$, $\exp(cd) = e$. Therefore, $cd = 2\pi i \sum_{j=1}^p n_j e_j$.

Now, the \mathfrak{M}_{1j} are disjoint in pairs, for suppose M is in $\mathfrak{M}_{1j} \cap \mathfrak{M}_{1k}$, $j \neq k$. Then $0 = (e_j e_k)(M) = e_j(M) e_k(M) = 1$, which is a contradiction. If $b = 2\pi i \cdot \sum_{j=1}^p n_j e_j$, we see that $b(M) = 2\pi i n_k \neq 0$ for M in $\cup_{j=1}^p \mathfrak{M}_{1j}$, $1 \leq k \leq p$. Hence

$(cd)(M) = b(M) \neq 0$ for M in $\cup_{j=1}^p \mathfrak{M}_{1j}$ and therefore $d(M) \neq 0, c(M) \neq 0$ for these M .

By the preceding lemma, c_j exists for $j = 1, \dots, p$. Let $d' = 2\pi i \sum_{j=1}^p n_j c_j e_j$. If M is not in $\cup_{j=1}^p \mathfrak{M}_{1j}$, then $d'(M) = 0$, and $(cd)(M) = b(M) = 0$. If $c(M) \neq 0$, then $d(M) = 0$. If $c(M) = 0$, then $d(M) = 0$ by the third line of this proof. Thus $d(M) = 0$ for all $M \notin \cup_{j=1}^p \mathfrak{M}_{1j}$. If M is in \mathfrak{M}_{1j} for some j , then $d'(M) = 2\pi i n_j c_j(M) = 2\pi i n_j / c(M)$. Since $(cd)(M) = 2\pi i n_j$ for this M , we have $d(M) = 2\pi i n_j / c(M)$. Thus $d(M) = d'(M)$ for all M . Since the radical is $\{0\}$, $d = d'$ as was to be shown.

Now suppose $b = 2\pi i \sum_{j=1}^p n_j e_j$ where $e_j^2 = e_j, e_j e_k = 0$ for $j \neq k$ and $|c(M)| > 0$ for M in $\cup_{j=1}^p \mathfrak{M}_{1j}$. Let K be the curve parametrized as follows: $z(t) = r \exp(2\pi i t b) + R(e - \sum_{j=1}^p e_j)$ where $0 < r < \min\{|c(M)|, M \text{ in } \cup_{j=1}^p \mathfrak{M}_{1j}\}$ and $R > r + \|c\|$. (Since $\cup_{j=1}^p \mathfrak{M}_{1j}$ is compact, $\min_{M \in \cup_{j=1}^p \mathfrak{M}_{1j}} \{|c(M)|\} > 0$.) We shall prove that $z(t)$ is in $G \cap (c + G)$ for $0 \leq t \leq 1$.

Let M be in $\cup_{j=1}^p \mathfrak{M}_{1j}$, say M in \mathfrak{M}_{1k} . Then $b(M) = 2\pi i n_k$ and $[z(t)](M) = r \exp(2\pi i n_k t) + R(1 - 1) = r \exp(2\pi i n_k t) \neq 0$. If M' is in $\cap_{j=1}^p \mathfrak{M}_{0j}$, then $e_j(M') = 0$ for $j = 1, \dots, p$. Hence $[z(t)](M') = r + R > 0$. Thus $[z(t)](M) \neq 0$ for all M in \mathfrak{M} ; i.e. $z(t)$ is in G for $0 \leq t \leq 1$.

Similarly, for M in $\cup_{j=1}^p \mathfrak{M}_{1j}$, $|[z(t) - c](M)| = |r \exp(2\pi i n_k t) - c(M)| \geq |c(M)| - r \geq \mu - r > 0$ where $\mu = \min\{|c(M)|, M \text{ in } \cup_{j=1}^p \mathfrak{M}_{1j}\}$. For M not in $\cup_{j=1}^p \mathfrak{M}_{1j}$, $|[z(t) - c](M)| = |r + R - c(M)| \geq R - r - |c(M)| \geq R - r - \|c\| > 0$. Thus $[z(t) - c]$ is in G for $0 \leq t \leq 1$.

The integrand $f(x)$ is therefore defined for all x on K . The integration proceeds as follows:

$$\begin{aligned} & \int_K f(x) dx \\ &= \int_0^1 \frac{2\pi i b r \exp(2\pi i t b) dt}{\left[r \exp(2\pi i t b) + R \left(e - \sum_1^p e_j \right) \right] \left[r \exp(2\pi i t b) + R \left(e - \sum_1^p e_j \right) - c \right]} \\ &= \sum_{j=1}^p e_j \int_0^1 \frac{2\pi i b r \exp(2\pi i t b) dt}{z(t) [z(t) - c]} = \sum_{j=1}^p e_j \int_K x^{-1} (x - c)^{-1} dx \\ &= \sum_{j=1}^p \int_K e_j x^{-1} (x - c)^{-1} dx = \sum_{j=1}^p \int_K c_j [(x - c)^{-1} - x^{-1}] dx \end{aligned}$$

since $c_j c = e_j$ and c_j exists by the Lemma 9.3.

Now let $I_0 = \int_K x^{-1} dx = \int_0^1 [2\pi i b r \exp(2\pi i t b)] \cdot [z(t)]^{-1} dt$. For M in $\cup_{j=1}^p \mathfrak{M}_{1j}$, $I_0(M) = \int_0^1 2\pi i n_k r \exp(2\pi i n_k t) / (r \exp(2\pi i n_k t)) dt = 2\pi i n_k$. For M not in $\cup_{j=1}^p \mathfrak{M}_{1j}$, $I_0(M) = \int_0^1 0 / (r + R) dt = 0$. Thus $I_0(M) = b(M)$ for all M . Since the radical is $\{0\}$, $I_0 = b$.

Let $I_c = \int_K (x - c)^{-1} dx = \int_0^1 2\pi i b r \exp(2\pi i t b) [z(t) - c]^{-1} dt$. For M in $\cup_{j=1}^p \mathfrak{M}_{1j}$,

$$I_c(M) = \int_0^1 \frac{2\pi i n_k r \exp(2\pi i n_k t)}{r \exp(2\pi i n_k t) - c(M)} dt.$$

Letting $\zeta(t) = r \exp(2\pi i n_k t)$, we have $d\zeta = 2\pi i n_k \zeta dt$ so that

$$I_c(M) = \int_{K'} \left(\frac{1}{\zeta - c(M)} \right) d\zeta$$

where K' is the closed curve $r \exp(2\pi i n_k t)$, $0 \leq t \leq 1$, in the complex plane. Since $|c(M)| \geq \mu > r$, $c(M)$ lies outside of K' for all M in $U_1^p \mathfrak{M}_{1j}$. Hence, $I_c(M) = 0$. Likewise, for M not in $U_1^p \mathfrak{M}_{1j}$, $I_c(M) = \int_0^1 0 / (R + r - c(M)) dt$ which equals 0. Thus $I_c = 0$.

Therefore, $\int_{K'} f(x) dx = - \sum_{j=1}^p c_j b$. Since $b = 2\pi i \sum_{j=1}^p n_j e_j$, and $c_j e_k = 0$ for $j \neq k$ (i.e. $[c_j e_k](M) = c_j(M) e_k(M) = 0$ since $c_j(M) = 0$ for M in \mathfrak{M}_{1k} and $e_k(M) = 0$ for M not in \mathfrak{M}_{1k}), we have

$$\int_{K'} f(x) dx = - 2\pi i \sum_{j=1}^p n_j c_j e_j.$$

Reversing the parametrization of K then yields the desired result.

We now demonstrate that all residues of $f(x)$ may be expressed as a sum of residues obtained by integrating on curves of the form $\{z_0 + \psi(t)e\}$ where z_0 is a pole of $f(x)$ and $\psi(t)$ is a continuous complex function, $0 \leq t \leq 1$. This is an analogue of the residues theorem for $f(x)$.

Let d be a residue of $f(x)$. Thus $d = 2\pi i \sum_{j=1}^p n_j c_j e_j$. Again, let $\mu = \min \{|c(M)|, M \text{ in } U \mathfrak{M}_{1j}\}$ and $\mu > r > 0, R > r + \|c\|$.

THEOREM 9.2. *The elements $z_j = R(e - e_j), j = 1, \dots, p$, are poles of $f(x)$ and the integral of $f(x)$ on the curve $K_j: \{z_j + r \exp 2\pi i n_j t, 0 \leq t \leq 1\}$ is equal to $-2\pi i n_j c_j$. Thus $\sum_{j=1}^p \int_{K_j} f(x) dx = -d$.*

To prove that z_j is a pole we must show that $f(z_j + \lambda e)$ is defined for $0 < |\lambda| < r + \epsilon$, where $r + \epsilon \|c\| < R, \epsilon > 0, r + \epsilon < \mu$.

If M is in \mathfrak{M}_{1j} , then $|(z_j + \lambda e)(M)| = |0 + \lambda| \neq 0$. If M is not in \mathfrak{M}_{1j} , $|(z_j + \lambda e)(M)| = |R + \lambda| \geq R - |\lambda| > 0$. Thus $(z_j + \lambda e)$ is in G for all λ such that $0 < |\lambda| < r + \epsilon$. Likewise, $|(z_j + \lambda e - c)(M)| = |\lambda - c(M)| \geq |c(M)| - |\lambda| \geq \mu - (r + \epsilon) > 0$ for M in \mathfrak{M}_{1j} and $|(z_j + \lambda e - c)(M)| = |R + \lambda - c(M)| \geq R - \|c\| - r - \epsilon > 0$ for M not in \mathfrak{M}_{1j} . Thus $(z_j + \lambda e)$ is in $(c + G)$ and z_j is a pole.

Let $d_j = \int_{K_j} f(x) dx = \int_0^1 2\pi i n_j r \exp(2\pi i n_j t) [z_j + r \exp 2\pi i n_j t]^{-1} [z_j + r \exp 2\pi i n_j t - c]^{-1} dt$. Then

$$d_j(M) = \int_0^1 \frac{2\pi i n_j r \exp(2\pi i n_j t) dt}{r \exp(2\pi i n_j t) [r \exp(2\pi i n_j t) - c(M)]} = \int_{K_j(M)} \frac{d\zeta}{\zeta [\zeta - c(M)]}$$

for M in \mathfrak{M}_{1j} . Here $K_j(M) = \{r \exp(2\pi i n_j t)\}$. Since $|c(M)| > r$ for M in \mathfrak{M}_{1j} , $d_j(M) = -2\pi i n_j / c(M)$. For M in \mathfrak{M}_{0j} , we have

$$d_j(M) = \int_0^1 \frac{2\pi i n_j r \exp(2\pi i n_j t) dt}{[R + r \exp 2\pi i n_j t][R + r \exp(2\pi i n_j t) - c(M)]}$$

$$= \int_{K_j(M)} \frac{d\zeta}{\zeta[\zeta - c(M)]} = 0$$

since $\zeta = R + r \exp(2\pi i n_j t)$ and $|\zeta| \geq R - r > \|c\| \geq |c(M)|$; i.e. $c(M)$ is exterior to $K_j(M) = \{R + r \exp(2\pi i n_j t)\}$. Thus $d_j(M)c(M) = -2\pi i n_j e_j(M)$ for all M . Since the radical is $\{0\}$, $d_j c = -2\pi i n_j e_j$, or $d_j = -2\pi i n_j c_j$. Clearly, $\sum_{j=1}^p d_j = -d$ or $\sum_{j=1}^p \int_{K_j} \dot{x}^{-1}(x-c)^{-1} dx = -d$ as was to be shown.

We conclude the discussion of the function $x^{-1}(x-c)^{-1}$ with some further remarks on the topological structure of its domain $G \cap (c+G)$.

If c is in G , then the component cG_1 of G intersects every component of $(c+G)$. In proof, let $(c+gG_1)$ be any component of $(c+G)$. Let g_1 be an element of G_1 such that $\|g_1\| \|g\| < 1/\|c^{-1}\|$. Then we have $\|(c+gg_1) - c\| = \|gg_1\| \leq \|g\| \|g_1\| < 1/\|c^{-1}\|$ and $(c+gg_1)$ is in cG_1 . Correspondingly, the component $(c+cG_1)$ intersects every component of G , for if g in G is of sufficiently small norm, then $g-c$ is in cG_1 and $c+(g-c) = g$ is in $(c+cG_1)$.

This by no means determines the set-theoretic structure of $G \cap (c+G)$ for it is not known whether a component gG_1 , other than cG_1 , intersects any component of $(c+G)$ except its translate $(c+gG_1)$ and $(c+cG_1)$, or whether these intersections are connected.

The discussion for $f(x) = x^{-1}(x-c)^{-1}$ may be extended to the case $[p_n(x)]^{-1}$ where $p_n(x)$ is a polynomial which is of degree n and splits into n linear factors. If $p_n(x)$ does not split into linear factors, the situation is more difficult. If $D(p_n)$ denotes the set of z in B such that $[p_n(z)]^{-1}$ exists and $O(p_n)$ is its complement, then we can state the following

THEOREM 9.3. *If $p_n(x) = a_0 + a_1x + \dots + a_nx^n$ is such that $O(p_n)$ and $D(p_n)$ are not empty, then $f_n(x) = [p_n(x)]^{-1}$ is a meromorphic function; i.e. it has no plane singularities other than poles. In fact, the poles are of order not greater than n , assuming that the radical is $\{0\}$.*

Proof. The domain of analyticity of $f_n(x)$ consists of the various components of $D(p_n)$. Suppose z_0 is a plane singularity. Then z_0 is in $O(p_n)$ and $z_0 + \lambda e$ is in $D(p_n)$ for all λ such that $0 < |\lambda| < r_1$. Let $f_n(x) = \sum_{-\infty}^{\infty} b_m(x-z_0)^m$ be the Laurent expansion around z_0 , where $b_m = (1/2\pi i) \int_K f_n(x)(x-z_0)^{-m-1} dx$, $m=0, \pm 1, \pm 2, \dots$, and K is the curve $\{z_0 + (r \exp 2\pi i t)e\}$, $0 < r < r_1$. For M in \mathfrak{M} , we have

$$b_m(M) = \frac{1}{2\pi i} \int_{K(M)} \frac{\lambda_t^{-m-1} d\lambda_t}{(a_0(M) + a_1(M)\zeta + \dots + a_N(M)\zeta^N)}$$

where $a_N(M) \neq 0$ and $0 < N \leq n$ and $\zeta = z_0(M) + \lambda_t$, $\lambda_t = r \exp(2\pi i t)$. Factoring the complex poly-

nomial, we obtain $a_0(M) + a_1(M)\zeta + \cdots + a_N(M)\zeta^N = [a^N(M)](\zeta - \lambda_1) \cdots (\zeta - \lambda_N)$. Thus

$$b_m(M) = \frac{1}{2\pi i} \int_{K(M)} \frac{\lambda_t^{-m-1} d\lambda_t}{a_N(M) [\lambda_t - (\lambda_1 - z_0(M))] \cdots [\lambda_t - (\lambda_N - z_0(M))]}.$$

If $\lambda_k - z_0(M) \neq 0$, then $|\lambda_k - z_0(M)| \geq r_1$ for $k=1, \dots, N$, for otherwise $z_0 + (\lambda_k - z_0(M))e$ is in $D(p_n)$. This is impossible because

$$[p_n(z_0 + (\lambda_k - z_0(M))e)](M) = a_N(M) [z_0(M) + \lambda_k - z_0(M) - \lambda_1] \cdots [z_0(M) + \lambda_k - z_0(M) - \lambda_N]$$

which equals zero. Thus for $m < -n$, the integrand in the expression for $b_m(M)$ is analytic interior to the curve $\lambda_t = r \exp(2\pi it)$ and also on the curve. Hence, for such m , $b_m(M) = 0$. This holds for all M . Since the radical is $\{0\}$, $b_m = 0$.

As a concluding remark, we observe that the theory of the general rational function $p_m(x)[p_n(x)]^{-1}$, where $p_n(x)$ and $p_m(x)$ are polynomials, is not immediately obtainable from the theory of $[p_n(x)]^{-1}$ because the domain of analyticity may possibly extend beyond $D(p_n)$.

BIBLIOGRAPHY

1. E. R. Lorch, *The theory of analytic functions in normed abelian vector rings*, Trans. Amer. Math. Soc. vol. 54 (1943) pp. 414-425.
2. ———, *The structure of normed abelian rings*, Bull. Amer. Math. Soc. vol. 50 (1944) pp. 447-463.
3. I. Gelfand, *Normierte Ringe*, Rec. Math. (Mat. Sbornik) N.S. vol. 9 (51) (1941) pp. 3-24.
4. E. Hille, *Functional analysis and semi-groups*, Amer. Math. Soc. Colloquium Publications, vol. 31, 1948.
5. N. Dunford, *Spectral theory*, Bull. Amer. Math. Soc. vol. 48 (1943) pp. 637-651.
6. A. E. Taylor, *Analysis in complex Banach spaces*, Bull. Amer. Math. Soc. vol. 49 (1943) pp. 652-669.
7. M. Nagumo, *Einige analytische Untersuchungen in linearen metrischen Ringen*, Jap. J. Math. (1936).
8. S. Mazur, *Sur les anneaux linéaires*, C. R. Acad. Sci. Paris vol. 207 (1938) p. 1025.
9. I. Gelfand and G. Silov, *Über verschiedene Methoden der Einführung der Topologie in die Menge der maximalen Ideale eines normierten Ringes*, Rec. Math. (Mat. Sbornik) N.S. vol. 9 (51) (1941) pp. 25-39.
10. C. E. Rickart, *B-algebras with an adjoint operation*, Ann. of Math. vol. 47 (1946) pp. 528-550.
11. ———, *Singular elements of a B-algebra*, Duke Math. J. vol. 14 (1947) pp. 1063-1077.
12. E. K. Blum, *The fundamental group of the principal component of a commutative Banach algebra*, Proc. Amer. Math. Soc. vol. 4 (1953) pp. 397-400.

COLUMBIA UNIVERSITY,
NEW YORK, N. Y.
UNIVERSITY OF MARYLAND,
COLLEGE PARK, MD.