

STUDIES IN FABER POLYNOMIALS. I

BY

JOSEPH L. ULLMAN

1. **Introduction.** The research on the following problem was suggested by Dr. G. Szegő, to whom much is owed, both for general encouragement and for many specific suggestions. In 1903, in his thesis, G. Faber⁽¹⁾ [1] introduced polynomials, which have since proved useful in Analysis, and have become known as Faber polynomials. The results of the following research concern the asymptotic distribution of the zeros of Faber polynomials.

DEFINITION 1.1. Let $f(z)$ be a power series of the type

$$(1.1) \quad z + a_0 + \frac{a_1}{z} + \cdots,$$

where the coefficients satisfy the condition that $\limsup |a_n|^{1/n} < \infty$. For each $n \geq 1$, form

$$(1.2) \quad [f(z)]^n = F_n(z) + R_n(z),$$

where $F_n(z)$ is a polynomial of degree n and $R_n(z)$ contains only negative powers of z . The polynomials $F_n(z)$ are the Faber polynomials associated with $f(z)$.

This is equivalent to Faber's original definition, except that he specified that $f(z)$ be the Taylor's series expansion about $z = \infty$ of the normalized exterior mapping function of a simple Jordan curve C in the z -plane. He referred to the polynomials as being associated with the curve C . In his thesis, he proved that an arbitrary analytic function $G(z)$, regular in the interior of C , could be expanded as a series of the Faber polynomials associated with C ,

$$(1.3) \quad G(z) = \sum_0^{\infty} a_n F_n(z),$$

such that the series converges to $G(z)$ uniformly in any closed subset of the interior of C . In proving this result, Faber derived and used the generating function for the polynomials. It is derived here by a method adapted to later applications.

Consider a series of type (1.1), with $\limsup |a_n|^{1/n} < \infty$. Because of the restriction on the coefficients, the series converges uniformly in a neighborhood of $z = \infty$. Indeed, a positive number γ can be chosen, so that in the closed exterior of the circle Γ , $|z| = \gamma$, the series defines a function that is called $w = f(z)$ without ambiguity, which is analytic except for a simple pole

Received by the editors February 25, 1959.

(¹) Numbers in brackets refer to the bibliography at the end of the paper.

at $z = \infty$, and performs a 1-1 mapping of the exterior of Γ onto the exterior of a simple analytic curve Γ' . These properties are preserved if γ is replaced by a larger number, so we can assume that $\gamma > |\zeta|$, where ζ is an arbitrary complex number. Consider the integral

$$(1.4) \quad I(\zeta) = \frac{1}{2\pi i} \int_{\Gamma^+} \frac{[f(z)]^n}{z - \zeta} dz.$$

Substitute (1.2) to obtain

$$(1.5) \quad I(\zeta) = \frac{1}{2\pi i} \int_{\Gamma^+} \frac{F_n(z)}{z - \zeta} dz + \frac{1}{2\pi i} \int_{\Gamma^+} \frac{R_n(z)}{z - \zeta} dz.$$

Since the series $f(z)$ represents an analytic function for z in the closed exterior of Γ , the same is true of $[f(z)]^n$, and therefore $R_n(z)$. Since $R_n(z)$ contains only negative powers of z , $R_n(z) = O(1/|z|)$. In the second integral on the right, the value is not changed by replacing Γ by a circle of radius ρ , where $\rho > \gamma$. Since $R_n(z) = O(1/|z|)$, the integral tends to zero as $\rho \rightarrow \infty$, and must therefore equal zero. Since ζ is in Γ , (1.5) becomes

$$(1.6) \quad I(\zeta) = F_n(\zeta) = \frac{1}{2\pi i} \int_{\Gamma^+} \frac{[f(z)]^n}{z - \zeta} dz.$$

Next, make the change of variable $z = g(w)$ in (1.6) where $g(w)$ is the inverse function of $f(z)$, obtaining

$$(1.7) \quad F_n(\zeta) = \frac{1}{2\pi i} \int_{\Gamma_1^+} \frac{w^n g'(w)}{g(w) - \zeta} dw.$$

The function $g'(w)/(g(w) - \zeta)$ has a simple zero at $w = \infty$, and therefore has an expansion of the type

$$(1.8) \quad \frac{g'(w)}{g(w) - \zeta} = \sum_0^{\infty} \frac{b_n(\zeta)}{w^{n+1}}.$$

Equation (1.7) is the Cauchy integral formula for the coefficient of $1/w^{n+1}$ in the right hand side of (1.8), hence $b_n(\zeta) = F_n(\zeta)$ and

$$(1.9) \quad \frac{g'(w)}{g(w) - \zeta} = \sum_0^{\infty} \frac{F_n(\zeta)}{w^{n+1}}.$$

This is the generating function of the Faber polynomials. The domain of convergence of the right hand side of (1.9) depends on ζ . The exact relationship between the radius of convergence of the series in (1.9) and the value of ζ is derived in §4.

2. **The zeros in a special case.** Let $\mathfrak{F}(z)$ be the branch of $(z^2 - 1)^{1/2}$ which is holomorphic in the z -plane cut along the real axis from $z = -1$ to $z = 1$, and

is positive for z real and greater than one. Let $f(z)$ also designate the Taylor's series expansion about $z = \infty$ of $\mathfrak{F}(z)$. We study the Faber polynomials associated with $f(z)$ in this section. The following results on the zeros of partial sums of Taylor's series are used. Let $h(z) = \sum_0^\infty a_k z^k$, $\limsup |a_k|^{1/k} = 1$. Let $h_n(z) = \sum_0^n a_k z^k$ have the zeros z_{ni} ($i=1, \dots, n$) which we designate by (z_{ni}) . Let $\{(z_{ni})\}$ denote the sequence of sets of points $(z_{1,i}), (z_{2,i}), \dots$. R. Jentzsch [3] proved that every point of the circle $|z| = 1$ is a limit point of the elements of $\{(z_{ni})\}$. Later G. Szegö [4] proved a generalization which we refer to as the Szegö-Jentzsch theorem. Consider only zeros of sections $h_{n_\nu}(z)$ for which $\lim |a_{n_\nu}|^{1/n_\nu} = 1$, indicating them by $\{(z_{n_\nu,i})\}$. These points are distributed with the following properties. Let $r(n)$ be a function which is $o(n)$. For any $\epsilon > 0$, the points $(z_{n_\nu,i})$ satisfy $||z_{n_\nu,i}| - 1| < \epsilon$, except for at most $r(n)$ values of i . Let $\arg z_{n_\nu,i} = \theta_{n_\nu,i}$, let α and β be numbers satisfying $0 \leq \alpha < \beta < 2\pi$ and let $\lambda_{n_\nu}(\alpha, \beta)$ be the number of $\theta_{n_\nu,i}$ satisfying $\alpha \leq \theta_{n_\nu,i} \leq \beta$, then

$$(2.1) \quad \lambda_{n_\nu}(\alpha, \beta) = n_\nu \left(\frac{\beta - \alpha}{2\pi} \right) + o(n_\nu).$$

Points $\{(z_{n_\nu,i})\}$ satisfying these two properties are said to be uniformly distributed about the circle $|z| = 1$. A greater degree of generality is required to discuss the zeros of the Faber polynomials of $f(z)$.

Let Γ be a curve consisting of finitely many analytic arcs $\Gamma_1, \dots, \Gamma_k$, touching only at their endpoints, and forming the boundary of a simply connected domain D which contains the point $z = \infty$. Let $w = F(z)$ be the holomorphic function which maps D conformally onto the exterior of the circle Σ , $|w| = 1$, in a one-to-one manner, taking $z = \infty$ into $w = \infty$ and satisfying the condition

$$\lim_{z \rightarrow \infty} \frac{F(z)}{z} = k,$$

where k is a positive constant. Let Γ' be the set of points of Γ at a distance greater than δ from the end points of the arcs Γ_i . Let D^* be the domain consisting of D and every other point in the z -plane whose distance from Γ is less than γ . For any δ, γ can be chosen so that D^* is simply covered, and so that $F(z)$ can be uniquely continued into D^* by the reflection principle and remain one-to-one. Call this holomorphic function $F^*(z)$.

A sequence of sets of points $\{(z_{n_\nu,i})\}$, ($n_1 < n_2 < \dots$; $i=1, \dots, n$) is said to be conformally distributed about the curve Γ if it has the following properties. Let $s(n)$ be a function which is $o(n)$. For any $\epsilon > 0$, the points $(z_{n_\nu,i})$ must be within a distance ϵ of Γ , except for at most $s(n)$ values of i . Let $(a, b)^\wedge$ be a closed subarc of Γ consisting of interior points of one of the arcs Γ_i , and whose image on Σ is $(e^{i\alpha}, e^{i\beta})^\wedge$. Choose δ so that a, b is contained

in Γ' and then choose γ so that $F^*(z)$ will be one-to-one in D^* . Let $\mu_{n_r}(a, b)$ be the number of $(z_{n_r, i})$ for which $F^*(z)$ is defined and satisfies $\alpha \leq \arg F^*(z_{n_r, i}) \leq \beta$. Then

$$(2.2) \quad \mu_{n_r}(a, b) = n_r \left(\frac{\beta - \alpha}{2\pi} \right) + o(n_r).$$

The Faber polynomials of $f(z)$ of even degree are given by

$$(2.3) \quad F_{2k}(z) = [f(z)]^{2k} = [(z^2 - 1)^{1/2}]^{2k} = (z^2 - 1)^k.$$

Thus the polynomial of degree $2k$ has a zero of multiplicity k at $z=1$ and of multiplicity k at $z=-1$. What we now prove is that the zeros of the polynomials of odd degree are all interior to the lemniscate Γ , $|z^2 - 1| = 1$, except that each has a simple zero at $z=0$, and that the zeros are conformally distributed on Γ . We use the generating function (1.9). The inverse of $w = \mathfrak{F}(z)$ is $z = \mathfrak{G}(w) = (w^2 + 1)^{1/2}$, where $\mathfrak{G}(w)$ designates the branch of $z = (w^2 + 1)^{1/2}$ which is holomorphic in the w -plane cut from i to $-i$, and is positive for positive values of w . Hence

$$(2.4) \quad \sum_0^\infty \frac{F_n(\zeta)}{w^{n+1}} = \frac{1}{(1 + w^{-2})^{1/2}} \cdot \frac{1}{w(1 + w^{-2})^{1/2} - \zeta}.$$

The generating function of the polynomials of odd degree is obtained by replacing w by $(-w)$ in (2.4) and adding the function obtained to (2.4) as follows,

$$(2.5) \quad \sum_0^\infty \frac{F_n(\zeta)}{w^{n+1}} + \sum_0^\infty \frac{F_n(\zeta)}{(-w)^{n+1}} = \frac{1}{(1 + w^{-2})^{1/2}} \cdot \left[\frac{1}{w(1 + w^{-2}) - \zeta} + \frac{1}{-w(1 + w^{-2})^{1/2} - \zeta} \right],$$

$$(2.6) \quad \sum_0^\infty \frac{F_{2k+1}(\zeta)}{w^{2k+2}} = \frac{\zeta}{(1 + w^{-2})^{1/2}} \cdot \frac{1}{w^2 + 1 - \zeta^2}.$$

Let $h(w^2)$ be the Taylor's expansion about $w = \infty$ of $(1 + w^{-2})^{-1/2}$, thus

$$(2.7) \quad h(w^2) = \sum_0^\infty u_k w^{-2k},$$

$$u_k = (-1)^k \frac{1 \cdot 3 \cdot \dots \cdot 2k - 1}{2 \cdot 4 \cdot \dots \cdot 2k}.$$

Let $h_n(w^2) = \sum_0^n u_k w^{-2k}$. We expand the right side of (2.6) in powers of w^{-2} , use the Cauchy rule for product of power series, and obtain

$$\begin{aligned}
 \frac{\zeta}{(1+w^{-2})^{1/2}} \cdot \frac{1}{w^2 - (\zeta^2 - 1)} &= \zeta \sum_0^\infty \frac{u_k}{w^{2k}} \cdot \sum_0^\infty \frac{(\zeta^2 - 1)^k}{w^{2k+2}} \\
 (2.8) \qquad \qquad \qquad &= \zeta \sum_{k=0}^\infty \frac{\sum_{j+l=k} u_j (\zeta^2 - 1)^l}{w^{2k+2}} \\
 &= \sum_{k=0}^\infty \frac{\zeta (\zeta^2 - 1)^k h_k (\zeta^2 - 1)}{w^{2k+2}}.
 \end{aligned}$$

Comparing coefficients of powers of w in (2.6) and (2.8), we obtain the identity

$$(2.9) \qquad F_{2k+1}(\zeta) = \zeta (\zeta^2 - 1)^k h_k (\zeta^2 - 1).$$

The coefficients of the polynomial

$$(2.10) \qquad t^k h_k(t) = \sum_{j=0}^k u_j t^{k-j}$$

decrease in magnitude, hence by a theorem of Kakeya-Eneström the k zeros t_{k1}, \dots, t_{kk} , or (t_{ki}) , all have modulus less than one. The roots of $F_{2k+1}(\zeta)$ are $\zeta_{2k+1,2k+1} = 0$ and the $2k$ solutions of $\zeta^2 - 1 = t_{ki}$ ($i = 1, \dots, k$). Let the two solutions corresponding to t_{ki} be indicated by $\zeta_{2k+1,2i-1}$ and $\zeta_{2k+1,2i}$. Hence $|\zeta_{2k+1,2i-1}^2 - 1| = |\zeta_{2k+1,2i}^2 - 1| = |t_{ki}| < 1$, ($i = 1, \dots, k$), and the zeros of $F_{2k+1}(\zeta)$ are inside the lemniscate Γ , except for the simple zero at $\zeta = 0$. Since $\lim |u_j|^{1/j} = 1$, using the Szegő-Jentzsch theorem, we find that the $\{(t_{ki})\}$ are uniformly distributed about the circle $|t| = 1$. Hence, for any $\rho, 0 < \rho < 1$, $|t_{ki}| > \rho$, except for at most $r(k)$ values of i . Now the lemniscates $\Gamma_\rho, |z^2 - 1| = \rho$, converge uniformly to Γ as ρ tends to one. Thus, for any ϵ, ρ can be chosen so that those points of $(\zeta_{2k+1,j})$ which lie in the annulus between Γ_ρ and Γ are at a distance less than ϵ from Γ . Since there are at most $s(2k+1) = 2r(k)$ points which do not lie in this annulus, the points $(\zeta_{2k+1,i})$ satisfy the first condition for conformal distribution about Γ . We now prove the second condition. The curve Γ consists of two arcs, Γ_1 and Γ_2 , one of which, say Γ_1 , is in the half-plane $\text{Re } z \geq 0$. Let $(a, b)^\wedge$ be a closed sub-arc of interior points of Γ_1 . Let $w = F(z)$ be the normalized exterior mapping function of Γ and let $(e^{i\alpha}, e^{i\beta})^\wedge$ be the image of $(a, b)^\wedge$, $-\pi < \alpha < \beta < \pi$. Of the two solutions of $\zeta^2 - 1 = t_{ki}$, say $\zeta_{2k+1,2i-1}$ is in the half-plane $\text{Re } z > 0$. Further, we note that the function $F(z)$ is the restriction of $\mathfrak{F}(z)$ to the exterior of Γ , which we denote by D . Also, $F^*(z)$ is the restriction of $\mathfrak{F}(z)$ to any properly chosen domain D^* , since the function $\mathfrak{F}(z)$ is one-to-one in the z -plane cut from $z = -1$ to $z = 1$. Now $\alpha \leq \arg \mathfrak{F}(\zeta_{2k+1,j}) \leq \beta$ implies that j is of the form $2i - 1$, and that the argument of a square root of t_{ki} lies between α and β , or $2\alpha \leq \arg t_{ki} \leq 2\beta$. Since $\{(t_{ki})\}$ is uniformly distributed on $|t| = 1$, there are by (2.1), $k((2\beta - 2\alpha)/2\pi) + o(k)$ solutions of this inequality. Furthermore, $F^*(z)$ is de-

finer and coincides with $\mathfrak{F}(z)$ for the $\zeta_{2k+1,j}$ lying in $\text{Re } z > 0$ which correspond to these values of $t_{k,i}$, except for at most $r(k)$ values. Hence

$$\begin{aligned}
 \mu_{2k+1}(a, b) &= \lambda_k(\alpha, \beta) + r(k) \\
 &= k \frac{2\beta - 2\alpha}{2\pi} + o(k) + r(k) \\
 (2.11) \qquad &= (2k + 1) \left(\frac{\beta - \alpha}{2\pi} \right) + o(k),
 \end{aligned}$$

and condition (2.2) is satisfied.

3. The zeros in the general case. Attention is now restricted to $f(z)$, an arbitrary but fixed series of type

$$(3.1) \qquad z + a_0 + \frac{a_1}{z} + \dots,$$

with $\limsup |a_n|^{1/n} < \infty$.

DEFINITION 3.1. Let $z_{n,i}$ ($i=1, \dots, n$) be the zeros of the Faber polynomial $F_n(z)$ associated with $f(z)$, and let Δ be the derived set of the $z_{n,i}$ ($n=1, \dots; i=1, \dots, n$).

To describe the location of Δ , the points of the z -plane are divided into classes, the method of classification depending on properties of $f(z)$ that are next to be discussed. Since the coefficients of (3.1) satisfy the condition $\limsup |a_n|^{1/n} < \infty$, the series (3.1) defines an analytic element of an analytic function. Designate the analytic function by $w=f(z)$. In a neighborhood of $z = \infty$, the series (3.1) defines a portion of $w=f(z)$ which is single valued and univalent. Therefore, the inverse of this portion of $w=f(z)$ is uniquely determined in a neighborhood of $w = \infty$, and has the expansion

$$(3.2) \qquad w + b_0 + \frac{b_1}{w} + \dots,$$

about the point $w = \infty$. The following notation will be used. For any $\eta \geq 0$, let σ_η be the circle $|w| = \eta$, Σ_η the domain $|w| > \eta$, and $\bar{\Sigma}_\eta$ the region $|w| \geq \eta$. If in (3.2), $\limsup |b_n|^{1/n} = \rho$, then (3.2) converges uniformly in any closed subset of Σ_ρ , defining a holomorphic function. Thus, a holomorphic function is defined in a neighborhood of every point of Σ_ρ .

DEFINITION 3.2. Let $z=g(w)$ be the holomorphic function defined by the series (3.2), in Σ_ρ .

Let ζ, ζ', \dots indicate complex constants in the z -plane, and ω, ω', \dots complex constants in the w -plane. Consider the equation

$$(3.3) \qquad g(w) - \zeta = 0.$$

Say $\rho' > \rho$. In $\Sigma_{\rho'}$, (3.3) has either no solution or finitely many solutions, hence

the total number of solutions of (3.3) in Σ_ρ is denumerable, and the solutions can be given an index $\omega_i, (i = 1, 2, \dots)$. Since the domain of definition of $z = g(w)$ is $\Sigma_\rho, |\omega_i| > \rho$. A value ω_i is called a ζ -point of $g(w)$. If $g'(\omega_i) \neq 0$, it is called a simple ζ -point, or a simple solution of (3.3).

DEFINITION 3.3. *The values of ζ for which (3.3) has no solutions are called 0-points, and belong to the set C_0 . Next, consider a value for which (3.3) has solutions. If, among the ζ -points of $g(w)$, there is one, say ω_1 , of greatest modulus, and if ω_1 is a simple ζ -point, then the value ζ is classified as a 1-point. The set of 1-points constitutes the set C_1 in the z -plane.*

The further classification of points in the z -plane is carried out in §4. Definition 3.3 is sufficient for the statement of Theorem 1, and for the proof of its first part.

THEOREM 1. *The set Δ (a) lies in the complement of C_1 , and (b) contains every boundary point of C_1 .*

Proof (a). It is first shown that C_1 contains a neighborhood of $z = \infty$, and that no point of Δ lies in this neighborhood.

LEMMA 3.1. *Given the function $z = g(w)$ of Definition 3.2 with domain of definition Σ_ρ (i) there is a $\rho' > \rho$ and a $P > 0$, such that if $|\zeta| \geq P, g(w) - \zeta = 0$ has one solution in $\Sigma_{\rho'}$, say ω , and it is a simple solution. Furthermore, (ii) the value ω corresponding to ζ is given by a holomorphic function, $w = f(z, \infty, P)$, defined for $|z| \geq P$.*

A proof of (i) is given here, and the proof of (ii) is found in [2, pp. 135-139]. Choose $\rho' > \rho$. Let $\max |g(w)| = M$ for w on $\sigma_{\rho'}$. Choose $P > M$. If ζ is chosen so that $|\zeta| \geq P$, then $|g(w)| < |\zeta|$ for w on $\sigma_{\rho'}$. The argument principle can then be used to show that $g(w) - \zeta = 0$ has a simple solution in $\Sigma_{\rho'}$, since $g(w)$ has a simple pole for $w = \infty$.

Now, if $|\zeta| \geq P$, only one solution of $g(w) - \zeta = 0$ is in $\Sigma_{\rho'}$; hence, any other solution has a smaller modulus. Thus ζ is a 1-point and the set $|z| \geq P$ is contained in C_1 . It is next shown that no point of Δ lies in the set $|z| \geq P$.

Consider the integral

$$(3.4) \quad I(\zeta) = \frac{1}{2\pi i} \int_{\sigma_{\rho'}} \frac{w^n g'(w)}{g(w) - \zeta} dw,$$

where ζ, ρ' are constants introduced in Lemma 3.1. By the lemma, $\omega = f(\zeta, \infty, P)$ is the only solution of $g(w) - \zeta = 0$ in $\Sigma_{\rho'}$. Let $\rho'' > |\omega|$, then

$$(3.5) \quad I(\zeta) = \frac{1}{2\pi i} \int_{\sigma_{\rho''} + \sigma_{\rho'}} \frac{w^n g'(w)}{g(w) - \zeta} dw + \frac{1}{2\pi i} \int_{\sigma_{\rho'}} \frac{w^n g'(w)}{g(w) - \zeta} dw.$$

The integrand has a simple pole at $w = \omega$ with residue $-(\omega)^n = -[f(\zeta, \infty, P)]^n$. The second integral, by (1.6), is equal to $F_n(\zeta)$, hence

$$(3.6) \quad I(\zeta) = F_n(\zeta) - [f(\zeta, \infty, P)]^n.$$

It follows that

$$(3.7) \quad \left| 1 - \frac{F_n(\zeta)}{f(\zeta, \infty, P)^n} \right| \leq \frac{1}{2\pi} \int_{\sigma_{\rho'}} \left| \frac{w}{f(\zeta, \infty, P)} \right|^n \left| \frac{g'(w)dw}{g(w) - \zeta} \right|.$$

Now, $|g(w) - \zeta| \geq |g(w)| - |\zeta| \geq P - M$, for w on $\sigma_{\rho'}$. Also, $\min |f(z, \infty, P)| > \rho' + \delta$, ($\delta > 0$) for $|z| \geq P$. In the contrary case, there would be a value ζ , such that $|\zeta| \geq P$, and $|\omega| = |f(\zeta, \infty, P)| = \rho'$. Hence, it would follow that $g(w) = \zeta$ and thus $|g(w)| \geq P$ for w on $\sigma_{\rho'}$, a contradiction.

Let $Q = \max |g'(w)|$ for w on $\sigma_{\rho'}$. It then follows that

$$(3.8) \quad \frac{1}{2\pi} \int_{\sigma_{\rho'}} \left| \frac{w}{f(\zeta, \infty, P)} \right|^n \left| \frac{g'(w)dw}{g(w) - \zeta} \right| \leq \left(\frac{\rho'}{\rho' + \delta} \right)^n \frac{Q_{\rho'}}{(P - M)}.$$

Hence, for $n > n_0$

$$(3.9) \quad \left| 1 - \frac{F_n(\zeta)}{f(\zeta, \infty, P)^n} \right| < 1.$$

The estimates used in (3.8) are independent of ζ , if $|\zeta| \geq P$, hence, (3.9) is true for $|\zeta| \geq P$. Thus, $F_n(\zeta) \neq 0$ for $|\zeta| \geq P$ and $n > n_0$, and, therefore, no point of Δ lies in the set $|z| \geq P$.

It is next shown that if ζ is a finite complex number in C_1 , then a neighborhood of ζ belongs to C_1 , and that no point of Δ lies in this neighborhood.

LEMMA 3.2. *Given the function $z = g(w)$ of Definition 3.2, with domain of definition Σ_{ρ} , let ζ be a simple 1-point, and ω the ζ -point of $g(w)$ of greatest modulus. (i) There is a $\rho' > \rho$ and a $\delta > 0$, such that if $|\zeta' - \zeta| \leq \delta$, $g(w) - \zeta' = 0$ has one solution ω in $\Sigma_{\rho'}$ and it is a simple solution. Furthermore, (ii) the values of ω are given by a holomorphic function $w = f(z, \zeta, \delta)$, defined for $|z - \zeta| \leq \delta$.*

It is sufficient to give a sketch of (i), since (ii) is proved in [2, pp. 135-139]. The number ρ' can be chosen so that $\rho < \rho' < |w|$, and so that no other ζ points of $g(w)$ lie in $\bar{\Sigma}_{\rho'}$. Say $\min |g(w) - \zeta| = 2\delta$ for w on $\sigma_{\rho'}$, with $\delta > 0$. Then, if $|\zeta' - \zeta| \leq \delta$, $g(w) - \zeta' = 0$ has one solution in $\Sigma_{\rho'}$, and it is simple. This is a consequence of the argument principle.

Thus, if $|\zeta' - \zeta| \leq \delta$, $g(w) - \zeta' = 0$ has one solution in $\Sigma_{\rho'}$; hence, any other solution has smaller modulus. Furthermore, the solution is simple. Hence, ζ' is a 1-point, and the set $|z - \zeta| \leq \delta$ belongs to C_1 .

Since ζ' is a simple 1-point, by Lemma 3.2, the ζ' -point of $g(w)$ of greatest modulus is given by $\omega' = f(\zeta', \zeta', \delta')$, δ' depending on ζ' . On the other hand, since $|\zeta' - \zeta| \leq \delta$, ζ' lies in the domain of definition of $f(z, \zeta, \delta)$, and hence $\omega' = f(\zeta', \zeta, \delta)$.

Consider the integral

$$(3.10) \quad I(\zeta) = \frac{1}{2\pi i} \int_{\sigma_\zeta} \frac{w^n g'(w)}{g(w) - \zeta} dw.$$

In a similar fashion to the treatment of (3.5), this becomes

$$(3.11) \quad I(\zeta) = F_n(\zeta) - [f(\zeta, \zeta, \delta)]^n.$$

If ζ' is a constant satisfying $|\zeta' - \zeta| \leq \delta$, then

$$(3.12) \quad \begin{aligned} I(\zeta') &= F_n(\zeta') - [f(\zeta', \zeta', \delta)]^n \\ &\geq F_n(\zeta') - [f(\zeta', \zeta, \delta)]^n. \end{aligned}$$

Thus, from (3.12), we obtain

$$(3.13) \quad \left| 1 - \frac{F_n(\zeta')}{[f(\zeta', \zeta, \delta)]^n} \right| \leq \frac{1}{2\pi} \int_{\sigma_{\zeta'}} \left| \frac{w}{f(\zeta', \zeta, \delta)} \right|^n \left| \frac{g'(w)dw}{g(w) - \zeta} \right|.$$

Now $|g(w) - \zeta'| = |g(w) - \zeta + \zeta - \zeta'| \geq |g(w) - \zeta| - |\zeta - \zeta'| \geq 2\delta - \delta = \delta$ for w on $\sigma_{\rho'}$, and $|\zeta' - \zeta| < \delta$. Also $f(\zeta', \zeta, \delta) \geq \rho' + \gamma$ for some $\gamma > 0$, since if $f(\zeta', \zeta, \delta) = \rho'$, then $g(w) = \zeta'$ for w on $\sigma_{\rho'}$, in contradiction to the inequality $|g(w) - \zeta'| \geq \delta$. Thus for $n > n_0$,

$$(3.14) \quad \left| 1 - \frac{F_n(\zeta')}{[f(\zeta', \zeta, \delta)]^n} \right| \leq \left| \frac{\rho'}{\rho' + \gamma} \right|^n \left| \frac{Q\rho'}{\delta} \right|,$$

where $Q = \max |g'(w)|$ for w on $\sigma_{\rho'}$. The estimates used are independent of ζ' for $|\zeta - \zeta'| < \delta$, hence for $|\zeta' - \zeta| < \gamma$ and $n > n_0$,

$$(3.15) \quad \left| 1 - \frac{F_n(\zeta')}{[f(\zeta', \zeta, \delta)]^n} \right| < 1.$$

Thus, Δ does not lie in the set $|z - \zeta| \leq \delta$.

This completes the proof that C_1 is an open set containing $z = \infty$, and that Δ lies in the complement of C_1 . The second part of Theorem 1 states that Δ contains all the boundary points of C_1 . The proof of this depends on the classification of the boundary points of C_1 and some properties of these classes. These are given in §4 and the proof of Theorem 1 is completed in §5.

4. The definition of p -points and their properties. Part (b) of Theorem 1 requires a classification of the boundary points of C_1 , and some properties of these classes. This material is contained in this section, and the proof of Theorem 2 is completed in the next section.

DEFINITION 4.1. Let $z = g(w)$ be as defined in Definition 3.2 with domain of definition Σ_ρ . Let ζ be a fixed complex number. If $g(w) - \zeta = 0$ has a solution ω such that $g^1(\omega) = \dots = g^{k-1}(\omega) = 0, g^k(\omega) \neq 0$, then ω is a ζ -point of multiplicity k . If $k = 1$, then ω is a simple ζ -point. Among the ζ -points of $g(w)$, a finite number, $\omega_1, \dots, \omega_n$ will have greatest modulus. Say $|\omega_1| = \dots = |\omega_n| = \rho'$; if

p is the total multiplicity of the ζ -points, then ζ is a p -point. If each ω_i is a simple ζ -point, then ζ is a simple p -point. Otherwise, it is called a multiple p -point.

LEMMA 4.1 (Property 1 and 2). Let ζ be a p -point of $g(w)$, then there is a $\delta > 0$, such that if $|\zeta' - \zeta| < \delta$, ζ' is a q -point, $q \leq p$. Furthermore it is impossible that each ζ' satisfying $|\zeta' - \zeta| < \delta$ is a p -point.

Proof. Let $\omega_1, \dots, \omega_n$ be the solutions of greatest modulus of $g(w) - \zeta = 0$, where $|\omega_1| = \dots = |\omega_n| = \rho'$. Since all other solutions have smaller modulus, ρ'' can be chosen so that $\rho < \rho'' < \rho'$ and no other ζ -points besides $\omega_1, \dots, \omega_n$ lie in $\Sigma_{\rho''}$. Let $\min |g(w) - \zeta| = 2\delta$ for w on $\sigma_{\rho''}$. Then, if $|\zeta' - \zeta| < \delta$, $g(w) - \zeta' = 0$ has solutions of total multiplicity p in $\Sigma_{\rho''}$. Since these need not be of the same modulus, ζ' is a q -point, $q \leq p$. We refer to this fact as property 1. The fact that $q = p$ for each ζ' in any δ neighborhood of ζ is impossible is called property 2, and is next proved.

We first note that only a finite number of ζ' satisfying $|\zeta' - \zeta| < \delta$ can be multiple p -points. To each multiple p -point, ζ' , there is a ζ' -point in $\Sigma_{\rho''}$, say ω' , such that $g'(\omega') = 0$. Now $g'(w) = 0$ has only finitely many solutions in $\Sigma_{\rho''}$, and is single valued, so our conclusion follows. Thus there is a value ζ_1 , and $\delta_1 > 0$ satisfying $|\zeta_1 - \zeta| < \delta$, $\delta_1 < \delta - |\zeta_1 - \zeta|$, such that if $|\zeta' - \zeta_1| < \delta_1$ is a simple p -point. The equation $g(w) = \zeta_1$ has p solutions $\omega_1, \dots, \omega_p$ of greatest modulus, and such that $g'(\omega_i) \neq 0$. Thus [2], inverse functions $w = f_i(z, \zeta_1, \delta_1)$ exist, each holomorphic and univalent for $|z - \zeta_1| < \delta_1$ and satisfying the condition $\omega_i = f_i(\zeta_1, \zeta_1, \delta_1)$, ($i = 1, \dots, p$). By hypothesis, each point z , $|z - \zeta_1| < \delta_1$ is a p -point, hence

$$(4.1) \quad \left| \frac{f_1(z, \zeta_1, \delta_1)}{f_2(z, \zeta_1, \delta_1)} \right| = 1$$

for $|z - \zeta_1| < \delta$. Therefore

$$(4.2) \quad f_1(z, \zeta_1, \delta_1) = e^{i\alpha} f_2(z, \zeta_1, \delta_1)$$

for $|z - \zeta_1| < \delta_1$ and α a constant. Now

$$(4.3) \quad g(f_1(z, \zeta_1, \delta_1)) = g(f_2(z, \zeta_1, \delta_1))$$

for $|z - \zeta_1| < \delta_1$, and therefore

$$(4.4) \quad g(w) = g(e^{i\alpha} w)$$

for w in an open set. Hence the functional relationship (4.4) must hold in Σ_ρ . However, $z = g(w)$ is univalent in a neighborhood of $w = \infty$, which leads to a contradiction.

LEMMA 4.2 (Property 3). Not every point in the neighborhood of a p -point is a 1- p -point.

has a unique solution, ζ' , satisfying

$$(4.8) \quad |\zeta' - \zeta| < \delta.$$

Also

$$(4.9) \quad (f')^{(a_1 \cdots a_k)/\alpha_i} = (\zeta' - \zeta)^{1/\alpha_i}.$$

Hence the functions (4.6) corresponding to i yield the α_i values of ω_j in c_i . Since ζ' is a 1-point, one of the p functions yields an image of largest modulus, and ζ' is said to be associated with this function. As in the first case, the 1-points in $|\zeta' - \zeta| < \delta$ associated with a given function form an open set, and hence all must be associated with the same function. But this, in the same manner, leads to a contradiction of the maximum modulus theorem.

5. Proof of Theorem 1 concluded. We start with the following remarks.

If ζ is a p -point, then there are $k\zeta$ -points of greatest modulus, $\omega_1, \dots, \omega_k$, ($k \leq p$), say of modulus ρ' . Let $F_n(z)$ be the Faber polynomial of degree n associated with $f(z)$, then

$$(5.1) \quad \limsup |F_n(\zeta)|^{1/n} = \rho'.$$

If ζ is a 1-point and ω is the ζ -point of $g(w)$ of greatest modulus, say $|\omega| = \rho'$, and $f(z, \zeta, \delta)$ is the inverse of $g(w)$ in the set $|z - \zeta| < \delta$ which satisfies $\omega = f(\zeta, \zeta, \delta)$, then

$$(5.2) \quad \lim (F_n(\zeta))^{1/n} = f(\zeta, \zeta, \delta),$$

where the root is chosen so that the expansion begins $\zeta + \dots$. The weaker form of this is sometimes used,

$$(5.3) \quad \lim |F_n(\zeta)|^{1/n} = \rho'.$$

Finally, if ζ is an 0-point

$$(5.4) \quad \limsup |F_n(\zeta)|^{1/n} = \rho,$$

where Σ_ρ is the domain of definition of $g(w)$. These limit relationships are obtained by applying the Cauchy-Hadamard formula for the radius of convergence of a power series to the generating function of the Faber polynomial. Consider (1.9)

$$(5.5) \quad \frac{g'(w)}{g(w) - \zeta} = \sum_0^\infty \frac{F_n(\zeta)}{w^{n+1}}$$

for the case ζ is a p -point, $p > 1$. The left side has poles of total multiplicity p on $\sigma_{\rho'}$, and no other singularities in $\Sigma_{\rho'}$. The formula for the radius of convergence yields (5.1). If ζ is a 1-point, (5.3) is obtained since the left side of (5.5) has a simple pole on $\sigma_{\rho'}$ and no singularities in $\Sigma_{\rho'}$. In this case, it is easily shown that (5.2) also holds. Finally if ζ is an 0-point, (5.3) is obtained,

since $g(w)$ has a singularity on σ_ρ , whereas in Σ_ρ the left side of (5.5) has no poles.

The notation $\sigma(z_1, r)$ denotes the closed circle of radius r and center z_1 . Let $\sigma(z_1, r)$ be in the complement of Δ , in which case it will be bounded from Δ . The functions $(F_n(z))^{1/n}$, for sufficiently large n , have all their branches regular in $\sigma(z, r)$ for otherwise $\sigma(z_1, r)$ would contain a point of Δ . For each n , choose one branch, then it will be shown that the sequence of functions forms a normal family in $\sigma(z_1, r)$. In the proof of Part (a) of Theorem 1, it was shown that Δ lies in the complement of C_1 , and that C_1 contains the domain $|z| \geq P, P > 0$. Hence for large $n, |z_{ni}| \leq P + \delta, (\delta > 0)$. But $\max |F_n(z)|^{1/n} \leq \max |z - z_{ni}| \leq \max |z| + |z_{ni}|$. For z in $\sigma(z_1, r)$ this is less than $|z_1| + r + P + \delta$. Hence $\{(F_n(z))^{1/n}\}$ forms a normal family in $\sigma(z_1, r)$, and a sequence can be extracted which converges uniformly to a holomorphic function in any interior subset of $\sigma(z_1, r)$.

To complete the proof of Part (b), it is sufficient to consider the following cases. Let b be a boundary point of C_1 .

CASE 1. The point b is a limit point of interior points of C_0 .

CASE 2. Case 1 does not hold, and b is an 0-point.

CASE 3. Case 1 does not hold, and b is a p -point, $p \neq 0$.

Proof of Case 1. Under the hypothesis of this case, it will be shown that b belongs to Δ . Suppose b is not in Δ . Since Δ is a closed set, there is an r such that $\sigma(b, r)$ does not intersect Δ . By hypothesis, $\sigma(b, r)$ contains an interior point of C_0 , say z_0 . Hence $\sigma(b, r)$ contains $\sigma(z_0, \gamma)$ where $\gamma, \gamma > 0$, is chosen so that $\sigma(z_0, \gamma)$ consists of 0-points. Since by (5.3), $\limsup |F_n(z_0)|^{1/n} = \rho$, there is a subsequence $\{n_\nu\}$ such that $|F_{n_\nu}(z_0)|^{1/n_\nu} = \rho$. Let $\{n_\mu\}$ be the indices of a subsequence of $(F_{n_\nu}(z))^{1/n_\nu}$ which converges to $F(z)$ in a subset of $\sigma(b, \delta)$ containing $\sigma(z_0, \gamma)$ and a 1-point ζ . Now $|F(z_0)| = \rho$, and $|F(z)| \leq \rho$ in $\sigma(z_0, \gamma)$ since points in this circle are 0-points. Hence $|F(z)| = \rho$ in $\sigma(z_0, \gamma)$ and therefore $F(z) = e^{i\theta_0}\rho$ where θ_0 is a real constant. Therefore $F(\zeta) = e^{i\theta_0}$, but since ζ is a 1-point, $\lim |F_{n_\mu}(\zeta)|^{1/n_\mu} = \rho_1$, where $\rho_1 > \rho$, (5.3). This contradiction shows that b belongs to Δ .

CASE 2. Let b be a boundary point of C_1 . From the hypothesis of this case, it follows that there is an r such that $\sigma(b, r)$ contains no point of Δ and no interior points of C_0 . Hence every point of $\sigma(b, r)$ is a limit of k -points ($k = 1, 2, \dots$). By property 1 and 2 of p -points, a k -point is a limit of j -points, $j < k$, hence a limit of 1-points. In other words, 1-points are dense in $\sigma(b, r)$. By hypothesis b is an 0-point, thus $\{n_\nu\}$ can be chosen so that $\lim_{\nu \rightarrow \infty} |F_{n_\nu}(b)|^{1/n_\nu} = \rho$. A further subsequence $\{n_\mu\}$ of $\{n_\nu\}$ exists such that in the subset $\sigma(b, \delta/2)$, $\lim_{\mu \rightarrow \infty} (F_{n_\mu}(z))^{1/n_\mu} = F(z)$, a holomorphic function. Since 1-points are dense in $\sigma(b, \delta/2)$, $|F(z)| \geq \rho$ at a dense set of points, strict inequality being attained at any 1-point and equality being attained at $b = 0$. Since $F(z)$ is continuous, $|F(z)| \geq \rho$ at each point in $\sigma(b, \delta/2)$, equality

being attained for $z=b$. Therefore $F(z)$ is a constant, which involves a contradiction.

CASE 3. In this case b is a boundary point of C_1 , a p -point, ($p \geq 2$), and by hypothesis not a point of Δ . A number r , $r > 0$, can be chosen so that $\sigma(b, r)$ lies in the complement of Δ . By property 3 of p points, not all points in $\sigma(b, \delta)$ are 1-points, hence b is a limit of 2-points. An argument is given in the proof of Lemma 4.3 showing that only finitely many are multiple points hence b is a limit of simple two points. Let ζ be one in $\sigma(b, \delta)$. There are two inverse functions, $f_1(z, \zeta, \gamma)$, $f_2(z, \zeta, \gamma)$, holomorphic for $|z-\zeta| \leq \gamma$, where $\sigma(\zeta, \gamma)$ is contained in $\sigma(b, \delta)$. These satisfy the conditions $\omega_1 = f_1(\zeta, \zeta, \gamma)$, $\omega_2 = f_2(\zeta, \zeta, \gamma)$, where ω_1, ω_2 are the ζ -points of greatest modulus. Since

$$\left| \frac{f_1(z, \zeta, \gamma)}{f_2(z, \zeta, \gamma)} \right| = 1$$

for $z=\zeta$, there are two sequences of points $\{z'_i\}$ and $\{z''_i\}$, both having limit ζ , such that

$$(5.6) \quad \left| \frac{f_1(z'_i, \zeta, \gamma)}{f_2(z'_i, \zeta, \gamma)} \right| > 1, \quad \left| \frac{f_1(z''_i, \zeta, \gamma)}{f_2(z''_i, \zeta, \gamma)} \right| < 1.$$

Now $(F_n(z))^{1/n}$ forms a normal family in $\sigma(b, r)$, hence a subsequence $\{n_\nu\}$ exists such that $\lim (F_{n_\nu}(z))^{1/n_\nu} = F(z)$, for z in $\sigma(b, \gamma/2)$, where $F(z)$ is a holomorphic function. Now, by (5.2), and (5.4),

$$(5.7) \quad \begin{aligned} F(z'_i) &= f_1(z'_i, \zeta, \gamma), \\ F(z''_i) &= f_2(z''_i, \zeta, \gamma). \end{aligned}$$

Hence $\lim_{z \rightarrow b} F(z) = \omega_1$ for one sequence and $\lim_{z \rightarrow b} F(z) = \omega_2$ for the other sequence. This contradiction shows that b belongs to Δ in Case 3, and this concludes the proof of Theorem 1.

BIBLIOGRAPHY

1. G. Faber, *Über Polynomische Entwicklungen*, Math. Ann. vol. 57 (1903) pp. 389-408.
2. A. Hurwitz and R. Courant, *Funktionentheorie*, Interscience Publishers, Inc., 1929.
3. R. Jentzsch, *Untersuchungen zur Theorie der Folgen analytischer Funktionen*, Acta Math. vol. 41 (1917) pp. 219-251.
4. G. Szegö, *Über die nullstellen von Polynomen, die in einem Kreise gleichmässig konvergieren*, Sitzungsberichte der Berliner Mathematische Gesellschaft vol. 21 (1922) pp. 59-64.

UNIVERSITY OF MICHIGAN,
ANN ARBOR, MICHIGAN