

HOMOGENEOUS MANIFOLDS OF ZERO CURVATURE

BY

JOSEPH A. WOLF⁽¹⁾

1. **Introduction.** It is well known⁽²⁾ that a connected Riemannian homogeneous manifold of constant curvature zero is isometric to a quotient R^n/D where R^n is Euclidean space and D is a discrete subgroup of the underlying vector group of R^n . We will extend that theorem to Lorentz manifolds (this is our main result), to compact manifolds with indefinite metric of arbitrary signature (in contrast to the affine case), and to indefinite-metric manifolds of dimension <5 . We will then give examples to show that the assumption "compact, or Riemannian, or Lorentz" is essential in dimensions 5 or more.

2. **Preliminaries.** In order to establish terminology and notation, we will recall some definitions. A *pseudo-Riemannian manifold* M_s^n is an n -dimensional differentiable manifold M with a differentiable family of symmetric bilinear forms Q_p on the tangentspaces M_p of M , where each Q_p is equivalent to the form $-\sum_1^s x_j y_j + \sum_{s+1}^n x_j y_j$. M_s^n has a unique affine connection—the Levi-Civita connection on its tangentbundle—with zero torsion and such that parallel translation is a linear isometry (preserves the Q_p) of tangentspaces. M_s^n is *complete* if the Levi-Civita connection is complete. M_s^n is *flat*, or has *constant curvature zero*, if the curvature tensor of the Levi-Civita connection is zero. A diffeomorphism of pseudo-Riemannian manifolds is an *isometry* if it induces linear isometries on the tangentspaces. The group of all isometries of M_s^n (onto itself) is denoted $\mathfrak{I}(M_s^n)$; M_s^n is *homogeneous* if $\mathfrak{I}(M_s^n)$ is transitive on the points of M_s^n . M_s^n is complete if it is homogeneous. A *pseudo-Riemannian covering* is a covering $p: N_s^n \rightarrow M_s^n$ of connected pseudo-Riemannian manifolds where p is a local isometry; then every deck transformation (homeomorphism d of N_s^n such that $p = p \cdot d$) is an isometry of N_s^n , and, if N_s^n is simply connected and D is the group of deck transformations of the covering, M_s^n is homogeneous if and only if the centralizer of D in $\mathfrak{I}(N_s^n)$ is transitive on N_s^n [7, Theorem 2.5]. The *holonomy group* of M_s^n at x is the group of linear transformations of the tangentspace $(M_s^n)_x$ obtained by parallel translation of tangentvectors about sectionally smooth closed curves based at x . We will make the convention that a Riemannian manifold is just a pseudo-Riemannian manifold $M^n = M_0^n$ or $M^n = M_n^n$, and a Lorentz manifold is either an M_1^n or an M_{n-1}^n .

Received by the editors October 27, 1961.

(1) The author thanks the National Science Foundation for fellowship support during the preparation of this paper.

(2) This is an easy consequence of the work of L. Bieberbach [5]. A somewhat different three-line proof is given in [6]. In addition, we will see that this follows directly from our Lemma 1.

The space V of real n -tuples, identified with its tangentspace at every point and endowed with the bilinear form $Q(x, y) = -\sum_1^s x_j y_j + \sum_{s+1}^n x_j y_j$, carries the structure of a pseudo-Riemannian manifold R_s^n . R_s^n is flat because its Levi-Civita connection is the usual affine connection on V , and inherits from V the structure of a real vectorspace. Subsets X and Y of R_s^n are called *orthogonal* (denoted $X \perp Y$) if $Q(X, Y) = 0$, and X^\perp denotes the largest subspace W of R_s^n such that $X \perp W$. An element $x \in R_s^n$ is *isotropic* if $Q(x, x) = 0$, and a subspace U of R_s^n is *totally isotropic* if $Q(U, U) = 0$. When speaking of R_s^n , we will often refer to Q , and will usually write $Q(x)$ for $Q(x, x)$. $\mathfrak{I}(R_s^n)$ is the group of all transformations $(A, a) : x \rightarrow Ax + a$ where $a \in R_s^n$ and A is a linear transformation of R_s^n which preserves Q ; thus R_s^n is homogeneous. (A, a) is a *translation* if $A = I$. The usefulness of R_s^n is due to the fact [7, Theorem 5] that M_s^n is connected, flat and complete if and only if it admits a pseudo-Riemannian covering by R_s^n .

A basis $\{v_i\}$ of $V = R_s^n$ is called *Q-orthonormal* if $Q(v_i, v_j) = -\delta_{ij}$ for $i \leq s$ and $Q(v_i, v_j) = \delta_{ij}$ for $i > s$, δ_{ij} being the Kronecker symbol. Now suppose that $s \leq n - s$ and that U is a totally isotropic subspace of R_s^n , say $\dim U = l$. One can then find a *Q-orthonormal* basis $\{v_i\}$ of R_s^n such that, defining $f_i = v_i - v_{n-l+i}$ and $e_i = v_i + v_{n-l+i}$ ($1 \leq i \leq l$), $\{e_i\}$ is a basis of U and $\{v_{l+1}, \dots, v_{n-l}\} \cup \{e_i\}$ is a basis of U^\perp . The basis

$$\{f_1, \dots, f_l; v_{l+1}, \dots, v_{n-l}; e_1, \dots, e_l\}$$

of R_s^n is called a *skew basis with respect to U*. One has the obvious definition of skew bases for the case $s \geq n - s$.

Let $p : R_s^n \rightarrow M_s^n$ be a pseudo-Riemannian covering, let D be the group of deck transformations, and let H be the holonomy group of M_s^n at $p(0)$. p gives an identification of R_s^n with the tangentspace to M_s^n at $p(0)$, so H may be viewed as acting on R_s^n . This gives us the *standard homomorphism* $\phi : D \rightarrow H$ by $d = (\phi_d, t_d) \in \mathfrak{I}(R_s^n)$; ϕ is onto, and the kernel $\text{Ker} \phi$ is the set of translations in D .

If A is a linear transformation, then $\text{Ker} A$ denotes the kernel and $\text{Im} A$ the image. If a and b are elements of a group, then $[a, b]$ denotes the commutator $aba^{-1}b^{-1}$.

3. Computational preparations. Let D be a subgroup of $\mathfrak{I}(R_s^n)$, let Z be the centralizer of D in $\mathfrak{I}(R_s^n)$, and suppose that Z acts transitively on R_s^n . We adopt the convention that d, d', d_i always represent arbitrary elements of D , $d = (R, t) = (I + N, t)$, $d' = (R', t') = (I + N', t')$, $d_i = (R_i, t_i) = (I + N_i, t_i)$ where I is the identity transformation of R_s^n .

In the proof of Theorem 14.1 of [7], it was seen that transitivity of Z implies:

LEMMA 1. *If $d \in D$, then $N^2 = 0$, $t \perp \text{Im} N$, and $\text{Im} N$ is totally isotropic.*

In the Riemannian case ($s = 0$ or $s = n$), there is no nonzero totally isotropic

subspace of R_s^n , whence the theorem on flat connected Riemannian homogeneous manifolds mentioned in the Introduction.

LEMMA 2. *If $d \in D$ and $x, y \in R_s^n$, then $Q(Nx, y) + Q(x, Ny) = 0$, $\text{Ker}.N = \text{Im}.N^\perp$, $\text{Im}.N = \text{Ker}.N^\perp$, and $N(t) = 0$.*

Proof. Lemma 1 gives us $0 = Q(Rx, Ry) - Q(x, y) = Q(Nx, y) + Q(x, Ny) + Q(Nx, Ny) = Q(Nx, y) + Q(x, Ny)$, proving the first statement. Thus $Q(Nx, y) = 0$ if $Ny = 0$, so $\text{Ker}.N \perp \text{Im}.N$, and the second and third statements follow from the fact that $\dim.\text{Ker}.N + \dim.\text{Im}.N = \dim.\text{Ker}.N + \dim.\text{Ker}.N^\perp = \dim.\text{Im}.N + \dim.\text{Im}.N^\perp = n$. Finally, $t \in \text{Ker}.N$ because $t \perp \text{Im}.N$ by Lemma 1. q.e.d.

An immediate consequence of Lemmas 1 and 2 is:

LEMMA 3. *If $d \in D$, then $d^m = (I + mN, mt)$ for every integer m .*

A useful tool for examining commutativity in D is:

LEMMA 4. *If $d, d', d_i \in D$, then $NN' + N'N = 0 = N_1N_2N_3$.*

Proof. Let $d'' = dd'$; then $N'' = N + N' + NN'$, and Lemma 1 gives

$$(*) \quad 0 = N''^2 = NN' + N'N + NN'N + N'NN' + NN'NN'.$$

Left multiplication of (*) by N and nonsingularity of $R' = I + N'$ gives $0 = NN'N$; right multiplication of (*) by N' and nonsingularity of $R = I + N$ gives $0 = N'NN'$. Thus (*) reduces to the first equality of the Lemma. It follows that $0 = N_3(N_1 + N_2 + N_1N_2) + (N_1 + N_2 + N_1N_2)N_3 = N_3N_1 + N_1N_3 = N_3N_2 + N_2N_3$, and the second equality follows. q.e.d.

Observe that Lemmas 2 and 4 give us $N'Nt' = -NN't' = 0$. Thus $d'' = dd'$ and Lemma 2 imply $(N + N' + NN')(t + t' + Nt') = 0 = Nt' + N't$. It follows that $dd'd^{-1} = (I + N, t)(I + N', t')(I - N, -t) = (I + N' + 2NN', t' + 2Nt')$ and $[d, d'] = (I + 2NN', 2Nt')$. This last implies that every translation in D is central, and that the third term in the lower central series of D consists of translations. In summary, we have just proved:

LEMMA 5. *If $d, d' \in D$, then $N'Nt' = 0 = NN't$, $Nt' + N't = 0$, $dd'd^{-1} = (I + N' + 2NN', t' + 2Nt')$, $[d, d'] = (I + 2NN', 2Nt')$. The set $T = \text{Ker}.\phi$ of all translations in D is central in D , and D is nilpotent of order 3.*

Here, of course, by nilpotent of order k we mean only that the $(k+1)$ st term of the lower central series is trivial, and do not exclude triviality of the k th term. Observe also that T is the full center of D if the translation parts of elements of D span R_s^n .

4. **Commutativity.** We retain the notation of §3, let $\phi: \mathfrak{F}(R_s^n) \rightarrow \mathcal{O}^*(n)$ be the canonical homomorphism, let $H = \phi(D)$, and define U_D to be the subspace of R_s^n spanned by $\{\text{Im}.\phi(d) - I : d \in D\}$.

PROPOSITION 1. R_s^n has a maximal totally isotropic subspace V_D which contains U_D and on which H acts trivially.

Proof. The Proposition is true for $n=1$, for we are then dealing with a Riemannian signature. Now assume $n>1$ and suppose the Proposition true in dimensions $<n$. We assume $D \neq I$. Then D has a central element $\neq I$ by Lemma 5, and transitivity of Z shows that the translation part of this element is nonzero. Let W be the subspace of R_s^n spanned by all translation parts of central elements of D ; then $W \neq 0$ and Lemma 5 shows that H acts trivially on W . We may assume $W^\perp \neq 0$, for, if not, then $H=I$ and the Proposition is trivial.

Suppose that $W \cap W^\perp = 0$. Then R_s^n is an orthogonal direct sum $W \oplus W^\perp$, this decomposition is preserved by $\phi(D)$ and $\phi(Z)$, and Q induces nondegenerate bilinear forms on W and W^\perp . Thus the Proposition follows by induction on n .

Now suppose that $X = W \cap W^\perp$ is nonzero. We choose a skew basis

$$\beta = \{f_1, \dots, f_l; v_{l+1}, \dots, v_{n-l}; e_1, \dots, e_l\}$$

of R_s^n with respect to X , $l = \dim X$. As H preserves and acts trivially on W , and thus on X , every element of H is of the form

$$h = \begin{pmatrix} h'_1 & h'_2 & h'_3 \\ 0 & h'_4 & h'_5 \\ 0 & 0 & I \end{pmatrix}$$

in block form relative to β . The process of restricting to X^\perp and passing to X^\perp/X shows, by induction on n , that the linear span of $\{v_{l+1}, \dots, v_{n-l}\}$ has a maximal totally isotropic subspace Y on which each h'_4 acts trivially and which contains the image of each $h'_4 - I$. We define $V_D = Y \oplus X$. $Y \subset X^\perp$ shows that V_D is totally isotropic, and then it is clear from dimensions that V_D is a maximal totally isotropic subspace of R_s^n which is H -invariant. Let β' be a skew basis of R_s^n with respect to V_D . Every $h \in H$ has form

$$h = \begin{pmatrix} h_1 & h_2 & h_3 \\ 0 & h_4 & h_5 \\ 0 & 0 & h_6 \end{pmatrix}$$

relative to $\beta' = \{f'_1, \dots, f'_m; v'_{m+1}, \dots, v'_{n-m}; e'_1, \dots, e'_m\}$. $h_2 = 0$ because (Lemma 1) $\text{Im}(h - I)$ is totally isotropic and the linear span S of $\{v'_{m+1}, \dots, v'_{n-m}\}$ is a positive or negative definite subspace; it follows that $h_5 = 0$ because $h \in \mathcal{O}^s(n)$. Similarly, $h_4 = I$. We wish to prove $h_1 = I$; then $h \in \mathcal{O}^s(n)$ will imply $h_6 = I$, we will have

$$h = \begin{pmatrix} I & 0 & h_3 \\ 0 & I & 0 \\ 0 & 0 & I \end{pmatrix}$$

relative to β' , and the Proposition will be proved. To prove $h_1 = I$, it suffices to prove $\text{Im.}(h - I) \subset V_D$. As $\text{Im.}(h - I)$ is totally isotropic and V_D is maximal totally isotropic, it suffices to show $\text{Im.}(h - I) \perp V_D$. It is clear that $\text{Im.}(h - I) \perp Y$, so we need only show $\text{Im.}(h - I) \perp X$. Looking at h in the basis β , this is clear because $h'_1 = I$. q.e.d.

PROPOSITION 2. *Let β be a skew basis of R_s^n with respect to U_D . Then every $h \in H$ is of the form*

$$h = \begin{pmatrix} I & 0 & \alpha_h \\ 0 & I & 0 \\ 0 & 0 & I \end{pmatrix}$$

relative to β , where α_h is skew-symmetric; H has an element h_0 such that α_{h_0} is nonsingular; the translation part of every element of D lies in U_D^\perp , and D is represented faithfully as a group of translations of U_D^\perp . In particular, D and H are torsion-free abelian groups.

Proof. The form of the elements $h \in H$ relative to β is clear from Proposition 1. We will write α_i for α_{h_i} . If $h_1 = h_2 h_3$, then $\alpha_1 = \alpha_2 + \alpha_3$; the existence of h_0 is now clear. Let $d_0 = (h_0, t_0) \in D$. Then $t_0 \in \text{Ker.}(h_0 - I) = U_D^\perp$ by Lemmas 1 and 2 and nonsingularity of α_0 . Now let $d_1 = (h_1, t_1)$ be any element of D , and observe that one can find integers u and v such that $u\alpha_0 + v\alpha_1$ is nonsingular. Let $d_2 = d_1^u d_0^v = (h_2, t_2)$; then $\alpha_2 = u\alpha_0 + v\alpha_1$ is nonsingular and $t_2 = vt_1 + ut_0 \in U_D^\perp$. Thus $t_1 \in U_D^\perp$. The Proposition follows. q.e.d.

5. **The main results.** Our main result is the Lorentz case of the following Theorem 1. As mentioned in the Introduction, the Riemannian case is known from the work of L. Bieberbach [5]. The compact case is interesting in relation of the work of L. Auslander on compact locally affine spaces ([1; 2; 3 and 4], for example), and shows that the metric plays a strong role.

THEOREM 1. *Let M_s^n be a connected flat homogeneous pseudo-Riemannian manifold, and suppose that*

1. M_s^n is compact; or
2. M_s^n is Riemannian, i.e., $s = 0$ or $s = n$; or
3. M_s^n is Lorentzian, i.e., $s = 1$ or $s = n - 1$; or
4. the dimension $n = \dim.M_s^n \leq 4$.

Then M_s^n is isometric to a quotient R_s^n/D where D is a discrete group of translations of R_s^n . This result is best possible in the sense that, if $n > 4$ and if $s \neq 0, 1, n - 1$ or n , then there is a connected noncompact flat homogeneous N_s^n with non-trivial holonomy group.

Proof. Let D be the group of deck transformations of the universal pseudo-Riemannian covering $\pi: R_s^n \rightarrow M_s^n$, let Z be the centralizer of D in $\mathfrak{S}(R_s^n)$, and let H be the holonomy group of M_s^n at $\pi(0)$. Z is transitive on R_s^n by homogeneity of M_s^n ; this allows us to use the results of §4.

We adopt the terminology of §4. If M_s^n is compact, then the translation parts of the elements of D span R_s^n , whence $U_D^1 = R_s^n$ by Proposition 2; thus $U_D = 0$. If M_s^n is Riemannian or Lorentzian, then every totally isotropic subspace of R_s^n has dimension < 2 ; thus $m = \dim U_D < 2$. If $m \neq 0$, then Proposition 2 gives us a skew nonsingular $m \times m$ matrix α_h ; it follows that $m = 0$; thus $U_D = 0$. If $n \leq 4$ and M_s^n is neither Riemannian nor Lorentzian, then $s = 2$ and $n = 4$. Then, if $U_D \neq 0$, Proposition 2 would give us $d = (h, t) \in D$ with $0 \neq t \in U_D = \text{Im.}(h - I)$. This gives $v \in R_s^n$ with $t = (h - I)v$, whence $d(-v) = h(-v) + (h - I)v = -v$, contradicting the fact that d has no fixed point.

Now $U_D = 0$ in all four cases, whence $H = I$ and D consists of translations. D is discrete because $M_s^n = R_s^n/D$ is a manifold.

The manifolds N_s^n will be constructed in §6, completing the proof of Theorem 1, in such a way as to show the bounds of Theorem 2 to be best possible.

THEOREM 2. *Let M_s^n be a connected flat homogeneous pseudo-Riemannian manifold, let D be the group of deck transformations of the universal pseudo-Riemannian covering $\pi: R_s^n \rightarrow M_s^n$, let H be the holonomy group of M_s^n , and let $\phi: D \rightarrow H$ be the standard homomorphism. Then D is free abelian on some number $m \leq n$ of generators, D is represented faithfully as a discrete group of translations of a linear subspace of R_s^n , D has a subgroup D' such that $D = D' \times \text{Ker.}\phi$, and $m \leq n - 2$ in case $H \neq I$. M_s^n has the homotopy type of an m -torus; its Euler-Poincaré characteristic is zero, and its integral cohomology is an exterior algebra on m generators⁽³⁾.*

Proof. The first part follows easily from Proposition 2, in the same manner as the proof of the first part of Theorem 1. To prove the second part, we observe that both M_s^n and an m -torus are Eilenberg-MacLane spaces $K(D, 1)$.

6. An example. In order to complete the proof of Theorem 1 and show the bound of Theorem 2 to be best possible, we take integers n and s with $2 \leq s \leq n - 2$ and $n > 4$, and will construct connected flat homogeneous pseudo-Riemannian manifolds $N_s^n = R_s^n/D$ with nontrivial holonomy group and with D free abelian on $m = n - 2$ generators.

Let $\{v_i\}$ be an "orthonormal" basis of R_s^n , and let U be a two-dimensional totally isotropic subspace of R_s^n such that we have a skew basis $\beta = \{f_1, f_2, v_3, \dots, v_{n-2}, e_1, e_2\}$ of R_s^n with $f_i = v_i + v_{n-2+i}$, $e_i = v_i - v_{n-2+i}$, and $\{e_1, e_2\}$ is a basis of U . Let

(³) Compare with [2; 3; 4], and the proof of [1, Lemma 1].

$$h = \begin{pmatrix} I & 0 & J \\ 0 & I & 0 \\ 0 & 0 & I \end{pmatrix}$$

relative to β , where

$$J = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix},$$

and observe that h is in the orthogonal group of R_s^n . We now define elements d_1, d_2, \dots, d_{n-2} of $\mathfrak{S}(R_s^n)$ by $d_1(x) = h(x) + v_3$, $d_2(x) = x + v_4, \dots, d_{n-4}(x) = x + v_{n-2}$, $d_{n-3}(x) = x + e_1$, and $d_{n-2}(x) = x + e_2$. Let D be the subgroup of $\mathfrak{S}(R_s^n)$ generated by the d_i , and let Z be the centralizer of D in $\mathfrak{S}(R_s^n)$. It is clear that D is free abelian on $n-2$ generators, that $N_s^n = R_s^n/D$ is a manifold, and that N_s^n has nontrivial holonomy group generated by h . Thus we need only prove that N_s^n is homogeneous, i.e., that Z is transitive on the points of R_p^n .

To prove transitivity of Z , let v be an arbitrary element of R_s^n ; we must find $(z, v) \in \mathfrak{S}(R_s^n)$ which commutes with each d_i . Write $v = a_1 f_1 + a_2 f_2^\perp + w$ where $w \in U^\perp$. To construct z , we define a linear transformation z' of U^\perp by $v_3 \rightarrow v_3 - a_2 e_1 + a_1 e_2$, $v_i \rightarrow v_i$ for $3 < i \leq n-2$, and $e_j \rightarrow e_j$. This preserves the bilinear form induced on U^\perp by Q , and thus extends to an element of the orthogonal group $O^*(n)$ of R_s^n , say z , by Witt's Theorem. As $z(v_i) = v_i$ for $3 < i \leq n-2$ and $z(e_j) = e_j$, it follows that (z, v) commutes with d_k for $k > 1$. To see that (z, v) commutes with d_1 , we must show that $zh = hz$ and $(z - I)v_3 = (h - I)v$. The second condition is satisfied by construction of z' . Now

$$z = \begin{pmatrix} z_1 & z_2 & z_3 \\ 0 & z_4 & z_5 \\ 0 & 0 & z_6 \end{pmatrix}$$

in block form relative to β . By construction of z' , $z_4 = I$ and $z_6 = I$. Then $z \in O^*(n)$ gives $z_1 = I$. Thus

$$zh = \begin{pmatrix} I & z_2 & J + z_3 \\ 0 & I & z_5 \\ 0 & 0 & I \end{pmatrix} = hz$$

in block form relative to β , proving transitivity of Z on R_s^n .

REFERENCES

1. L. Auslander, *On the group of affinities of locally affine spaces*, Proc. Amer. Math. Soc. 9 (1958), 471-473.

2. ———, *On the Euler characteristic of compact locally affine spaces*, Comment. Math. Helv. **35** (1961), 25–29.
3. ———, *On the Euler characteristic of compact locally affine spaces. II*, Bull. Amer. Math. Soc. **67** (1961), 405–406.
4. ———, *On radicals of discrete subgroups of Lie groups, with applications to locally affine spaces*, to appear.
5. L. Bieberbach, *Über die Bewegungsgruppen der Euklidischen Räume*, Math. Ann. **70** (1911), 297–336; II, Math. Ann. **72** (1912), 400–412.
6. J. A. Wolf, *Sur la classification des variétés riemanniennes homogènes à courbure constante*, C. R. Acad. Sci. Paris **250** (1960), 3443–3445.
7. ———, *Homogeneous manifolds of constant curvature*, Comment. Math. Helv. **36** (1961), 112–147.

THE INSTITUTE FOR ADVANCED STUDY,
PRINCETON, NEW JERSEY