

ON SPHERICAL CHARACTERISTIC COHOMOLOGY⁽¹⁾

BY
A. W. ADLER

The theory of characteristic classes, primarily a topological discipline, can be (and, at its inception, was) treated geometrically by means of mappings into classifying spaces and invariant polynomials in the curvature forms of these spaces. The theory goes something like this. Suppose that n and p are positive integers, that G is a closed subgroup of the orthogonal group $O(n)$, and that S is an invariant symmetric q -linear functional on the Lie algebra of G . Let Ω denote the curvature form of the *canonical connection of the second kind* on $O(n+p)$ as a bundle over the *classifying space* $B_G = O(n+p)/G \times O(p)$. The $2q$ -form

$$\text{Alternation } S(\Omega, \dots, \Omega)$$

on $O(n+p)$ then induces a closed $2q$ -form Ω^S on B_G ; the $2q$ th real cohomology group of B_G is, for any $2q < p$, generated by such forms Ω^S . If M is an n -dimensional manifold and \bar{B} a principal G -bundle over M , then \bar{B} can be induced by a mapping g of M into B_G (with p sufficiently large); the *characteristic $2q$ -forms* of M with respect to the bundle \bar{B} are then the forms $g^*(\Omega^S)$. The resulting *characteristic cohomology groups* of M turn out to be independent of the choice of p and g .

An analogous *spherical characteristic cohomology* of M can be defined by means of a more carefully chosen bundle B_G^+ and mapping g . If p is large, it can be shown as a consequence of Nash's imbedding theorem that there exists an isometric imbedding of M in the unit sphere S^{n+p-1} . The *spherical image* of this imbedding (defined in §2) then gives rise to a mapping g of M into the space $B_G^+ = O(n+p)/G \times O(p-1)$. The forms $g^*(\Omega^S)$ are nothing new: they generate the characteristic cohomology groups of M . But now one can choose instead to consider the curvature form Ω' of the canonical connection on $O(n+p)$ as a bundle over S^{n+p-1} . The $2q$ -forms $\text{Alternation } S(\Omega', \dots, \Omega')$ on $O(n+p)$ then induce $2q$ -forms Ω'_S on B_G^+ , and hence induce $2q$ -forms $\Omega_S = g^*(\Omega'_S)$ on M . The question, of course, is whether the forms Ω_S are closed, whether they are independent of p and g , and whether they generate some noncharacteristic cohomology in M . This paper is a preliminary attempt to supply some answers. For obvious reasons we will call the forms Ω_S the *spherical characteristic forms* of M with respect to the bundle \bar{B} .

Received by the editors June 14, 1962.

(1) Supported by a National Science Foundation Fellowship.

First, the conditions under which the forms Ω_S are closed. It is easy to show that the characteristic forms of M are closed, because the forms Ω^S are closed in B_G ; but such is not the case for the spherical characteristic forms, because the forms Ω'_S are in general not closed in B_G^+ . It can be proved using the methods of §2 that Ω_S is closed whenever the restriction of the form Alternation $S(\Omega', \dots, \Omega')$ to $g(M)$ is invariant under the right-action of $O(n)$, but such a condition is quite unsatisfactory. A better condition uses Chern's notion of a G -manifold: An n -dimensional Riemannian manifold M is a G -manifold if its bundle of frames can be reduced to a principal G -bundle B which admits a torsionless connection. It is proved in §2 that the spherical characteristic forms of M with respect to such a bundle B are closed.

The proof that the spherical characteristic classes are independent of p and of g is, on the other hand, much simpler than the corresponding proof for characteristic classes (§3). For characteristic classes, independence is established by a theorem of Weil. For spherical characteristic classes, independence is essentially a consequence of the fact that the forms Ω'_S are horizontal over S^{n+p-1} (this, of course, is due to the fact that the form Ω' is horizontal over S^{n+p-1} ; the forms Ω and Ω^S are horizontal only over B_G). Part C of Theorem 2 is the heart of the proof.

In the special case n even and G the unitary group $U(n/2)$, the G -manifolds M are the *almost-Kähler manifolds*. There exists a nonvanishing invariant 1-linear functional S_0 on the Lie algebra of $U(n/2)$: The elements of the Lie algebra are certain $n \times n$ matrices $A = (a_{ij})$, and

$$S_0(A) = \sum_{k=1}^n a_{k, k+n/2} .$$

The resulting characteristic 2-form $(2\pi)^{-1} \cdot g^*(\Omega^{S_0})$ on M is then the *Chern form* c_1 of the almost-Kähler metric of M . c_1 and the *fundamental form* of the metric will in general differ; c_1 is an integral form, while the fundamental form is integral only if the metric is Hodge. The fundamental form in fact will generally not be found among the characteristic forms of M . It is proved in [1], however, that the fundamental form is a spherical characteristic form of M : It is the form $(2\pi)^{-1} \cdot g^*(\Omega'_{S_0})$. This example suggests that it is exactly the differences $g^*(\Omega^S) - g^*(\Omega'_S)$ between the characteristic forms and the spherical characteristic forms which give the most fundamental information about the geometry of G -manifolds.

The forms Alternation $S(\Omega, \dots, \Omega)$ on $O(n + p)$ are exact (although the induced forms Ω^S on B_G are not). Analogously, it would be useful to know when the forms

$$\text{Alternation } S(\Omega', \dots, \Omega')$$

are exact over images $g(M)$ of G -manifolds (they cannot be exact on all of $O(n + p)$ since they are not even closed there). The following result has been proved (§4).

The integrals over the group $SO(n)$ of the forms Alternation $S(\Omega', \dots, \Omega')$ are exact forms on $O(n + p)$.

Exactness can, in general, be expected to have quite strong consequences. It is shown in Theorem 4 of [1], for example, that for the case $G = U(n/2)$ the following conclusion can be drawn:

If the spherical characteristic form Alternation $S_0(\Omega')$ is exact over $g(M)$, then M is an algebraic manifold.

The important topological properties of compact Kähler manifolds are consequences of the fact that the fundamental form is never exact: the $n/2$ th power of the fundamental form is the volume element. It is tempting to conjecture that this is a special case of some property shared by all spherical characteristic forms on all compact G -manifolds. For this reason we consider in §6 the set $s(O(n)/G)$ of all invariant symmetric multilinear functionals on the Lie algebra of G which *cannot* be written in the form

$$\sum_i S_i T_i,$$

with S_i and T_i invariant symmetric multilinear functionals of positive degree on the Lie algebra of G such that each S_i is extendable to an invariant symmetric multilinear functional on the Lie algebra of $O(n)$. It is proved that, for G -manifolds, the algebraic homomorphism $S \rightarrow \Omega_S$ induces an algebraic homomorphism θ of $s(O(n)/G)$ onto the spherical characteristic cohomology ring of M . Since the previously-defined linear functional S_0 on the Lie algebra of $U(n/2)$ cannot be extended to Lie algebra of $O(n)$, the generalization of the nonexactness property of fundamental forms should be of the following form:

If M is compact, then θ is an isomorphism.

This proposition remains to be proved. Since $s(O(n)/G)$ is isomorphic to the characteristic cohomology ring of $O(n)/G$, it would follow that the spherical characteristic cohomology ring of a compact G -manifold is isomorphic with the characteristic ring of $O(n)/G$.

If ϕ is a mapping and \bar{X} a vector, then $\phi(\bar{X})$ will always denote the vector often denoted by $d\phi(\bar{X})$. A form on a bundle will be called *horizontal* if it vanishes whenever one of its arguments is a vertical vector. If H is some connection (concept of horizontality) on the bundle, then a form will be called *H-horizontal* (resp. *V-vertical*) if it vanishes whenever one of its arguments is a vertical (resp. *H-horizontal*) vector. (Thus horizontal and *H-horizontal* mean the same thing.)

1. **The forms Ω'_i .** Let G be a closed subgroup of $O(n)$ and let p be a large positive integer. Let $B_G^+ = O(n + p)/G \times O(p - 1)$, and let

$$B_{O(n)}^+ = O(n + p)/O(n) \times O(p - 1).$$

Then we have the following collection of bundles and mappings (σ, τ, ∂ , and α denote the natural projection mappings, and S^{n+p-1} is the unit sphere $O(n+p)/O(n+p-1)$ of Euclidean space $E(n+p)$).

$$\begin{array}{c}
 O(n+p) \\
 \sigma \downarrow \\
 O(n+p)/O(p-1) \\
 \tau \downarrow \\
 B_G^+ \\
 \partial \downarrow \\
 B_{O(n)}^+ \\
 \alpha \downarrow \\
 S^{n+p-1}
 \end{array}$$

Let $o(n+p)$, $o(n)$ and $o(p-1)$ denote, respectively, the Lie algebras of $O(n+p)$, $O(n)$ and $O(p-1)$; let \mathfrak{g} denote the Lie algebra of G . Let ω be the 1-form on $o(n+p)$ which assigns to each element of $o(n+p)$ its $o(n)$ -component (with respect to the Killing form). Let ω^0 be the 1-form which assigns to each element of $o(n+p)$ its \mathfrak{g} -component. Finally, let ω' be the 1-form which assigns to each element of $o(n+p)$ its component in the Lie algebra of $O(n+p-1)$. (The forms ω , ω^0 , and ω' are the *canonical connections of the second kind* on $O(n+p)$ as a bundle over $O(n+p)/O(n)$, $O(n+p)/G$, and $O(n+p)/O(n+p-1)$.)

Let S be a symmetric q -linear functional on the Lie algebra of G . If P denotes the projection of $o(n+p-1)$ onto \mathfrak{g} (with respect to the Killing form), then the functional $P^*(S)$ is an extension of S to the Lie algebra of $O(n+p-1)$. For simplicity we denote it also by S . Let

$$\Omega' = d\omega' + (1/2) \cdot \omega' \wedge \omega'$$

be the curvature form of ω' . Let $\varepsilon(j_1, \dots, j_{2q})$ denote the sign of the permutation taking $(1, \dots, 2q)$ into (j_1, \dots, j_{2q}) . Then a real-valued $2q$ -form Λ_S can be defined on $O(n+p)$ in the following way:

If x_1, \dots, x_{2q} are tangent vectors to $O(n+p)$ and if X_1, \dots, X_{2q} denote their extensions to elements of $o(n+p)$, then

$$\Lambda_S(x_1, \dots, x_{2q}) = ((2q)!)^{-1} \cdot \sum \varepsilon(j_1, \dots, j_{2q}) \cdot S(\Omega'(X_{j_1}, X_{j_2}), \dots, \Omega'(X_{j_{2q-1}}, X_{j_{2q}}))$$

(the summation extends over all permutations of $1, \dots, 2q$).

More concisely, one can write

$$\Lambda_S = \text{Alternation } S(\Omega', \dots, \Omega').$$

Let H' denote the operator which assigns to each vector of $O(n + p)$ its component in the nullspace of ω' . It follows immediately (from the fact that Ω' is H' -horizontal) that Λ_S is an H' -horizontal form on $O(n + p)$. It is of course left-invariant under $O(n + p)$, and invariant under the right-action of $G \times O(p - 1)$. So there exists a $2q$ -form Ω'_S on B_G^+ with $(\tau \circ \sigma)^*(\Omega'_S) = \Lambda_S$. The form Ω'_S is also invariant under left-translation by $O(n + p)$.

2. G -manifolds. Let M be a G -manifold of dimension n . Let F be an isometric imbedding of M in S^{n+p-1} , for p sufficiently large. For each point m of M , let $\pi(F(m))$ denote the n -plane through the origin of $E(n + p)$ parallel to the tangent plane to $F(M)$ at $F(m)$. Then π is a mapping of $F(M)$ into $B_{O(n)}^+$ (called the *spherical image mapping*) and $f = \pi \circ F$ is a mapping of M into $B_{O(n)}^+$. Since M is a G -manifold, there exists a mapping g of M into B_G^+ satisfying the following conditions:

1. $\partial \circ g = f$;
2. g induces (as a characteristic mapping) the principal G -bundle B over M to which the frame bundle of M can be reduced.

Let Ω_S denote the $2q$ -form $g^*(\Omega'_S)$ on M .

THEOREM 1. Ω_S is closed.

Proof. Let $M' = g(M)$. Let $\delta = \tau \circ \sigma$. Since $\delta^*(\Omega'_S) = \Lambda_S$, it suffices to prove that Λ_S is a closed form on the submanifold $\delta^{-1}(M')$ of $O(n + p)$. This is proved in the following lemmas.

LEMMA 1. The restrictions of ω and ω^0 to $\delta^{-1}(M')$ coincide.

Proof. This is proved in Theorem 1' of [1]. What is involved in the proof is this: In order for the Riemannian connection of M to reduce to a connection on the bundle B , the manifold $M' = g(M)$ must be horizontal in B_G^+ as a bundle over $B_{O(n)}^+$.

LEMMA 2. The restrictions of $d\Omega'$ and $\Omega' \wedge \omega$ to $\delta^{-1}(M')$ have identical $o(n)$ -components.

Proof. The Bianchi identity for Ω' can be written in the form

$$d\Omega' = \Omega' \wedge \omega'.$$

So it must be shown that the restrictions of $\Omega' \wedge \omega'$ and $\Omega' \wedge \omega$ to $\delta^{-1}(M')$ have identical $o(n)$ -components. Let \bar{x} , \bar{y} , and \bar{z} be tangent vectors to $\delta^{-1}(M')$; it will be shown in A—C, below, that the components in $o(n)$ of

$$[\Omega'(\bar{x}, \bar{y}), \omega'(\bar{z})] \quad \text{and} \quad [\Omega'(\bar{x}, \bar{y}), \omega(\bar{z})]$$

coincide (and so the lemma is proved).

A. Let \bar{x}_1 and \bar{x}_2 be tangent vectors to $O(n + p)$, and let \bar{X}_1 and \bar{X}_2 denote their extensions to left-invariant vector fields on $O(n + p)$. So

$$\Omega'(\bar{X}_1, \bar{X}_2) = -(1/2) \cdot \omega' [H'(\bar{X}_1), H'(\bar{X}_2)].$$

The sphere S^{n+p-1} is a symmetric homogeneous space; that is, $[H'(\bar{X}_1), H'(\bar{X}_2)]$ is in $o(n+p-1)$ for any left-invariant vector fields \bar{X}_1 and \bar{X}_2 . So

$$\omega'[H'(\bar{X}_1), H'(\bar{X}_2)] = [H'(\bar{X}_1), H'(\bar{X}_2)],$$

and we conclude that

$$(*) \quad \Omega'(\bar{X}_1, \bar{X}_2) = -(1/2) \cdot [H'(\bar{X}_1), H'(\bar{X}_2)].$$

B. The vector fields \bar{X}_1 and \bar{X}_2 are elements of $o(n+p)$ and so can be considered skew-symmetric $(n+p) \times (n+p)$ matrices:

$$\bar{X}_i = \left[\begin{array}{c|c|c} A_i & C_i & U_i \\ \hline -C_i^t & B_i & V_i \\ \hline -U_i & -V_i & O \end{array} \right], \quad i = 1, 2.$$

Here

- (a) A_i is an $n \times n$ matrix with $A_i^t = -A_i$,
- (b) B_i is an $(p-1) \times (p-1)$ matrix with $B_i^t = -B_i$,
- (c) C_i is a $(p-1) \times n$ matrix,
- (d) U_i is a $1 \times n$ matrix, and V_i is a $1 \times (p-1)$ matrix.

Note that

$$H'(\bar{X}_i) = \left[\begin{array}{c|c|c} O & O & U_i \\ \hline O & O & V_i \\ \hline -U_i & -V_i & O \end{array} \right]$$

and, furthermore, that Lemma 1 is the statement that A_i lies in \mathfrak{g} if \bar{x}_i is tangent to $\delta^{-1}(M')$.

The manifold M' satisfies the condition

$$\partial(M') = \pi(F(M));$$

that is, $\partial(M')$ is the spherical image of a submanifold of S^{n+p-1} . Hence if \bar{x}_1 and \bar{x}_2 are tangent to $\delta^{-1}(M')$, then the fields \bar{X}_1 and \bar{X}_2 satisfy the condition

$$(**) \quad V_1 = V_2 = 0.$$

It follows from (*) and (**) that $\Omega'(\bar{X}_1, \bar{X}_2)$ is in $o(n)$ whenever \bar{x}_1 and \bar{x}_2 are tangent to $\delta^{-1}(M')$. Hence we conclude:

$$\Omega' \text{ is } o(n)\text{-valued on } \delta^{-1}(M').$$

C. Suppose U is an element of $o(n)$ and V an element of $o(n+p-1)$. Let $V_{o(n)}$ denote the $o(n)$ -component of V . It follows immediately from matrix-multiplication that

$$o(n)\text{-component of } [U, V] = o(n)\text{-component of } [U, V_{o(n)}].$$

Note that $\omega(\bar{z})$ is the $o(n)$ -component of $\omega'(\bar{z})$. So the substitution $U = \Omega'(\bar{x}, \bar{y})$ and $V = \omega'(\bar{z})$ proves the assertion of the lemma.

LEMMA 3. $d\Lambda_S = 0$ on $\delta^{-1}(M')$.

Proof. Using Lemmas 1 and 2 we find that (letting \bar{S} denote Alternation S)

$$\begin{aligned} d\Lambda_S &= d(\bar{S}(\Omega', \dots, \Omega')) \\ &= \bar{S}(d\Omega', \Omega', \dots, \Omega') + \dots + \bar{S}(\Omega', \dots, \Omega', d\Omega') \\ &= \bar{S}(\Omega' \wedge \omega', \Omega', \dots, \Omega') + \dots + \bar{S}(\Omega', \dots, \Omega', \Omega' \wedge \omega') \\ &= \bar{S}(\Omega' \wedge \omega, \Omega', \dots, \Omega') + \dots + \bar{S}(\Omega', \dots, \Omega', \Omega' \wedge \omega) \\ &= \bar{S}(\Omega' \wedge \omega^0, \Omega', \dots, \Omega') + \dots + \bar{S}(\Omega', \dots, \Omega', \Omega' \wedge \omega^0). \end{aligned}$$

S is invariant under the group G , and ω^0 is \mathfrak{g} -valued; hence, the last line of the above string of equalities equals zero.

3. Dependence on the G -structure.

THEOREM 2. Ω_S depends only on the G -structure of M .

Proof. Let B again denote the principal G -bundle over M defined by the G -structure on M (that is, the bundle to which the frame bundle of M can be reduced). For $i = 1, 2$, let F^i be an isometric imbedding of M in S^{n+p-1} , let $f^i = \pi \circ F^i$, and let g^i be a mapping of M into B_G^+ satisfying the conditions

1. $\partial \circ g^i = f^i$,
2. g^i induces the bundle B over M .

It must be proved that $g^{1*}(\Omega'_S) = g^{2*}(\Omega'_S)$. Let ϕ and Φ be the mappings defined by

1. $F^2(m) = \phi(F^1(m))$ for each point m of M ,
2. $g^2(m) = \Phi(g^1(m))$ for each point m of M .

So it suffices to prove that $\Phi^*(\Omega'_S) = \Omega'_S$.

The proof that $\Phi^*(\Omega'_S) = \Omega'_S$ proceeds as follows. Let $M'^1 = g^1(M)$, let m^1 be a point of M'^1 , and let $n^1 = \partial(m^1)$. Suppose that $(n^1, f^1_1, \dots, f^1_n)$ is a point of $O(n+p)/O(p-1)$ with $\tau(n^1, f^1_1, \dots, f^1_n) = m^1$ (note that f^1_1, \dots, f^1_n is an orthonormal basis of the n -plane n^1). Let F^1_1, \dots, F^1_n be vectors tangent to M'^1 at m^1 satisfying the condition

$$(\alpha \circ \partial)(F^1_k) = f^1_k, \quad k = 1, \dots, n.$$

We will show that

$$\Omega'_S(\Phi(F^1_1), \dots, \Phi(F^1_{2q})) = \Omega'_S(F^1_1, \dots, F^1_{2q}),$$

thus proving the theorem.

A. B_G^+ a nonprincipal bundle over S^{n+p-1} ; its associated principal bundle is the bundle $O(n+p)$ over S^{n+p-1} . Thus the connection H' induces a connection H , (concept of horizontal vectors) on B_G^+ as a bundle over S^{n+p-1} . We will show in C, below, that there exists an element h in $O(n+p)$ satisfying the condition

$$H_*(\Phi(F^1_k)) = H_*(L_h F^1_k), \quad k = 1, \dots, n$$

(L_h denotes left-translation by the element h).

It then follows from the fact that Ω'_S is H ,-horizontal and invariant under the left-action of $O(n + p)$ that

$$\begin{aligned} \Omega'_S(\Phi(F_1^1), \dots, \Phi(F_{2q}^1)) &= \Omega'_S(H, \Phi(F_1^1), \dots, H, \Phi(F_{2q}^1)) \\ &= \Omega'_S(H, L_h(F_1^1), \dots, H, L_h(F_{2q}^1)) \\ &= \Omega'_S(L_h F_1^1, \dots, L_h F_{2q}^1) \\ &= (L_h^* \Omega'_S)(F_1^1, \dots, F_{2q}^1) \\ &= \Omega'_S(F_1^1, \dots, F_{2q}^1), \end{aligned}$$

as required.

B. We first show that $(\partial(\Phi(m^1)), \phi(f_1^1), \dots, \phi(f_n^1))$ is a point of $O(n + p)/O(p - 1)$ which projects onto the point $\Phi(m^1)$ under τ .

Since g^1 induces the bundle B , there exists a point b of B which maps into $(n^1, f_1^1, \dots, f_n^1)$ under g^1 . B is a subbundle of the bundle of frames of M ; hence $b = (m, e_1, \dots, e_n)$, with m a point of M and with e_1, \dots, e_n an orthonormal basis of the tangent space to M at m . Thus

1. $f^1(m) = n^1$,
2. $F^1(e_k) = f_k^1, k = 1, \dots, n$.

The mapping g^2 also induces the bundle B over M , and so the point

$$(f^2(m), F^2(e_1), \dots, F^2(e_n))$$

is in $O(n + p)/O(p - 1)$ and projects onto the point $g^2(m)$ under τ . Since $f^2(m) = \partial(\Phi(m^1))$ and $F^2(e_k) = \phi(f_k^1)$ (see a and b below) the assertion of **B** is proved.

- a. $\partial(\Phi(m^1)) = \partial(\Phi(g^1(m))) = \partial(g^2(m)) = f^2(m)$,
- b. $\phi(f_k^1) = \phi(F^1(e_k)) = F^2(e_k), k = 1, \dots, n$.

C. $O(n + p)$ acts transitively on $O(n + p)/O(p - 1)$ by left-translation. So there exists an element h of $O(n + p)$ with

$$(\partial(\Phi(m^1)), \phi(f_1^1), \dots, \phi(f_n^1)) = L_h(\partial(m^1), f_1^1, \dots, f_n^1).$$

Hence $L_h(f_k^1) = \phi(f_k^1), k = 1, \dots, n$, and we have:

- a. $(\alpha \circ \partial)(L_h F_k^1) = L_h(\alpha \circ \partial)(F_k^1) = L_h(f_k^1) = \phi(f_k^1), k = 1, \dots, n$;
- b. F_k^1 is the tangent vector to M'^1 at m^1 which projects onto f_k^1 under $\alpha \circ \partial$; hence $\Phi(F_k^1)$ is the tangent vector to $M'^2 = \Phi(M'^1)$ at $m^2 = \Phi(m^1)$ which projects onto $\phi(f_k^1)$ under $\alpha \circ \partial, k = 1, \dots, n$.

Hence $L_h F_k^1$ and $\Phi(F_k^1)$ are vectors at m^2 both of which project onto $\phi(f_k^1)$ under $\alpha \circ \partial$. Hence their H ,-horizontal components $H_*(L_h F_k^1)$ and $H_*(\Phi(F_k^1))$ coincide.

4. Invariant integration. If β is a left-invariant r -form on $O(n + p)$ and if g is an element of $SO(n + p)$, let β^g denote the right-translate $R_g^*(\beta)$ of β . Let $I(\beta)$ denote the following r -form on $O(n + p)$:

If x is a point of $O(n + p)$ and if the integration is taken with respect to the normalized Haar measure on $SO(n)$, then

$$I(\beta)(x) = \int_{SO(n)} \beta^g(x) dg.$$

The integral I has the following properties;

1. $I \circ I = I$;
2. $d \circ I = I \circ d$;
3. if β is invariant under the right-action of $SO(n)$, then $I(\beta) = \beta$.

For a detailed discussion and proofs, see [2].

DEFINITION. If \bar{x} is a tangent vector of $O(n + p)$, we will denote by $V'(\bar{x})$ the unique vector satisfying the condition

$$\bar{x} = H'(\bar{x}) + V'(\bar{x}).$$

THEOREM 3. $I(\Lambda_S)$ is an exact form on $O(n + p)$.

Proof. The remainder of this section is devoted to the proof of this theorem. The following formula for differentiation will be used:

If Δ is a left-invariant $(q - 1)$ -form on a Lie group, and if X_1, \dots, X_q are left-invariant vector fields on the group, then

$$(1) \quad q \cdot d\Delta(X_1, \dots, X_q) = \sum_{r < s} (-1)^{r+s+1} \cdot \Delta([X_r, X_s], X_1, \dots, \hat{X}_r, \dots, \hat{X}_s, \dots, X_q).$$

We first show, in (1)-(8') below, that $I(\Lambda_S)$ is closed.

The Bianchi identity for Ω' can be written in the form $d\Omega' = \Omega' \wedge \omega'$. Hence

$$\begin{aligned} dI(\Lambda_S) &= I(dS(\Omega', \dots, \Omega')) \\ &= I(S(d\Omega', \Omega', \dots, \Omega')) + \dots + I(S(\Omega', \dots, \Omega', d\Omega')) \\ &= I(S(\Omega' \wedge \omega', \Omega', \dots, \Omega')) + \dots + I(S(\Omega', \dots, \Omega', \Omega' \wedge \omega')). \end{aligned}$$

Since Ω' is H' -horizontal and ω' is V' -vertical, it follows that it suffices to prove that

$$dI(\Lambda_S)(\bar{v}, \bar{h}_1, \dots, \bar{h}_{2q}) = 0$$

for any V' -vertical vector \bar{v} and any H' -horizontal vectors $\bar{h}_1, \dots, \bar{h}_{2q}$.

We extend $\bar{v}, \bar{h}_1, \dots, \bar{h}_{2q}$ to left-invariant vector fields on $O(n + p)$, to be denoted by $\bar{V}, \bar{H}_1, \dots, \bar{H}_{2q}$. The sphere S^{n+p-1} is a symmetric homogeneous space, that is, the brackets $[\bar{H}_i, \bar{H}_j]$ are V' -vertical for all $1 \leq i, j \leq 2q$. Formula (1) thus shows that

$$(2) \quad \begin{aligned} &(2q + 1) \cdot dI(\Lambda_S)(\bar{V}, \bar{H}_1, \dots, \bar{H}_{2q}) \\ &= I(\Lambda_S)([\bar{V}, \bar{H}_1], \bar{H}_2, \dots, \bar{H}_{2q}) + \dots + I(\Lambda_S)(\bar{H}_1, \dots, \bar{H}_{2q-1}, [\bar{V}, \bar{H}_{2q}]). \end{aligned}$$

Let us for the moment consider one of the terms

$$I(\Lambda_S)(\bar{H}_1, \dots, \bar{H}_{r-1}, [\bar{V}, \bar{H}_r], \bar{H}_{r+1}, \dots, \bar{H}_{2q}).$$

It follows from the definition of I and of Λ_S that this term is equal to

$$(3) \quad (2q)! \sum_{j_s; s} \varepsilon \cdot \delta_{r, j_s} \cdot \int_{SO(n)} S(\Omega'(R_g \bar{H}_{j_1}, R_g \bar{H}_{j_2}), \dots, \Omega'(R_g [\bar{V}, \bar{H}_{j_s}], R_g \bar{H}_{j_{s+1}}), \dots, \Omega'(R_g \bar{H}_{j_{2q-1}}, R_g \bar{H}_{j_{2q}})) dg,$$

where ε is the sign of the permutation taking $(1, \dots, 2q)$ into (j_1, \dots, j_{2q}) , and where the sum extends

1. over all such permutations, and
2. over all s for each fixed choice of (j_1, \dots, j_{2q}) . It follows from the symmetry of S^{n+p-1} that

$$\Omega'(R_g \bar{H}_\alpha, R_g \bar{H}_\beta) = -(1/2) \cdot R_g [\bar{H}_\alpha, \bar{H}_\beta];$$

and so the expression (3) becomes

$$(4) \quad 2^{-q} (2q)! \sum_{j_s; s} \varepsilon \cdot \delta_{r, j_s} \cdot I(S)([\bar{H}_{j_1}, \bar{H}_{j_2}], \dots, [[\bar{V}, \bar{H}_{j_s}], \bar{H}_{j_{s+1}}], \dots, [\bar{H}_{j_{2q-1}}, \bar{H}_{j_{2q}}]).$$

Application of the Jacobi identity

$$[[\bar{V}, \bar{H}_{j_s}], \bar{H}_{j_{s+1}}] - [[\bar{V}, \bar{H}_{j_{s+1}}], \bar{H}_{j_s}] = [\bar{V}, [\bar{H}_{j_s}, \bar{H}_{j_{s+1}}]],$$

then shows that

$$(5) \quad I(\Lambda_S)(\bar{H}_1, \bar{H}_2, \dots, [\bar{V}, \bar{H}_r], \dots, \bar{H}_{2q}) = 2^{-q} (2q)! \sum_{j_s; s} \varepsilon \cdot \delta_{r, j_s} \cdot I(S)([\bar{H}_{j_1}, \bar{H}_{j_2}], \dots, [\bar{V}, [\bar{H}_{j_s}, \bar{H}_{j_{s+1}}]], \dots, [\bar{H}_{j_{2q-1}}, \bar{H}_{j_{2q}}]).$$

We thus conclude from (2) and (5) that

$$(6) \quad 2^q (2q + 1) ((2q)!)^{-1} \cdot dI(\Lambda_S)(\bar{V}, \bar{H}_1, \dots, \bar{H}_{2q}) = \sum_{j; } \varepsilon \cdot I(S)([\bar{H}_{j_1}, \bar{H}_{j_2}], \dots, [\bar{V}, [\bar{H}_{j_s}, \bar{H}_{j_{s+1}}]], \dots, [\bar{H}_{j_{2q-1}}, \bar{H}_{j_{2q}}]).$$

We next show that the right side of the equality (6) is equal to

$$(7) \quad (q + 1) \sum \varepsilon \cdot dI(S)(\bar{V}, [\bar{H}_{j_1}, \bar{H}_{j_2}], \dots, [\bar{H}_{j_{2q-1}}, \bar{H}_{j_{2q}}]).$$

This can be done in the following way. With the aid of (1), the expression (7) becomes

$$(8) \quad \sum_{j_s; s} \varepsilon \cdot I(S)([\bar{H}_{j_1}, \bar{H}_{j_2}], \dots, [\bar{V}, [\bar{H}_{j_s}, \bar{H}_{j_{s+1}}]], \dots, [\bar{H}_{j_{2q-1}}, \bar{H}_{j_{2q}}]) + \sum_{r < s} \varepsilon \cdot \sum (-1)^{\beta(r, s)} \cdot I(S)([[\bar{H}_{j_r}, \bar{H}_{j_{r+1}}], [\bar{H}_{j_s}, \bar{H}_{j_{s+1}}]], [\bar{H}_{j_1}, \bar{H}_{j_2}], \dots),$$

where $\beta(r, s) = 1 + (1/2)((r + 1) + (s + 1))$.

For fixed r and s , however, we have:

- a. $\varepsilon(\dots, j_r, j_{r+1}, \dots, j_s, j_{s+1}, \dots) = \varepsilon(\dots, j_s, j_{s+1}, \dots, j_r, j_{r+1}, \dots)$;
- b. $[[\bar{H}_{j_r}, \bar{H}_{j_{r+1}}], [\bar{H}_{j_s}, \bar{H}_{j_{s+1}}]] = - [[\bar{H}_{j_s}, \bar{H}_{j_{s+1}}], [\bar{H}_{j_r}, \bar{H}_{j_{r+1}}]]$.

Hence (8) reduces to

$$(8') \quad \sum_{j_s; s} \varepsilon \cdot I(S)([\bar{H}_{j_1}, \bar{H}_{j_2}], \dots, [\bar{V}, [\bar{H}_{j_s}, \bar{H}_{j_{s+1}}]], \dots, [\bar{H}_{j_{2q-1}}, \bar{H}_{j_{2q}}]) + 0.$$

This is, of course, the right side of the equality (6).

It is now a simple matter to show that $dI(\Lambda_S) = 0$. In view of (6) and (7), it suffices to show that $dI(S) = 0$. The form $I(S)$ is an invariant q -form on $SO(n)$, hence, (see [2]) it is closed: $dI(S)$ vanishes identically on the Lie algebra of $SO(n)$. Since the fields $[\bar{H}_{j_k}, \bar{H}_{j_{k+1}}]$ and $[[V, H_{j_k}], H_{j_{k+1}}]$ all lie in this Lie algebra, we see that $dI(\Lambda_S) = 0$.

The proof of exactness proceeds as follows. $I(\Lambda_S)$ is a horizontal form on $O(n + p)$ as a bundle over $O(n + p)/O(p - 1)$, and is invariant under the right-action of $O(p - 1)$. Hence, there exists a form β on $O(n + p)/O(p - 1)$ with $\sigma^*(\beta) = I(\Lambda_S)$. The form β is closed because $I(\Lambda_S)$ is; hence, the form β is exact, since for p much larger than q the q th cohomology group of $O(n + p)/O(p - 1)$ vanishes. So $\beta = d\mathcal{L}$ for some form \mathcal{L} , and so $I(\Lambda_S) = d(\sigma^*(\mathcal{L}))$.

5. Homomorphisms.

LEMMA. Suppose $S = \sum_i S_i \cdot T_i$, with T_i invariant by G and S_i invariant by $O(n)$. Then $\Omega_S = 0$.

Proof. It suffices to show that $\Omega_S = 0$ if S invariant by $O(n)$. For $i = 1, \dots, n$, let \bar{X}_i be the $(n + p) \times (n + p)$ matrix with $(i, n + p)$ th entry 1, $(n + p, i)$ th entry -1 , and all other entries 0. As before (part B, Lemma 2, §2) the tangent planes to $g(M)$ are spanned by vectors $\bar{x}_1, \dots, \bar{x}_n$ with $H_i(\bar{x}_i) = \delta(\bar{X}_i)$; and so it suffices to show that

$$\Lambda_S(\bar{X}_{i_1}, \dots, \bar{X}_{i_2}) = 0$$

for any $1 \leq i_1, \dots, i_{2q} \leq n$ and any invariant symmetric q -linear functional S on the Lie algebra of $O(n)$.

If g is an element of $O(n)$, then

$$\Lambda_S(\bar{X}_{i_1}, \dots, \bar{X}_{i_{2q}}) = \Lambda_S(R_g \bar{X}_{i_1}, \dots, R_g \bar{X}_{i_{2q}}).$$

If we choose g in $O(n)$ with

1. $R_g \bar{X}_{i_1} = -\bar{X}_{i_1}$,
2. $R_g \bar{X}_{i_k} = \bar{X}_{i_k}$, $k = 2, \dots, 2q$,

we thus have

$$\Lambda_S(\bar{X}_{i_1}, \dots, \bar{X}_{i_{2q}}) = -\Lambda_S(\bar{X}_{i_1}, \dots, \bar{X}_{i_2}),$$

as required.

BIBLIOGRAPHY

1. A. W. Adler, *Classifying spaces for Kähler metrics*, Math. Ann. (to appear).
2. C. Chevalley and S. Eilenberg, *Cohomology theory of Lie groups and Lie algebras*, Trans. Amer. Math. Soc. **63** (1948), 85-124.

PURDUE UNIVERSITY,
LAFAYETTE, INDIANA