

FACTORIZATION OF POLYNOMIALS OVER BANACH ALGEBRAS⁽¹⁾

BY
JOHN A. LINDBERG, JR.

Introduction. A Banach algebra in this paper will be understood to mean a commutative, semi-simple Banach algebra with multiplicative unit e . By the carrier space Φ_A of the Banach algebra A , we mean the space of multiplicative linear functionals on A to C , the complex numbers, and it is to be endowed with the usual weak* topology (cf. [6]). For $a \in A$, \hat{a} denotes the Gelfand transform of a defined on Φ_A and \hat{A} will denote the collection of such functions. We set the following notation. x will be used to denote an indeterminate over A as well as over \hat{A} and C . If $\alpha(x) = \sum_{i=0}^n \alpha_i x^i$ is a polynomial over A , let $\hat{\alpha}(x)$ and $\alpha_h(x)$ denote, respectively, $\sum_{i=0}^n \hat{\alpha}_i x^i$ and $\sum_{i=0}^n \hat{\alpha}_i(h) x^i$, $h \in \Phi_A$.

The set $Z(\alpha(x), A) = \{(h, \lambda) \in \Phi_A \times C : \alpha_h(x) \neq 0 \text{ and } \alpha_h(\lambda) = 0\}$, $\alpha(x) \in A[x]$, plays an important role in the present paper. $Z(\alpha(x), A)$ is topologized with the relative product topology from $\Phi_A \times C$. The mapping π is defined by $\pi(h, \lambda) = h$, $(h, \lambda) \in Z(\alpha(x), A)$. The multiplicity function M of $\alpha(x)$ is defined as follows: for $(h, \lambda) \in Z(\alpha(x), A)$, $M(h, \lambda)$ is equal to the multiplicity of λ as a root of $\alpha_h(x) = 0$.

In §1 we introduce the concept of M -neighborhood of a point in $Z(\alpha(x), A)$. We say that $W \subset Z(\alpha(x), A)$ is a M -neighborhood of $(h_0, \lambda_0) \in Z(\alpha(x), A)$ if W is a neighborhood in $Z(\alpha(x), A)$ of (h_0, λ_0) and if, for each $h \in \pi(W)$, $M(h_0, \lambda_0)$ is equal to the sum of the values of M at the points (h, λ) in $\pi^{-1}(h) \cap W$. Proposition 1.1 states that M -neighborhoods exist and that they form a base for the neighborhood system at each point of $Z(\alpha(x), A)$. The remainder of this section contains most of the topological lemmas needed for our work on factorization. Particular attention is paid to the case where $Z(\alpha(x), A)$ contains a compact open subset K ($\pi(K) = \Phi_A$) on which M is constant. When this condition obtains, K and Φ_A decompose topologically and this decomposition in turn forces $\alpha(x)$ to factor.

The main factorization theorem (2.1) says that if $\alpha(x) \in A[x]$ and if K ($\pi(K) = \Phi_A$) is a compact open subset of $Z(\alpha(x), A)$, then there exists a monic polynomial $\beta(x) \in A[x]$ such that $\beta(x)$ is a factor of $\alpha(x)$, $Z(\beta(x), A) = K$ and $Z(\alpha(x)/\beta(x), A) = Z(\alpha(x), A) \sim K$. A more detailed description of the factorization of monic polynomials follows. For example, it is shown that if A is indecomposable

Received by the editors February 5, 1962 and, in revised form, April 3, 1963.

(1) This paper is based on portions of the author's Doctoral dissertation which was prepared under the guidance of Professor B. R. Gelbaum, University of Minnesota.

and if M is locally constant on $Z(\alpha(x), A)$, then the factorization of $\alpha(x)$ into irreducible factors is unique (to within units).

§3 deals with the problem of finding zeros of a given polynomial over A in A and in extensions of A which are also Banach algebras. Theorem 3.2 extends, for semi-simple Banach algebras, a theorem (4.3 [1]) of Arens and Calderón. Our theorem states that if $\alpha(x) \in A[x], f \in C(\Phi_A)$ such that $\hat{\alpha}(f) = 0$ and $M(\cdot, f(\cdot))$ is locally constant on Φ_A , then $f \in \hat{A}$.

1. **Some topological results.** Let A be a Banach algebra and

$$\alpha(x) = \sum_{i=0}^n \alpha_i x^i \in A[x].$$

(The results of this section do not depend on the semi-simplicity of A .) The topology in Φ_A is to be the usual weak* topology. Then the topology in $Z(\alpha(x), A)$ induced by the product topology in $\Phi_A \times C$ is precisely the weakest topology which renders the functions $\sum_{i=0}^m \hat{a}_i f^i, a_i \in A$, continuous on $Z(\alpha(x), A)$, where $f(h, \lambda) \equiv \lambda$ for $(h, \lambda) \in Z(\alpha(x), A)$. From its definition, it is easily seen that π is continuous with respect to the prescribed topologies and it follows from Proposition 1.1 that π is also an open mapping. $\pi(Z(\alpha(x), A))$ need not be all of Φ_A ; however, it is an open subset of Φ_A .

DEFINITION. A neighborhood W in $Z(\alpha(x), A)$ of (h_0, λ_0) is called a M -neighborhood of (h_0, λ_0) if, for each $h \in \pi(W)$,

$$M(h_0, \lambda_0) = \sum_{(h, \lambda) \in \pi^{-1}(h) \cap W} M(h, \lambda),$$

that is, $M(h_0, \lambda_0)$ is equal to the sum of the values of M at the points in $\pi^{-1}(h) \cap W$.

It follows from the above definition that if W is a M -neighborhood of (h, λ) and if V is a neighborhood in Φ_A of h , then $\pi^{-1}(V) \cap W$ is also a M -neighborhood of (h, λ) .

PROPOSITION 1.1. *Let $\alpha(x) = \sum_{i=0}^n \alpha_i x^i \in A[x]$ and $(h_0, \lambda_0) \in Z(\alpha(x), A)$. If W is a neighborhood in $Z(\alpha(x), A)$ of (h_0, λ_0) , then there exists a M -neighborhood W_0 of (h_0, λ_0) which is contained in W .*

Proof. It suffices to prove the proposition for the case where W is of the form $(V \times \{\lambda \in C : |\lambda - \lambda_0| < \varepsilon\}) \cap Z(\alpha(x), A)$, where $\varepsilon > 0$ and V is a neighborhood in Φ_A of h_0 . Consider the polynomial $p(z_0, z_1, \dots, z_n, w) = z_0 + z_1 w + \dots + z_n w^n$. Then λ_0 is a root of $p(\hat{\alpha}_0(h_0), \hat{\alpha}_1(h_0), \dots, \hat{\alpha}_n(h_0), w) = 0$. Suppose that λ_0 is a root of multiplicity m . From the Weierstrass preparation theorem, it follows that there exists $s > 0$ and $t > 0$ such that if $|z_i - \hat{\alpha}_i(h_0)| < t, i = 0, 1, \dots, n$, then $p(z_0, z_1, \dots, z_n, w) = 0$ possesses exactly m roots (multiplicities counted) in $\{w \in C : |w - \lambda_0| < s\}$ (cf. [5]). Furthermore, s can be so chosen that it is less than ε . Now, $V_0 = \{h \in \Phi_A : |\hat{\alpha}_i(h) - \hat{\alpha}_i(h_0)| < t, i = 0, 1, \dots, n\}$ is a neighborhood in Φ_A of h_0 so that $V \cap V_0$ is also a neighborhood in Φ_A of h_0 . Let W_0

denote the neighborhood $((V \cap V_0) \times \{\lambda \in C : |\lambda - \lambda_0| < s\}) \cap Z(\alpha(x), A)$. Then $W_0 \subset W$ and W_0 is a M -neighborhood of (h_0, λ_0) . Q.E.D.

The above proposition tells us that the M -neighborhoods of a point in $Z(\alpha(x), A)$ form a base for the neighborhood system of that point.

From the construction in the proof of the proposition we have the following important result:

1.2. π is an open mapping.

This can be seen in the following way. Let W be an open subset of $Z(\alpha(x), A)$ and let $(h, \lambda) \in W$. Then let W_0 be the M -neighborhood of (h, λ) contained in W which was constructed above. It has the property that $\pi(W_0)$ is a neighborhood in Φ_A of h . Thus it follows that $\pi(W)$ is open and that π must be an open mapping.

Since the topology in $Z(\alpha(x), A)$ is Hausdorff, we can conclude from the proposition that any two distinct points of $Z(\alpha(x), A)$ possess disjoint M -neighborhoods. If the points have the same first coordinate, then M -neighborhoods can be found which have the same image under π . For if W_1 and W_2 are the disjoint M -neighborhoods of these points, then $V = \pi(W_1) \cap \pi(W_2)$ is a neighborhood so that $\pi^{-1}(V) \cap W_1$ and $\pi^{-1}(V) \cap W_2$ are the desired M -neighborhoods.

In the next section we concern ourselves with subsets K of $Z(\alpha(x), A)$ which are compact and open, and such that $\pi(K) = \Phi_A$. The next proposition shows such sets behave, with respect to Φ_A , in a manner which is reminiscent of covering spaces (cf. [3]). If $h \in \Phi_A$, let $r(h)$ denote the cardinality of $\pi^{-1}(h) \cap K$ and let $(h, \lambda_1), (h, \lambda_2), \dots, (h, \lambda_{r(h)})$ denote the distinct points in $\pi^{-1}(h) \cap K$.

PROPOSITION 1.3. *Let $\alpha(x) = \sum_{i=0}^n \alpha_i x^i \in A[x]$ and K be a compact open subset of $Z(\alpha(x), A)$ such that $\pi(K) = \Phi_A$. Then for each $h \in \Phi_A$, there are disjoint M -neighborhoods $V_k(h)$ of the points (h, λ_k) and positive numbers (depending on h) t and $s_k, k = 1, 2, \dots, r(h)$ such that*

(i) *for each $k, V_k(h)$ is contained in $\{(h', \lambda') \in K : |\hat{\alpha}_j(h') - \hat{\alpha}_j(h)| < t, j = 0, 1, \dots, n \text{ and } |\lambda - \lambda_k| < s_k\}$,*

(ii) *for each $k, \pi(V_k(h)) = \pi(V(h_1))$, and*

(iii) $\pi^{-1}(\pi(V_1)) \cap K = \bigcup_{k=1}^{r(h)} V_k(h)$.

Furthermore, if $U(h) = \{(z_0, z_1, \dots, z_n) \in C^{n+1} : |z_j - \hat{\alpha}_j(h)| < t, j = 0, 1, \dots, n\}$, then

(iv) $p(z_0, z_1, \dots, z_n, w) = \sum_{i=0}^n z_i w^i$ has exactly $M(h, \lambda_k)$ roots (each distinct root repeated according to its multiplicity) in $\{\lambda \in C : |\lambda - \lambda_k| < s_k\}$ if $(z_0, z_1, \dots, z_n) \in U(h)$.

Proof. Let $W_1, \dots, W_{r(h)}$ be disjoint neighborhoods of the points $(h, \lambda_1), \dots, (h, \lambda_{r(h)})$. In view of Proposition 1.1 (and its proof) and the fact that K is open, we may and do assume that the W_k 's are M -neighborhoods satisfying (i) for an appropriate choice of $t > 0$ and $s_k > 0, k = 1, 2, \dots, r(h)$, and such that (iv) is satisfied for this choice. As seen earlier, the openness of π allows us to assume that

$\pi(W_k) = \pi(W_1)$, $k = 1, 2, \dots, r(h)$. Now let $W = \pi^{-1}(\pi(W_1)) \cap K \sim \bigcup_{k=1}^{r(h)} W_k$ and consider the closure \bar{W} of W . Since K is compact, \bar{W} is compact and $\pi(\bar{W})$ is a compact subset of Φ_A ; hence, $\pi(\bar{W})$ is a closed subset of Φ_A . Since $(h, \lambda_k) \notin \bar{W}$, for each k , $h \notin \pi(\bar{W})$ so that there is a neighborhood V in Φ_A of h such that $\pi(W) \cap V = \emptyset$; thus, $\pi^{-1}(V) \cap W = \emptyset$. If we take $V_k = W_k \cap \pi^{-1}(V)$, then the above properties are satisfied with t and the s_k 's chosen as above. Q.E.D.

By M locally constant (π a local homeomorphism) at $(h, \lambda) \in Z(\alpha(x), A)$, we mean that there exists a neighborhood W in $Z(\alpha(x), A)$ of (h, λ) on which M is constant ($\pi|_W : W \rightarrow \pi(W)$ is a homeomorphism).

LEMMA 1.4. *Let $\alpha(x) \in A[x]$. The following are equivalent:*

- (i) M is locally constant at $(h, \lambda) \in Z(\alpha(x), A)$,
- (ii) π is a local homeomorphism at (h, λ) , and
- (iii) there exists a neighborhood V in Φ_A of h and a function $f \in C(V)$ such that $\alpha_h(f(h')) = 0$ for each $h' \in V$, $M(\cdot, f(\cdot))$ is constant on V and $f(h) = \lambda$.

Proof. We show (iii) \Rightarrow (ii) \Rightarrow (i) \Rightarrow (iii).

(iii) \Rightarrow (ii). Let W be a M -neighborhood of (h, λ) and let V_0 be a neighborhood in Φ_A of h such that $V_0 \subset V$ and $h' \in V_0$ implies that $(h', f(h')) \in W$. Then $\pi^{-1}(V_0) \cap W$ is a M -neighborhood of (h, λ) and hence we assume that $W = W \cap \pi^{-1}(V_0)$. Now, if $\pi|_W$ is not a homeomorphism, it must be the case that there is a point $(h', \lambda') \in W$ such that $\lambda' \neq f(h')$ since π is an open mapping. But this means that $M(h', f(h')) < M(h', f(h')) + M(h', \lambda') \leq M(h, f(h))$. This is a contradiction since $M(\cdot, f(\cdot))$ is constant on V and so it follows that $\pi|_W$ must be a homeomorphism.

(ii) \Rightarrow (i). Suppose that W is a neighborhood in $Z(\alpha(x), A)$ of (h, λ) such that $\pi|_W$ is a homeomorphism. It may be assumed that W is a M -neighborhood since W contains such a neighborhood. Now, M must be constant on W . For if it is not, then there are two points (h', λ') and (h', λ'') in W , $\lambda' \neq \lambda''$. But this is a contradiction so that M is constant on W .

(i) \Rightarrow (iii). Let W be a neighborhood in $Z(\alpha(x), A)$ of (h, λ) on which M is constant. We assume that W is a M -neighborhood of (h, λ) . Thus, the cardinality of $\pi^{-1}(h') \cap W$ is always equal to one if $h' \in \pi(W)$. This means that $\pi|_W$ is a homeomorphism so that if f is defined to be the function on $V = \pi(W)$ such that $(h', f(h')) \in W$, $h' \in V$, then f is continuous on V . Furthermore, $\alpha_h(f(h')) = 0$, $h' \in V$, $f(h) = \lambda$ and $M(\cdot, f(\cdot))$ is constant on V . Q.E.D.

If K is a compact open subset of $Z(\alpha(x), A)$ such that $\pi(K) = \Phi_A$, then we define $m_K(h)$ to be the sum of the values of M at the points of $\pi^{-1}(h) \cap K$. (If we are dealing in any given situation with only one such set K , we will simply denote m_K by m .)

LEMMA 1.5. *If K is a compact open subset of $Z(\alpha(x), A)$ such that $\pi(K) = \Phi_A$, then there exists disjoint open subsets X_i , $i = 1, 2, \dots, s$, of Φ_A such that $\bigcup_{i=1}^s X_i = \Phi_A$ and the function $m(\cdot)$ is constant on each X_i .*

Proof. For each $h \in \Phi_A$, let $(h, \lambda_1(h)), \dots, (h, \lambda_{r(h)}(h))$ denote the points in $\pi^{-1}(h) \cap K$ and let $X_i = \{h \in \Phi_A : m(h) = i\}$. Then the X_i are open sets in Φ_A . For if $h \in X_i$, for some i , then let $V_k, k = 1, 2, \dots, r(h)$, be the neighborhoods of the (h, λ_k) 's which were given in Proposition 1.3. Then $h' \in \pi(V_1)$ implies that

$$\begin{aligned} m(h') &= \sum_{i=1}^{r(h')} M(h', \lambda(h')) = \sum_{i=1}^{r(h)} \left(\sum_{(h', \lambda') \in \pi^{-1}(h') \cap V_i} M(h', \lambda') \right) \\ &= \sum_{i=1}^{r(h)} M(h, \lambda_i(h)) = m(h) \end{aligned}$$

so that $h' \in X_i$ and hence $\pi(V_1) \subset X_i$. Thus, each X_i is open in Φ_A . Clearly, the X_i 's are disjoint. Q.E.D.

We now turn our attention to the case where $\alpha(x)$ is a monic polynomial. In this case, $Z(\alpha(x), A)$ is compact and its image under π is all of Φ_A . ($Z(\alpha(x), A)$ is, for monic polynomials, the carrier space of the Banach algebra $A[x]/(\alpha(x))$; see [2] for details. If M is locally constant on $Z(\alpha(x), A)$, then $Z(\alpha(x), A)$ and Φ_A must both decompose topologically in the way specified in

LEMMA 1.6. *Suppose that $\alpha(x) \in A[x]$ is a monic polynomial. If M is locally constant on $Z(\alpha(x), A)$, then there exist disjoint open sets $X_i, i = 1, 2, \dots, s$, in Φ_A and disjoint open sets Y_{ij} in $Z(\alpha(x), A), j = 1, 2, \dots, s_i; i = 1, 2, \dots, s$, such that (i) $\Phi_A = \bigcup_{i=1}^s X_i$, (ii) $\pi^{-1}(X_i) = \bigcup_{j=1}^{s_i} Y_{ij}$, (iii) $\text{card } \pi^{-1}(h) \cap Y_{ij}$ is constant on X_i for each i and j and (iv) M is constant on each Y_{ij} .*

Proof. Let F_j denote the set $\{(h, \lambda) \in Z(\alpha(x), A) : M(h, \lambda) = j\}$. Since M is locally constant on $Z(\alpha(x), A)$, the F_j 's are open subsets of $Z(\alpha(x), A)$ and hence closed since they are disjoint. Now, let G_{ij} denote the set $\{h \in \Phi_A : \text{card}(\pi^{-1}(h) \cap F_j) = i\}$. We will show that G_{ij} is both open and closed by showing that if h is in the closure \bar{G}_{ij} of G_{ij} , then h is in the interior of G_{ij} . To this end, let $h \in \Phi_A$ and let $(h, \lambda_k), k = 1, 2, \dots, r(h)$, denote the points in $\pi^{-1}(h)$. We can find disjoint neighborhoods V_k in $Z(\alpha(x), A)$ of the points (h, λ_k) such that for each k (i) $\pi|_{V_k} : V_k \rightarrow \pi(V_k)$ is a homeomorphism, (ii) M is constant on V_k and (iii) $\pi(V_k) = \pi(V_1)$. Furthermore, (iv) $\pi^{-1}(\pi(V_1)) = \bigcup_{k=1}^{r(h)} V_k$. To see this, take M -neighborhoods W_k of the (h, λ_k) which satisfy conditions (ii) and (iv) of Proposition 1.3. In view of Lemma 1.4, we know there exists, for each k , a neighborhood U_k in $Z(\alpha(x), A)$ of (h, λ_k) such that M is constant on U_k and $\pi|_{U_k} : U_k \rightarrow \pi(U_k)$ is a homeomorphism. Let $V = \bigcap_{k=1}^{r(h)} \pi(U_k)$. If V_k is taken to be $\pi^{-1}(V) \cap W_k$, then the above conditions (i)–(iv) are satisfied.

Now, if $h \in \bar{G}_{ij}$, then there exists a $h' \in \pi(V_1) \cap G_{ij}$ and hence there are distinct points $(h', \lambda'_1), \dots, (h', \lambda'_i)$ in F_j . Each point $(h', \lambda'_t), 1 \leq t \leq i$, belongs to one and only one of the V_k 's. Denote the subscript by $k(t)$. Then $M \equiv j$ on $V_{k(t)}$. Furthermore, if $M \neq j$ on V_k , then $V_k \cap V_{k(t)} = \emptyset, t = 1, 2, \dots, i$. Thus, $\text{card}(\pi^{-1}(h) \cap F_j) = i$ and h must be in G_{ij} . Also, by the same argument, $\pi(V_1) \subset G_{ij}$ so that G_{ij} is both open and closed.

For each $h \in \Phi_A$, let $V(h) = \bigcap_{h \in G_{ij}} G_{ij}$. Since there are only finitely many G_{ij} 's the $V(h)$'s are both open and closed in Φ_A . We next show that if $h' \in V(h)$, then $V(h') = V(h)$. For clearly, $V(h') \subset V(h)$. Let $G_{i_1, j_1}, \dots, G_{i_n, j_n}$ be the set of all G_{ij} 's whose intersection is $V(h)$. Then $\sum_{k=1}^n i_k j_k = m(h)$ ($m(\cdot)$ defined above). If the intersection $V(h')$ is smaller than $V(h)$, then

$$m(h) = \sum_{k=1}^n i_k j_k < \sum_{(h', \lambda') \in \pi^{-1}(h')} M(h', \lambda').$$

But the right-hand side is equal to $m(h')$. Observing that $m(\cdot)$ is constant on Φ_A , we have a contradiction. Thus, $V(h) = V(h')$.

Since the sets $V(h)$ are open and either identical or disjoint, the compactness of Φ_A implies that there are points h_1, h_2, \dots, h_s in Φ_A such that $\Phi_A = \bigcup_{i=1}^s V(h_i)$ and $V(h_i) \cap V(h_j) = \emptyset$ if $i \neq j$. Let $X_i = V(h_i)$ and $Y_{ij} = \pi^{-1}(X_i) \cap F_j$. With this choice, conditions (i)–(iv) of the lemma are satisfied. Q.E.D.

2. Factorization of polynomials. Let us recall that we are assuming that our Banach algebras are semi-simple. Examples can be given which show that this assumption is a necessary one.

Suppose that $\alpha(x), \alpha_i(x) \in A[x]$, $i = 1, 2$, and that $\alpha(x) = \alpha_1(x)\alpha_2(x)$. Then $Z(\alpha(x), A) = Z(\alpha_1(x), A) \cup Z(\alpha_2(x), A)$. (We are assuming that not all the coefficients of $\hat{\alpha}(x)$ vanish at the same points of Φ_A .) Associated with each set $Z(\alpha_i(x), A)$ we have the multiplicity function M_i of the polynomial $\alpha_i(x)$, and if we agree that M_i is to be zero at those points of $Z(\alpha(x), A)$ for which it is not defined, then $M_1(h, \lambda) + M_2(h, \lambda) = M(h, \lambda)$, $(h, \lambda) \in Z(\alpha(x), A)$. Furthermore, the sets $Z(\alpha_i(x), A)$ are closed in the prescribed topology in $Z(\alpha(x), A)$.

Suppose now that there exist closed subsets X_1 and X_2 of $Z(\alpha(x), A)$ such that $Z(\alpha(x), A) = X_1 \cup X_2$ and functions M_1 and M_2 which behave like multiplicity functions on X_1 and X_2 , respectively, and $M_1 + M_2 = M$. Thus, we ask: is this enough to force a factorization over A of $\alpha(x)$, with factors $\alpha_1(x)$ and $\alpha_2(x)$ such that $Z(\alpha_i(x), A) = X_i$, $i = 1, 2$. The answer is yes if $A = C(\Omega)$ but is no in general. In fact, if A is regular and self-adjoint the above conditions are still not sufficient. Witness the following example. Let $A = L_1(I)$, $I = \text{integers}$, let $f \in A$ such that $|f| \notin \hat{A}$, and let $\alpha(x) = x^2 - g$, $\hat{g} = |f|^2$. Then the above conditions are satisfied if $X_i = \{(h, \lambda) \in \Phi_A \times C : \lambda = (-1)^i |f(h)|\}$ and $M_i \equiv 1$, $i = 1, 2$, but $\alpha(x)$ fails to factor into a product $\alpha_1(x)\alpha_2(x)$, $Z(\alpha_i(x), A) = X_i$, $i = 1, 2$, since this would imply that $|f| \in \hat{A}$. Thus, we see that the factorization of a polynomial $\alpha(x)$ over A is not, in general, a topological property of the space $Z(\alpha(x), A)$. However, if $X_1 \cap X_2 = \emptyset$, then $\alpha(x)$ must factor as indicated above. Before proving this (Theorem 2.1) we set the following notation.

Let $\alpha(x) = \sum_{i=0}^n \alpha_i x^i \in A[x]$. We will use (z) for (z_0, z_1, \dots, z_n) , $((z), \lambda)$ for $(z_0, z_1, \dots, z_n, \lambda)$ and $(z(h))$ for $(\hat{\alpha}_0(h), \hat{\alpha}_1(h), \dots, \hat{\alpha}_n(h))$, $h \in \Phi_A$. If K is a compact open subset of $Z(\alpha(x), A)$ such that $\pi(K) = \Phi_A$, then for each $h \in \Phi_A$, let

$\pi^{-1}(h) \cap K = \{(h, \lambda_1(h)), \dots, (h, \lambda_r(h)(h))\}$. Let $V_i(h) \subset K$ denote the M -neighborhood of $(h, \lambda_i(h))$ and $U(h)$ the neighborhood in C^{n+1} of $(z(h))$ which were specified in Proposition 1.3. Furthermore, let $U_i(h)$ denote the set

$$\{(z), \lambda) \in U(h) \times C : p(z_0, z_1, \dots, z_n, \lambda) = \sum_{i=0}^n z_i \lambda^i = 0, \text{ and } |\lambda - \lambda_i| < s_i\}$$

(see Proposition 1.3 for the definition of s_i ; s_i depends on h), i varying from one to $r(h)$.

THEOREM 2.1. *Let $\alpha(x) = \sum_{i=0}^n \alpha_i x^i \in A[x]$. If K is a compact open subset of $Z(\alpha(x), A)$ such that $\pi(K) = \Phi_A$, then there are mutually orthogonal idempotents e_i in A and polynomials $\beta_i(x), Q_i(x) \in A[x]$, $i = 1, 2, \dots, s$, such that (i) $e = \sum_{i=1}^s e_i$, (ii) $e_i \beta_i(x)$ is monic over $e_i A$, (iii) $K = \bigcup_{i=1}^s Z(e_i \beta_i(x), e_i A)$, and (iv) $\alpha(x) = (\sum_{i=1}^s e_i \beta_i(x)) (\sum_{i=1}^s e_i Q_i(x))$.*

Proof. For simplicity (of notation), let us first assume that the function $m(\cdot)$ (defined in the last section) is constant on Φ_A . We use m itself to denote this constant.

For each $h \in \Phi_A$, let $f_j(h)$ be defined by

$$f_j(h) = (-1)^{m-j} (\lambda_1(h) \lambda_2(h) \cdots \lambda_{m-j}(h) + \lambda_1(h) \lambda_2(h) \cdots \lambda_{m-j-1}(h) \lambda_{m-j+1}(h) + \cdots)$$

for $j = 0, 1, \dots, m - 1$. Clearly $f_j \in C(\Phi_A)$ for each j . The next step is to show that $f_j \in \hat{A}$.

For each $h \in \Phi_A$, let $F_{j,h}$ denote the function defined by

$$F_{j,h}(z) = (-1)^{m-j} (\mu_1(z) \mu_2(z) \cdots \mu_{m-j}(z) + \mu_1(z) \mu_2(z) \cdots \mu_{m-j-1}(z) \mu_{m-j+1}(z) + \cdots)$$

for $j = 0, 1, \dots, m - 1$, where $(z) \in U(h)$ and $\mu_k(z)$ denotes the roots of $p(z_0, z_1, \dots, z_n, \mu) = 0$, each repeated according to its multiplicity, such that $((z), \mu_k(z)) \in \bigcup_{i=1}^{r(h)} U_i(h)$. Each $F_{j,h}$ is analytic in $U(h)$ (cf., for example, [4]). Furthermore, if $h' \in V(h)$, then $F_{j,h} = F_{j,h'}$ in $U(h) \cap U(h')$, and $F_{j,h}(z(h)) = f_j(h)$ for each $h \in \Phi_A$. (This says that $h \rightarrow F_{j,h}$ is a continuous mapping in the sense described in [1].) Hence by a theorem (6.2, [1]) of Arens and Calderón, $f_j \in \hat{A}$.

Now let $\beta_i \in A$ be the unique elements such that $\hat{\beta}_j = f_j$ for each j . If $\beta(x)$ denotes the monic polynomial $\sum_{i=0}^{m-1} \beta_i x^i + x^m$, then $Z(\beta(x), A) = K$ and the multiplicity function of $\beta(x)$ is precisely $M | K$, where M is the multiplicity function of $\alpha(x)$. Since $\beta(x)$ is monic, there are polynomials $Q(x), R(x) \in A[x]$ such that $\alpha(x) = \beta(x) Q(x) + R(x)$ and the degree of $R(x)$ is less than that of $\beta(x)$. Now, for each $h \in \Phi_A$, we have that $\alpha_h(x) = \beta_h(x) Q_h(x) + R_h(x)$ so that $R_h(x) = 0$ since $\beta_h(x) = 0$ has m roots (including multiplicities). Hence $R(x) = 0$ and $\alpha(x) = \beta(x) Q(x)$.

We now turn to the more general situation of the theorem where m is not necessarily constant on Φ_A . However, in this case m is locally constant on Φ_A .

(cf. Lemma 1.5). Let $X_i, i = 1, 2, \dots, s$, be the (disjoint) sets of constancy of the function m . By a theorem of Šilov (cf. [6]), there exist mutually perpendicular idempotents e_1, e_2, \dots, e_s , such that $e = \sum_{i=1}^s e_i$ and $\hat{e}_i(h) = 1$ if and only if $h \in X_i$. Also, $\Phi_{e_i A}$ is (identifiable with) X_i . Applying the first part of the proof to $e_i \alpha(x)$ over $e_i A$, for each i , we have polynomials $\beta_i(x)$ and $Q_i(x)$ over A such that $e_i \alpha(x) = e_i \beta_i(x) Q_i(x)$, $e_i \beta_i(x)$ monic over $e_i A$. Thus $(\sum_{i=1}^s e_i \beta_i(x))(\sum_{i=1}^s e_i Q_i(x)) = \sum_{i=1}^s e_i \alpha(x) = \alpha(x)$. It is clear that $Z(\sum_{i=1}^s e_i \beta_i(x), A) = K$. Q.E.D.

The factorization demonstrated above is not unique. For example, when $s = 2$,

$$\begin{aligned} \alpha(x) &= (e_1 \beta_1(x) + e_2 \beta_2(x))(e_1 Q_1(x) + e_2 Q_2(x)) \\ &= (e_1 \beta_1(x) + e_2 Q_2(x))(e_1 Q_1(x) + e_2 \beta_2(x)). \end{aligned}$$

We will take up this question of uniqueness in Theorem 2.4.

We now turn to the question of when a factor appears as a multiple factor. If $\alpha(x), \beta(x) \in A[x]$ and if k is a positive integer such that $\beta(x)^k = \alpha(x)$, then $M(h, \lambda)/k$ is an integer for every $(h, \lambda) \in Z(\beta(x), A)$ ($Z(\beta(x), A)$ is a subset of $Z(\alpha(x), A)$). Without any assumptions on $\alpha(x)$, the converse is false, that is, the existence of such a k need not imply that there exists a polynomial $\beta(x)$ such that $\beta(x)^k = \alpha(x)$. Witness the following example. Let $\Delta = \{\lambda \in C : |\lambda| \leq 1\}$ and $A = \{f \in C(\Delta) : f \text{ analytic in the interior of } \Delta \text{ and } f'(0) = 0\}$. If $\alpha(x) = z^2 x^2 - 2z^3 x + z^4$, then M is identically equal to two on $Z(\alpha(x), A)$ but $\alpha(x)$ is not the square of any other polynomial. This difficulty vanishes if we assume that $\alpha(x)$ is monic over A . That this is true follows from the corollary below, which is stated in a more general setting for later use. In the remainder of this section, we restrict our attention to monic polynomials.

In the following corollary, $Z(\alpha(x), A)$ is assumed to be the union of disjoint compact open subsets $X_i, i = 1, 2, \dots, t$. The functions $m_{X_i}(\cdot)$ defined earlier will be denoted by $m_i(\cdot)$.

COROLLARY 2.2. *Let $\alpha(x)$ be a monic polynomial over A and X_1, X_2, \dots, X_t , be disjoint open subsets of $Z(\alpha(x), A)$, the union of which is $Z(\alpha(x), A)$. Assume that each $m_i(\cdot)$ is constant on Φ_A and denote its constant value by m_i . If there are positive integers k_1, k_2, \dots, k_t , such that for each i and every $(h, \lambda) \in X_i, M(h, \lambda)/k_i$ is an integer, then there are monic polynomials $\alpha_i(x), i = 1, 2, \dots, t$, of degree m_i/k_i over A such that $Z(\alpha_i(x), A) = X_i$ and $\alpha(x) = \prod_{i=1}^t \alpha_i(x)^{k_i}$.*

Proof. By repeated applications of the theorem, there are monic polynomials $\beta_1(x), \beta_2(x), \dots, \beta_t(x) \in A[x]$ such that $\alpha(x) = \prod_{i=1}^t \beta_i(x)$ and $Z(\beta_i(x), A) = X_i$. The degree of $\beta_i(x)$ is m_i and the multiplicity function of $\beta_i(x)$ is $M_i = M|_{X_i}$. Recall that the coefficient $\hat{\beta}_{ij}$ of X_j in $\beta_i(x)$ satisfies

$$\hat{\beta}_{ij}(h) = (-1)^{m_i-j} (\lambda_1^{(i)}(h) \dots \lambda_{m_i-j}^{(i)}(h) + \dots),$$

where $\lambda_1^{(i)}(h), \dots, \lambda_{m_i}^{(i)}(h)$ are all the roots of $(\beta_i)_h(x) = 0$, each distinct root repeated

according to its multiplicity. Let $\mu_1^{(i)}(h), \dots, \mu_{n_i}^{(i)}(h)$, $n_i = m_i/k_i$, denote all the roots of $(\beta_i)_h(x) = 0$ but now each distinct $\mu_j^{(i)}(h)$ is to be repeated only $M(h, \mu_j^{(i)}(h))/k_i$ times. Then the functions $\alpha_{i,j}$ defined by

$$\alpha_{i,j}(h) = (-1)^{n_i-j} (\mu_1^{(i)}(h) \cdots \mu_{n_i-j}^{(i)}(h) + \cdots), \quad j = 0, 1, 2, \dots, n_i - 1,$$

are continuous on Φ_A . Furthermore, if $\alpha_i(x) = \sum_{j=0}^{n_i-1} \alpha_{i,j} x^j + x^{n_i}$, then $\alpha_i(x)^{k_i} = \sum_{i=0}^{m_i} \beta_{ij} x^i$. But this means that the coefficients of $\alpha_i(x)$ must lie in \hat{A} ; for the coefficient of x^{m_i-l} in $\alpha_i(x)^{k_i}$, $n_i = m_i/k_i \geq l \geq 1$, is of the form

$$k_i(\alpha_{i, n_i-l}) + (\text{terms in } \alpha_{ij}, n_i > j > n_i - l)$$

so that by induction the coefficients of $\alpha_i(x)$ must lie in \hat{A} . Hence, $\alpha(x) = \prod_{i=1}^s \alpha_i(x)^{k_i}$. Since A is semi-simple, we may and do take $\alpha_i(x)$ to be a polynomial over A . This proves the corollary since the $\alpha_i(x)$ are monic, of degree $n_i = m_i/k_i$ and $Z(\alpha_i(x), A) = X_i$.

We now turn our attention to the case where M is locally constant on $Z(\alpha(x), A)$, with $\alpha(x)$ monic.

THEOREM 2.3. *Let $\alpha(x)$ be a monic polynomial over A . If M is locally constant on $Z(\alpha(x), A)$, then there are mutually orthogonal idempotents $e_i \in A$, positive integers k_{ij} and polynomials $\alpha_{ij}(x) \in A[x]$, $j = 1, 2, \dots, s_i$; $i = 1, 2, \dots, s$, such that $e_i \alpha_{ij}(x)$ is monic over $e_i A$, the discriminant of $\prod_{j=1}^{s_i} e_i \alpha_{ij}(x)$ is invertible in $e_i A$ and*

$$\alpha(x) = \sum_{i=1}^s e_i \prod_{j=1}^{s_i} \alpha_{ij}(x)^{k_{ij}}.$$

Furthermore, if $\gamma(x)$ is any other polynomial over A such that $\gamma_h(\lambda) = 0$ for each $(h, \lambda) \in Z(\alpha(x), A)$, then $\gamma(x)$ is a multiple of $\beta(x) = \sum_{i=1}^s e_i \prod_{j=1}^{s_i} \alpha_{ij}(x)$.

Proof. Let X_i, Y_{ij} , $j = 1, 2, \dots, s_i$; $i = 1, 2, \dots, s$ be the sets constructed in Lemma 1.6. Let k_{ij} be equal to the constant value of M on Y_{ij} . It follows from Theorem 2.1 and its corollary that there are mutually orthogonal idempotents e_i, \dots, e_s and polynomials $\alpha_{ij}(x)$ over A such that $\sum_{i=1}^s e_i = e$,

$$e_i \alpha(x) = \prod_{j=1}^{s_i} e_i \alpha_{ij}(x)^{k_{ij}}$$

and $Y_{ij} = \{(h, \lambda) \in X_i \times C : (\alpha_{ij})_h(\lambda) = 0\}$, $i = 1, 2, \dots, s$. The degree of $e_i \alpha_{ij}(x)$ is equal to

$$\sum_{(h, \lambda) \in \pi^{-1}(h) \cap Y_{ij}} (M(h, \lambda)/k_{ij})$$

which in turn is equal to the number of points in $\pi^{-1}(h) \cap Y_{ij}$, $h \in X_i$. Hence there are as many distinct roots of $(\alpha_{ij})_h(x) = 0$, $h \in X_i$, as the degree of $e_i \alpha_{ij}(x)$

so that the discriminant of $e_i\alpha_{ij}(x)$ is invertible in e_iA . Since the Y_{ij} are disjoint, the discriminant of $e_i\prod_{j=1}^{s_i}\alpha_{ij}(x)$ is invertible in e_iA .

Now suppose that $\gamma_h(\lambda)=0$ for all $(h, \lambda)\in Z(\alpha(x), A)$. Then $e_i\gamma(x) = e_i\beta(x)Q_i(x) + e_iR_i(x)$ for some $Q_i(x)$ and $R_i(x)$ in $A[x]$, where the degree of $e_iR_i(x)$ is less than the degree of $e_i\beta(x)$ (defined above). Since $\beta_h(x) = 0$ has no multiple roots for $h \in X_i$, $(R_i)_h(x) = 0$ has a greater number of distinct roots than its degree so that the coefficients of $e_iR_i(x)$ belong to every maximal ideal of e_iA . Hence $R_i(x) = 0$ since A is semi-simple and

$$\gamma(x) = \left(\sum_{i=1}^s e_i Q_i(x) \right) \beta(x). \tag{Q.E.D.}$$

The last assertion of the above theorem says that, in other words, if M is locally constant on $Z(\alpha(x), A)$, with $\alpha(x)$ monic, then the radical $A[x]/(\alpha(x))$ is a principal ideal generated by the coset $\beta(x) + (\alpha(x))$ (cf. [2]).

We return now to the question of uniqueness of the factorization. Our earlier comments showed that if A is the direct sum of two of its ideals e_1A and e_2A , $e_i e_j = \delta_{ij} e_i$, $e = e_1 + e_2$, then the factorization is not unique. If A is not decomposable, the factorization is still not necessarily unique. In this case M is not locally constant on $Z(\alpha(x), A)$, as following theorem shows.

THEOREM 2.4. *Suppose that A is an indecomposable algebra. If $\alpha(x)$ is a monic polynomial over A and if its multiplicity function M is locally constant on $Z(\alpha(x), A)$, then a factorization of $\alpha(x)$ into irreducible factors is unique (to within units).*

Proof. Recall first that $Z(\alpha(x), A)$ is compact since $\alpha(x)$ is monic. In this situation it is well known that the mapping π from $Z(\alpha(x), A)$ onto Φ_A is a closed mapping.

Since A is indecomposable, we know that Φ_A is connected (cf. [7]). Hence $Z(\alpha(x), A)$ must be the union of disjoint open connected subsets X_i such that $\pi(X_i) = \Phi_A$. For if $Z(\alpha(x), A) = Q_1 \cup Q_2$, where Q_1 and Q_2 are disjoint and open, and hence closed, then $\pi(Q_i)$ is both open and closed in Φ_A so that $\pi(Q_i) = \Phi_A$ or $Q_i = \emptyset$. This process of decomposition, when repeated, must terminate after a finite number of steps so that there are disjoint open connected sets X_1, X_2, \dots, X_t in $Z(\alpha(x), A)$ such that $\bigcup_{i=1}^t X_i = \Phi_A$ and $\pi(X_i) = \Phi_A$ for each i . Furthermore, the assumption that M is locally constant on $Z(\alpha(x), A)$ implies that M is constant on each X_i . This assumption together with the fact that X_i is connected implies that the number of points in $\pi^{-1}(h) \cap X_i$, for each i , is constant on Φ_A . Let k_i denote the constant value of M on X_i and n_i the $\text{card}(\pi^{-1}(h) \cap X_i)$. Applying Theorem 2.3, we have that there are monic polynomials $\alpha_i(x) \in A[x]$ such that the degree of $\alpha_i(x)$ is n_i , $\alpha(x) = \prod_{i=1}^t \alpha_i(x)^{k_i}$ and $Z(\alpha_i(x), A) = X_i$. Also, the discriminant of each $\alpha_i(x)$ is invertible in A . That the factors $\alpha_i(x)$ are irreducible follows from the argument given below.

Suppose that $\alpha(x) = \beta_1(x)\beta_2(x)\cdots\beta_n(x)$. We can assume without loss of gen-

erality that $n = 2$. Then $Z(\alpha(x), A) = Z(\beta_1(x), A) \cup Z(\beta_2(x), A)$ and the $Z(\beta_i(x), A)$ are closed in $Z(\alpha(x), A)$. We now show that each $Z(\beta_i(x), A)$ is also open. Let M_i denote the function on $Z(\alpha(x), A)$ which is equal to the multiplicity function of $\beta_i(x)$ on $Z(\beta_i(x), A)$ and zero otherwise. Then $M_1(h, \lambda) + M_2(h, \lambda) \equiv M(h, \lambda)$. Each point $(h, \lambda) \in Z(\beta_1(x), A)$ possesses a neighborhood W_0 in $Z(\alpha(x), A)$ on which M_1 is constant. To see this, choose a neighborhood W_0 in $Z(\alpha(x), A)$ of (h, λ) , on which M is constant, so that $W_0 \cap Z(\alpha(x), A)$ is a M -neighborhood (relative to the polynomial $\beta_1(x)$) in $Z(\beta_1(x), A)$ of (h, λ) and $W_0 \cap Z(\beta_2(x), A)$ (relative to the polynomial $\beta_2(x)$) is a M -neighborhood in $Z(\beta_2(x), A)$ of (h, λ) if $(h, \lambda) \in Z(\beta_2(x), A)$ but $W_0 \cap Z(\beta_2(x), A) = \emptyset$ if $(h, \lambda) \notin Z(\beta_2(x), A)$. Then $M_i(h', \lambda) \leq M_i(h, \lambda)$ for each $(h', \lambda') \in W_0$ and $i = 1, 2$. If M_1 is not constant on W_0 , then there is a point $(h', \lambda') \in W_0$ such that $M_1(h', \lambda') < M_1(h, \lambda)$. By the above restrictions on W_0 , $M_2(h', \lambda') \leq M_2(h, \lambda)$ so that $M(h', \lambda') = M_1(h', \lambda') + M_2(h', \lambda') < M_1(h, \lambda) + M_2(h, \lambda) = M(h, \lambda)$. This is a contradiction and so it must be the case that M_i is constant on W_0 . Since $M_1(h, \lambda) > 0$, W_0 is a subset of $Z(\beta_1(x), A)$. Thus, $Z(\beta_1(x), A)$ is open, as well as closed, in $Z(\alpha(x), A)$. This means that $Z(\beta_1(x), A)$ is the union of some of the X_i , say X_1, X_2, \dots, X_s (after possible relabeling) and fails to intersect the other X_i 's. $\prod_{i=1}^s \alpha_i(x)$ is a monic polynomial with an invertible discriminant since the $Z(\alpha_i(x), A)$'s are disjoint. It follows from the previous theorem that $\prod_{i=1}^s \alpha_i(x)$ is a factor of $\beta_1(x)$.

By continuing this process we have that

$$\beta_j(x) = Q_j \prod_{i=1}^t \alpha_i(x)^{k_{ij}}, \quad j = 1, 2,$$

for an appropriate choice of non-negative integers k_{ij} and $Q_j \in A$. Since $Q_1 Q_2 = e$, the Q_i are invertible in A . Q.E.D.

3. Zeros of polynomials. In this section we deal with sufficient conditions under which $\alpha(x) \in A[x]$ has a zero in (i) a Banach algebra extension of A , that is, a Banach algebra B in which A is embedded isomorphically and isometrically, and (ii) in A itself. In [2] it is shown that if A is a Banach algebra and if $\beta(x) \in A[x]$ has an invertible leading coefficient, then $A[x]/(\beta(x))$ possesses a norm which makes it a Banach algebra extension of A , the isomorphism being the canonical one.

It is also shown in [2] that if A is semi-simple and if R denotes the radical of $A[x]/(\beta(x))$, then $(A[x]/(\beta(x)))/R$ possessess a norm so that it is a Banach algebra extension of A . Again the isomorphism is the canonical one. We now make use of these propositions in proving

THEOREM 3.1. *Let A be a semi-simple Banach algebra and let $\alpha(x)$ be a polynomial over A . If K is a compact open subset of $Z(\alpha(x), A)$ such that $\pi(K) = \Phi_A$, then there exists a Banach algebra extension $A[b]$ of A such that $\alpha(b) = 0$.*

Proof. By Theorem 2.1, $\alpha(x) = (\sum_{i=1}^s e_i \alpha_i(x)) (\sum_{i=1}^s e_i Q_i(x))$ where the e_i are mutually orthogonal idempotents such that $e = \sum_{i=1}^s e_i$. Furthermore, the $e_i \alpha_i(x)$ are monic over $e_i A$ and $Z(\sum_{i=1}^s e_i \alpha_i(x), A) = K$. Let m denote the least common multiple of the n_i , $n_i = \text{degree of } e_i \alpha_i(x)$, and let $\beta(x) = \sum_{i=1}^s e_i \alpha_i(x)^{m/n_i}$. Then $\beta(x)$ is a monic polynomial over A and hence has a solution in $A[x]/(\beta(x))$. Now $(A[x]/(\beta(x)))/R$ is isomorphic to $(A[x]/(\beta(x)))^\wedge$, the Gelfand representation of $A[x]/(\beta(x))$. Thus, $(A[x]/(\beta(x)))^\wedge$ is of the form $\hat{A}[b]$, where $b = (x + (\beta(x)))^\wedge$, and $\sum_{i=1}^s e_i \hat{\alpha}_i(b) = 0$. By the above comments, $(A[x]/(\beta(x)))/R$ is a Banach algebra extension of A ; hence, $\hat{A}[b]$ is a Banach algebra extension of A . Q.E.D.

Our next theorem gives a sufficient condition that $\alpha(x) = 0$ has a solution in A , where $\alpha(x) \in A[x]$, and generalizes a theorem (in case A is semi-simple, which we assume here) due to Arens and Calderón.

THEOREM 3.2. *Let A be a semi-simple Banach algebra. If $\alpha(x) \in A[x]$, if $f \in C(\Phi_A)$ such that $\hat{\alpha}(f) = 0$ and if $M(\cdot, f(\cdot))$ is locally constant on Φ_A , then $f \in \hat{A}$, that is, there is an $a \in A$ such that $\alpha(a) = 0$ and $\hat{a} = f$.*

Proof. Let K denote the graph of f . Then K is a compact subset of $Z(\alpha(x), A)$. By Lemma 1.4, K is an open subset of $Z(\alpha(x), A)$ so that by Theorem 2.1 and its corollary,

$$\alpha(x) = \sum_{i=1}^s e_i \alpha_i(x)^{k_i} Q_i(x)$$

where $K = Z(\sum_{i=1}^s e_i \alpha_i(x), A)$ and the degree of $e_i \alpha_i(x)$ is one. If $e_i \alpha_i(x) = e_i x - e_i b_i$, then $\sum_{i=1}^s (e_i b_i)^\wedge$ must be f . Thus $f \in \hat{A}$, or equivalently, there exists $a \in A$ such that $\alpha(a) = 0$ and $\hat{a} = f$. Q.E.D.

The Arens-Calderón theorem (4.1, [1]) assumes that $M(\cdot, f(\cdot))$ is identically one but does not require that A be semi-simple. Precisely: if $\sum_{i=0}^n \hat{a}_i(h)(f(h))^i = 0$ and $M(h, f(h)) = 1$ for each $h \in \Phi_A$, then there exists $a \in A$ such that $\sum_{i=0}^n \alpha_i a^i = 0$ and $\hat{a} = f$. When $M(\cdot, f(\cdot))$ is just locally constant on Φ_A and A not semi-simple, we can not draw the stronger conclusion that there is an $a \in A$ such that $\alpha(a) = 0$ and $\hat{a} = f$.

REMARK. Some of the main theorems of the Arens-Calderón paper [1] deal with functions $G(z_0, z_1, \dots, z_n, w)$ which are analytic in a neighborhood of $K = \{(\hat{\alpha}_0(h), \hat{\alpha}_1(h), \dots, \hat{\alpha}_n(h), f(h)) : h \in \Phi_A\}$. Specifically, they show that if $G(z_0, z_1, \dots, z_n, w) = 0$, $G_w(z_0, z_1, \dots, z_n, w) \neq 0$ for each $(z_0, z_1, \dots, z_n, w) \in K$, then $f \in \hat{A}$ (cf. Theorem 6.8, [1]). The condition on G_w can be replaced by the condition that the multiplicity of $f(h)$ as a root of $G(\hat{\alpha}_0(h), \hat{\alpha}_1(h), \dots, \hat{\alpha}_n(h), \lambda) = 0$ is locally constant on Φ_A . If so, then f once again must belong to \hat{A} . The proof of Theorem 3.2 can be extended so as to give a proof of the above assertion.

REFERENCES

1. R. Arens and A. P. Calderón, *Analytic functions of several Banach algebra elements*, Ann. of Math. (2) 6 (1955), 204-216.

2. R. Arens and K. Hoffman, *Algebraic extensions of normed algebras*, Proc. Amer. Math. Soc. 7 (1956), 203–210.
3. C. Chevalley, *Theory of Lie groups*, Princeton Univ. Press, Princeton, N. J., 1946.
4. S. Bochner and W. T. Martin, *Several complex variables*, Princeton Univ. Press, Princeton, N. J., 1948.
5. W. F. Osgood, *Lehrbuch der Funktionentheorie*, Vol. 2 (1st part) Teubner, Leipzig, 1929.
6. C. E. Rickart, *General theory of Banach algebras*, Van Nostrand, New York, 1960.
7. G. E. Šilov. *Decomposition of a commutative normed ring* Mat. Sb. (N. S) 32 (74) (1953), 353–364; Amer. Math. Soc. Transl. (2) 5 (1955), 37–48.

YALE UNIVERSITY,
NEW HAVEN, CONNECTICUT
SYRACUSE UNIVERSITY,
SYRACUSE, NEW YORK