

p -VALENT CLOSE-TO-CONVEX FUNCTIONS

BY

A. E. LIVINGSTON⁽¹⁾

1. **Introduction.** Let $S(p)$ denote the class of functions, which are regular and p -valently star-like in $|z| < 1$. A function

$$f(z) = a_1z + a_2z^2 + \dots \quad (|z| < 1)$$

is a member of $S(p)$, if there exists a positive number ρ such that for $\rho < |z| < 1$

$$(1.1) \quad \operatorname{Re} \left[\frac{zf'(z)}{f(z)} \right] > 0$$

and

$$(1.2) \quad \int_0^{2\pi} \operatorname{Re} \left[\frac{zf'(z)}{f(z)} \right] d\theta = 2p\pi.$$

The class $S(p)$ has been studied previously by Goodman [4], Robertson [9] and others. Goodman [4] has shown that a function in $S(p)$ is p -valent and has exactly p roots in $|z| < 1$.

Goodman [4] also defined the class of p -valent convex functions, which we will refer to as $C(p)$. A function

$$f(z) = a_1z + a_2z^2 + \dots \quad (|z| < 1)$$

is said to be in $C(p)$, if there exists a ρ such that for $\rho < |z| < 1$

$$(1.3) \quad 1 + \operatorname{Re} \left[\frac{zf''(z)}{f'(z)} \right] > 0$$

and

$$(1.4) \quad \int_0^{2\pi} \left[1 + \operatorname{Re} \frac{zf''(z)}{f'(z)} \right] d\theta = 2p\pi.$$

A function in $C(p)$ is at most p -valent and has $(p - 1)$ critical points in $|z| < 1$. $S(p)$ and $C(p)$ are related to each other in the same way as $S(1)$ and $C(1)$. Namely, $f(z)$ is in $C(p)$ if and only if $zf'(z)$ is in $S(p)$.

Kaplan [5] defined the class of close-to-convex functions. A function $F(z)$,

Received by the editors August 23, 1963 and, in revised form, December 5, 1963.

⁽¹⁾ This paper is part of a doctoral thesis written under the direction of Professor M. S. Robertson.

regular for $|z| < 1$, with $F(0) = 0$ and $F'(0) \neq 0$ is said to be close-to-convex if there exists $\phi(z)$ in $C(1)$ such that

$$\operatorname{Re} \left[\frac{F'(z)}{\phi'(z)} \right] > 0 \quad (|z| < 1).$$

Notice that we may rewrite the last inequality to read

$$\operatorname{Re} \left[\frac{zF'(z)}{f(z)} \right] > 0 \quad (|z| < 1)$$

for some function $f(z)$ in $S(1)$.

Umezawa [13] extended this definition to the case of p -valent functions. According to Umezawa, a function

$$F(z) = z^q + a_{q+1}z^{q+1} + \dots \quad (|z| < 1)$$

is p -valently close-to-convex, if there exists

$$\phi(z) = z^q + b_{q+1}z^{q+1} + \dots \quad (|z| < 1)$$

in $C(p)$ such that

$$(1.5) \quad \operatorname{Re} \left[\frac{F'(z)}{\phi'(z)} \right] > 0 \quad (|z| < 1).$$

It is known that a function in this class is at most p -valent in $|z| < 1$ [13].

However, Umezawa's definition requires that the zeros of $F'(z)$ and $\phi'(z)$ have the same positions and multiplicities. We will redefine the concept of a close-to-convex function by requiring that (1.5) should hold only in some range $\rho < |z| < 1$. Furthermore, we will not require that our functions be normalized.

DEFINITION. We shall say that a function

$$F(z) = a_1z + a_2z^2 + \dots \quad (|z| < 1),$$

regular for $|z| < 1$, is p -valently close-to-convex, or is in $\mathcal{H}(p)$, if it satisfies one of the following conditions.

(A) There exists a function $f(z)$ in $S(p)$ and a positive number ρ such that

$$(1.6) \quad \operatorname{Re} \left[\frac{zF'(z)}{f(z)} \right] > 0 \quad (\rho < |z| < 1).$$

(B) $F(z)$ is regular on $|z| = 1$ and there exists a function $f(z)$ in $S(p)$, also regular on $|z| = 1$, such that (1.6) holds on $|z| = 1$.

Notice that if $F(z)$ satisfies (A), then there exists a δ such that $G(z) = F(\beta z)$ satisfies (B) for $\delta < \beta < 1$.

If $F(z)$ is in $S(p)$, then taking $f(z) = F(z)$, we see that $F(z)$ is in $\mathcal{H}(p)$. Also, if $F(z)$ is in $C(p)$, then taking $f(z) = zF'(z)$, we see that $F(z)$ is in $\mathcal{H}(p)$.

In §2 we will show that a function in $\mathcal{H}(p)$ is at most *p*-valent in $|z| < 1$. We are also able to obtain sufficient conditions for a function $F(z)$ to be in $\mathcal{H}(p)$, provided $F(z)$ is regular on $|z| = 1$: If $F(z)$ has *p* zeros at the origin, then we are able to remove the condition of regularity on $|z| = 1$.

Considerable interest has been shown in the coefficient problem for functions, which are at most *p*-valent in $|z| < 1$. Goodman [3] has conjectured that if

$$F(z) = a_1z + a_2z^2 + \dots \quad (|z| < 1)$$

is regular and at most *p*-valent in $|z| < 1$, then

$$|a_n| < \sum_{k=1}^p \frac{2k(n+p)!}{(p+k)!(p-k)!(n-p-1)!(n^2-k^2)} |a_k|$$

for $n > p$.

The conjecture was proven by Goodman and Robertson [2] for a function in $S(p)$, in case all its coefficients are real and by Robertson [9] for $F(z)$ in $S(p)$, in case $a_1 = a_2 = \dots = a_{p-2} = 0$, the remaining coefficients being complex. In §3 we will prove the conjecture for the (*p* + 1)st coefficient of an arbitrary function in $\mathcal{H}(p)$. This is the largest class of *p*-valent functions for which the exact bound on the (*p* + 1)st coefficient is known. We also obtain some sharp upper and lower bounds on $|F'(z)|$ for $F(z)$ in $\mathcal{H}(p)$.

§4 deals with the radii of close-to-convexity and convexity for a function in $\mathcal{H}(p)$. If

$$F(z) = a_qz^q + a_{q+1}z^{q+1} + \dots \quad (|z| < 1)$$

is in $\mathcal{H}(p)$, then we obtain a $r_q < 1$ such that $F(z)$ is *q*-valently close-to-convex in $|z| < r_q$ and $\beta_q < 1$ such that $F(z)$ is *q*-valently convex in $|z| < \beta_q$. The numbers r_q and β_q depend upon the nonzero critical points of $F(z)$. We are able to show that the number β_q gives us the best possible result. However, we are not able to show this for the number r_q .

2. The class $\mathcal{H}(p)$. We will make use of the following lemma due to Umezawa [12].

LEMMA 1. Let $f(z)$ be regular for $|z| \leq r$ and $f'(z) \neq 0$ on $|z| = r$. Suppose that for $z = re^{i\theta}$

$$\int_0^{2\pi} d \arg df(z) = \int_0^{2\pi} \frac{\partial}{\partial \theta} [\arg zf'(z)] d\theta = \int_0^{2\pi} \operatorname{Re} \left[1 + \frac{zf''(z)}{f'(z)} \right] d\theta = 2p\pi^{(2)}.$$

If, furthermore,

⁽²⁾ Geometrically this says that the angle that the tangent to the image of $|z| = r$ makes with the positive *x*-axis goes through a change of $2p\pi$ as z traverses $|z| = r$. In other words, the image of $|z| = r$, under $w = f(z)$, makes *p*-loops.

$$\int_{\theta_1}^{\theta_2} d \arg df(z) = \int_{\theta_1}^{\theta_2} \frac{\partial}{\partial \theta} [\arg zf'(z)] d\theta > -\pi \quad \text{for } \theta_1 < \theta_2,$$

then $f(z)$ is at most p -valent in $|z| < r$.

THEOREM 1. *If $F(z)$ is in $\mathcal{S}(p)$, then $F(z)$ is at most p -valent in $|z| < 1$.*

Proof. There exists $f(z)$ in $S(p)$ and $\rho < 1$ such that

$$(2.1) \quad \operatorname{Re} \left[\frac{zF'(z)}{f(z)} \right] > 0 \quad (\rho < |z| < 1).$$

Since $zF'(z)/f(z) \neq 0$ and $zF'(z) \neq 0$ for $|z| = r$ ($\rho < r < 1$), we may define $\arg [zF'(z)/f(z)]$ and $\arg [zF'(z)]$ to be single-valued and continuous on $|z| = r$. Since $f(z) = [f(z)/zF'(z)][zF'(z)]$, then $\arg f(z) = \arg [zF'(z)] - \arg [zF'(z)/f(z)]$ will be uniquely determined and by (2.1) we have for $z = re^{i\theta}$ ($\rho < r < 1$),

$$-\frac{\pi}{2} < \arg zF'(z) - \arg f(z) < \frac{\pi}{2}.$$

Let $\theta_1 < \theta_2$, then

$$(2.2) \quad -\frac{\pi}{2} < \arg re^{i\theta_2}F'(re^{i\theta_2}) - \arg f(re^{i\theta_2}) < \frac{\pi}{2}$$

and

$$(2.3) \quad -\frac{\pi}{2} < -\arg re^{i\theta_1}F'(re^{i\theta_1}) + \arg f(re^{i\theta_1}) < \frac{\pi}{2}.$$

Combining (2.2) and (2.3), we obtain

$$(2.4) \quad \begin{aligned} & -\pi + \arg f(re^{i\theta_2}) - \arg f(re^{i\theta_1}) \\ & < \arg [re^{i\theta_2}F'(re^{i\theta_2})] - \arg [re^{i\theta_1}F'(re^{i\theta_1})] \\ & < \pi + \arg f(re^{i\theta_2}) - \arg f(re^{i\theta_1}) \end{aligned}$$

or

$$(2.5) \quad \begin{aligned} -\pi + \int_{\theta_1}^{\theta_2} d \arg f(re^{i\theta}) & < \int_{\theta_1}^{\theta_2} d \arg dF(re^{i\theta}) \\ & < \pi + \int_{\theta_1}^{\theta_2} d \arg f(re^{i\theta}). \end{aligned}$$

Since $f(z)$ is in $S(p)$,

$$\int_{\theta_1}^{\theta_2} d \arg f(re^{i\theta}) > 0.$$

Thus the left side of (2.5) gives

$$(2.6) \quad \int_{\theta_1}^{\theta_2} d \arg dF(re^{i\theta}) > -\pi.$$

Taking $\theta_1 = 0$ and $\theta_2 = 2\pi$ in (2.5) and using the fact that

$$\int_0^{2\pi} d \arg f(re^{i\theta}) = 2p\pi$$

we obtain

$$(2.7) \quad (2p - 1)\pi < \int_0^{2\pi} d \arg dF(re^{i\theta}) < (2p + 1)\pi.$$

However, the integral in (2.7) is an integral multiple of 2π . Therefore,

$$(2.8) \quad \int_0^{2\pi} d \arg dF(re^{i\theta}) = 2p\pi.$$

Thus, by Lemma 1, $F(z)$ is at most p -valent for $|z| < r$. Since r was arbitrary ($\rho < r < 1$), $F(z)$ is at most p -valent for $|z| < 1$.

Since (2.8) holds for any function in $\mathcal{H}(p)$ for some range $\rho < |z| < 1$, we easily obtain the following corollary.

COROLLARY. *If $F(z)$ is in $\mathcal{H}(p)$, then $F'(z)$ has exactly $(p - 1)$ zeros in $|z| < 1$.*

Necessary and sufficient conditions for a function $F(z)$, regular in $|z| < 1$, with $F(0) = 0$ and $F'(z) \neq 0$ to be in $\mathcal{H}(1)$ have been given by Kaplan [5]. We see from the proof of Theorem 1 that necessary conditions for $F(z)$ to be in $\mathcal{H}(p)$ are that (2.6) and (2.8) hold in some range $\rho < |z| < 1$. We will now show these conditions to be sufficient in two particular cases. The method of proof used is that established by Kaplan [5].

LEMMA 2. *Let*

$$F(z) = a_p z^p + a_{p+1} z^{p+1} + \dots$$

be regular for $|z| \leq 1$. If

$$(2.9) \quad \int_0^{2\pi} d \arg dF(z) = 2p\pi$$

and

$$(2.10) \quad \int_{\theta_1}^{\theta_2} d \arg dF(z) > -\pi \quad (\theta_1 < \theta_2)$$

for $|z| = 1$, then $F(z)$ is in $\mathcal{H}(p)$.

REMARK. We will show that there exists a function $f(z)$ in $S(p)$ with all its zeros at the origin, which is regular for $|z| < 1 + \epsilon$ for some $\epsilon > 0$, and

such that $\operatorname{Re}[zF'(z)/f(z)] > 0$ for $|z| < 1 + \epsilon$. This is actually more than we need to prove the lemma, but it is needed in the proof of Theorem 3.

Proof. Since $F(z)$ is regular on $|z| = 1$, it is regular in some circle containing $|z| \leq 1$. By continuity we then have the existence of some $\epsilon > 0$ such that (2.9) and (2.10) hold for $1 \leq |z| \leq (1 + \epsilon)$. Now, the function $z^{(1-p)}F'(z)$ is free of zeros in $|z| \leq (1 + \epsilon)$. Hence, we may define $\arg z^{(1-p)}F'(z)$ to be single-valued and continuous in $|z| \leq 1 + \epsilon$.

Let

$$p(r, \theta) = \arg[(re^{i\theta})^{(1-p)}F'(re^{i\theta})] \quad (r \leq 1 + \epsilon)$$

and

$$P(r, \theta) = p(r, \theta) + p\theta.$$

Then, since (2.9) and (2.10) hold for $|z| = 1 + \epsilon$, we have

$$P(1 + \epsilon, \theta + 2\pi) - P(1 + \epsilon, \theta) = 2p\pi,$$

$$P(1 + \epsilon, \theta_2) - P(1 + \epsilon, \theta_1) > -\pi \quad \text{for } \theta_1 < \theta_2.$$

Using an argument identical to Kaplan's [5], we may show the existence of a function $S(1 + \epsilon, \theta)$, which is increasing in θ and such that

$$(2.11) \quad S(1 + \epsilon, \theta + 2\pi) - S(1 + \epsilon, \theta) = 2p\pi$$

and

$$(2.12) \quad |S(1 + \epsilon, \theta) - P(1 + \epsilon, \theta)| \leq \frac{\pi}{2}.$$

Let

$$(2.13) \quad q(r, \theta) = \frac{1}{2\pi} \int_0^{2\pi} \frac{[(1 + \epsilon)^2 - r^2][S(1 + \epsilon, \alpha) - p\alpha] d\alpha}{(1 + \epsilon)^2 + r^2 - 2(1 + \epsilon)r \cos(\alpha - \theta)}.$$

Then, $q(r, \theta)$ is harmonic for $r < 1 + \epsilon$.

Let $Q(r, \theta) = q(r, \theta) + p\theta$ for $r < 1 + \epsilon$. Using the fact that $S(1 + \epsilon, \alpha) - p\alpha$ has period 2π , we obtain for $r < 1 + \epsilon$ and $\theta_1 < \theta_2$,

$$\begin{aligned} Q(r, \theta_2) - Q(r, \theta_1) \\ = \int_0^{2\pi} \frac{[(1 + \epsilon)^2 - r^2][S(1 + \epsilon, \alpha + \theta_2) - S(1 + \epsilon, \alpha + \theta_1)] d\alpha}{(1 + \epsilon)^2 + r^2 - 2(1 + \epsilon)r \cos \alpha}. \end{aligned}$$

Since $S(1 + \epsilon, \alpha)$ is increasing

$$Q(r, \theta_2) - Q(r, \theta_1) \geq 0.$$

Thus $(\partial/\partial\theta) Q(r, \theta) \geq 0$ for $r < 1 + \epsilon$.

Let $h(z)$ be a function, regular for $|z| < 1 + \epsilon$, such that $\operatorname{Im}[h(re^{i\theta})] = q(r, \theta)$ and let

$$f(z) = z^p e^{h(z)} = b_p z^p + \dots \quad (|z| < 1 + \epsilon).$$

For $|z| < 1 + \epsilon$,

$$\operatorname{Re} \left[\frac{zf'(z)}{f(z)} \right] = \frac{\partial}{\partial \theta} \arg f(z) = \frac{\partial}{\partial \theta} (p\theta + q(r, \theta)) = \frac{\partial}{\partial \theta} Q(r, \theta) \geq 0.$$

But $zf'(z)/f(z)$ is regular for $|z| < 1 + \epsilon$. Thus,

$$\operatorname{Re} \left[\frac{zf'(z)}{f(z)} \right] > 0 \quad \text{for } |z| < 1 + \epsilon.$$

Since $f(z)$ has p zeros, all of them at the origin,

$$\int_0^{2\pi} \operatorname{Re} \left[\frac{zf'(z)}{f(z)} \right] d\theta = 2p\pi \quad (|z| < 1 + \epsilon).$$

Hence, $f(z)$ is p -valently star-like for $|z| < 1 + \epsilon$.

Now, for $z = re^{i\theta}$, $r < 1 + \epsilon$, we have

$$\begin{aligned} \left| \arg \frac{zF'(z)}{f(z)} \right| &= |\arg zF'(z) - \arg f(z)| \\ &= |P(r, \theta) - q(r, \theta) - p\theta| \\ &= |p(r, \theta) - q(r, \theta)|. \end{aligned}$$

Since $p(r, \theta)$ is harmonic for $|z| < 1 + \epsilon$, we may write

$$(2.14) \quad p(r, \theta) = \frac{1}{2\pi} \int_0^{2\pi} \frac{[(1 + \epsilon)^2 - r^2]p(1 + \epsilon, \alpha)}{(1 + \epsilon)^2 + r^2 - 2(1 + \epsilon)r \cos(\alpha - \theta)} d\alpha.$$

Then, using (2.12), (2.13) and (2.14), we obtain

$$\begin{aligned} \left| \arg \frac{zF'(z)}{f(z)} \right| &= |p(r, \theta) - q(r, \theta)| \\ &= \frac{1}{2\pi} \left| \int_0^{2\pi} \frac{[(1 + \epsilon)^2 - r^2][P(1 + \epsilon, \alpha) - S(1 + \epsilon, \alpha)] d\alpha}{(1 + \epsilon)^2 + r^2 - 2(1 + \epsilon)r \cos(\alpha - \theta)} \right| \\ &\leq \frac{\pi}{2}. \end{aligned}$$

Thus $\operatorname{Re}[zF'(z)/f(z)] \geq 0$ for $|z| < 1 + \epsilon$. Hence, either $\operatorname{Re}[zF'(z)/f(z)] > 0$ for $|z| < 1 + \epsilon$, in which case $F(z)$ is in $\mathcal{S}(p)$, or $zF'(z)/f(z)$ reduces to a constant for $|z| < 1 + \epsilon$. In the second case $F(z)$ is in $C(p) \subset \mathcal{S}(p)$.

THEOREM 2. *Let*

$$F(z) = a_p z^p + a_{p+1} z^{p+1}, \dots \quad (|z| < 1)$$

be regular for $|z| < 1$. If (2.9) and (2.10) hold for some range $\rho < |z| < 1$, then $F(z)$ is in $\mathcal{S}(p)$.

Proof. Let $\rho < \delta < 1$. Then the function $G_\delta(z) = F(\delta z)$ is regular on $|z| = 1$

and satisfies (2.9) and (2.10) on $|z| = 1$. Hence, by Lemma 2, $G_\delta(z)$ is in $\mathcal{H}(p)$ and there exists

$$f_\delta(z) = b_p z^p + \dots \quad (|z| < 1)$$

in $S(p)$ such that

$$(2.15) \quad \operatorname{Re} \left[\frac{zG'_\delta(z)}{f_\delta(z)} \right] > 0 \quad (|z| < 1).$$

We may assume that $|b_p| = 1$. Cartwright [1] has shown that the family of p -valent functions with the moduli of the first p coefficients fixed is a normal family. Thus we may choose a sequence δ_n tending to 1, such that the sequence of functions $f_{\delta_n}(z)$ tends to a function $f(z)$ in $S(p)$. Since $zG'_{\delta_n}(z)$ tends to $zF'(z)$, we obtain from (2.15) that

$$\operatorname{Re} \left[\frac{zF'(z)}{f(z)} \right] \geq 0 \quad \text{for } |z| < 1.$$

This implies that $F(z)$ is in $\mathcal{H}(p)$.

THEOREM 3. *Let*

$$F(z) = a_q z^q + \dots \quad (1 \leq q \leq p)$$

be regular for $|z| \leq 1$. If (2.9) and (2.10) hold on $|z| = 1$, then $F(z)$ is in $\mathcal{H}(p)$.

Proof. By condition (2.9) $F'(z)$ has $(p - 1)$ zeros in $|z| < 1$, $(q - 1)$ of them at the origin. Let $\alpha_1, \alpha_2, \dots, \alpha_{p-q}$ be the nonzero roots of $F'(z)$ and let

$$G(z) = \int_0^z \frac{z^{p-q} F'(z) dz}{\prod_{i=1}^{p-q} (z - \alpha_i)(1 - \bar{\alpha}_i z)} = d_p z^p + \dots$$

$G(z)$ is regular for $|z| \leq 1$ and

$$zG'(z) = \frac{z^{p-q} z F'(z)}{\prod_{i=1}^{p-q} (z - \alpha_i)(1 - \bar{\alpha}_i z)}.$$

Since

$$\arg \left[\frac{z^{p-q}}{\prod_{i=1}^{p-q} (z - \alpha_i)(1 - \bar{\alpha}_i z)} \right] = 0 \quad \text{for } |z| = 1,$$

$\arg zG'(z) = \arg zF'(z)$ for $|z| = 1$.

Thus, $G(z)$ satisfies (2.9) and (2.10) on $|z| = 1$. Hence, by Lemma 2, $G(z)$ is in $\mathcal{H}(p)$ and there exists $f(z)$ in $S(p)$, regular for $|z| \leq 1$, such that

$$\operatorname{Re} \left[\frac{zG'(z)}{f(z)} \right] > 0 \quad (|z| \leq 1).$$

But using the same reasoning as above, we have

$$\arg \left[\frac{zG'(z)}{f(z)} \right] = \arg \left[\frac{zF'(z)}{f(z)} \right] \quad \text{on } |z| = 1.$$

Hence,

$$\operatorname{Re} \left[\frac{zF'(z)}{f(z)} \right] > 0 \quad \text{for } |z| = 1.$$

Thus, $F(z)$ is in $\mathcal{K}(p)$.

Theorem 3 immediately gives us the following lemma, which will prove useful in obtaining a bound for the $(p + 1)$ st coefficient of a function in $\mathcal{K}(p)$.

LEMMA 3. *If $F(z)$ is regular in $|z| \leq 1$ and in $\mathcal{K}(p)$, then there exists*

$$f(z) = b_p z^p + \dots \quad (|b_p| = 1)$$

regular and in $S(p)$ for $|z| \leq 1$, such that

$$\operatorname{Re} \left[\frac{zF'(z)}{f(z)} \right] > 0 \quad \text{on } |z| = 1.$$

3. Some extremal problems for the class $\mathcal{K}(p)$. The following lemma has been proven by Royster [11]. However, the proof we give, which was communicated to me by Professor M. S. Robertson, seems to be different.

LEMMA 4. *Let $f(z) = [h(z)]^{-p}$, where $h(z)$ is in $S(1)$, $h(0) = 0$, $h'(0) = 1$ and let*

$$f(z) = \sum_{n=-p}^{\infty} C_n z^n \quad (0 < |z| < 1, C_{-p} = 1),$$

then

$$|C_n| \leq \binom{2p}{n+p} \quad (n = -p, \dots, 1),$$

and these inequalities are sharp.

Proof. We write

$$(3.1) \quad z^p f(z) = z^p [h(z)]^{-p} = \sum_{n=0}^{\infty} d_n z^n \quad (|z| < 1, d_0 = 1).$$

The lemma will then be proven, if we can show

$$|d_n| \leq \binom{2p}{n} \quad (n \leq p + 1).$$

Taking the logarithm of both sides of (3.1), differentiating and multiplying through by z , we obtain

$$-\frac{zf'(z)}{pf(z)} = \frac{zh'(z)}{h(z)}.$$

Thus, we have for $|z| < 1$

$$(3.2) \quad \operatorname{Re} \left[-\frac{zf'(z)}{pf(z)} \right] = \operatorname{Re} \left[\frac{zh'(z)}{h(z)} \right] > 0 \quad (|z| < 1).$$

Let

$$P(z) = -\frac{zf'(z)}{pf(z)},$$

then

$$\operatorname{Re} \left[\frac{1}{P(z)} \right] > 0 \quad \text{for } |z| < 1.$$

Let

$$\begin{aligned} \frac{1}{P(z)} &= 1 + \sum_{n=1}^{\infty} \mu_n z^n, \\ \frac{1}{P(z)} &= -\frac{pf(z)}{zf'(z)} = -\frac{pz^p f(z)}{z^{p+1} f'(z)}, \\ -\frac{1}{P(z)} z^{p+1} f'(z) &= pz^p f(z), \end{aligned}$$

or

$$\left[-\sum_{m=0}^{\infty} \mu_m z^m \right] \left[\sum_{s=0}^{\infty} (s-p) d_s z^s \right] = p \sum_{n=0}^{\infty} d_n z^n.$$

Equating coefficients, we obtain

$$\begin{aligned} pd_n &= \sum_{r=0}^n (p-r) d_r \mu_{n-r}, \\ nd_n &= \sum_{r=0}^{n-1} (p-r) d_r \mu_{n-r}. \end{aligned}$$

Since $|\mu_{n-r}| \leq 2$, we obtain

$$(3.3) \quad n|d_n| \leq 2 \sum_{r=0}^{n-1} (p-r) |d_r|$$

provided $p - r \geq 0$. That is, provided $n \leq p + 1$. Using (3.3) and a simple induction argument, we have

$$|d_n| \leq \binom{2p}{n} \quad \text{for } n \leq p + 1.$$

That the inequalities are sharp is shown by the function

$$f(z) = \left[\frac{z}{(1+z)^2} \right]^{-p}.$$

THEOREM 4. *Let*

$$F(z) = \sum_{n=1}^{\infty} a_n z^n \quad (|z| < 1)$$

be regular and in $\mathcal{S}(p)$ for $|z| < 1$, then

$$(3.4) \quad |a_{p+1}| \leq \sum_{k=1}^p \frac{2k(2p+1)!}{(p+k)!(p-k)![(p+1)^2 - k^2]} |a_k|$$

and this inequality is sharp in all the variables $|a_1|, \dots, |a_p|$.

REMARK. This theorem was first proven for $p = 1$ by Reade [8].

Proof. We may assume without loss of generality that $F(z)$ is regular for $|z| \leq 1$. Then, by Lemma 3 there exists a function

$$f(z) = b_p z^p + \dots \quad (|b_p| = 1),$$

regular for $|z| \leq 1$ and in $S(p)$, such that

$$(3.5) \quad \operatorname{Re} \left[\frac{zF'(z)}{f(z)} \right] > 0 \quad (|z| = 1).$$

We may assume that $b_p = 1$ since $\arg[b_p]$ is not involved in the inequality to be obtained. Thus we may write $f(z)$ in the form $[\phi(z)]^p$, where

$$\phi(z) = z + \sum_{n=2}^{\infty} h_n z^n$$

is regular for $|z| < 1$ and in $S(1)$.

We may then write (3.5) in the form

$$\operatorname{Re}[zF'(z)[\phi(z)]^{-p}] > 0 \quad \text{on } |z| = 1.$$

Let

$$[\phi(z)]^{-p} = \sum_{n=-p}^{\infty} C_n z^n \quad (0 < |z| < 1, C_{-p} = 1).$$

Then

$$\begin{aligned} zF'(z) [\phi(z)]^{-p} &= \left[\sum_{n=1}^{\infty} na_n z^n \right] \left[\sum_{n=-p}^{\infty} C_n z^n \right] \\ &= \sum_{k=-(p-1)}^{\infty} d_k z^k, \end{aligned}$$

where

$$d_k = \sum_{n=1}^{p+k} C_{-(n-k)} na_n \quad (k = -(p-1), \dots).$$

Consider the function $G(z)$ given by

$$\begin{aligned} (3.6) \quad G(z) &= zF'(z) [\phi(z)]^{-p} - \sum_{k=-(p-1)}^{-1} d_k z^k \\ &\quad + \sum_{k=-(p-1)}^{-1} \bar{d}_k z^{-k}. \end{aligned}$$

Since $\bar{z} = z^{-1}$ for $|z| = 1$, the last two terms in (3.6) add up to a purely imaginary number for $|z| = 1$. Thus,

$$\operatorname{Re}[G(z)] = \operatorname{Re}[zF'(z) [\phi(z)]^{-p}] > 0 \quad \text{for } |z| = 1.$$

But $G(z)$ is regular for $|z| \leq 1$. Therefore,

$$\operatorname{Re}[G(z)] > 0 \quad \text{for } |z| \leq 1.$$

Now

$$G(z) = d_0 + (d_1 + \bar{d}_{-1})z + \dots \quad (|z| \leq 1).$$

Hence

$$\begin{aligned} |d_1 + \bar{d}_{-1}| &\leq 2 \operatorname{Re}[d_0] \leq 2|d_0|, \\ \left| \sum_{n=1}^{p+1} C_{-(n-1)} na_n + \sum_{n=1}^{p-1} \bar{C}_{-(n+1)} n\bar{a}_n \right| &\leq 2 \left| \sum_{n=1}^p C_{-n} na_n \right|, \\ (p+1)|a_{p+1}| &\leq \sum_{n=1}^{p-1} [2n|C_{-n}| + n|C_{-(n-1)}| + n|C_{-(n+1)}|] |a_n| \\ &\quad + [2p|C_{-p}| + p|C_{-(p-1)}|] |a_p|. \end{aligned}$$

By Lemma 4

$$|C_{-k}| \leq \binom{2p}{p-k} \quad (k = 1, 2, \dots, p).$$

Therefore,

$$\begin{aligned}
 (p + 1)|a_{p+1}| &\leq \sum_{n=1}^{p-1} \left[2n \binom{2p}{p-n} + n \binom{2p}{p-n+1} + n \binom{2p}{p-n-1} \right] |a_n| \\
 &\quad + \left[2p + p \binom{2p}{1} \right] |a_p| \\
 &= (p + 1) \sum_{n=1}^p \frac{2n(2p + 1)!}{(p + n)!(p - n)![(p + 1)^2 - n^2]} |a_n|
 \end{aligned}$$

which is (3.4).

We remark that the inequality is sharp, since it is known to be sharp for $f(z)$ in $S(p)$ with real coefficients [2], [4].

In order to obtain bounds on $|F'(z)|$ for $F(z)$ in $\mathcal{H}(p)$, we will make use of the following lemma.

LEMMA 5. *Let*

$$F(z) = a_q z^q + \dots \quad (|z| \leq 1)$$

be regular and in $\mathcal{H}(p)$ for $|z| \leq 1$. Let $\alpha_1, \alpha_2, \dots, \alpha_{p-q}$ be the nonzero critical points of $F'(z)$ in $|z| < 1$. Then the function

$$H(z) = \int_0^z z^{p-q} F'(z) \left[\prod_{i=1}^{p-q} \left(\frac{\alpha_i}{|\alpha_i|} - \frac{z}{|\alpha_i|} \right) (\bar{\alpha}_i z - 1) \right]^{-1} dz$$

is regular for $|z| \leq 1$ and in $\mathcal{H}(p)$.

Proof. By Lemma 3, there exists

$$h(z) = b_p z^p + \dots \quad (|b_p| = 1),$$

regular and in $S(p)$ for $|z| \leq 1$, such that

$$\operatorname{Re} \frac{zF'(z)}{h(z)} > 0 \quad \text{for } |z| = 1.$$

$$\frac{zH'(z)}{h(z)} = \frac{z^{p-q} z F'(z) \left[\prod_{i=1}^{p-q} \left(\frac{\alpha_i}{|\alpha_i|} - \frac{z}{|\alpha_i|} \right) (\bar{\alpha}_i z - 1) \right]^{-1}}{h(z)}.$$

But,

$$\arg \left(z^{p-q} \left[\prod_{i=1}^{p-q} \left(\frac{\alpha_i}{|\alpha_i|} - \frac{z}{|\alpha_i|} \right) (\bar{\alpha}_i z - 1) \right]^{-1} \right) = 0 \quad \text{on } |z| = 1.$$

Thus,

$$\frac{zH'(z)}{h(z)} = M \frac{zF'(z)}{h(z)}, \quad M > 0 \quad \text{on } |z| = 1.$$

Hence,

$$\operatorname{Re} \left[\frac{zH'(z)}{h(z)} \right] > 0 \quad \text{for } |z| = 1.$$

Therefore, $H(z)$ is in $\mathcal{H}(p)$.

THEOREM 5. *Let*

$$F(z) = a_q z^q + \dots \quad (|z| < 1),$$

be regular and in $\mathcal{H}(p)$ for $|z| < 1$. Let $\alpha_1, \alpha_2, \dots, \alpha_{p-q}$ be the nonzero critical points of $F(z)$ and let $\rho = \max |\alpha_i|$ and $\rho^ = \min |\alpha_i|$. Then*

$$(3.7) \quad |F'(re^{i\theta})| \leq \frac{(1+r)r^{q-1}}{(1-r)^{2p+1}} q|a_q| \left[\prod_{i=1}^{p-q} \left(1 + \frac{r}{|\alpha_i|} \right) (1+r|\alpha_i|) \right] \quad (r < 1),$$

$$(3.8) \quad |F'(re^{i\theta})| \geq \frac{(1-r)r^{q-1}}{(1+r)^{2p+1}} q|a_q| \left[\prod_{i=1}^{p-q} \left(\frac{r}{|\alpha_i|} - 1 \right) (1-r|\alpha_i|) \right] \quad (\rho < r < 1),$$

$$(3.9) \quad |F'(re^{i\theta})| \geq \frac{(1-r)r^{q-1}}{(1+r)^{2p+1}} q|a_q| \left[\prod_{i=1}^{p-q} \left(1 - \frac{r}{|\alpha_i|} \right) (1-r|\alpha_i|) \right] \quad (r < \rho^*).$$

All these inequalities are sharp, equality being attained by the function

$$F_0(z) = \int_0^z \frac{(1+z)z^{q-1}}{(1-z)^{2p+1}} q|a_q| \prod_{i=1}^{p-q} \left(1 + \frac{z}{|\alpha_i|} \right) (1+z|\alpha_i|) dz.$$

Note that inequality (3.7) was obtained by Umezawa [13] for his class of p -valent close-to-convex functions.

Proof. We may assume without loss of generality that $F(z)$ is regular for $|z| \leq 1$. Consider the functions $H(z)$ and $h(z)$, given in Lemma 5 and in its proof.

$$\frac{zH'(z)}{h(z)} = d_0 + d_1 z + \dots \quad (|z| \leq 1),$$

where

$$d_0 = \frac{qa_q}{b_p} \left[\prod_{i=1}^{p-q} (-e^{i \arg \alpha_i}) \right]^{-1}.$$

Then

$$\frac{1}{\operatorname{Re}[d_0]} \left[\frac{zH'(z)}{h(z)} - i \operatorname{Im}[d_0] \right] = P(z),$$

where $\operatorname{Re} P(z) > 0$ for $|z| < 1$ and $P(0) = 1$. Thus,

$$\left| \frac{P(z) - 1}{P(z) + 1} \right| \leq |z|.$$

Hence

$$\left| \frac{\frac{zH'(z)}{h(z)} - d_0}{\frac{zH'(z)}{h(z)} + \bar{d}_0} \right| \leq |z| = r,$$

$$(1 - r) \left| \frac{zH'(z)}{h(z)} \right| \leq (1 + r) |d_0| = (1 + r)q|\alpha_q|.$$

Using the known bound

$$|h(z)| \leq \frac{r^p}{(1 - r)^{2p}} \text{ for } |z| = r$$

and using the definition of $H(z)$, we obtain

$$|F'(re^{i\theta})| \leq \frac{(1 + r)}{(1 - r)r^{p-q+1}} q|\alpha_q| |h(z)| \left| \prod_{i=1}^{p-q} \left(\frac{\alpha_i}{|\alpha_i|} - \frac{z}{|\alpha_i|} \right) (\bar{\alpha}_i z - 1) \right|$$

$$\leq \frac{(1 + r)r^{q-1}}{(1 - r)^{2p+1}} q|\alpha_q| \prod_{i=1}^{p-q} \left(1 + \frac{r}{|\alpha_i|} \right) (1 + r|\alpha_i|),$$

which is (3.7).

To obtain (3.8) and (3.9), we notice that for $z = re^{i\theta}$

$$\left| \frac{P(z) + 1}{P(z) - 1} \right| \geq \frac{1}{r},$$

$$|h(z)| \geq \frac{r^p}{(1 + r)^{2p}},$$

$$\left| \frac{\alpha_i}{|\alpha_i|} - \frac{z}{|\alpha_i|} \right| |\bar{\alpha}_i z - 1| \geq \left(\frac{r}{|\alpha_i|} - 1 \right) (1 - r|\alpha_i|) \quad (|\alpha_i| < r),$$

and

$$\left| \frac{\alpha_i}{|\alpha_i|} - \frac{z}{|\alpha_i|} \right| |\bar{\alpha}_i z - 1| \geq \left(1 - \frac{r}{|\alpha_i|} \right) (1 - r|\alpha_i|) \quad (r < |\alpha_i|).$$

Going through the same type of argument as before, we obtain (3.8) and (3.9).

The function $F_0(z)$ is in $\mathcal{S}(p)$ relative to

$$f(z) = \frac{z^q}{(1 - z)^{2p}} \prod_{i=1}^{p-q} \left(1 + \frac{z}{|\alpha_i|} \right) (1 + z|\alpha_i|).$$

Equality in (3.7) is attained by $F'_0(r)$, in (3.8) by $F'_0(-r)$, $r > \rho$, and in (3.9) by $F'_0(-r)$, $r < \rho^*$.

4. **Radii of close-to-convexity and convexity for functions in $\mathcal{H}(p)$.** Goodman [4] has proven that if

$$f(z) = a_q z^q + \dots \quad (|z| < 1)$$

is in $S(p)$, then

$$(4.1) \quad \operatorname{Re} \frac{zf'(z)}{f(z)} \geq J_q(r) \quad \text{for } r < \rho,$$

where

$$J_q(r) = q - r \left[\frac{2p}{1+r} + \sum_{i=1}^{p-q} \frac{1}{|\alpha_i| - r} + \frac{|\alpha_i|}{1 - |\alpha_i|r} \right],$$

$\alpha_1, \dots, \alpha_{p-q}$ being the nonzero roots of $f(z)$ and $\rho = \min |\alpha_i|$. $J_q(r)$ is a decreasing function of r for $r < \rho$, is positive for $r = 0$ and tends to $-\infty$ as r tends to ρ . Thus, $J_q(r)$ has a least positive root r_q and $J_q(r) > 0$ for $r < r_q$.

We thus have that $f(z)$ is q -valently star-like for $|z| < r_q$. This estimate is sharp, since (4.1) was shown to be sharp [4], equality being attained at $z = -r$ by the function

$$(4.2) \quad f(z) = z^q (1 - z)^{-2p} \prod_{i=1}^{p-q} \left(1 + \frac{z}{|\alpha_i|} \right) (1 + z |\alpha_i|).$$

THEOREM 6. *Let*

$$F(z) = a_q z^q + \dots \quad (|z| < 1)$$

be in $\mathcal{H}(p)$. Let $\alpha_1, \dots, \alpha_{p-q}$ be the nonzero roots of $F'(z)$ and let r_q be the least positive root of $J_q(r)$, defined in (4.1). Then $F(z)$ is q -valently close-to-convex for $|z| < r_q$.

Proof. We first prove the theorem for $F(z)$, regular on $|z| = 1$. Then there exists

$$f(z) = b_p z^p + \dots \quad (|z| \leq 1),$$

regular and in $S(p)$ for $|z| \leq 1$, such that

$$\operatorname{Re} \left[\frac{zF'(z)}{f(z)} \right] > 0 \quad \text{on } |z| = 1.$$

Since

$$\arg \left(z^{p-q} \left[\prod_{i=1}^{p-q} (z - \alpha_i)(1 - \bar{\alpha}_i z) \right]^{-1} \right) = 0 \quad \text{on } |z| = 1,$$

we have

$$\operatorname{Re} \left[\frac{z^{p-q} z F'(z)}{\prod_{i=1}^{p-q} (z - \alpha_i)(1 - \bar{\alpha}_i z) \cdot f(z)} \right] > 0 \quad \text{for } |z| \leq 1.$$

Let

$$g(z) = z^{q-p} \left[\prod_{i=1}^{p-q} (z - \alpha_i)(1 - \bar{\alpha}_i z) \right] f(z).$$

Then, $g(z)$ is in $S(p)$ since $\operatorname{Re}[zg'(z)/g(z)] > 0$ on $|z| = 1$. But $g(z)$ has nonzero roots at $\alpha_1, \alpha_2, \dots, \alpha_{p-q}$. Therefore, $g(z)$ is q -valently star-like for $|z| < r_q$. Since

$$\operatorname{Re} \left[\frac{z F'(z)}{g(z)} \right] > 0 \quad \text{for } |z| \leq r_q,$$

$F(z)$ is q -valently close-to-convex for $|z| < r_q$.

If $F(z)$ is not regular on $|z| = 1$, there exists a $\rho^* < 1$ such that for $\rho^* < \delta < 1$ the function $G_\delta(z) = F(\delta z)$ is in $\mathcal{S}(p)$ and regular on $|z| = 1$. $G'_\delta(z) = 0$ for $z = \alpha_i/\delta$. Thus, $G_\delta(z)$ is q -valently close-to-convex for $|z| < r_{q,\delta}$, where $r_{q,\delta}$ is the least positive root of

$$J_{q,\delta}(r) = q - r \left[\frac{2p}{1+r} + \sum_{i=1}^{p-q} \frac{\delta}{|\alpha_i| - r\delta} + \frac{|\alpha_i|}{\delta - |\alpha_i|r} \right].$$

Thus, there exists

$$f_\delta(z) = C_q z^q + \dots \quad (|z| < r_{q,\delta}, |C_q| = 1)$$

q -valently star-like for $|z| < r_{q,\delta}$, such that

$$\operatorname{Re} \left[\frac{z G'_\delta(z)}{f_\delta(z)} \right] > 0 \quad \text{for } |z| < r_{q,\delta}.$$

But $r_{q,\delta} \geq r_q$, since $J_{q,\delta}(r) \geq J_q(r)$ for $r < \min |\alpha_i|$. Thus $f_\delta(z)$ is q -valently star-like for $|z| < r_q$.

By a result of M. Cartwright [1] the family of q -valent functions $f(z) = a_q z^q + \dots$ ($|a_q| = 1$) is a normal family. Thus we may choose an increasing sequence δ_i tending to 1, such that the functions $f_{\delta_i}(z)$ tend to a function $f(z)$, which is q -valently star-like for $|z| < r_q$. Since for each i

$$\operatorname{Re} \left[\frac{z G'_{\delta_i}(z)}{f_{\delta_i}(z)} \right] > 0 \quad \text{for } |z| < r_q$$

and since $z G'_{\delta_i}(z)$ tends to $z F'(z)$, we have

$$\operatorname{Re} \left[\frac{z F'(z)}{f(z)} \right] \geq 0 \quad \text{for } |z| < r_q.$$

Thus either $\operatorname{Re}[zF'(z)/f(z)] > 0$ for $|z| < r_q$, in which case $F(z)$ is q -valently close-to-convex for $|z| < r_q$, or $[zF'(z)/f(z)]$ reduces to a constant for $|z| < r_q$. In the second case $F(z)$ is q -valently convex and hence q -valently close-to-convex for $|z| < r_q$.

THEOREM 7. *Let*

$$F(z) = a_q z^q + \dots \quad (|z| < 1),$$

be in $\mathcal{H}(p)$, then $F(z)$ is q -valently convex for $|z| < \beta_q$, where β_q is the least positive root of

$$K_q(r) = J_q(r) - \frac{2r}{1-r^2}$$

and this estimate is the best possible.

Proof. Let us first assume that $F(z)$ is regular on $|z| = 1$. Then, as we have seen before, there exists

$$g(z) = b_q z^q + \dots \quad (|z| < 1),$$

which is in $S(p)$ for $|z| < 1$, such that

$$\operatorname{Re} \left[\frac{zF'(z)}{g(z)} \right] > 0 \quad \text{for } |z| \leq 1.$$

Let

$$\begin{aligned} \frac{zF'(z)}{g(z)} &= P(z), \quad \operatorname{Re}[P(z)] > 0 \quad \text{for } |z| \leq 1, \\ 1 + \frac{zF''(z)}{F'(z)} &= \frac{zP'(z)}{P(z)} + \frac{zg'(z)}{g(z)}. \end{aligned}$$

Now $g(z)$ has the same zeros as $F'(z)$. Therefore,

$$\operatorname{Re} \left[\frac{zg'(z)}{g(z)} \right] \geq J_q(r) \quad \text{for } r < \min |\alpha_i|.$$

By a result, obtained independently by Libera [6], MacGregor [7] and Robertson [10], we have

$$\operatorname{Re} \left[\frac{zP'(z)}{P(z)} \right] \geq -\frac{2r}{1-r^2}.$$

Thus

$$\operatorname{Re} \left[1 + \frac{zF''(z)}{F'(z)} \right] \geq -\frac{2r}{1-r^2} + J_q(r) = K_q(r)$$

for $r < \min |\alpha_i|$.

Thus, if $|z| < \beta_q$

$$\operatorname{Re} \left[1 + \frac{zF''(z)}{F'(z)} \right] > 0.$$

Since $F'(z)$ has $(q-1)$ zeros in $|z| < \beta_q$, all of them at the origin,

$$\int_0^{2\pi} \operatorname{Re} \left[1 + \frac{zF''(z)}{F'(z)} \right] d\theta = 2q\pi \quad (|z| < \beta_q).$$

Thus $F(z)$ is q -valently convex for $|z| < \beta_q$.

Arguing as in Theorem 6, we may remove the assumption of regularity on $|z| = 1$.

The function

$$F(z) = \int_0^z \frac{(1+z)^{q-1}}{(1-z)^{2p+1}} \prod_{i=1}^{p-q} \left(1 + \frac{z}{|\alpha_i|} \right) (1+z|\alpha_i|) dz$$

shows that the radius found is sharp, since

$$1 + \frac{zF''(z)}{F'(z)} = K_q(r)$$

for $z = -r$, $r < \min |\alpha_i|$.

REFERENCES

1. M. Cartwright, *Some inequalities in the theory of functions*, Math. Ann. 111 (1935), 98-118.
2. A. W. Goodman and M. S. Robertson, *A class of multivalent functions*, Trans. Amer. Math. Soc. 70 (1951), 127-136.
3. A. W. Goodman, *On some determinants related to p -valent functions*, Trans. Amer. Math. Soc. 63 (1948), 175-192.
4. ———, *On the Schwarz-Christoffel transformation and p -valent functions*, Trans. Amer. Math. Soc. 68 (1950), 204-223.
5. W. Kaplan, *Close-to-convex schlicht functions*, Michigan Math. J. 1 (1952), 169-185 (1953).
6. R. J. Libera, *Radius of convexity problems*, Thesis, Rutgers, The State University, New Brunswick, N. J., 1962.
7. T. H. MacGregor, *Functions whose derivative has a positive real part*, Trans. Amer. Math. Soc. 104 (1962), 532-537.
8. Maxwell O. Reade, *The coefficients of close-to-convex functions*, Duke Math. J. 23 (1956), 459-462.
9. M. S. Robertson, *Multivalently star-like functions*, Duke Math. J. 20 (1953), 539-550.
10. ———, *Extremal problems for analytic functions with positive real part and applications*, Trans. Amer. Math. Soc. 106 (1963), 236-253.
11. W. C. Royster, *Meromorphic starlike multivalent functions*, Trans. Amer. Math. Soc. 107 (1963), 300-308.
12. T. Umezawa, *On the theory of univalent functions*, Tôhoku Math. J. 7 (1955), 212-223.
13. ———, *Multivalently close-to-convex functions*, Proc. Amer. Math. Soc. 8 (1957), 869-874.

LAFAYETTE COLLEGE,
EASTON, PENNSYLVANIA