

CONVOLUTION MEASURE ALGEBRAS WITH GROUP MAXIMAL IDEAL SPACES⁽¹⁾

BY
JOSEPH L. TAYLOR

Let G denote a locally compact abelian topological group (an l.c.a. group) with dual group G^\wedge . We will denote by $M(G)$ the Banach algebra of bounded regular Borel measures on G under convolution multiplication and by $L(G)$ the algebra of bounded measures absolutely continuous with respect to Haar measure on G (for discussions of these Banach algebras cf. [1], [2], and [5]).

In this paper we shall be concerned with closed subalgebras \mathfrak{M} of $M(G)$ with the following two properties:

- (1) if $\mu \in \mathfrak{M}$ and ν is absolutely continuous with respect to μ , then $\nu \in \mathfrak{M}$;
- (2) the maximal ideal space of \mathfrak{M} is G^\wedge , where the Gelfand transform μ^\wedge of $\mu \in \mathfrak{M}$ is given by $\mu^\wedge(\chi) = \int \chi d\mu$ for $\chi \in G^\wedge$; i.e., the Gelfand transform coincides with the Fourier-Stieltjes transform on \mathfrak{M} .

Any closed subalgebra of $M(G)$ satisfying (1) will be called an L -subalgebra of $M(G)$. It is well known that $L(G)$ satisfies (1) and (2) (cf. [5, Chapter 1]). In Theorem 1 we show that any L -subalgebra \mathfrak{M} of $M(G)$, with $L(G) \subset \mathfrak{M} \subset (L(G))^{1/2}$, also satisfies (2), where $(L(G))^{1/2}$ is the intersection of all maximal ideals of $M(G)$ containing $L(G)$. We conjecture that the converse is also true; i.e., if \mathfrak{M} satisfies (1) and (2) then $L(G) \subset \mathfrak{M} \subset (L(G))^{1/2}$. In Theorem 2 we prove that this is true provided G contains no copy of R^n for $n > 1$.

The problem arises in the following way: in [6] we define the concept of abstract convolution measure algebra and prove that any such algebra \mathfrak{M} , provided it is commutative and semisimple, may be represented as an L -subalgebra of $M(S)$, where S is a compact topological semigroup called the structure semigroup of \mathfrak{M} . $M(G)$ and $L(G)$ are convolution measure algebras as is any L -subalgebra of the measure algebra on a semigroup. The map $\mu \rightarrow \mu_S$ which embeds \mathfrak{M} in $M(S)$ is an isometry as well as an algebraic isomorphism and it preserves the order theoretic properties of \mathfrak{M} as a space of measures. The maximal ideal space of \mathfrak{M} may be identified as S^\wedge , the set of all semicharacters of S , where the Gelfand transform of $\mu \in \mathfrak{M}$ is given by $\mu^\wedge(f) = \int f d\mu_S$ for $f \in S^\wedge$ (a semicharacter of S is a continuous homomorphism of S into the unit disc which is not identically zero). Under pointwise multiplication S^\wedge is a semigroup provided \mathfrak{M} has an approximate identity.

Received by the editors March 14, 1966.

⁽¹⁾ Portions of these results were presented to the meeting of the Society held January 24–29, 1966, under the same title. This research was supported in part by the United States Army Research Office (Durham).

Now, given this representation of \mathfrak{M} , it is natural to try to relate the algebraic structures of S and S^\wedge to the structure of \mathfrak{M} . In particular, if S and S^\wedge are groups, what can be said about \mathfrak{M} ? More generally, if G_1 is a maximal group contained in S , then its dual group G_1^\wedge is embedded in S^\wedge (cf. [6, §3]) and, if G_1 is not a set of measure zero for \mathfrak{M} , then G_1^\wedge is locally compact in the Gelfand topology of S^\wedge and, with this topology, is the maximal ideal space of \mathfrak{M}_1 , the algebra of measures in \mathfrak{M} which are concentrated on G_1 . It follows that G_1^\wedge , with the Gelfand topology (as opposed to the discrete topology it inherits from the compact group G_1), is the dual group G^\wedge of some l.c.a. group G (with Bohr compactification G_1) and that \mathfrak{M}_1 may be embedded as an L -subalgebra of $M(G)$. Thus \mathfrak{M}_1 is a subalgebra of $M(G)$ satisfying (1) and (2); i.e., by studying the relationship between maximal groups of S and the structure of \mathfrak{M} we are led to the problem posed in this paper.

Our problem may also be viewed as an attempt to extend results of Rieffel in [4]. Rieffel proves that if A is a commutative semisimple Banach algebra which is Tauberian and in which every multiplicative linear functional is L' -inducing, then A is $L(G)$ for some l.c.a. group G . The hypothesis that every multiplicative linear functional is L' -inducing is equivalent to the existence of an l.c.a. group G such that A is a closed subalgebra of $M(G)$ satisfying (1) and (2). The Tauberian hypothesis is extremely strong; it means that those elements of A whose transforms have compact support on G^\wedge are dense in A . If our conjecture is true, then without Rieffel's Tauberian hypothesis we may still conclude that $L(G) \subset A \subset (L(G))^{1/2}$. In Theorem 5.1 of [4] Rieffel mentions an example which shows that $L(G)$ and $(L(G))^{1/2}$ may be distinct.

We now proceed with our development of Theorems 1 and 2. If B is any commutative Banach algebra and I is a closed ideal of B , then $I^{1/2}$ denotes the intersection of all maximal ideals of B which contain I . Under the natural map $x \rightarrow \xi_x$ from B to B/I , $I^{1/2}$ is the inverse image of $(0)^{1/2}$; but $\xi_x \in (0)^{1/2}$ if and only if $\lim_n \|\xi_x^n\|^{1/n} = 0$ (cf. [2, 24B]). Hence, $x \in I^{1/2}$ if and only if $\lim_n \inf_{y \in I} \|x^n + y\|^{1/n} = 0$. In Lemma 1 we use this fact to obtain a measure theoretic characterization of the radical of an L -ideal, where an L -ideal \mathfrak{N} of a convolution measure algebra \mathfrak{M} is an ideal such that whenever $\mu \in \mathfrak{N}$ and $\nu \in \mathfrak{M}$ with ν absolutely continuous with respect to μ , then $\nu \in \mathfrak{N}$.

If two measures μ and ν are mutually singular, then we write $\mu \perp \nu$. If \mathfrak{N}_1 and \mathfrak{N}_2 are sets of measures, then $\mathfrak{N}_1 \perp \mathfrak{N}_2$ means $\mu \perp \nu$ for every $\mu \in \mathfrak{N}_1$ and $\nu \in \mathfrak{N}_2$. By $|\mu|$ we mean the measure defined by the total variation of μ .

LEMMA 1. *If \mathfrak{N} is an L -ideal in a convolution measure algebra \mathfrak{M} , then $\mathfrak{N}^{1/2}$ is also an L -ideal and $\mu \in \mathfrak{N}^{1/2}$ if and only if $\mu \perp \{\nu \in \mathfrak{M} : |\nu|^n \perp \mathfrak{N} \text{ for } n = 1, 2, \dots\}$.*

Proof. Let S be the structure semigroup of \mathfrak{M} as in [6] and $\mu \rightarrow \mu_S$ the embedding of \mathfrak{M} into $M(S)$. Let A be the smallest closed subset of S on which each μ_S for $\mu \in \mathfrak{N}$ is concentrated. A maximal regular ideal of \mathfrak{M} containing \mathfrak{N} corresponds to a semicharacter $f \in S^\wedge$ such that $\int f d\mu_S = 0$ for each $\mu \in \mathfrak{N}$. Since \mathfrak{N} is an L -ideal it

follows that such an f must be identically zero on A . Let J be the set of all such semicharacters f together with the identically zero function on S ; then J is closed under conjugation and pointwise multiplication. Let $A' = \{s \in S : f(s) = 0 \text{ for all } f \in J\}$. Since S^\wedge separates points in S (cf. [6, Theorem 2.2]), J separates points in $S \setminus A'$ and, thus, the Stone-Weierstrass theorem implies that the closed linear span of J in $C(S)$ consists of those continuous functions which vanish on A' . Now $\mathfrak{N}^{1/2}$ consists of those measures μ in \mathfrak{M} for which $\int f d\mu_S = 0$ for all $f \in J$; i.e., those measures $\mu \in \mathfrak{M}$ for which μ_S is concentrated on A' . It follows that $\mathfrak{N}^{1/2}$ is an L -ideal.

Since $\mathfrak{N}^{1/2}$ is an L -ideal, $\mu \in \mathfrak{N}^{1/2}$ if and only if $|\mu| \in \mathfrak{N}^{1/2}$. Therefore, it suffices to prove the second part of Lemma 1 for positive μ . Let $\mu \in \mathfrak{M}$, $\mu \geq 0$, and $\nu \in \mathfrak{M}$ such that $|\nu|^n \perp \mathfrak{N}$ for $n = 1, 2, \dots$; then if μ and ν are not mutually singular, we may write $\mu = \mu_1 + \mu_2$ where $\mu_1, \mu_2 \geq 0$ and $0 \neq \mu_2 \leq |\nu|$. Then $\|\omega + \mu^n\| \geq \|\mu_2\|^n$ for every $\omega \in \mathfrak{M}$, since $\mu_2^n \leq |\nu|^n \perp \mathfrak{N}$. Hence $\lim_n \inf_{\omega \in \mathfrak{M}} \|\omega + \mu^n\|^{1/n} \geq \|\mu_2\| \neq 0$, and $\mu \notin \mathfrak{N}^{1/2}$. Thus if $\mu \in \mathfrak{N}^{1/2}$ then $\mu \perp \nu$ whenever $\nu \in \mathfrak{M}$ and $|\nu|^n \perp \mathfrak{N}$ for $n = 1, 2, \dots$

To see the converse, note that if $\mu \notin \mathfrak{N}^{1/2}$ then there is a semicharacter f in the set J described above for which $\int f d\mu_S \neq 0$. We may write $\mu = \omega + \nu$ where ω_S is concentrated on the set where f is zero and ν_S is concentrated on the set where f is nonzero. Since f is multiplicative, ν_S^n is also concentrated on the set where f is nonzero and, hence, $|\nu|^n \perp \mathfrak{N}$ for each n . This completes the proof, since μ and ν are not mutually singular.

THEOREM 1. *If \mathfrak{M} is an L -subalgebra of $M(G)$ and $L(G) \subset \mathfrak{M} \subset (L(G))^{1/2}$ then the maximal ideal space of \mathfrak{M} is G^\wedge ; i.e., \mathfrak{M} satisfies (2).*

Proof. It follows from the characterization of the radical in Lemma 1, that if \mathfrak{M}_2 is an L -subalgebra of a convolution measure algebra \mathfrak{M}_1 and if \mathfrak{N} is an L -ideal of both \mathfrak{M}_1 and \mathfrak{M}_2 , then the radical of \mathfrak{N} in \mathfrak{M}_2 is the intersection with \mathfrak{M}_2 of the radical of \mathfrak{N} in \mathfrak{M}_1 . Thus, if $L(G) \subset \mathfrak{M} \subset (L(G))^{1/2}$, then the radical of $L(G)$ in \mathfrak{M} is \mathfrak{N} . It follows that $L(G)$ and \mathfrak{M} have the same maximal ideal space, namely G^\wedge .

Before considering the converse problem we require a lemma concerning closed semigroups in l.c.a. groups.

LEMMA 2. *If T is a closed proper subsemigroup of the l.c.a. group G which generates G (i.e., the closed subgroup generated by T is G), then there is a continuous homomorphism γ of G into the additive real numbers R such that $0 \neq \gamma(T) \subset R^+$, the set of nonnegative reals.*

Proof. If T has an interior point x such that $-x \notin T$, then the lemma follows from results of Rieffel (cf. [4, Theorem 6.4]). We shall prove the lemma directly in the case where $G = K \times R^n$ for a compact group K , and use this to reduce the general case to Rieffel's result.

We prove the lemma for G of the form $K \times R^n$ by induction on n . If $n = 0$ then G is compact and the lemma holds vacuously, since a closed subsemigroup of a

compact group must be a group (cf. [7, p. 99]) and, thus, T generates G would imply $T=G$. We now assume the lemma holds for $n=p-1$ and let $G=K \times R^p$ for some compact group K . Since T is not a group, there is some $x \in T$ for which $-x \notin T$. Since K is compact, $T \cap K$ is a group; hence, $x \notin K$. Note that without loss of generality we may assume that $0 \notin T$ for, if $0 \in T$, we may replace T by $T'=x+T$; then $0 \notin T'$ and if γ is a homomorphism of G into R such that $\gamma(T') \subset R^+$, then $\gamma(T) \subset R^+ - \gamma(x)$; but $\gamma(T)$ is a semigroup, so $\gamma(T) \subset R^+$.

In view of the above considerations, we may, without loss of generality, write G as $H \times R$, where $H=K \times R^{p-1}$, x is the element $(0, 1) \in H \times R$, and $(0, n) \in T$ for positive integers n but not for nonpositive integers n .

Let $G^+ = \{(h, r) \in H \times R : r \geq 0\}$ and $G^- = \{(h, r) \in H \times R : r \leq 0\}$. If $T \subset G^+$ then the projection of $H \times R$ on R is the required γ . If $T \not\subset G^+$ we set $T^- = T \cap G^-$, $Z = \{(0, n) : n = 0, \pm 1, \pm 2, \dots\}$, and let ϕ be the natural map from G to G/Z . Note that $(0, 1) \in T$ implies that $\phi(T^-) = \phi(T_0^-)$, where

$$T_0^- = \{(h, r) \in H \times R : -1 \leq r \leq 0\} \cap T^-,$$

and since $\phi(T_0^-)$ is closed, so is $\phi(T^-)$. Also, $Z \cap T^-$ is empty so $0 \notin \phi(T^-)$ and $\phi(T^-)$ is a closed proper subsemigroup of G/Z . To apply the induction hypothesis we need only show that $\phi(T^-)$ generates G/Z . If not, then $\phi(T^-) \subset J$ where J is a proper closed subgroup of G ; if $y \in T$ and $w \in T^-$, then $y + w^n \in T^- \subset \phi^{-1}(J)$ for sufficiently large n , and, hence, $y \in \phi^{-1}(J)$; i.e., $T \subset \phi^{-1}(J)$ which contradicts the assumption that T generates G . Note that if $p=1$ then T must be contained in G^+ , since G/Z is compact. Thus the lemma is proven in this case.

Now if $p > 1$ the noncompact part of G/Z has dimension $p-1 \neq 0$, and we may apply the induction hypothesis and conclude that there is a continuous homomorphism f of G/Z into R such that $0 \neq f(\phi(T^-)) \subset R^+$. We now define a map g of G into $R \times R$ by $g(h, r) = (f(\phi(h, r)), r)$; then g is a continuous homomorphism and $g(T) \cap \{(t, r) : t < 0, r < 0\}$ is empty; i.e., $g(T)$ misses the open third quadrant. From this and the fact that $g(T)$ is a semigroup, it follows readily that the convex hull of $g(T)$ also misses the third quadrant and may thus be separated from it by a straight line l passing through the origin. The line l may contain $g(T)$, in which case we have reduced the problem to the one-dimensional case and may apply the last comment of the previous paragraph. If $g(T) \not\subset l$, then there is a continuous homomorphism α of $R \times R$ onto R , with kernel l , such that $0 \neq \alpha(g(T)) \subset R^+$. Then α composed with g is the required map γ . This completes the induction.

To prove the lemma for G a general l.c.a. group, we use the structure theorem for l.c.a. groups (cf. [5, Theorem 2.4.1]) which says that G contains an open subgroup G_1 of the form $K \times R^n$ for a compact group K . Let β be the natural map from G to G/G_1 . If $\beta(T)$ is proper in G/G_1 , we may apply the above mentioned result of Rieffel to obtain γ , since G/G_1 is discrete. If $\beta(T) = G/G_1$ then every coset of G_1 in G contains an element of T . Then $T \cap G_1$ must be a proper generating subsemigroup of G_1 if T is to be a proper generating subsemigroup of G . Thus, we

may apply the result of the previous paragraph to obtain a continuous homomorphism γ_1 of G_1 into R such that $0 \neq \gamma(T \cap G_1) \subset R^+$. We write

$$G_1^+ = \{x \in G_1 : \gamma_1(x) \geq 0\}.$$

Then $G_1^+ + T$ is proper in G , for if $y \in G_1$ and $y = x + t$ with $x \in G_1^+$ and $t \in T$, then $y - x \in G_1 \cap T$ and $\gamma_1(y - x) = \gamma_1(y) - \gamma_1(x) \geq 0$ so that $y \in G_1^+$ also. Clearly, $G_1^+ + T$ is closed, generates G , and has an interior point x such that $-x \notin G_1^+ + T$ (e.g., any $x \in G_1$ for which $\gamma_1(x) > 0$). We again apply Rieffel's result and the proof is complete.

Note that Lemma 2 implies that on a semigroup T of the above type there is a proper semicharacter, i.e., a semicharacter f for which $|f| \neq 1$; in fact, $f(x) = e^{-s\gamma(x)}$ for $s > 0$ and $x \in T$ is such a semicharacter.

In what follows, \mathfrak{M} will denote a closed subalgebra of $M(G)$ which satisfies (1) and (2). Lemmas 3 and 4 hold for arbitrary l.c.a. groups G , while the remainder of the proof of Theorem 2 requires special assumptions on G .

LEMMA 3. *\mathfrak{M} is weak-* dense in $M(G)$ and \mathfrak{M}^\wedge is uniformly dense in $C_0(G^\wedge)$.*

Proof. Let $T = \text{carrier } (\mathfrak{M})$ be the smallest closed subset of G on which each $\mu \in \mathfrak{M}$ is concentrated. Since \mathfrak{M} is an L -subalgebra of $M(G)$, T is a closed subsemigroup of G , and \mathfrak{M} is weak-* dense in $M(G)$ if and only if $T = G$. Since \mathfrak{M} separates points in G^\wedge , T cannot be contained in any closed proper subgroup of G ; i.e., T generates G . If $T \neq G$ we may apply Lemma 2 and obtain a continuous homomorphism γ of G into R such that $0 \neq \gamma(T) \subset R^+$. Then $\mu \rightarrow \int e^{-\gamma(x)} d\mu(x)$ is a complex homomorphism of \mathfrak{M} which clearly does not correspond to any character of G . This contradicts the fact that \mathfrak{M} satisfies (2). We conclude that $T = G$ and \mathfrak{M} is weak-* dense in $M(G)$.

Let $\lambda \in M(G^\wedge)$; then $\lambda^\wedge(x) = \int \bar{\chi}(x) d\lambda(x)$ is continuous on G , so there exists $\mu \in \mathfrak{M}$ for which $\int \lambda^\wedge d\mu = \int \mu^\wedge d\lambda \neq 0$. Hence \mathfrak{M}^\wedge is a dense subspace of $C_0(G^\wedge)$. This completes the proof.

LEMMA 4. *If $\mathfrak{M} \cap L(G) \neq (0)$ then $L(G) \subset \mathfrak{M} \subset (L(G))^{1/2}$.*

Proof. $\mathfrak{M} \cap L(G)$ is an L -subalgebra of $L(G)$ and an L -ideal of \mathfrak{M} . If $\nu \in L(G)$ then Lemma 3 implies that ν is the weak-* limit in $M(G)$ of a net $\{\mu_\alpha\}$ of elements of \mathfrak{M} . If $f \in L_\omega(G)$ and $\omega \in \mathfrak{M} \cap L(G)$, then $g(x) = \int f(x+y) d\omega(y) \in C(G)$ and, hence, $\int f d\omega \cdot \mu_\alpha$ converges to $\int f d\nu = \int f d\omega \cdot \nu$; i.e., $\{\omega \cdot \mu_\alpha\}$ is a net in $\mathfrak{M} \cap L(G)$ which converges weakly to $\omega \cdot \nu$ and so $\omega \cdot \nu \in \mathfrak{M} \cap L(G)$. It follows that $\mathfrak{M} \cap L(G)$ is an L -ideal of $L(G)$. However, the only L -ideals of $L(G)$ are (0) and $L(G)$. Hence $L(G) \subset \mathfrak{M}$. Also $\mathfrak{M} \subset (L(G))^{1/2}$, otherwise there would be a complex homomorphism of $M(G)$ which was zero on $L(G)$ and nonzero on \mathfrak{M} , contradicting the fact that the maximal ideal space of \mathfrak{M} is G^\wedge .

LEMMA 5. *If G contains no copy of R , then $L(G) \subset \mathfrak{M} \subset (L(G))^{1/2}$.*

Proof. Under this hypothesis, G and G^\wedge each contain an open-compact subgroup, by the principal structure theorem (cf. [5, Theorem 2.4.1]). Let K be an open-compact subgroup of G^\wedge . Since G^\wedge is the maximal ideal space of \mathfrak{M} , Shilov's theorem (cf. [3, Theorem 3.6.3]) implies the existence of $\mu \in \mathfrak{M}$ such that $\mu^\wedge(\chi) = 0$ for $\chi \notin K$ and $\mu^\wedge(\chi) = 1$ for $\chi \in K$. Then μ is Haar measure on an open-compact subgroup H of G ($H = \{x \in G : \chi(x) = 1 \text{ for } x \in K\}$). Thus $\mu \in L(G)$ and the conclusion follows from Lemma 4.

LEMMA 6. *If G is of the form $K \times R$ where K is compact, and*

$$G^+ = \{(k, r) \in K \times R : r > 0\},$$

then there exists a nonnegative measure $\mu \in \mathfrak{M}$ concentrated on G^+ with compact support and $\|\mu\| > 1$ such that $\mu^\wedge(\chi) \neq 1$ for each $\chi \in G^\wedge$.

Proof. It follows from Lemma 3 that it is enough to show the existence of a $\nu \in L(G)$ with the above properties. An example of such a measure ν is $\nu = 2(\rho \times \omega)$, where ρ is Haar measure on K and ω is Lebesgue measure on $[0, 1]$ in R .

THEOREM 2. *If G contains no copy of R^n for $n > 1$, then $L(G) \subset \mathfrak{M} \subset (L(G))^{1/2}$.*

Proof. If G contains no copy of R , then Lemma 5 gives us the result; otherwise, the hypothesis, together with the structure theorem for l.c.a. groups, implies that G has an open subgroup G_1 of the form $K \times R$ with K compact. Let \mathfrak{M}_1 be the subalgebra of \mathfrak{M} consisting of all measures in \mathfrak{M} which are concentrated on G_1 . It is easily seen that \mathfrak{M}_1 is an L -subalgebra of $M(G_1)$ with maximal ideal space G_1^\wedge . Thus the preceding lemmas hold for \mathfrak{M}_1 as a subalgebra of $M(G_1)$.

Now let μ be the element of \mathfrak{M}_1 given by Lemma 6. Since μ^\wedge never assumes the value one, μ has an adverse (cf. [2, §21]); i.e., there is an element $\nu \in \mathfrak{M}_1$ such that $\mu\nu = \mu + \nu$. If μ had norm less than one, its adverse would be $\omega = -\sum_{n=1}^\infty \mu^n$. Since μ is concentrated on G_1^+ , the series $\sum \mu^n$ converges on each compact subset of G_1 , but since $\|\mu\| > 1$ and $\mu \geq 0$, it converges to an unbounded measure.

The Laplace-Stieltjes transform, $\bar{\rho}$, for a measure ρ on $G_1 = K \times R$ is defined by $\bar{\rho}(\chi, z) = \int_{K \times R} e^{-zt} \bar{\chi}(k) d\rho(k, t)$, for $\chi \in K^\wedge$ and z a complex number, whenever this integral converges. Note that since μ has compact support, $\bar{\mu}$ exists and is analytic in z for all χ , and $\mu^\wedge(\chi, y) = \bar{\mu}(\chi, iy)$, for y real, is the Fourier-Stieltjes transform of μ . Since ν is a bounded measure, $\bar{\nu}$ exists for z imaginary and

$$\bar{\nu}(\chi, iy) = \nu^\wedge(\chi, y) = \mu^\wedge(\chi, y)[\mu^\wedge(\chi, y) - 1]^{-1}.$$

Also, for sufficiently large positive r , $\int e^{-rt} d\mu(k, t) < 1$ and $\int e^{-rt} d\omega(k, t)$ is finite. Thus, for $x \geq r$, $\bar{\omega}(\chi, x + iy)$ exists and equals $\bar{\mu}(\chi, x + iy)[\bar{\mu}(\chi, x + iy) - 1]^{-1}$.

The function $f(\chi, z) = \bar{\mu}(\chi, z)[\bar{\mu}(\chi, z) - 1]^{-1}$ is analytic in z except where $\bar{\mu}(\chi, z) = 1$, and at such points it has simple poles. Also, since $\bar{\mu}(\chi, z)$ approaches zero at infinity for $\text{Re}(z) \geq 0$, there can be at most finitely many poles of f in the region $\text{Re}(z) \geq 0$.

Let $\{\rho_\alpha\}$ be a weak-* approximate identity for $M(G)$ consisting of elements of $L(G_1)$ with compact support and norm one, whose Laplace transforms $\hat{\rho}_\alpha$ have the property that $\hat{\rho}_\alpha(\chi, a + iy)$ is integrable in (χ, y) for each $a \in R$. Let $g_\alpha(k, t)$ be the Radon-Nikodym derivative of ρ_α for each α ; then, the inversion formula (cf. [5], Theorem 1.5.1) implies that

$$g_\alpha * \nu(k, t) = \frac{1}{2\pi i} \int_{-i\infty}^{i\infty} \left[\sum_{\chi \in K^\wedge} e^{zt} \chi(k) f(\chi, z) \hat{\rho}_\alpha(\chi, z) \right] dz$$

and

$$g_\alpha * \omega(k, t) = \frac{1}{2\pi i} \int_{x-i\infty}^{x+i\infty} \left[\sum_{\chi \in K^\wedge} e^{zt} \chi(k) f(\chi, z) \hat{\rho}_\alpha(\chi, z) \right] dz,$$

for $x \geq r$. It follows that

$$g_\alpha * (\omega - \nu)(k, t) = \frac{1}{2\pi i} \sum_{\chi \in P} \int_{\Gamma} e^{zt} \chi(k) f(\chi, z) \hat{\rho}_\alpha(\chi, z) dz$$

where P is the finite set of points $\chi \in K^\wedge$ for which $f(\chi, z)$ has a pole in the region $\text{Re}(z) \geq 0$, and Γ is any simple closed curve in $\text{Re}(z) \geq 0$ enclosing all such poles. Then $g_\alpha * (\omega - \nu)$ converges uniformly on compact subsets of G_1 to the function

$$\phi(k, t) = \frac{1}{2\pi i} \sum_{\chi \in P} \int_{\Gamma} e^{zt} \chi(k) f(\chi, z) dz,$$

while $\rho_\alpha \cdot (\omega - \nu)$ converges weakly, relative to continuous functions with compact support, to the measure $\omega - \nu$. It follows that $\omega - \nu$ is absolutely continuous on each compact set with Radon-Nikodym derivative ϕ . Thus, if λ is the restriction of $\omega - \nu$ to any compact subset of G_1 , we have $\lambda \in L(G_1) \cap \mathfrak{M}_1 \subset L(G) \cap \mathfrak{M}$. In view of Lemma 4, the proof is complete.

Added in proof. We have recently obtained a proof of Theorem 2 without the condition on G . Thus, the conjecture mentioned at the beginning of this paper is true.

BIBLIOGRAPHY

1. E. Hewitt and K. A. Ross, *Abstract harmonic analysis*, Springer-Verlag, Berlin, 1963.
2. L. H. Loomis, *An introduction to abstract harmonic analysis*, Van Nostrand, New York, 1953.
3. C. E. Rickart, *General theory of Banach algebras*, Van Nostrand, Princeton, N. J., 1960.
4. M. A. Rieffel, *A characterization of commutative group algebras and measure algebras*, Trans. Amer. Math. Soc. **116** (1965), 32-65.
5. W. Rudin, *Fourier analysis on groups*, Interscience, New York, 1962.
6. J. L. Taylor, *The structure of convolution measure algebras*, Trans. Amer. Math. Soc. **119** (1965), 150-166.
7. A. D. Wallace, *The structure of topological semigroups*, Bull. Amer. Math. Soc. **61** (1955), 95-112.

UNIVERSITY OF UTAH,
SALT LAKE CITY, UTAH