

A-TRANSFORMS AND HILBERT FUNCTIONS IN LOCAL LATTICES

BY
E. W. JOHNSON

0. Introduction. In [1], Dilworth introduced the notion of a principal element of a multiplicative lattice and used it to define a class of lattices in which the Noether decomposition theorems hold. He called these Noether lattices and showed, among other things, that Noether lattices satisfy the Krull Intersection Theorem and the Krull Principal Ideal Theorem.

Noether lattices in general, however, are lacking in many important ring-like properties. For example, in a local Noether lattice of altitude d , there need not exist d principal elements with join an element which is primary for the maximal element, so that systems of parameters may be much larger than one would hope.

In this paper it is shown that if P is the maximal element of a local Noether lattice, then for large n the number of principal elements required to generate P^n is given by a polynomial $B^*(P, x)$ (Corollary 3.3). It seems reasonable then to consider the degree of this polynomial as a second measure of the number of elements in a system of parameters, or possibly as an alternative definition of the dimension of the lattice. Lemma 3.6 and Theorem 3.9 establish that the degree of $B^*(P, x)$ is $d-1$ where d is the height of P .

§2 is concerned with the introduction of a graded multiplicative lattice, denoted $\mathcal{R}(\mathcal{L}, A)$. This lattice is analogous to the A -transform of a Noetherian ring introduced by D. Rees and serves as the vehicle in §3 for obtaining the polynomial $B^*(P, x)$. The main result of this section is that, if \mathcal{L} is a Noether lattice, then $\mathcal{R}(\mathcal{L}, A)$ is a Noether lattice (Theorem 2.11). A determination is also made of the heights of the maximal primes of $\mathcal{R}(\mathcal{L}, A)$ in terms of A and \mathcal{L} (Lemma 2.16).

In §4 regular local lattices are considered. It is shown that the 0-element of a regular local lattice is prime (Theorem 4.1) and the results of §2 are applied to show that the degree function is a valuation (Corollary 4.4).

The author expresses his gratitude to Professor Louis J. Ratliff for many helpful suggestions.

Throughout the remainder of this paper, the symbol \mathcal{L} will denote a Noether lattice.

1. The following lemma is a useful generalization of Lemma (6.3) in [1].

LEMMA 1.1. *Let A, B and C be elements of \mathcal{L} with A principal, $B \leq C$ and $(0 : A) \wedge C \leq B$. Then the correspondence $D \rightarrow AD$ of the quotient $C|B$ into the quotient $CA|BA$ is a lattice isomorphism.*

Received by the editors February 8, 1967 and, in revised form, December 8, 1967.

Proof. Clearly the correspondence is order preserving. Let D and E be elements of C/B . Then since A is principal and $(0 : A) \wedge C \leq B$, $D = D \vee ((0 : A) \wedge C) = C \wedge (D \vee (0 : A)) = C \wedge (DA : A)$, and similarly, $E = C \wedge (EA : A)$. Hence if $DA \leq EA$, then $D \leq E$, and if $DA = EA$, then $D = E$. Now, let F be any element of CA/BA . Then $(F : A) \wedge C$ is an element of C/B and $((F : A) \wedge C)A = F \wedge CA = F$, since A is principal. Hence $C/B \cong CA/BA$. Q.E.D.

If B, C are elements of \mathcal{L} with $B < C$, then C will be called a cover for B if $B \leq X < C$ implies $B = X$. The following theorem characterizes the covering relation in \mathcal{L} .

THEOREM 1.2. *Let B and C be elements of \mathcal{L} with $B < C$. Then C is a cover for B if, and only if, there exists a principal element $A \in \mathcal{L}$ with $B \vee A = C$ such that $B : A$ is maximal in \mathcal{L} .*

Proof. If A is principal in \mathcal{L} , then by Lemma 1.1, $(B \vee A)/B \cong AI/(B \wedge A) = AI/(B : A)A \cong I/(B : A)$. The theorem follows immediately from this. Q.E.D.

In [1] it was shown that Noether lattices satisfy an abstract "Krull Intersection Theorem." An immediate but useful consequence is stated for ease of reference in the following

LEMMA 1.3. *If \mathcal{L} is a local lattice and if B, C , and D are elements of \mathcal{L} with $D < I$, then $B \leq C \vee BD$ implies $B \leq C$.*

Proof. $B \leq C \vee BD$ implies $B \leq C \vee BD^n$ for all n , so that necessarily $B \leq C$. Q.E.D.

If an element $B \in \mathcal{L}$ is the join of principal elements A_1, \dots, A_k then the elements A_i will be called a *base* or *basis* for B . If this is the case and if no proper subcollection of the A_i also form a basis for B , then the A_i will be said to be a *minimal* base for B . As for rings, it is clear that any base can be reduced to a minimal base.

THEOREM 1.4. *Let \mathcal{L} be a local lattice with maximal element P and let $A \leq P$ be an element of \mathcal{L} . Then*

1. *the quotient A/AP is finite dimensional and*
2. *the number of elements in any minimal base for A is equal to the dimension of A/AP .*

Proof. Let A_1, \dots, A_r be a minimal base for A . Set $S_0 = AP$ and for $1 \leq i \leq r$, let $S_i = S_{i-1} \vee A_i$. Then $PA_i \leq PA \leq S_{i-1}$, and hence $S_{i-1} : A_i \geq P$ ($1 \leq i \leq r$). Since P is maximal, it follows that either $S_{i-1} : A_i = P$ or $S_{i-1} : A_i = I$ and hence that either $S_i = S_{i-1} \vee A_i$ is a cover for S_{i-1} (Theorem 1.2) or that $S_i = S_{i-1}$. It is immediate then that $AP = S_0 \leq S_1 \leq \dots \leq S_r = A$ can be reduced to a composition chain from AP to A and hence that the quotient A/AP is finite dimensional with dimension (say) $d \leq r$.

Assume now that $d < r$. Then for some i , $1 \leq i \leq r$, $S_i = S_{i-1}$. Hence

$$\begin{aligned} A &= S_i \vee A_{i+1} \vee \cdots \vee A_r \\ &= S_{i+1} \vee A_{i+1} \vee \cdots \vee A_r \\ &= AP \vee A_1 \vee \cdots \vee A_{i-1} \vee A_{i+1} \vee \cdots \vee A_r. \end{aligned}$$

But then by Lemma 1.3, $A = A_1 \vee \cdots \vee A_{i-1} \vee A_{i+1} \vee \cdots \vee A_r$, which contradicts the minimality of the base A_1, \dots, A_r . It follows, therefore, that the number of elements in any minimal base for A is the dimension of the quotient A/AP . Q.E.D.

COROLLARY 1.5. *Let \mathcal{L} be a local lattice and let \mathcal{C} be a collection of principal elements with the property that for every element $A \in \mathcal{L}$ there exist elements A_1, \dots, A_k (depending on A) in \mathcal{C} such that $A = A_1 \vee \cdots \vee A_k$. Then \mathcal{C} is the complete collection of principal elements of \mathcal{L} .*

Proof. If B and A_1, \dots, A_k are principal elements of \mathcal{L} with $B = A_1 \vee \cdots \vee A_k$, then A_1, \dots, A_k can be reduced to a minimal base for B . Since B is itself principal, it follows that $B = A_i$, for some i . Q.E.D.

COROLLARY 1.6. *Let R be a local ring, and let \mathcal{L} be the lattice of the ideals of R . Then the principal elements of \mathcal{L} are the principal ideals of R .*

Proof. It was established in [1] that the principal ideals of R are principal elements of \mathcal{L} , and it follows from Corollary 1.5 that these are the only principal elements. Q.E.D.

If $B \in \mathcal{L}$, then as shown in [1], the quotient I/B becomes a Noether lattice, denoted \mathcal{L}/B , if multiplication is defined by $C \circ D = CD \vee B$. The principal elements of \mathcal{L}/B include the elements $A \vee B$ where A is principal in \mathcal{L} . Since some confusion will naturally result about the multiplication when more than one quotient lattice \mathcal{L}/B is under consideration at the same time, the notation C/B will be used to denote the element $C \vee B$ of \mathcal{L}/B when it is necessary to distinguish. With this notation, if A is principal in \mathcal{L} , then A/B is principal in \mathcal{L}/B .

COROLLARY 1.7. *If an element $B \in \mathcal{L}$ is such that \mathcal{L}/B is local, then the principal elements of \mathcal{L}/B are the elements A/B with A principal in \mathcal{L} .*

Proof. The corollary is immediate from Corollary 1.5. Q.E.D.

If an element P of \mathcal{L} is *prime* (i.e., $AB \leq P$ implies $A \leq P$ or $B \leq P$), then the *height* of P , denoted $\text{ht } P$, is defined as the supremum of all integers n for which there exists a prime chain $P_0 < P_1 < \cdots < P_n = P$ in \mathcal{L} .

LEMMA 1.8. *Let P be a prime element of \mathcal{L} and let A be a principal element with $A \leq P$. Then*

$$\text{ht } P \geq \text{ht } P/A \geq \text{ht } P - 1.$$

If $A \not\leq P'$ for every minimal associated prime $P' \leq P$ of 0 , then $\text{ht } P/A = \text{ht } P - 1$.

Proof. It was shown in [1] that an element C/A is prime or primary in \mathcal{L}/A if, and only if, the element $C \vee A$ is respectively prime or primary in \mathcal{L} . In particular then, the element $P=P/A$ is prime in \mathcal{L}/A . Now, let $d = \text{ht } P$ and let $P_0 < P_1 < \dots < P_d = P$ be a prime chain in \mathcal{L} . Then (by Lemma 6.4 of [1]), there exists a prime chain $P_0^* < P_1^* < \dots < P_d^* = P$ in \mathcal{L} with $P_0 = P_0^*$ and $A \leq P_1^*$. It is immediate from this that $\text{ht } P/A \geq d - 1$. Since $\text{ht } P/A \leq \text{ht } P$ is clear, the first statement is proved. Now, if $\text{ht } P/A = d$, then there is a prime sequence $P'_0 < P'_1 < \dots < P'_d = P$ in \mathcal{L}/A and therefore also in \mathcal{L} , with $A \leq P'_0$. Since P'_0 is necessarily a minimal associated prime of 0 in \mathcal{L} , the second statement follows. Q.E.D.

2. Let \mathcal{L} denote a Noether lattice with least element 0 and greatest element I , and let A be an arbitrary element of \mathcal{L} . In this section a lattice analogue of D. Rees' A -transform of a Noetherian ring R by an ideal A [4] will be introduced. It is this transform, denoted $\mathcal{R}(\mathcal{L}, A)$, which will form the basis for the later proof of the existence of Hilbert Characteristic Polynomials.

Let $\mathcal{F}(A)$ denote the collection of all formal sums $\sum_{i=-\infty}^{\infty} B_i$ of elements of \mathcal{L} such that

$$(2.1) \quad A^i \geq B_i \geq B_{i+1} \geq AB_i, \text{ for all } i, \text{ where for } i \leq 0, A^i = I.$$

The elements of $\mathcal{F}(A)$ will generally be denoted by capitals B, C, \dots , and if $B \in \mathcal{F}(A)$, then B_i will denote the i th coordinate of B . For elements $B, C \in \mathcal{F}(A)$ the following definitions will be used:

$$(2.2) \quad B \leq C \text{ if } B_i \leq C_i \text{ for all } i,$$

$$(2.3) \quad B \vee C = \sum (B_i \vee C_i),$$

$$(2.4) \quad B \wedge C = \sum (B_i \wedge C_i),$$

$$(2.5) \quad B \cdot C = \sum_i (\bigvee_{r+s=i} B_r C_s).$$

It is seen that the collection $\mathcal{F}(A)$ forms a complete, modular, lattice under the relation \leq with resulting join and meet given by (2.3) and (2.4), and that (2.5) defines a commutative, associative multiplication on $\mathcal{F}(A)$ which distributes over the join operation (2.3). The resulting lattice, together with the multiplication (2.5) will be denoted by the symbol $\mathcal{R}(\mathcal{L}, A)$ and will be called the A -transform of \mathcal{L} . It is clear then from the defining relations that $\mathcal{R}(\mathcal{L}, A)$ has greatest element $I^* = \sum A^i$. The element I^* is also a multiplicative identity, since by (2.1) and (2.5) it is seen that if B is any element of $\mathcal{R}(\mathcal{L}, A)$, then the i th coordinate of BI^* is $\bigvee_{r+s=i} B_r I_s^* = \bigvee_{r+s=i} B_r A^s = B_i$. These comments are summarized in the following

LEMMA 2.1. $\mathcal{R}(\mathcal{L}, A)$ is a complete, modular, multiplicative lattice.

DEFINITION 2.2. Let C be an element of \mathcal{L} with $C \leq A^r$. Then $C^{(r)}$ is the least element D of $\mathcal{R}(\mathcal{L}, A)$ such that $C \leq D_r$. The notation $C_A^{(r)}$ will also be used when it is necessary to indicate that $C^{(r)}$ is to be computed in $\mathcal{R}(\mathcal{L}, A)$.

The following lemmas will be useful in computations.

LEMMA 2.3. If $C \leq A^r$ in \mathcal{L} , then $C^{(r)} = \sum CA^{i-r}$. If also $D \leq A^r$, then $C^{(r)} \vee D^{(r)} = (C \vee D)^{(r)}$.

Proof. Clearly $\sum CA^{i-r} \in \mathcal{R}(\mathcal{L}, A)$, so in view of (2.1) it follows that $C^{(r)} = \sum CA^{i-r}$. If also $D \leq A^r$, so that $D^{(r)} = \sum DA^{i-r}$, then by (2.3)

$$C^{(r)} \vee D^{(r)} = \sum (C \vee D)A^{i-r} = (C \vee D)^{(r)}. \quad \text{Q.E.D.}$$

LEMMA 2.4. *If $B \in \mathcal{R}(\mathcal{L}, A)$ and if $C \in \mathcal{L}$ is such that $C \leq A^r$, then $BC^{(r)} = \sum B_{i-r}C$.*

Proof. $BC^{(r)}$ has i th coordinate $\bigvee_{s+t=i} B_sC^{(r)}_t$. Let $s+t=i$. If $s > i-r$ so that $t < r$, then $B_sC^{(r)}_t = B_sC \leq B_{i-r}C$ by (2.1). On the other hand, if $s < i-r$ so that $t > r$, then $B_sC^{(r)}_t = B_sCA^{t-r} \leq B_{s+t-r}C = B_{i-r}C$. If $s = i-r$ so that $t = r$, then $B_sC^{(r)}_t = B_{i-r}C$. Hence

$$B_{i-r}C \leq \bigvee_{s+t=i} B_sC^{(r)}_t \leq B_{i-r}C. \quad \text{Q.E.D.}$$

COROLLARY 2.5. *If $C \leq A^r$ and $D \leq A^s$ in \mathcal{L} , then $C^{(r)}D^{(s)} = (CD)^{(r+s)}$.*

Proof. By Lemmas 2.3 and 2.4, $C^{(r)}D^{(s)}$ has i th coordinate $(C^{(r)})_{i-s}D = CA^{i-s-r}D = CDA^{i-(s+r)}$, which is the i th coordinate of $(CD)^{(s+r)}$. Q.E.D.

LEMMA 2.6. *If $C \in \mathcal{R}(\mathcal{L}, A)$ and if $B \leq A^r$ in \mathcal{L} , then $C : B^{(r)} = \sum_i [(C_{i+r} : B) \wedge A^i]$.*

Proof. Let D be any element of $\mathcal{R}(\mathcal{L}, A)$ such that $DB^{(r)} \leq C$. Then $D_{i-r}B \leq C_i$, for all i ; and hence $D_{i-r} \leq (C_i : B) \wedge A^{i-r}$, for all i . Now, set $E_i = (C_{i+r} : B) \wedge A^i$, for all i . Then $A^i \geq E_i \geq E_{i+1} \geq AE_i$, for all i , so that $E = \sum E_i \in \mathcal{R}(\mathcal{L}, A)$. Since $(EB^{(r)})_i = E_{i-r}B = ((C_i : B) \wedge A^{i-r})B \leq C_i$, for all i , it now follows that $E = C : B^{(r)}$. Q.E.D.

THEOREM 2.7. *Let B be a principal element of \mathcal{L} such that $B \leq A^r$. Then $B^{(r)}$ is a principal element of $\mathcal{R}(\mathcal{L}, A)$.*

Proof. Let $B \leq A^r$ be a principal element of \mathcal{L} and let C and D be any elements of $\mathcal{R}(\mathcal{L}, A)$. Then by Lemmas 2.4 and 2.6,

$$\begin{aligned} ((C \wedge (D : B^{(r)}))B^{(r)})_i &= (C \wedge (D : B^{(r)}))_{i-r}B \\ &= (C_{i-r} \wedge (D_i : B) \wedge A^{i-r})B = (C_{i-r} \wedge (D_i : B))B \\ &= C_{i-r}B \wedge D_i = (CB^{(r)} \wedge D)_i, \end{aligned}$$

for all i . Similarly,

$$\begin{aligned} ((C \vee DB^{(r)}) : B^{(r)})_i &= ((C_{i+r} \wedge D_iB) : B) \wedge A^i = ((C_{i+r} : B) \vee D_i) \wedge A^i \\ &= ((C_{i+r} : B) \wedge A^i) \vee D_i = ((C : B^{(r)}) \vee D)_i, \end{aligned}$$

for all i . Hence $B^{(r)}$ is principal. Q.E.D.

The following theorem will be instrumental in establishing that $\mathcal{R}(\mathcal{L}, A)$ satisfies the ascending chain condition.

THEOREM 2.8. *Let B be any element of $\mathcal{R}(\mathcal{L}, A)$. Then there exists an integer N such that $B_{N+i} = B_NA^i$, for all $i \geq 0$.*

Proof. Let \mathcal{F} be the family of all elements B in $\mathcal{R}(\mathcal{L}, A)$ for which there fails to exist an integer N such that $B_{N+i} = B_NA^i$, for all $i \geq 0$. Assume $\mathcal{F} \neq \emptyset$. Let C be any

chain of elements of \mathcal{F} . Then it is easily seen that $S = \bigvee C \in \mathcal{F}$, for if not, then since \mathcal{L} is Noetherian, there exist integers N and K such that $S_i = S_K$ for $i \leq K$ and $S_i = S_N A^{i-N}$, for $i \geq N$, and for each i , $K \leq i \leq N$, there exists an element $C(i) \in C$ such that $C(i)_i = S_i$. Let M be the maximum of the elements $C(i)$. Then $M \in C$ and $M_i = S_i$ for $K \leq i \leq N$. Furthermore, if $i > N$, then $S_i \geq M_i \geq M_N A^{i-N} = S_N A^{i-N} = S_i$; and if $i < K$, then $S_i \geq M_i \geq M_K = S_K = S_i$. Hence $S_i = M_i$, for all i , so that $S = M \in \mathcal{F}$, which is a contradiction.

Hence if $\mathcal{F} \neq \emptyset$, then there exists a maximal element F of \mathcal{F} . Then $F_1 < A$ since $F_1 = A$ implies $A^n \geq F_n \geq F_1 A^{n-1} = A^n$, for all $n \geq 1$, which contradicts $F \in \mathcal{F}$. Let E be a principal element of \mathcal{L} such that $F_1 < F_1 \vee E \leq A$. Then $F < F \vee E^{(1)}$, so $F \vee E^{(1)} \notin \mathcal{F}$ and there exists an integer N such that $(F \vee E^{(1)})_n A = (F \vee E^{(1)})_{n+1}$, for all $n \geq N$. Hence $(F_n \vee EA^{n-1})A = F_n A \vee EA^n = F_{n+1} \vee EA^n$; and therefore $F_{n+1} \leq F_n A \vee EA^n$, for $n \geq N$. Then since \mathcal{L} is modular and E is principal in \mathcal{L} ,

$$(2.6) \quad \begin{aligned} F_{n+1} &= F_{n+1} \wedge (F_n A \vee EA^n) = F_n A \vee (F_{n+1} \wedge EA^n) \\ &= F_n A \vee (A^n \wedge (F_{n+1} : E))E, \end{aligned}$$

for $n \geq N$. Now, by Lemma 2.6, $A^n \wedge (F_{n+1} : E)$ is the n th coordinate of the element $F : E^{(1)}$ of $\mathcal{R}(\mathcal{L}, A)$; so if $F : E^{(1)} = F$, then $F_n = A^n \wedge (F_{n+1} : E)$, for all n , so that by (2.6), $F_{n+1} = F_n A \vee F_n E = F_n A$, for all $n \geq N$, in contradiction to the assumption that $F \in \mathcal{F}$. Hence $F : E^{(1)} > F$, so $F : E^{(1)} \notin \mathcal{F}$, and there exists an integer $N' \geq N$ such that

$$(2.7) \quad A^n \wedge (F_{n+1} : E) = (A^{n-1} \wedge (F_n : E))A, \quad \text{for all } n \geq N'.$$

But then by (2.6) and (2.7), for $n > N'$, $F_{n+1} = F_n A \vee (A^n \wedge (F_{n+1} : E))E = F_n A \vee (A^{n-1} \wedge (F_n : E))AE = F_n A$, which again contradicts the assumption that $F \in \mathcal{F}$ and shows that $\mathcal{F} = \emptyset$. Q.E.D.

COROLLARY 2.9. *If A and B are elements of \mathcal{L} , then there exists an integer N such that $B \wedge A^{N+n} = (B \wedge A^N)A^n$, for all $n \geq 0$.*

LEMMA 2.10. *If $C \in \mathcal{R}(\mathcal{L}, A)$, then C is the join of finitely many principal elements of the form $B^{(r)}$ where $B \leq A^r$ is principal in \mathcal{L} .*

Proof. Choose integers N and K ($K \leq N$) such that $C_{N+i} = C_N A^i$, for $i \geq 0$, and $C_{K+i} = C_K$, for $i \leq 0$. Then clearly $C = \bigvee_{K \leq i \leq N} (C_i)^{(1)}$. For each i , $K \leq i \leq N$, choose finitely many principal elements $B_{i,j}$ in \mathcal{L} such that $\bigvee_j B_{i,j} = C_i$. Then

$$C = \bigvee_{i,j} (B_{i,j})^{(1)}. \quad \text{Q.E.D.}$$

THEOREM 2.11. *$\mathcal{R}(\mathcal{L}, A)$ is a Noether lattice.*

Proof. By Lemmas 2.1 and 2.10, it is only necessary to show that $\mathcal{R}(\mathcal{L}, A)$ satisfies the ascending chain condition. Hence, let $F(1) \leq F(2) \leq \dots$ be an ascending chain in $\mathcal{R}(\mathcal{L}, A)$. Set $S = \bigvee_j F(j)$ so that, for each i , $S_i = \bigvee_j (F(j))_i$. Choose integers N and K ($K \leq N$) so that $S_{N+i} = S_N A^i$, for $i \geq 0$, and $S_{K+i} = S_K$, for $i \leq 0$. For

each $i, K \leq i \leq N$, choose $F(\emptyset(i))$ so that $(F(\emptyset(i)))_i = S_i$; and let M be the maximum of the elements $F(\emptyset(i))$ thus chosen. Then, as in the proof of Theorem 2.8, $S = M = F(j)$, for some j . Q.E.D.

It is natural to define the *altitude* of a Noether lattice \mathcal{L} to be the supremum of the heights of the prime elements P of \mathcal{L} ($P < I$). As for rings, the altitude of \mathcal{L} will be denoted by $\text{alt } \mathcal{L}$.

The altitude of $\mathcal{R}(\mathcal{L}, A)$ will be determined presently in terms of the altitude of \mathcal{L} and certain properties of A . However, before doing this, it is convenient first to investigate some of the basic properties of $\mathcal{R}(\mathcal{L}, A)$.

If B is any element of \mathcal{L} , then it is clear that for every $i, A^i \geq B \wedge A^i \geq B \wedge A^{i+1} \geq (A(B \wedge A^i))$, so that $\sum (B \wedge A_i)$ is an element of $\mathcal{R}(\mathcal{L}, A)$. This element will be denoted by B^* .

LEMMA 2.12. *The correspondence $B \rightarrow B^*$ of \mathcal{L} into $\mathcal{R}(\mathcal{L}, A)$ is 1-1, isotone and meet-preserving. If $P \in \mathcal{L}$ is prime and $Q \in \mathcal{L}$ is P -Primary, then Q^* is P^* -primary in $\mathcal{R}(\mathcal{L}, A)$.*

Proof. The first statement is immediate from the definition of B^* . Hence assume that P is a prime element of \mathcal{L} and that Q is P -primary. If D and E are any elements of $\mathcal{R}(\mathcal{L}, A)$ with $DE \leq Q^*$ and $D \not\leq Q^*$, then there exists an integer s such that $D_s \not\leq Q \wedge A^s$ and therefore also such that $D_s \not\leq Q$. Now, for every choice of $r, D_s E_r \leq (DE)_{s+r} \leq Q \wedge A^{s+r} \leq Q$, so it follows from the assumption that Q is P -primary in \mathcal{L} that $E_r \leq P$, for all r , so that $E_r \leq P \wedge A^r$, for all r . Hence $E \leq P^*$. Now, choose n so that $P^n \leq Q \leq P$ in \mathcal{L} . It is easily seen that for every r and $s, (B \wedge A^r) \cdot (C \wedge A^s) \leq BC \wedge A^{r+s}$, and it follows from this that $B^* C^* \leq (BC)^*$. Hence, in particular, $(P^*)^n \leq (P^n)^*$ and therefore $(P^*)^n \leq (P^n)^* \leq Q^* \leq P^*$, so that [1] P^* is prime and Q^* is P^* -primary. Q.E.D.

COROLLARY 2.13. *Let $B = Q_1 \wedge \dots \wedge Q_k$ be a normal decomposition of B in \mathcal{L} in which Q_i is P_i -primary. Then $B^* = Q_1^* \wedge \dots \wedge Q_k^*$ is a normal decomposition of B^* in $\mathcal{R}(\mathcal{L}, A)$ in which Q_i^* is P_i^* -primary. Furthermore, if P_1, \dots, P_r are the minimal associated primes of B , then P_1^*, \dots, P_r^* are the minimal associated primes of B^* .*

Proof. This is immediate from Lemma 2.12. Q.E.D.

LEMMA 2.14. *Let P be a prime element of $\mathcal{R}(\mathcal{L}, A)$ with $I^{(-1)} \not\leq P$. Then $P = (P_0)^*$.*

Proof. Assume $I^{(-1)} \not\leq P$. Choose s so that $P_0 \wedge A^{s+i} = (P_0 \wedge A^s)A^i$ for all $i \geq 0$ (Corollary 2.9). Then $(P_0)^* I^{(-s)} = P_0^* (I^{(-1)})^s \leq P$ (Corollary 2.5), so $(P_0)^* \leq P$. Now, if $i \geq 0$, then $P_i \leq P_0 \wedge A^i$; and if $i < 0$, then $(P_i)^{(0)} I^{(i)} = (P_i)^{(0)} (I^{(-1)})^{-i} \leq P$, so that $P_i \leq P_0$. It follows therefore that also $P \leq P_0^*$ and hence that $P = (P_0)^*$. Q.E.D.

LEMMA 2.15. *Let P denote a maximal element of \mathcal{L} . If $P \geq A$, then $P^* \vee I^{(-1)}$ is maximal in $\mathcal{R}(\mathcal{L}, A)$. If $P \not\geq A$, then P^* is maximal in $\mathcal{R}(\mathcal{L}, A)$. Furthermore, if M is any maximal element of $\mathcal{R}(\mathcal{L}, A)$, then M_0 is maximal in \mathcal{L} and either $M = (M_0)^* \vee I^{(-1)}$ or $M = (M_0)^*$, depending on whether $A \leq M_0$ or $A \not\leq M_0$.*

Proof. It is easily seen that if an element M of $\mathcal{R}(\mathcal{L}, A)$ is maximal, then the conditions $M \geq I^{(-1)}$ and $M_0 \geq A$ are equivalent. Note that if $B \in \mathcal{R}(\mathcal{L}, A)$, then $B_0 = I$ if, and only if, $B = I^*$. Now, assume that P is maximal in \mathcal{L} and that $P \not\geq A$. Then $(P^* \vee I^{(-1)})_0 = P \vee A = I$, so that if $P^* \leq T < I^*$, then also $I^{(-1)} \not\leq T$. Furthermore, if $P^* \leq T < I^*$, then $P = (P^*)_0 = T_0$, so that, if T is maximal, then (Lemma 2.14) $T = (T_0)^* = P^*$. Assume now that $P \geq A$. Then $(P^* \vee I^{(-1)})_0 = P$, so $P^* \vee I^{(-1)} < I^*$, and $(P^* \vee I^{(-1)})_i = A_i$ for all $i \neq 0$, so that $P^* \vee I^{(-1)}$ is clearly maximal in $\mathcal{R}(\mathcal{L}, A)$.

Assume now that M is maximal in $\mathcal{R}(\mathcal{L}, A)$. If $M_0 \geq A$, then $M \geq I^{(-1)}$ and $M_i = A^i$ for all $i \neq 0$, so that $M = (M_0)^* \vee I^{(-1)}$. If M_0 were not maximal in \mathcal{L} , say $M_0 < P < I$, then the inequality $M = (M_0)^* \vee I^{(-1)} < P^* \vee I^{(-1)} < I^*$ would hold in $\mathcal{R}(\mathcal{L}, A)$, in contradiction to the maximality of M . If $M_0 \not\geq A$, then $M \not\geq I^{(-1)}$ and $M = M_0^*$, by Lemma 2.14. In this case, M_0 is maximal in \mathcal{L} since $M_0 < T$ would imply $(M_0)^* < T^*$. Q.E.D.

If P is a maximal element of \mathcal{L} , denote by $m(P)$ the associated maximal element of $\mathcal{R}(\mathcal{L}, A)$.

LEMMA 2.16. *Let P be a maximal element of \mathcal{L} . Then $\text{ht } m(P) = \text{ht } P + 1$ if $P \geq A$ and $\text{ht } m(P) = \text{ht } P$ if $P \not\geq A$.*

Proof. Let $P_0 < P_1 < \dots < P_d = P$ be a prime chain in \mathcal{L} ($d = \text{ht } P$). Then in $\mathcal{R}(\mathcal{L}, A)$, $P_0^* < P_1^* < \dots < P_d^* < P^* \vee I^{(-1)} \leq I$, with the equality $P^* \vee I^{(-1)} = I$ holding if, and only if, $P \not\geq A$. It follows that $\text{ht } m(P) \geq \text{ht } P$ and that $\text{ht } m(P) \geq \text{ht } P + 1$ if $P \geq A$.

Now, if $m(P) = P^*$, then it follows from Lemma 2.14 that any prime chain for $m(P)$ is the image under the map $B \rightarrow B^*$ of a prime chain of the same length for P in \mathcal{L} , and hence that $\text{ht } m(P) \leq \text{ht } P$. It can be assumed therefore that $m(P) \neq P^*$, so that (Lemmas 2.14 and 2.15) $I^{(-1)} \leq m(P)$ and $P \geq A$. Now, let A_1, \dots, A_s be principal elements of \mathcal{L} with $A = A_1 \vee \dots \vee A_s$. The inequality $\text{ht } m(P) \leq \text{ht } P + 1$ will now be established by induction on s . Hence assume $s = 1$, and let $P(0) < P(1) < \dots < P(d+i) = m(P)$ be a prime chain in $\mathcal{R}(\mathcal{L}, A)$ with $d = \text{ht } P$ and $d+i = \text{ht } m(P)$. If $A^{(1)} \leq P(0)$, then by Theorem 2.7 above and Lemma 6.4 in [1], it can be assumed that $I^{(-1)} \leq P(1)$ (be rechoosing $P(1)$ if necessary). Then $P(1)_0 < \dots < P(d+i)_0 = P$ is a prime chain in \mathcal{L} ; so, in this case, $d+i-1 \leq d$. If, on the other hand, $A^{(1)} \not\leq P(0)$ it can be assumed that $A^{(1)} \leq P(1)$. Now, if also $I^{(-1)} \leq P(1)$ then $P(0)_0 < P(1)_0 < \dots < P(d+i)_0 = P$ is a prime chain in \mathcal{L} , in contradiction to the assumption that $\text{ht } P = d$. Hence, $I^{(-1)} \not\leq P(1)$, so $d+i \geq 2$; and it can be assumed that $I^{(-1)} \leq P(2)$. Then $P(0)_0 < P(2)_0 < \dots < P(d+i)_0 = P$ is a prime chain in \mathcal{L} , so again $d+i-1 \leq d$. Therefore, if $s = 1$, then $\text{ht } m(P) \leq \text{ht } P + 1$. Assume now that $s > 1$. If $A_j^{(1)} \leq P(0)$, for all j , then, as before (by rechoosing $P(1)$ if necessary), it can be assumed that $I^{(-1)} \leq P(1)$, so that $P(0)_0 < P(2)_0 < \dots < P(d+i)_0 = P$ in \mathcal{L} ; and therefore $d+i-1 \leq d$. Assume, therefore, that $A_1^{(1)} \not\leq P(0)$. Then, as above, it can also be assumed that $A_1^{(1)} \leq P(1)$; and by continuing, it can be assumed that $A_j^{(1)} \leq P(h)$, for all j , and that for $1 \leq j \leq h$, $A_j^{(1)} \leq P(j)$ and $A_j^{(1)} \not\leq P(j-1)$. Also, it can be assumed that $I^{(-1)} \leq P(r)$, $I^{(-1)}$

$\not\leq P(r-1)$, and that $1 \leq r \leq h+1$. Now, set $B = \bigvee_{j \neq 1} A_j$. Then the elements $P'_j = \sum (P(j)_i \wedge B^i)$ of $\mathcal{R}(\mathcal{L}, B)$ are prime ($j=0, \dots, d+i$). To see this, assume C and D are elements of \mathcal{L} with $C \leq B^r$, $D \leq B^s$ and $C_B^{(r)} D_B^{(s)} = (CD)_B^{(r+s)} \leq P'_j$. Then $CD \leq (P'_j)_{r+s} = P(j)_{r+s} \wedge B^{r+s}$, so $(CD)_A^{(r+s)} = C_A^{(r)} D_A^{(s)} \leq P(j)$ and therefore either $C_A^{(r)} \leq P(j)$ or $D_A^{(s)} \leq P(j)$. Assume $C_A^{(r)} \leq P(j)$. Then $C \leq P(j)_r$, and therefore $C \leq P(j)_r \wedge B^r = (P'_j)_r$, so $C_B^{(r)} \leq P'_j$. By Lemma 2.10 it now follows that P'_j is prime. Further, P'_{d+1} is the maximal element $m(P)$ determined in $\mathcal{R}(\mathcal{L}, B)$ by P . It will now be shown that also $P'_0 < P'_1 < \dots < P'_{d+1}$. Since either $P(1) \geq I^{(-1)}$ and $P(0) \not\geq I^{(-1)}$ or else $P(1)$ and $P(0)$ are the images of distinct primes in \mathcal{L} under the map $C \rightarrow C^*$ of \mathcal{L} into $\mathcal{R}(\mathcal{L}, A)$, it follows that $P'_0 \neq P'_1$. Also $P'_j \neq P'_{j+1}$ for $1 \leq j \leq h-1$, since, in this case, $A_{j+1} \leq B$, $A_{j+1}^{(1)} \leq P(j+1)$, and $A_{j+1}^{(1)} \not\leq P(j)$, so that $(P(j))_1 \wedge B \neq (P(j+1))_1 \wedge B$. Now, $P'_h \neq P'_{h+1}$, since either $P(h+1) \geq I^{(-1)}$ and $P(h) \not\geq I^{(-1)}$, or else $P(h+1)_0 \neq P(h)_0$. And $P'_j \neq P'_{j+1}$ for $j > h$, since necessarily $(P(j))_0 \neq (P(j+1))_0$.

Hence, $P'_0 < P'_1 < \dots < P'_{d+i} = m(p)$ is a prime chain in $\mathcal{R}(\mathcal{L}, B)$; so by the inductive hypothesis, $d+i \leq d+1$. Therefore, $\text{ht } m(p) \leq \text{ht } P+1$. Q.E.D.

These results are summarized for local lattices by the following

THEOREM 2.17. *If \mathcal{L} is a local Noether lattice of altitude d , and if $A \in \mathcal{L}$ is contained in the maximal element, then $\mathcal{R}(\mathcal{L}, A)$ is a local Noether lattice of altitude $d+1$.*

3. Hereafter \mathcal{L} will denote a local Noether lattice with maximal element P .

In this section it will be shown that if Q is an element of \mathcal{L} which is P -primary, then there exists a polynomial $D^*(Q, x)$ such that for all n sufficiently large, $D^*(Q, n)$ is the lattice dimension of the quotient I/Q^n . Also, it will be shown that, if A is any element of \mathcal{L} , then there exists a polynomial $B^*(A, x)$ such that for all sufficiently large n , $B^*(A, n)$ is the number of elements in a minimal base for A^n .

LEMMA 3.1. *Let M denote the maximal element of $\mathcal{R}(\mathcal{L}, A)$, and let B be an element of $\mathcal{R}(\mathcal{L}, A)$, such that M_1/B_1 is finite dimensional. Then M_n/B_n is finite dimensional for all $n \geq 1$.*

Proof. Since $M_n = A^n \geq B_n \geq B_1 A^{n-1}$, it suffices to show the finite dimensionality of $A^n/B_1 A^{n-1}$, for $n \geq 2$. Let C_1, \dots, C_K be elements of \mathcal{L} such that $B_1 = C_1 < C_2 < \dots < C_K = A$, and such that for each $i = 1, \dots, K-1$, C_{i+1} is a cover for C_i . Then $A^{n-1} B_1 = A^{n-1} C_1 \leq A^{n-1} C_2 \leq \dots \leq A^{n-1} C_K = A^n$, so it suffices to show that each of the quotients $A^{n-1} C_{i+1} / A^{n-1} C_i$ is finite dimensional. Let i be fixed, $1 \leq i \leq K-1$. Then by Theorem 1.2, there exists a principal element $C \in \mathcal{L}$ with $C_i \vee C = C_{i+1}$ and $C_i : C = P$. Let A_1, \dots, A_r be a basis for A^{n-1} . Then

$$A^{n-1} C_i \leq A^{n-1} C_i \vee A_1 C \leq \dots \leq A^{n-1} C_i \vee A_1 C \vee \dots \vee A_r C = A^{n-1} C_{i+1}$$

and

$$P(A^{n-1} C_i \vee A_1 C \vee \dots \vee A_{j+1} C) \leq (A^{n-1} C_i \vee A_1 C \vee \dots \vee A_j C)$$

for $j = 1, \dots, r-1$, so that (Theorem 1.2) there exists a composition chain for the quotient $A^{n-1} C_{i+1} / A^{n-1} C_i$. Q.E.D.

THEOREM 3.2. *Let M and B be as in Lemma 3.1. Then there exists a polynomial $p(x)$ with rational coefficients such that for all n sufficiently large, $p(n)$ is the lattice dimension $d(M_n/B_n)$ of the quotient M_n/B_n .*

Proof. Let \mathcal{F} be the family of all elements $B \in \mathcal{R}(\mathcal{L}, A)$ for which M_1/B_1 is finite dimensional and for which there exists no polynomial with the desired properties. If $\mathcal{F} \neq \emptyset$, then \mathcal{F} has a maximal element C , since $\mathcal{R}(\mathcal{L}, A)$ is Noetherian. Clearly, $C_1 < M_1$, so there exists a principal element $E \in \mathcal{L}$ with $C < C \vee E^{(1)} \leq M$. Then by the choice of C and E , $C \vee E^{(1)} \notin \mathcal{F}$ and there exists a polynomial $p'(x)$ for $C \vee E^{(1)}$. Set $D = C : E^{(1)}$ so that (Lemma 2.6) $D_n = (C_{n+1} : E) \wedge A^n$, for all n . Then $D \geq C$, and

$$\begin{aligned} d(M_n/(C \vee E^{(1)})_n) &= d(M_n/(C_n \vee EA^{n-1})) \\ &= d(M_n/C_n) - d((C_n \vee EA^{n-1})/C_n) \\ &= d(M_n/C_n) - d(EA^{n-1}/(C_n \wedge EA^{n-1})) \\ &= d(M_n/C_n) - d(A^{n-1}/(C_n : E) \wedge A^{n-1}) \\ &= d(M_n/C_n) - d(M_{n-1}/D_{n-1}), \end{aligned}$$

by Lemma 1.1. Now, if $D > C$, then also $D \notin \mathcal{F}$ and there exists a polynomial $p''(x)$ for D . But then

$$\begin{aligned} d(M_n/C_n) &= d(M_n/(C \vee E^{(1)})_n) + d(M_{n-1}/D_{n-1}) \\ &= p'(n) + p''(n-1) \end{aligned}$$

for all n sufficiently large, which is clearly a contradiction. Hence $D = C$, so $D_n = (C_{n+1} : E) \wedge A^n = C_n$ and $d(M_n/(C \vee E^{(1)})_n) = d(M_n/C_n) - d(M_{n-1}/C_{n-1})$ for all n . Choose N so that $d(M_n/(C \vee E^{(1)})_n) = p'(n)$ for $n \geq N$. Then

$$\sum_{n=N}^w p'(n) = d(M_w/C_w) - d(M_{N-1}/C_{N-1}) \quad \text{for } w > N,$$

so

$$d(M_w/C_w) = \sum_{n=N}^w p'(n) + d(M_{N-1}/C_{N-1}) \quad \text{for } w > N.$$

But this again contradicts the choice of C and hence shows that $\mathcal{F} = \emptyset$. Q.E.D.

Before proceeding further, it is convenient to introduce the following notation:

(3.1) If $C \in \mathcal{L}$, then for $n \geq 1$, $B(C, n)$ is the number of elements in a minimal base for C^n .

(3.2) If Q is a P -primary element of \mathcal{L} , then for $n \geq 0$, $F(Q, n)$ is the dimension of the quotient Q^n/Q^{n+1} .

(3.3) If Q is a P -primary element of \mathcal{L} , then for $n \geq 1$, $D(Q, n)$ is the dimension of the quotient I/Q^n .

COROLLARY 3.3. *Let C be an element of \mathcal{L} . Then there exists a polynomial $B^*(C, X)$ such that $B^*(C, n) = B(C, n)$ for all n sufficiently large.*

Proof. Set $B=(PC)^{(1)}$ in $\mathcal{R}(\mathcal{L}, C)$. Then $B_n=PC^n$ for $n \geq 1$ and $d(M_n/B_n) = d(C^n/PC^n) = B(C, n)$, by Theorem 1.4. The Corollary now follows from Theorem 3.2. Q.E.D.

COROLLARY 3.4. *Let Q be a P -primary element of \mathcal{L} . Then there exists a polynomial $F^*(Q, X)$ such that $F^*(Q, n) = F(Q, n)$ for all n sufficiently large.*

Proof. Set $B = Q^{(0)}$ in $\mathcal{R}(\mathcal{L}, Q)$. Q.E.D.

COROLLARY 3.5. *Let Q be a P -primary element of \mathcal{L} . Then there exists a polynomial $D^*(Q, X)$ such that $D^*(Q, n) = D(Q, n)$ for all n sufficiently large.*

Proof. This is immediate from Corollary 3.4. Q.E.D.

Let $\partial p(x)$ denote the degree of a polynomial $p(x)$. It will presently be shown that if \mathcal{L} has altitude d , then for every choice of the P -primary element Q , $\partial B^*(Q, X) = \partial F^*(Q, X) = d - 1$ and $\partial D^*(Q, X) = d$. Of course, the equality $\partial D^*(Q, X) = \partial F^*(Q, X) + 1$ is clear, since for all sufficiently large n , $D^*(Q, n + 1) - D^*(Q, n) = F^*(Q, n)$. And since $P^K \leq Q \leq P$ for some K , it is immediate that $\partial D^*(Q, X) = \partial D^*(P, X)$ and hence also that $\partial F^*(Q, X) = \partial F^*(P, X)$. These remarks establish part of the following

LEMMA 3.6. *If Q is a P -primary element, then*

$$\begin{aligned} \partial B^*(Q, X) &= \partial B^*(P, X) = \partial F^*(Q, X) = \partial F^*(P, X) \\ &= \partial D^*(Q, X) - 1 = \partial D^*(P, X) - 1. \end{aligned}$$

Proof. Since $B^*(P, X) = F^*(P, X)$, it is necessary only to show that $\partial B^*(Q, X) = \partial F^*(Q, X)$. Choose K so that $P^K \leq Q$ and let $h = B(P, 1)$. Then

$$\begin{aligned} F(Q, n) &= d(Q^n/Q^{n+1}) = d(Q^n/Q^n P) + d(Q^n P/Q^{n+1}) \\ &\leq d(Q^n/Q^n P) + d(Q^n P/Q^n P^K) = d(Q^n/Q^n P) + \sum_{r=1}^{K-1} d(Q^n P^r/Q^n P^{r+1}) \\ &\leq d(Q^n/Q^n P) + \sum_{r=1}^{K-1} d(Q^n/Q^n P) h^r \leq B(Q, n) + B(Q, n)(K-1)h^{(K-1)} \\ &= B(Q, n)(1 + (K-1)h^{(K-1)}). \end{aligned}$$

Since also $B(Q, n) \leq F(Q, n)$, it follows that $\partial B^*(Q, X) \leq \partial F^*(Q, X) \leq \partial B^*(Q, X)$. Q.E.D.

COROLLARY 3.7. *If Q is a P -primary element of \mathcal{L} with $B(Q, 1) = h$, then $\partial D^*(Q, X) \leq h$.*

Proof. If $B(Q, 1) = h$, then clearly

$$B(Q, n) \leq \binom{n+h-1}{h-1}$$

It follows therefore that $\partial B^*(Q, X) = \partial D^*(Q, X) - 1 \leq h - 1$. Q.E.D.

LEMMA 3.8. *If Q is a P -primary element of \mathcal{L} , then $\partial D^*(Q, X) \geq \text{alt } \mathcal{L}$.*

Proof. By Lemma 3.6, it is sufficient to show that $\partial D^*(P, X) \geq \text{alt } \mathcal{L}$. If $\text{alt } \mathcal{L} = 0$, then $P^n = 0$ for all large n , so $\partial D^*(P, X) = 0$. Hence assume \mathcal{L} has altitude $d > 0$ and let $P_0 < P_1 < \dots < P_d = P$ be a prime chain in \mathcal{L} . Then $\text{alt } \mathcal{L}/P_0 = \text{alt } \mathcal{L}$ and $D^*(P/P_0, X) \leq D^*(P, X)$. Let $A \leq P$ be a principal element of \mathcal{L} with $A \not\leq P_0$ so that A/P_0 is principal in \mathcal{L}/P_0 with $A/P_0 \neq 0$. Choose K so that

$$(P/P_0)^{K+n} \wedge (A/P_0) = [(P_0/P_0)^K \wedge (A/P_0)](P/P_0)^n, \text{ for all } n \geq 0$$

(Corollary 2.9). Then

$$((P/P_0)^{K+n} : (A/P_0))(A/P_0) = ((P/P_0)^K : (A/P_0))(P/P_0)^n(A/P_0),$$

and hence

$$(P/P_0)^{K+n} : (A/P_0) = ((P/P_0)^K : (A/P_0))(P/P_0)^n \leq (P/P_0)^n,$$

for all $n \geq 0$ (Lemma 1.1). Now, since $A \leq P$, $(P/P_0)^{K+n-1} \leq ((P/P_0)^{K+n} : A/P_0)$. It follows that

$$\begin{aligned} D(P/P_0, K+n-1) &\geq \dim (I[(P/P_0)^{K+n} : (A/P_0)]) && \text{(Lemma 1.1)} \\ &= \dim ((A/P_0)/[(P/P_0)^{K+n} : (A/P_0)](A/P_0)) \\ &= \dim ((A/P_0)/(P/P_0)^{K+n} \wedge (A/P_0)) \\ &= \dim (((P/P_0)^{n+K} \vee A)/(P/P_0)^{n+K}) \\ &= D(P/P_0, n+K) - D(P/(P_0 \vee A), n+K) \geq D(P/P_0, n), \end{aligned}$$

so

$$\begin{aligned} D(P/P_0, n+K-1) &\geq D(P/P_0, n+K) - D(P/(P_0 \vee A), n+k) \\ &\geq D(P/P_0, n) \text{ for all } n \geq 0. \end{aligned}$$

Hence for n sufficiently large,

$$D^*(P/(P_0 \vee A), n+K) \geq D^*(P/P_0, n+K) - D^*(P/P_0, n+K-1)$$

and

$$D^*(P/P_0, n+K) - D^*(P/P_0, n) \geq D^*(P/(P_0 \vee A), n+K),$$

so $\partial D^*(P/(P_0 \vee A), X) = \partial D^*(P/P_0, X) - 1$. Now, it is clear that $\text{alt } \mathcal{L}/(P_0 \vee A) \leq d-1$ and therefore (Lemma 1.8) that $\text{alt } \mathcal{L}/(P_0 \vee A) = d-1$. Hence, by the inductive hypothesis, $\partial D^*(P/P_0 \vee A) \geq d-1$ so that

$$\begin{aligned} \partial D^*(P, X) &\geq \partial D^*(P/P_0, X) = 1 + \partial D^*(P/(P_0 \vee A), X) \\ &\geq \text{alt } \mathcal{L}/(P_0 \vee A) + 1 = \text{alt } \mathcal{L}/P_0 = \text{alt } \mathcal{L}. \end{aligned} \quad \text{Q.E.D.}$$

THEOREM 3.9. *If Q is a P -primary element of the local lattice \mathcal{L} , then*

$$\partial D^*(Q, X) = \text{alt } \mathcal{L}.$$

Proof. Let $A \neq 0$ be a principal element of \mathcal{L} with $A \leq Q$. Choose s so that $Q^{n+s} \wedge A = Q^n(Q^s \wedge A)$, for all $n \geq 0$. Then $(Q^{n+s} : A)A \leq Q^n A$, so $Q^{n+s} : A \leq Q^n \vee (0 : A)$, since A is principal. Now,

$$\begin{aligned} \dim(I/Q^n \vee A) &= \dim(I/Q^n) - \dim(Q^n \vee A/Q^n) \\ &= \dim(I/Q^n) - \dim(A/Q^n \wedge A) \\ &= \dim(I/Q^n) - \dim(I/Q^n : A) \quad (\text{Lemma 1.1}), \end{aligned}$$

so

$$\dim(I/Q^n) - \dim(I/Q^{n-1} \vee (0 : A)) \leq \dim(I/Q^n \vee A)$$

and

$$\dim(I/Q^n \vee A) \leq \dim(I/Q^n) - \dim(I/Q^{n-s} \vee (0 : A)).$$

Hence

$$D(Q, n) - D(Q/(0 : A), n-1) \leq D(Q/A, n)$$

and

$$D(Q/A, n) \leq D(Q, n) - D(Q/(0 : A), n-s),$$

so that

$$D^*(Q, n) \leq D^*(Q/A, n) + D^*(Q/(0 : A), n-1)$$

and

$$D^*(Q, n) \geq D^*(Q/A, n) + D^*(Q/(0 : A), n-s),$$

for all n sufficiently large. It follows that one of the polynomials $D^*(Q/A, X)$ and $D^*(Q/(0 : A), X)$ has the same degree as $D^*(Q, X)$.

Now, let A_1, \dots, A_h be principal elements in \mathcal{L} with $Q = A_1 \vee \dots \vee A_h$; and set $d = \text{alt } \mathcal{L}$. By Lemma 3.6 it can be assumed that no fewer than h principal elements have join a P -primary element. If $d=0$, then the calculation is immediate, so assume $d>0$. Choose K so that $0 : (A_h)^K = 0 : (A_h)^{K+i}$ for all $i \geq 0$, and set $A = (A_h)^K$. Then $A_1 \vee \dots \vee A_{h-1} \vee A$ is P -primary element of \mathcal{L} , so (Lemma 3.6) it can be assumed $Q = A_1 \vee \dots \vee A_{h-1} \vee A$. Now, if $h-d=0$, then (Corollary 3.7 and Lemma 3.8) $d \leq \partial D^*(Q, X) \leq h=d$. Hence it can be assumed $h-d>0$.

Now assume, $\partial D^*(Q, X) = \partial D^*(Q/A, X)$. If $\text{alt } \mathcal{L}/A < d$, then it follows by the induction on altitude that $\partial D^*(Q/A, X) = \text{alt } \mathcal{L}/A$ so that $\partial D^*(Q, X) = \partial D^*(Q/A, X) \leq d$. On the other hand, if $\text{alt } \mathcal{L}/A = d$, then it follows by the induction on the number of generators that $\partial D^*(Q/A, X) = \text{alt } \mathcal{L}/A$ and hence that $\partial D^*(Q, X) = d$. Hence if $\partial D^*(Q, X) = \partial D^*(Q/A, X)$, then $\partial D^*(Q, X) = d$.

Assume now that $\partial D^*(Q, X) = \partial D^*(Q/(0 : A), X)$. By the choice of A , $A/(0 : A)$ is not a zero divisor in $\mathcal{L}/(0 : A)$. Hence, by the earlier part of the proof,

$$D^*(Q/(0 : A), n) \leq D^*(Q/(0 : A) \vee A, n) + D^*(Q/(0 : A), n-1)$$

and

$$D^*(Q/(0 : A), n) \geq D^*(Q/(0 : A) \vee A, n) + D^*(Q/(0 : A), n-s);$$

so

$$\partial D^*(Q/(0 : A) \vee A, X) = \partial D^*(Q/(0 : A), X) - 1 = \partial D^*(Q, X) - 1.$$

It is easily seen that $\mathcal{L}/(0 : A) \vee A$ has altitude $< d$, so it follows by the inductive assumption on altitude that

$$\partial D^*(Q/(0 : A) \vee A, X) = \partial D^*(Q, X) - 1 < d,$$

and hence (Lemma 3.8) that $\partial D^*(Q, X) = d$. Q.E.D.

4. Regular local lattices. A local lattice \mathcal{L} of altitude d is said to be a *regular local lattice* if the maximal element $P \in \mathcal{L}$ is the join of d -principal elements.

It is noted that by Theorem 1.4, the local lattice \mathcal{L} is regular if, and only if, $\dim(P/P^2) = \text{alt}(\mathcal{L})$.

THEOREM 4.1. *If \mathcal{L} is a regular local lattice, then 0 is prime.*

Proof. If $\text{alt}(\mathcal{L}) = 0$, then the result is clear. The proof will proceed by induction. Assume $\text{alt} \mathcal{L} = d \geq 1$. Let $P_0 < P_1 < \dots < P_d = P$ be a prime chain in \mathcal{L} and let A_1, \dots, A_d be a principal element of \mathcal{L} with $P = A_1 \vee \dots \vee A_d$. Then \mathcal{L}/P_0 is again regular with maximal element $P = P/P_0 = A_1/P_0 \vee \dots \vee A_d/P_0$. Since P/P_0 is not the join of fewer than d -principal elements [1], it is clear that $A_i/P_0 \neq 0/P_0$ for $i = 1, \dots, d$. Then $(\mathcal{L}/P_0) \mid (A_1/P_0) = \mathcal{L} \mid (A_1 \vee P_0)$ has altitude $d-1$ (Lemma 1.8) and is regular since $P/(A_1 \vee P_0) = A_2/(A_1 \vee P_0) \vee \dots \vee A_d/(A_1 \vee P_0)$. Hence by the inductive hypothesis, $A_1 \vee P_0 = 0/(A_1 \vee P_0)$ is prime in $\mathcal{L}/(A_1 \vee P_0)$, and therefore also in \mathcal{L} .

Now, $P = P/A_1$ is the join of the $d-1$ principal elements $A_2/A_1, \dots, A_d/A_1$ in \mathcal{L}/A_1 , so \mathcal{L}/A_1 has altitude $\leq d-1$ [1] and therefore $= d-1$, (Lemma 1.8). Hence, by the inductive hypothesis, $A_1 = 0/A_1$ is prime in \mathcal{L}/A_1 , and therefore also in \mathcal{L} . Since $A_1 \leq A_1 \vee P_0$, and since \mathcal{L}/A_1 and $\mathcal{L}/(A_1 \vee P_0)$ both have altitude $d-1$, so that A_1 and $A_1 \vee P_0$ have the same depth in \mathcal{L} , it is immediate that $A_1 = A_1 \vee P_0$, whence $P_0 \leq A_1$. Then $P_0 = P_0 \wedge A_1 = (P_0 : A_1)A_1 = P_0A_1$ (since P_0 is prime $A_1 \not\leq P_0$), so $P_0 = 0$ by Lemma 1.3. Q.E.D.

COROLLARY 4.2. *Let \mathcal{L} be a regular local lattice of altitude d , and let A_1, \dots, A_d be principal elements with $P = A_1 \vee \dots \vee A_d$. Then $A_1 \vee \dots \vee A_i$ is prime for $i = 0, \dots, d$.*

Proof. By Lemma 1.8, it follows (by induction on i) that $\mathcal{L}/(A_1 \vee \dots \vee A_i)$ is regular of altitude $d-i$, so that $0/(A_1 \vee \dots \vee A_i) = A_1 \vee \dots \vee A_i$ is prime in $\mathcal{L}/(A_1 \vee \dots \vee A_i)$ and therefore also in \mathcal{L} . Q.E.D.

COROLLARY 4.3. *Let A be a principal element of the regular local lattice \mathcal{L} , with $A \leq P$ and $A \not\leq P^2$. Then A is prime in \mathcal{L} .*

Proof. Let \mathcal{L} be a regular of altitude d , so that $\dim(P/P^2)=d$, (Theorem 1.4). Now, $(A \vee P^2)/P^2 \cong A/(P^2 \wedge A) = A/(P^2 : A)A \cong I/(P^2 : A)$ (Lemma 1.1) so $(A \vee P^2)/P^2$ has dimension 1 and $P/(A \vee P^2)$ has dimension $d-1$. Let $A \vee P^2 = C_1 < C_2 < \dots < C_d = P$ be a composition chain for $P/(A \vee P^2)$. For each $i=2, \dots, d$ let A_i be a principal element of \mathcal{L} such that $A_i < C_i$ and $A_i \not\leq C_{i-1}$. Then $C_{i-1} \vee A_i = C_i$, for $i=2, \dots, d$. Set $A = A_1$. Then $P = C_1 \vee A_2 \vee \dots \vee A_d = P^2 \vee A_1 \vee \dots \vee A_d$, so $P = A_1 \vee \dots \vee A_d$ by Lemma 1.3, so A_1 is prime by Corollary 4.2. Q.E.D.

COROLLARY 4.4. *Let A, B be elements of a regular local lattice \mathcal{L} , with $A \leq P^n$, $A \not\leq P^{n+1}$ and $B \leq P^m$, $B \not\leq P^{m+1}$. Then $AB \leq P^{n+m}$ and $AB \not\leq P^{n+m+1}$.*

Proof. Let \mathcal{L} be regular of altitude d , with $P = A_1 \vee \dots \vee A_d$ (A_i principal). Then the maximal element of $\mathcal{R}(\mathcal{L}, P)$ is the join of the $d+1$ principal elements $I^{(-1)}$, $A_1^{(1)}$, \dots , $A_d^{(1)}$ (Lemma 2.3 and Theorem 2.7), so $\mathcal{R}(\mathcal{L}, P)$ is a regular local lattice of altitude $d+1$ (Theorem 2.17). Hence $I^{(-1)}$ is prime (Corollary 4.2). Now, assume $A, B \in \mathcal{L}$, $A \leq P^n$, $A \not\leq P^{n+1}$, and $B \leq P^m$, $B \not\leq P^{m+1}$. Then $I^{(-1)}$ has i th coordinate P^{i+1} , so $A^{(n)} \not\leq I^{(-1)}$ and $B^{(m)} \not\leq I^{(-1)}$. Hence $(AB)^{(n+m)} = A^{(n)}B^{(m)} \not\leq I^{(-1)}$, and therefore $AB \not\leq P^{n+m+1}$. Q.E.D.

Hence the degree function on a regular local lattice is a valuation.

It is noted that, as for local rings, if \mathcal{L} is regular of altitude 1, then Theorem 2.7 can be strengthened to say that every element $A \neq 0$ is a power P^n of the maximal element ($I = P^0$). For in this case, P is principal, so if n is the largest i such that $A \leq P^i$, then $A = A \wedge P^n = (A : P^n)P^n$, so that if $P^n \neq A$, then $A \leq PP^n = P^{n+1}$. Hence if \mathcal{L} is a regular local lattice of altitude 1, then \mathcal{L} is isomorphic to the lattice of ideals of a regular local ring, and $\mathcal{L}^* = \mathcal{L} - \{0\}$ is a Gaussian semigroup. This second property also holds in the semigroup of principal elements of a regular local lattice of altitude 2, as is easily seen.

REFERENCES

1. R. P. Dilworth, *Abstract commutative ideal theory*, Pacific J. Math. **12** (1962), 481-498.
2. M. Nagata, *Local rings*, Interscience, New York, 1962.
3. D. G. Northcott, *Ideal theory*, Cambridge Univ. Press, New York, 1963.
4. D. Rees, *A-transforms of local rings and a theorem on multiplicities of ideals*, Proc. Cambridge Philos. Soc. **57** (1961), 8-17.
5. O. Zariski and P. Samuel, *Commutative algebra*, Vol. II, Van Nostrand, Princeton, N. J., 1960.

UNIVERSITY OF IOWA,
IOWA CITY, IOWA