

# FAMILIES OF ARCS IN $E^3$

BY  
R. B. SHER<sup>(1)</sup>

**1. Introduction.** In [6], R. H. Bing announced the result that Euclidean 3-space does not contain an uncountable collection of pairwise disjoint wild surfaces. J. Stallings, in [19], gave an example of an uncountable collection of pairwise disjoint wild disks in 3-space. In [17] Stallings' construction has been modified so as to obtain an uncountable collection of pairwise disjoint inequivalently imbedded disks. In §4 of this paper we consider arcs in 3-space which are locally tame modulo a compact 0-dimensional set. Examples of such arcs have been studied in [1], [3], [9], [10], [11], and [12]. It is shown (assuming the Continuum Hypothesis) that an uncountable collection of pairwise disjoint arcs of this type exists which contains a representative of each imbedding class (under space homeomorphism) of such arcs.

In §5 we answer, in the affirmative, the following question raised in [8] by Bing: Given a 2-sphere  $S$  in  $E^3$ , is there a family  $F$  of mutually exclusive tame arcs such that for each point of  $S$  at which  $S$  can be pierced by a tame arc, there is a member of  $F$  piercing there?

**2. Definitions and notation.** We use  $E^n$  to denote Euclidean  $n$ -space, coordinatized in the usual way by a set of mutually perpendicular axes  $X_1, X_2, \dots$ , and  $X_n$ , and with points denoted by their coordinates  $(x_1, x_2, \dots, x_n)$ . If  $p = (x_1, x_2, \dots, x_n)$  and  $q = (y_1, y_2, \dots, y_n)$ , the distance between  $p$  and  $q$  is  $\rho(p, q) = (\sum_{i=1}^n (x_i - y_i)^2)^{1/2}$ . If  $p \in E^n$  and  $\epsilon$  is a positive number,  $N(p, \epsilon)$  is the set  $\{p' \in E^n \mid \rho(p, p') < \epsilon\}$ . If  $X$  is a bounded subset of  $E^n$ , the diameter of  $X$  is the least upper bound of the set of numbers  $\{\rho(p, q) \mid p, q \in X\}$ . An  $\epsilon$ -set is a subset of  $E^n$  whose diameter is no more than  $\epsilon$ . We use  $D_{\max}(X)$  to denote the least upper bound of the set of numbers which are diameters of components of  $X$ . If  $X$  is a subset of the  $x_1x_2$ -coordinate plane in  $E^3$ ,  $X \times [a, b]$  denotes the set  $\{(x_1, x_2, x_3) \mid (x_1, x_2, 0) \in X \text{ and } a \leq x_3 \leq b\}$ . The homeomorphism  $h: E^n \rightarrow E^n$  is an  $\epsilon$ -homeomorphism if  $\rho(p, h(p)) < \epsilon$  for each  $p \in E^n$ .

An arc  $A$  in  $E^3$  is *tame* if there is a homeomorphism of  $E^3$  onto itself taking  $A$  onto a polygonal arc. Otherwise,  $A$  is *wild*. The arc  $A$  is *locally tame* (*locally polygonal*) at the point  $p \in A$  if  $p$  lies in the interior of a tame subarc (polygonal subarc) of  $A$ . Otherwise  $A$  is *locally wild* (*locally nonpolygonal*) at  $p$ . If  $X$  is a closed subset of  $A$  and  $A$  is locally tame (locally polygonal) at each point of  $A - X$ , then

---

Presented to the Society, January 25, 1967 under the title *A family of inequivalently imbedded arcs in  $E^3$* ; received by the editors January 4, 1967.

(<sup>1</sup>) This work was supported by the National Science Foundation under grant GP-6016.

$A$  is said to be *locally tame (locally polygonal) modulo  $X$* . For generalizations of the above concepts, see §1 of [4].

A compact 0-dimensional subset  $M$  of  $E^3$  is *tame* if there is a homeomorphism of  $E^3$  onto itself taking  $M$  into a polygonal arc. Otherwise  $M$  is *wild*. See [5] for characterizations of tame compact 0-dimensional subsets of  $E^3$ .

The closure of a set  $X$  will be denoted by  $\text{Cl}(X)$ . The notation  $\text{Int}(X)$  may mean point set interior, combinatorial interior, or bounded complementary domain. Similarly,  $\text{Bd}(X)$  may mean point set boundary or combinatorial boundary. When seen in context, the usage will be clear in each case.

The setting for the results included here, with one exception, is  $E^3$ . The exception is Theorem 5 which generalizes Theorem 4 to  $E^n$ ,  $n > 3$ .

**3. Preliminary results.** In this section some results will be obtained dealing with imbeddings of compact 0-dimensional sets in  $E^3$ , one of which, Theorem 1, will find application in the following sections.

Suppose  $C$  is a polyhedral cube with  $n \geq 0$  handles in  $E^3$ . Then  $C$  is the homeomorphic image of a closed neighborhood of a bouquet of  $n$  circles in  $E^3$ . If  $n=0$ , interpret "bouquet of 0 circles" as "point". The image of the bouquet under such a homeomorphism shall be referred to as a *center* of  $C$ . We shall refer to any bouquet of  $n$  simple closed curves in  $C$  which is equivalently imbedded in  $C$  (under a homeomorphism of  $C$  onto itself fixed on  $\text{Bd}(C)$ ) to a center of  $C$  as a *central curve* of  $C$ . Notice that if  $K$  is a closed subset of  $\text{Int}(C)$ ,  $B$  is a central curve of  $C$ , and  $\varepsilon > 0$ , then there is a homeomorphism of  $C$  onto itself, fixed on  $\text{Bd}(C)$ , carrying  $K$  into  $N(B, \varepsilon)$ .

**LEMMA 1.** *If  $C$  is a polyhedral cube with  $n$  handles, and  $A$  is an arc, then there is a central curve  $S$  of  $C$  so that  $A \cap S = \emptyset$ .*

**Proof.** There is a homeomorphism  $h$  of  $C$  into  $E^3$  taking  $B$ , a center of  $C$ , into the  $x_1x_2$ -coordinate plane. Let  $\varepsilon > 0$  be a number so small that  $(h(B) \times [0, \varepsilon]) \subset \text{Int}(h(C))$ . There is a positive number  $\delta \leq \varepsilon$  so that the plane  $x_3 = \delta$  meets the set  $h(A \cap C)$  in at most a set of dimension 0, for an arc cannot contain uncountably many pairwise disjoint open intervals. Using elementary facts from plane topology, one is able to construct a central curve for  $h(C)$ , lying in the plane  $x_3 = \delta$ , which misses  $h(A \cap C)$ . The inverse image of this curve, under  $h$ , is a central curve of  $C$  which misses  $A$ .

**THEOREM 1.** *Suppose  $M$  is a compact 0-dimensional subset of  $E^3$ ,  $\{A_i\}_{i=1}^{\infty}$  is a countable collection of arcs, and  $\varepsilon > 0$ . Then there is an  $\varepsilon$ -homeomorphism  $h: E^3 \rightarrow E^3$  such that  $h(M) \cap (\bigcup_{i=1}^{\infty} A_i) = \emptyset$ .*

**Proof.** The set  $M$  is definable by cubes with handles (Lemma 4 of [2] or Theorem 2 of [15]). In other words, there is a sequence  $M_1, M_2, M_3, \dots$  of compact 3-manifolds with boundary such that

(1) for each positive integer  $i$ ,  $M_i$  is the union of a finite number of pairwise disjoint polyhedral cubes with handles,

(2) for each positive integer  $i$ ,  $M_{i+1} \subset \text{Int}(M_i)$ , and

(3)  $M = \bigcap_{i=1}^{\infty} M_i$ .

The homeomorphism  $h$  will be the limit of homeomorphisms  $h_1, h_2, h_3, \dots$ . We begin the construction of the sequence in this paragraph with the construction of  $h_1$ . Since  $M$  is totally disconnected, there is a positive integer  $i(1)$  such that  $D_{\max}(M_{i(1)}) < \max(1, \epsilon)$ . Denote the components of  $M_{i(1)}$  by  $C_1^1, \dots, C_{n(1)}^1$ . If  $j=1, 2, \dots$ , or  $n(1)$ ,  $C_j^1$  is a polyhedral cube with handles. By Lemma 1,  $C_j^1$  has a central curve  $S_j^1$  such that  $S_j^1 \cap A_1 = \emptyset$ . Let  $f_j^1$  be a homeomorphism of  $C_j^1$  onto itself, fixed on  $\text{Bd}(C_j^1)$ , which moves  $M_{i(1)+1} \cap C_j^1$  so close to  $S_j^1$  that  $f_j^1(M_{i(1)+1} \cap C_j^1) \cap A_1 = \emptyset$ . Let  $h_1: E^3 \rightarrow E^3$  be defined by  $h_1(p) = p$  if  $p \in E^3 - M_{i(1)}$ ,  $h_1(p) = f_j^1(p)$  if  $p \in C_j^1$ .

Now suppose that homeomorphisms  $h_1, \dots, h_m$  of  $E^3$  onto itself have been defined so that

(4) there are positive integers  $i(1) < i(2) < \dots < i(m)$  such that  $h_j|_{E^3 - M_{i(j)}} = h_{j-1}|_{E^3 - M_{i(j)}}$ , where  $j=1, 2, \dots$ , or  $m$  and  $h_0$  is the identity on  $E^3$ ,

(5)  $D_{\max}(h_j(M_{i(j)})) < 1/j$ , if  $j=1, 2, \dots$ , or  $m$ , and

(6)  $h_j(M_{i(j)+1}) \cap (\bigcup_{i=1}^j A_i) = \emptyset$ , if  $j=1, 2, \dots$ , or  $m$ .

Let  $i(m+1)$  be an integer greater than  $i(m)$  such that  $D_{\max}(h_m(M_{i(m+1)})) < 1/m+1$ . Denote the components of  $h_m(M_{i(m+1)})$  by  $C_1^{m+1}, \dots, C_{n(m+1)}^{m+1}$ . If  $j=1, 2, \dots$ , or  $n(m+1)$ ,  $C_j^{m+1}$  is a cube with handles and  $C_j^{m+1} \cap (\bigcup_{i=1}^m A_i) = \emptyset$ . By Lemma 1,  $C_j^{m+1}$  has a central curve  $S_j^{m+1}$  such that  $S_j^{m+1} \cap A_{m+1} = \emptyset$ . Let  $f_j^{m+1}$  be a homeomorphism of  $C_j^{m+1}$  onto itself, fixed on  $\text{Bd}(C_j^{m+1})$ , which moves  $h_m(M_{i(m+1)+1}) \cap C_j^{m+1}$  so close to  $S_j^{m+1}$  that  $f_j^{m+1}(h_m(M_{i(m+1)+1}) \cap C_j^{m+1}) \cap A_{m+1} = \emptyset$ . Let  $h_{m+1}: E^3 \rightarrow E^3$  be defined by  $h_{m+1}(p) = h_m(p)$  if  $p \in E^3 - M_{i(m+1)}$ ,  $h_{m+1}(p) = f_j^{m+1}(h_m(p))$  if  $p \in h_m^{-1}(C_j^{m+1})$ . The homeomorphisms  $h_1, \dots, h_{m+1}$  satisfy conditions (4)–(6) above, so the construction of the sequence may be continued in the fashion described here.

It is not difficult to show that  $h = \lim \{h_i\}$  is a homeomorphism. The set  $M$  is carried by  $h$  onto the set  $\bigcap_{i=1}^{\infty} h(M_i)$ . For each positive integer  $j$ ,  $h(M_{i(j)+1}) = h_j(M_{i(j)+1})$  fails to intersect  $\bigcup_{i=1}^j A_i$ , so  $h(M)$  fails to intersect  $\bigcup_{i=1}^{\infty} A_i$ . Since  $h$  acts as the identity on  $E^3 - M_{i(1)}$  and  $D_{\max}(M_{i(1)}) < \epsilon$ ,  $h$  is an  $\epsilon$ -homeomorphism.

**COROLLARY 1.** *Suppose  $M$  is a compact 0-dimensional subset of  $E^3$ ,  $\epsilon > 0$ , and  $A$  is a subset of  $E^3$  which is contained in the union of a countable collection of arcs. Then there is an  $\epsilon$ -homeomorphism  $h: E^3 \rightarrow E^3$  such that  $h(M) \cap A = \emptyset$ .*

An example of a set  $A$  satisfying the hypothesis of Corollary 1 is the union of all straight lines which are parallel to a coordinate axis and intersect a coordinate plane in a point all of whose coordinates are rational. Applying Corollary 1, we obtain the following result.

**COROLLARY 2.** *If  $M$  is a compact 0-dimensional subset of  $E^3$ , and  $\varepsilon > 0$ , then there is an  $\varepsilon$ -homeomorphism  $h: E^3 \rightarrow E^3$  such that if  $(x_1, x_2, x_3) \in h(M)$ , then at most one of the numbers  $x_1, x_2, x_3$  is rational.*

One might wonder if the conclusion of Corollary 2 could be strengthened to conclude that all of the numbers  $x_1, x_2, x_3$  are irrational. The following shows that, in general, it can not be.

**THEOREM 2.** *If  $M$  is a compact 0-dimensional subset of  $E^3$ , then  $M$  is tame if and only if there is a homeomorphism  $h: E^3 \rightarrow E^3$  such that if  $(x_1, x_2, x_3) \in h(M)$ , then the numbers  $x_1, x_2, x_3$  are irrational.*

**Proof.** If  $M$  is tame, there is a homeomorphism  $h_1: E^3 \rightarrow E^3$  such that  $h_1(M) \subset L$ , where  $L$  is the straight line through the point  $(\pi, \pi, 0)$  parallel to the  $x_3$ -axis. By the analog of Theorem 1 on the real line, there is a homeomorphism  $h_2: L \rightarrow L$  such that if  $(\pi, \pi, x_3) \in h_2 h_1(M)$ , then  $x_3$  is irrational. Let  $h_3$  be a homeomorphism of  $E^3$  onto itself such that  $h_3$  restricted to  $L$  is  $h_2$ . Then  $h = h_3 h_1$  carries  $M$  onto a set of points of the form  $(\pi, \pi, x_3)$  with  $x_3$  irrational.

Now suppose  $M$  is compact and 0-dimensional and that  $h$  is a homeomorphism of  $E^3$  onto itself such that  $(x_1, x_2, x_3) \in h(M)$  implies  $x_1, x_2, x_3$  are irrational. Let  $(x_1, x_2, x_3) \in h(M)$  and  $\varepsilon$  be an arbitrary positive number. If  $i$  is 1, 2 or 3, let  $r_1^i$  and  $r_2^i$  be rational numbers such that  $x_i - \varepsilon/4 < r_1^i < x_i < r_2^i < x_i + \varepsilon/4$ . The set  $C = \{(x'_1, x'_2, x'_3) \mid r_1^i \leq x'_i \leq r_2^i, i = 1, 2, 3\}$  is a polyhedral cube of diameter less than  $\varepsilon$  containing  $(x_1, x_2, x_3)$  in its interior and having no point of  $h(M)$  on its boundary. By Theorem 3.1 of [5],  $h(M)$ , and hence  $M$ , is tame.

**COROLLARY 3.** *If  $M$  is a compact 0-dimensional subset of  $E^3$  and the set of points of  $M$  which have a rational coordinate is closed and tame, then  $M$  is tame.*

**Proof.** Let  $N = \{(x_1, x_2, x_3) \in M \mid x_1 \text{ or } x_2 \text{ or } x_3 \text{ is rational}\}$ . By Theorem 2,  $M$  is locally tame modulo  $N$ . By Theorem 6.1 of [5],  $M$  is tame.

**LEMMA 2.** *Suppose  $\{M_i\}_{i=1}^\infty$  is a countable collection of compact 0-dimensional sets in  $E^2$ ,  $a$  and  $b$  are points of  $E^2 - \bigcup_{i=1}^\infty M_i$ , and  $J$  is a simple closed curve containing  $a$  and  $b$ . Then there is an arc  $ab$  such that  $\text{Int}(ab) \subset \text{Int}(J)$  and  $ab \cap (\bigcup_{i=1}^\infty M_i) = \emptyset$ .*

**REMARK.** The proof of Lemma 2 to be given here involves the concept of simple chains. For definitions of this and other terms, one is referred to Chapter 3, pp. 105–119, of [14].

**Proof.** By the classical Schoenflies Theorem, we may suppose that  $J$  is the unit circle in  $E^2$ . Let  $A_1$  be a polygonal arc from  $a$  to  $b$  so that  $\text{Int}(A_1) \subset \text{Int}(J)$  and  $A_1 \cap M_1 = \emptyset$ . As a convention, let us suppose that all arcs with end points  $a$  and  $b$  are linearly ordered from  $a$  to  $b$ . Then  $A_1$  is the union of a finite collection  $A_1^1, \dots, A_{n(1)}^1$ , of straight line intervals such that

- (1) if  $j=1, 2, \dots$ , or  $n(1)$ , then the diameter of  $A_j^1$  is less than 1,
  - (2) the intersection of any pair of intervals  $A_1^1, \dots, A_{n(1)}^1$  is either empty or an end point of each, and
  - (3) if  $1 \leq j < k \leq n(1)$ , then  $p \in \text{Int}(A_j^1)$  implies that  $p$  precedes each point of  $A_k^1$ .
- If  $j=1, 2, \dots$ , or  $n(1)$ , there is an open disk  $d_j^1$  containing  $A_j^1$  such that
- (4)  $C_1: d_1^1, \dots, d_{n(1)}^1$  is a simple 1-chain from  $a$  to  $b$ ,
  - (5) if  $j=2, 3, \dots$ , or  $n(1)-1$ ,  $d_j^1 \subset \text{Int}(J)$ ,
  - (6)  $(\bigcup_{i=1}^{n(1)} d_i^1) \cap M_1 = \emptyset$ , and
  - (7) if  $j=1, 2, \dots$ , or  $n(1)-1$ ,  $d_j^1 \cap d_{j+1}^1$  is an open disk.

There is a polygonal arc  $A_2$  from  $a$  to  $b$  such that

- (8)  $\text{Int}(A_2) \subset \text{Int}(J)$ ,
- (9)  $A_2 \subset \bigcup_{i=1}^{n(1)} d_i^1$ ,
- (10) if  $p \in A_2$  precedes  $q \in A_2$  and  $p$  and  $q$  are in the same link of  $C_1$ , then every point of  $A_2$  between  $p$  and  $q$  also lies in the link, and
- (11)  $A_2 \cap M_2 = \emptyset$ .

The arc  $A_2$  is the union of a finite collection  $A_1^2, \dots, A_{n(2)}^2$  of straight line intervals such that

- (12) if  $j=1, 2, \dots$ , or  $n(2)$ , then the diameter of  $A_j^2$  is less than  $1/2$ ,
- (13) the intersection of any pair of intervals  $A_1^2, \dots, A_{n(2)}^2$  is either empty or an end point of each,
- (14) if  $1 \leq j < k \leq n(2)$ , then  $p \in \text{Int}(A_j^2)$  implies that  $p$  precedes each point of  $A_k^2$ , and

(15) if  $j=1, 2, \dots$ , or  $n(2)$ , there is an integer  $k$  such that  $A_j^2 \subset d_k^1$ .

If  $j=1, 2, \dots$ , or  $n(2)$ , there is an open disk  $d_j^2$  containing  $A_j^2$  such that

- (16)  $C_2: d_1^2, \dots, d_{n(2)}^2$  is a simple  $1/2$ -chain from  $a$  to  $b$ ,
- (17) if  $j=2, 3, \dots$ , or  $n(2)-1$ ,  $d_j^2 \subset \text{Int}(J)$ ,
- (18)  $(\bigcup_{i=1}^{n(2)} d_i^2) \cap M_2 = \emptyset$ ,
- (19) if  $j=1, 2, \dots$ , or  $n(2)-1$ ,  $d_j^2 \cap d_{j+1}^2$  is an open disk, and
- (20)  $C_2$  is a refinement of  $C_1$  running straight through  $C_1$ .

Continuing in this manner, we construct simple chains  $C_1, C_2, C_3, \dots$  such that

- (21)  $C_k: d_1^k, \dots, d_{n(k)}^k$  is a simple  $1/k$ -chain from  $a$  to  $b$ ,
- (22) if  $j=2, 3, \dots$ , or  $n(k)-1$ ,  $d_j^k \subset \text{Int}(J)$ ,
- (23)  $(\bigcup_{i=1}^{n(k)} d_i^k) \cap M_k = \emptyset$ , and
- (24)  $C_{k+1}$  is a refinement of  $C_k$  running straight through  $C_k$ . By the proof of Theorem 3-15, pp. 116-117 of [14],  $ab = \bigcap_{k=1}^{\infty} (\bigcup_{i=1}^{n(k)} d_i^k)$  is an arc from  $a$  to  $b$ . By condition (22) satisfied by the simple chains  $C_1, C_2, C_3, \dots$ ,  $\text{Int}(ab) \subset \text{Int}(J)$ , while by condition (23),  $ab \cap (\bigcup_{i=1}^{\infty} M_i) = \emptyset$ .

The proof of Lemma 2 shows that, in  $E^n (n \geq 2)$ , the complement of a countable union of compact sets of dimension less than  $n-1$  is arcwise connected. An example given by R. H. Bing in the case  $n=3$  (Theorem 6.3 of [5]) shows that there may be no polygonal arcs in this set.

**4. Arcs locally tame modulo a compact 0-dimensional set.** If  $A$  is an arc in  $E^3$ , the *wild set* of  $A$ , denoted by  $W(A)$ , is the set  $\{p \in A \mid A \text{ is locally wild at } p\}$ . The *tame set* of  $A$  is the set  $T(A) = A - W(A)$ . Note that  $W(A)$  is closed,  $T(A)$  is open (relative to  $A$ ) and  $A$  is locally tame modulo  $W(A)$ . In this section we shall consider only those arcs  $A$  for which  $W(A)$  is a compact 0-dimensional subset of  $E^3$ .

Subsets  $H$  and  $K$  of  $E^3$  are said to be *equivalently imbedded* in  $E^3$  if there is a homeomorphism  $h: E^3 \rightarrow E^3$  such that  $h(H) = K$ . This notion induces an equivalence relation on the nonempty subsets of  $E^3$ , where the sets  $H$  and  $K$  are equivalent if and only if  $H$  and  $K$  are equivalently imbedded in  $E^3$ . We use  $C(H)$  to denote the equivalence class of  $H$  under this equivalence relation. It is easy to see that if  $A$  is an arc and  $B \in C(A)$ , then  $W(B) \in C(W(A))$ . The converse is, in general, false. By [12], there are uncountably many equivalence classes of arcs which are locally tame modulo an end point. By [11], there are uncountably many equivalence classes of arcs which are locally tame modulo an interior point. Since every Cantor set  $M$  lies on an arc which is locally tame modulo  $M$ , it follows from [18] that there are uncountably many equivalence classes of arcs which are locally tame modulo a Cantor set.

The result to be proved in this section is that there is a collection  $\{A_\alpha\}$  of pairwise disjoint arcs in  $E^3$  so that if  $A$  is an arc in  $E^3$  with a compact 0-dimensional wild set, then some member of  $\{A_\alpha\}$  is a member of  $C(A)$ .

**THEOREM 3.** *There is a collection  $\{A_\alpha\}$  of pairwise disjoint arcs in  $E^3$  such that if  $A$  is an arc in  $E^3$  with  $W(A)$  a compact 0-dimensional set, then there is an  $A' \in \{A_\alpha\}$  such that  $C(A') = C(A)$ .*

**Proof.** The theorem is to be proved by giving a constructive process for building the collection  $\{A_\alpha\}$ . Assuming the Continuum Hypothesis, the disjoint equivalence classes of arcs locally tame modulo a compact 0-dimensional set can be well-ordered so that each class is preceded in the ordering by only countably many classes. The construction of  $\{A_\alpha\}$  is begun by simply choosing a representative of the first class in the well-ordered sequence. To show that the construction may be continued through the sequence, suppose  $C(B)$  is an equivalence class of arcs locally tame modulo a compact 0-dimensional set and that the arcs  $\{A_i\}_{i=1}^\infty$  have been constructed as pairwise disjoint representatives of the classes preceding  $C(B)$ ; a member of  $C(B)$  shall be constructed in  $E^3 - (\bigcup_{i=1}^\infty A_i)$ . The element of  $C(B)$  in  $E^3 - (\bigcup_{i=1}^\infty A_i)$  will be constructed by adjusting  $B$  as described below.

By Theorem 1, there is no loss of generality in supposing that  $W(B) \subset E^3 - (\bigcup_{i=1}^\infty A_i)$ . By Theorem 1 of [16] we may also suppose that  $B$  is locally polygonal modulo  $W(B)$ . Then  $T(B)$  is the union of a countable number of straight line intervals. It is not hard to show that there is a homeomorphism of  $E^3$  onto itself which is fixed on  $W(B)$ , moves intervals of  $T(B)$  onto intervals, and end points of intervals of  $T(B)$  into  $E^3 - (\bigcup_{i=1}^\infty A_i)$ . We denote the image of  $B$  under this homeomorphism by  $B'$ .

Denote the intervals of  $T(B')$  by  $I_1, I_2, I_3, \dots$ . There is a null-sequence of polyhedral 2-spheres  $S_1, S_2, S_3, \dots$  such that

- (1)  $S_n \cap B'$  consists of the end points of  $I_n$ ,
- (2)  $\text{Int}(I_n) \subset \text{Int}(S_n)$ , and
- (3)  $S_n \cap S_m = \emptyset$  if  $I_n \cap I_m = \emptyset$ ,  $S_n \cap S_m$  is the common end point of  $I_n$  and  $I_m$  if  $I_n \cap I_m \neq \emptyset$ .

For each  $n$ , there is a plane  $P_n$  containing  $I_n$  and intersecting each element of  $\{A_i\}_{i=1}^\infty$  in at most a compact 0-dimensional set. Denote the end points of  $I_n$  by  $a_n$  and  $b_n$ . Let  $H_n$  be an arc from  $a_n$  to  $b_n$  lying in  $P_n \cap (\text{Int}(S_n) \cup \{a_n, b_n\})$  and so that  $(\text{Int}(H_n)) \cap I_n = \emptyset$ . By Lemma 2, there is an arc  $K_n$  in  $P_n$  so that (1)  $K_n \cap (\bigcup_{i=1}^\infty A_i) = \emptyset$ , and (2)  $\text{Int}(K_n) \subset \text{Int}(I_n \cup H_n)$ . It is clear that the arc  $(B - \bigcup_{n=1}^\infty I_n) \cup (\bigcup_{n=1}^\infty K_n)$  lies in  $C(B)$  and misses  $\bigcup_{i=1}^\infty A_i$ .

**5. Piercing 2-spheres with tame arcs.** The 2-sphere  $S$  in  $E^3$  is said to be *pierced* by the arc  $ab$  at the point  $p \in S$  if

- (1)  $S \cap (ab) = p$ ,
- (2)  $p \in \text{Int}(ab)$ , and
- (3)  $a$  and  $b$  lie in different complementary domains of  $S$ .

Since each point of  $S$  is arcwise accessible from both  $\text{Int}(S)$  and  $\text{Ext}(S)$ ,  $S$  can be pierced by an arc at each of its points. However, an example described in [10] shows that there may be points of  $S$  at which it is impossible to pierce  $S$  with a tame arc.

In [8], Bing showed that each 2-sphere in  $E^3$  can be pierced by a tame arc. David Gillman showed, in [13], that the set of points at which a 2-sphere cannot be pierced by a tame arc is a 0-dimensional  $F_\sigma$  set.

Theorem 4 below provides an affirmative answer to a question raised by Bing in [8].

**THEOREM 4.** *Given a 2-sphere  $S$  in  $E^3$ , there is a family  $F$  of mutually exclusive tame arcs such that for each point of  $S$  at which  $S$  can be pierced by a tame arc, there is a member of  $F$  piercing there.*

**Proof.** The collection  $F$  is to be constructed in much the same way as the collection  $\{A_\alpha\}$  was constructed in the proof of Theorem 3. The crucial step is in proving that the following statement is true: If  $\{A_i\}_{i=1}^\infty$  is a countable collection of pairwise disjoint tame arcs each of which pierces  $S$ , and  $p$  is a point of  $S - (\bigcup_{i=1}^\infty A_i)$  at which  $S$  can be pierced by a tame arc, then there is a tame arc  $A \subset E^3 - (\bigcup_{i=1}^\infty A_i)$  which pierces  $S$  at  $p$ .

Under the above conditions, let  $ab$  be a tame arc which pierces  $S$  at  $p$  and whose end points,  $a$  and  $b$ , lie in  $E^3 - (\bigcup_{i=1}^\infty A_i)$ . The point  $a$  is supposed, without loss of generality, to lie in  $\text{Ext}(S)$ . Let  $h: E^3 \rightarrow E^3$  be a homeomorphism of  $E^3$  onto itself carrying  $ab$  onto a straight line interval. There is a plane  $P$  containing  $h(ab)$  so that  $P$  intersects each element of  $\{A_i\}_{i=1}^\infty$  in at most a 0-dimensional set.

Let  $A'$  be an arc in  $P \cap (\text{Ext}(h(S)))$  from  $h(a)$  to  $h(p)$  so that  $A' \cap (\bigcup_{i=1}^{\infty} h(A_i)) = \emptyset$  and let  $A''$  be an arc in  $P \cap (\text{Int}(h(S)))$  from  $h(b)$  to  $h(p)$  so that  $A'' \cap (\bigcup_{i=1}^{\infty} h(A_i)) = \emptyset$ . Since  $A' \cup A''$  lies in  $P$ ,  $A' \cup A''$  is a tame arc which pierces  $h(S)$  at  $h(p)$  and which lies in  $E^3 - (\bigcup_{i=1}^{\infty} h(A_i))$ . Let  $A = h^{-1}(A' \cup A'')$ .

If  $S$  is an  $(n-1)$ -sphere in  $E^n$ , and  $n > 3$ , then  $S$  can be pierced by a tame arc at each of its points. From the above proof we obtain the following

**THEOREM 5.** *If  $S$  is an  $(n-1)$ -sphere in  $E^n$ , and  $n > 3$ , then there is a family  $F$  of mutually exclusive tame arcs such that  $S$  is pierced at each of its points by a member of  $F$ .*

#### REFERENCES

1. W. R. Alford and B. J. Ball, *Some almost polyhedral wild arcs*, Duke Math. J. **30** (1963), 33–38.
2. Steve Armentrout, *Decompositions of  $E^3$  with a compact 0-dimensional set of nondegenerate elements*, Trans. Amer. Math. Soc. **123** (1966), 165–177.
3. B. J. Ball, "Penetration indices and applications" in *Topology of 3-manifolds and related topics*, Prentice-Hall, Englewood Cliffs, N. J., 1962, pp. 37–39.
4. R. H. Bing, *Locally tame sets are tame*, Ann. of Math. (2) **59** (1954), 145–158.
5. ———, *Tame Cantor sets in  $E^3$* , Pacific J. Math. **11** (1961), 435–446.
6. ———,  *$E^3$  does not contain uncountably many mutually exclusive wild surfaces*, Abstract 63-801t, Bull. Amer. Math. Soc. **63** (1957), 404.
7. ———, *Each disk in  $E^3$  contains a tame arc*, Amer. J. Math. **84** (1962), 583–590.
8. ———, *Each disk in  $E^3$  is pierced by a tame arc*, Amer. J. Math. **84** (1962), 591–599.
9. Ralph H. Fox, *A remarkable simple closed curve*, Ann. of Math. (2) **50** (1949), 264–265.
10. Ralph H. Fox and Emil Artin, *Some wild cells and spheres in three-dimensional space*, Ann. of Math. (2) **49** (1948), 979–990.
11. R. H. Fox and O. G. Harrold, "The wilder arcs" in *Topology of 3-manifolds and related topics*, Prentice-Hall, Englewood Cliffs, N. J., 1962, pp. 184–187.
12. C. H. Giffen, *Uncountably many inequivalent nearly tame arcs*, Abstract 607-19, Notices Amer. Math. Soc. **10** (1963), 666.
13. David S. Gillman, *Side approximation, missing an arc*, Amer. J. Math. **85** (1963), 459–476.
14. J. G. Hocking and G. S. Young, *Topology*, Addison-Wesley, Reading, Mass., 1961.
15. H. W. Lambert and R. B. Sher, *Point-like 0-dimensional decompositions of  $S^3$* , Pacific J. Math. **24** (1968), 511–518.
16. E. E. Posey, *Proteus forms of wild and tame arcs*, Duke Math. J. **31** (1964), 63–72.
17. R. B. Sher, *A note on an example of Stallings*, Proc. Amer. Math. Soc. **19** (1968), 619–620.
18. ———, *Concerning wild Cantor sets in  $E^3$* , Proc. Amer. Math. Soc. **19** (1968), 1195–1200.
19. John R. Stallings, *Uncountably many wild disks*, Ann. of Math. (2) **71** (1960), 185–186.

THE UNIVERSITY OF GEORGIA,  
ATHENS, GEORGIA