

TWO NOTES ON RECURSIVE FUNCTIONS AND REGRESSIVE ISOLS⁽¹⁾

BY
JOSEPH BARBACK

Introduction. The theory of regressive isols was introduced by J. C. E. Dekker in [7]. The results that we wish to present in this paper belong to this theory and is a continuation of some of our studies in [1], [3] and [4]. We will assume that the reader is familiar with the terminology and some of the main results of the papers listed as references. We let E denote the collection of all nonnegative integers (*numbers*), Λ the collection of all isols, Λ^* the collection of all isolic integers, and $\Lambda_{\mathbb{R}}$ the collection of all regressive isols. If f is a function from a subset of E into E then δf will denote its domain and ρf its range. Let u_n and v_n be two one-to-one functions from E into E . Then $u_n \leq^* v_n$, if there is a partial recursive function f such that

$$(1) \quad \rho u \subseteq \delta f \quad \text{and} \quad (\forall n)[f(u_n) = v_n].$$

In addition, u_n and v_n are said to be *recursively equivalent* (denoted $u_n \simeq v_n$), if there is a one-to-one partial recursive function f such that (1) holds. It is easy to see that

$$u_n \simeq v_n \Rightarrow \rho u_n \simeq \rho v_n.$$

Also, it can be shown [8], that

$$(2) \quad u_n \simeq v_n \Leftrightarrow (u_n \leq^* v_n \text{ and } v_n \leq^* u_n).$$

Let α and β be two infinite subsets of E . Then $\alpha \leq^* \beta$, if there is a partial recursive function f such that $\alpha \subseteq \delta f$, $f(\alpha) = \beta$ and f is one-to-one on α . If α and β are each isolated sets then the following is true [8, Proposition P9.(b)],

$$(3) \quad (\alpha \leq^* \beta \text{ and } \beta \leq^* \alpha) \Leftrightarrow \alpha \simeq \beta.$$

Let α and β be infinite and isolated regressive sets, and let a_n and b_n be any regressive functions that range over α and β respectively. Then

$$(4) \quad \alpha \leq^* \beta \Leftrightarrow a_n \leq^* b_n,$$

Received by the editors April 23, 1968 and, in revised form, October 2, 1968.

⁽¹⁾ The author gratefully acknowledges conversations about some of the topics in the paper with Professors Dekker, Ellentuck, Hasset, Myhill, Nerode and Sansone. The author is an NIH Postdoctoral Fellow in the Department of Statistics (Division of Biostatistics), Stanford University, Stanford, California. Special thanks are expressed for comments of the referee.

and

$$(5) \quad \alpha \simeq \beta \Leftrightarrow a_n \simeq b_n.$$

Property (4) is [8, Proposition P11], and (5) follows directly from (2), (3) and (4). Let A and B be two isols. Then $A \leq^* B$, if either A is a finite isol and $A \leq B$, or both A and B are infinite isols and there are sets α and β belonging to A and B respectively, such that $\alpha \leq^* \beta$. It is readily seen that if both A and B are infinite isols, then $A \leq^* B$ if and only if $\alpha \leq^* \beta$ for any sets α and β belonging to A and B respectively. Combining this property with (4), it follows that in the special case both A and B are infinite regressive isols, then $A \leq^* B$ if and only if $a_n \leq^* b_n$ for all regressive functions a_n and b_n that range over sets belonging to A and B respectively.

A. Nerode associated with each recursive function $f: E^n \rightarrow E$, a function $D_f: \Lambda^n \rightarrow \Lambda^*$, called the *canonical extension of f to Λ^n* . A discussion of this extension procedure is given in [9, §2] and [15, §4]. A function $f: E \rightarrow E$ is said to be *increasing*, if for any numbers x and y , $x < y$ implies $f(x) \leq f(y)$. In the special case that a recursive function $f: E \rightarrow E$ is combinatorial, then $D_f: \Lambda \rightarrow \Lambda$, [15, Theorem 6]; and if f is increasing, then $D_f: \Lambda_R \rightarrow \Lambda_R$, [1, Theorem 4]. In [8], J. C. E. Dekker associated with each pair of regressive isols (A, B) , a particular regressive isol $\min(A, B)$ called the *minimum of A and B* . The minimum function has the following properties,

$$(6) \quad \min(A, B) = \min(B, A),$$

$$(7) \quad \min(A, B) = A \Leftrightarrow A \leq^* B.$$

Because there exist regressive isols that are not comparable relative to the relation \leq^* , it can happen that $\min(A, B)$ assumes neither the value A nor the value B . Let $D_{\min}(A, B)$ denote the canonical extension to Λ^2 of the ordinary $\min(a, b)$ function from E^2 into E . Then, by [4, p. 15],

$$(8) \quad \min(A, B) = D_{\min}(A, B), \quad \text{for } A, B \in \Lambda_R.$$

Properties of regressive isols that are related to the function $\min(A, B)$ and the relation \leq^* were first studied in [8], and later in [3]. In the first note of this paper we wish to present some additional properties of this kind. We also consider the ordinary recursive function $\text{maximum}(a, b)$ for pairs of numbers, and its canonical extension to Λ^2 . Some of the main results of the first note are as follows: Let A and B be two regressive isols. Let $f: E \rightarrow E$ be an increasing recursive function, and let D_f be its canonical extension to Λ . Let $\max(A, B)$ denote the canonical extension to Λ^2 of the recursive function $\text{maximum}(a, b)$. Then

$$\min(A, B) \leq^* A + B,$$

$$D_f(\min(A, B)) = \min(D_f(A), D_f(B)),$$

$$A \leq^* B \Rightarrow D_f(A) \leq^* D_f(B).$$

It need not be true that $\max(A, B)$ will be a regressive isol. However, in the special case that $A+B$ is a regressive isol then $\max(A, B)$ will also be a regressive isol. In this special case the following properties are also shown to be true,

$$\begin{aligned} \min(A, B) &\leq A+B, \\ D_f(\max(A, B)) &= \max(D_f(A), D_f(B)), \\ D_f(A) + D_f(B) &= D_f(\min(A, B)) + D_f(\max(A, B)), \\ D_f(A) \cdot D_f(B) &= D_f(\min(A, B)) \cdot D_f(\max(A, B)). \end{aligned}$$

The last two identities turn out to be of interest for the following reason; it is possible to choose regressive isols A and B such that $A+B$ is a regressive isol and such that the four (regressive) isols $D_f(A)$, $D_f(B)$, $D_f(\min(A, B))$, and $D_f(\max(A, B))$ are pairwise distinct.

In part II of the paper we study some properties dealing with regressive isols that are prime. Prime isols were first introduced and studied by J. C. E. Dekker and J. Myhill in [10]. Let $w(x)$ denote the strictly increasing recursive function that ranges over the collection of all prime numbers, and let D_w denote its canonical extension to Λ . Because w is an increasing recursive function, D_w will map Λ_R into Λ_R . In the second note we show that D_w maps regressive isols onto regressive isols that are prime, and also that there are prime regressive isols that do not belong to $D_w(\Lambda_R)$. Our main result gives the existence of four distinct prime regressive isols U_1 , U_2 , V_1 and V_2 such that

$$U_1, U_2, V_1, V_2 \in D_w(\Lambda_R), \quad U_1 \cdot U_2 \in \Lambda_R, \quad \text{and} \quad U_1 \cdot U_2 = V_1 \cdot V_2.$$

It follows therefore, that for regressive isols a unique factorization into primes is not possible.

We wish to acknowledge that throughout the paper an especially important and useful tool is the application of some of the well-known metatheorems, concerning recursive functions and their canonical extensions and isols, of A. Nerode [17].

I. ON PROPERTIES OF REGRESSIVE ISOLS

1. **A property of the relation \leq^* .** In [16], A. Nerode introduced an extension procedure for recursive relations on numbers to relations on isols. This extension procedure is also discussed by E. Ellentuck in [11]. In this section, we wish to prove that between two regressive isols, the extension to the isols of the ordinary relation \leq between two numbers, is equivalent to the relation \leq^* . We first describe how the extension to the isols of a recursive binary relation can be characterized. Let $\Delta \subseteq E^2$ be a recursive binary relation, and let $f, g: E^2 \rightarrow E$ be any recursive functions such that

$$\Delta = \{(x, y) \mid x, y \in E \text{ and } f(x, y) = g(x, y)\}.$$

Then the extension of the relation Δ to Λ can be characterized as the set

$$(1.1) \quad \Delta_{\Lambda} = \{(X, Y) \mid X, Y \in \Lambda \text{ and } D_r(X, Y) = D_g(X, Y)\}.$$

Let Δ_R denote the binary relation on Λ_R that is defined by

$$(1.2) \quad \Delta_R = \Delta_{\Lambda} \cap \Lambda_R^2.$$

THEOREM 1.1. *Let \leq_{Λ} denote the extension to Λ of the ordinary relation \leq between two numbers. Let \leq_R denote the relation on Λ_R obtained from \leq_{Λ} by (1.2). Then for all regressive isols A and B ,*

$$A \leq_R B \Leftrightarrow A \leq^* B.$$

Proof. Let Δ denote the relation \leq . Then

$$\begin{aligned} \Delta &= \{(x, y) \mid x, y \in E \text{ and } x \leq y\} \\ &= \{(x, y) \mid x, y \in E \text{ and } \min(x, y) = x\}. \end{aligned}$$

Therefore

$$(1.3) \quad \Delta_{\Lambda} = \{(X, Y) \mid X, Y \in \Lambda \text{ and } D_{\min}(X, Y) = X\}.$$

Combining (7), (8), (1.2) and (1.3) we see that

$$\begin{aligned} \Delta_R &= \{(A, B) \mid A, B \in \Lambda_R \text{ and } D_{\min}(A, B) = A\} \\ &= \{(A, B) \mid A, B \in \Lambda_R \text{ and } \min(A, B) = A\} \\ &= \{(A, B) \mid A, B \in \Lambda_R \text{ and } A \leq^* B\}. \end{aligned}$$

The desired result follows directly from the last equality, and this completes the proof.

2. Some fundamental properties of $\min(A, B)$. In [8], Dekker proved the following three properties;

$$(2.1) \quad A \leq B, \quad B \in \Lambda_R \Rightarrow A \in \Lambda_R,$$

$$(2.2) \quad A + B \in \Lambda_R \Leftrightarrow A \cdot B \in \Lambda_R, \quad \text{when } A, B \geq 1,$$

$$(2.3) \quad (\exists A, B \in \Lambda_R)[A + B \notin \Lambda_R].$$

In contrast to the isols, the regressive isols are neither closed under addition nor multiplication. One of the main goals of [3] was to try to obtain useful necessary conditions and sufficient conditions on two regressive isols for their sum to be a regressive isol. The next two theorems that we wish to prove are in a manner related to this goal.

THEOREM 2.1. *Let A and B be any regressive isols. Then*

$$(a) \quad \min(A, B) \leq^* A + B,$$

$$(b) \quad A + B \in \Lambda_R \Rightarrow \min(A, B) \leq A + B.$$

THEOREM 2.2. *There are (infinite) regressive isols A and B , for which the relation $\min(A, B) \leq A + B$ is not true.*

REMARK. Part (a) of Theorem 2.1 is a result due to W. Richter. Part (b) of Theorem 2.1 was stated without proof in [3], and its proof is related to the proof of [3, Theorem 1.2].

Proof of Theorem 2.1. If either of the regressive isols A or B is finite, then the value of $\min(A, B)$ will equal A or equal B . In this event both parts of the theorem are readily seen to be true.

Let us assume now that both A and B are infinite regressive isols. Let a_n and b_n be regressive functions that range over the sets α and β respectively, and with $\alpha \in A$, $\beta \in B$ and $\alpha | \beta$. Then $\alpha + \beta \in A + B$.

Re (a). Let $d_n = j(a_n, b_n)$ for $n \in E$. Then d_n is a regressive function and will range over a set in $\min(A, B)$. Let ∂ denote the range of d_n . Consider the mapping from ∂ onto $\alpha + \beta$ given by

$$(*) \quad \begin{aligned} d_n = j(a_n, b_n) &\rightarrow a_k, & \text{if } n = 2k, \\ &\rightarrow b_k, & \text{if } n = 2k + 1. \end{aligned}$$

Taking into account that j is a one-to-one recursive function and that each of the functions a_n , b_n and d_n is regressive, it is readily seen that the mapping denoted by (*) has a partial recursive extension. This property implies that $\partial \leq^* \alpha + \beta$, and therefore also that $\min(A, B) \leq^* A + B$.

Re (b). Assume $A + B \in \Lambda_R$. Then $\alpha + \beta$ will be an (infinite) regressive set. Let c_n be a regressive function that ranges over $\alpha + \beta$, and let $p(x)$ be a (partial recursive) regressing function for c_n . Let p^* be a partial recursive function associated with p that has the property, $p^*(c_n) = n$ for $n \in E$. We note that for each number $n \in E$, exactly one of the two relations $p^*(a_n) < p^*(b_n)$ or $p^*(b_n) < p^*(a_n)$ will be true. Let

$$\delta = \{x \mid (x = a_n \text{ and } p^*(b_n) < p^*(a_n)) \vee (x = b_n \text{ and } p^*(a_n) < p^*(b_n))\}.$$

Then $\delta \subseteq \alpha + \beta$ and for each number $n \in E$, exactly one of the two numbers a_n or b_n will belong to δ .

We now verify,

$$(b.1) \quad \delta | (\alpha + \beta) - \delta,$$

$$(b.2) \quad \delta \in \min(A, B).$$

For (b.1). Let $x \in \alpha + \beta$. We wish to determine whether or not $x \in \delta$. Since α and β are separated sets we can find out whether $x \in \alpha$ or $x \in \beta$. First suppose that $x \in \alpha$. Because a_n and c_n are regressive functions, we can compute the numbers u and v such that $x = a_u = c_v$. In view of the definition of δ , we see that

$$(2.4) \quad a_u \in \delta \Leftrightarrow b_u \in (c_0, \dots, c_{v+1}) = \{p^r(x) \mid 1 \leq r \leq v\}.$$

The members of the set on the right side of (2.4) can be effectively obtained from x , since p is a partial recursive function. In addition, using again the separability of the sets α and β , and the regressive property of the function b_n , we can determine

whether or not the number b_u belongs to the set (c_0, \dots, c_{v-1}) . In view of (2.4), it follows that we can therefore also determine whether or not the number x belongs to δ . We recall that we had assumed that $x \in \alpha$. It can be readily verified in a similar fashion that one can determine whether or not $x \in \delta$, in the event $x \in \beta$. We can conclude from these remarks that the two sets δ and $(\alpha + \beta) - \delta$ are separated.

For (b.2). Let the functions u_n and v_n be defined by

$$\begin{aligned} u_n &= a_n, & \text{if } a_n \in \delta, \\ &= b_n, & \text{if } b_n \in \delta; \end{aligned}$$

and

$$\begin{aligned} v_n &= a_n, & \text{if } a_n \notin \delta, \\ &= b_n, & \text{if } b_n \notin \delta. \end{aligned}$$

Then each of the functions u_n and v_n is everywhere defined and one-to-one; also u_n ranges over the set δ and v_n ranges over the set $(\alpha + \beta) - \delta$. Let f be the partial function with domain δ and defined by, $f(u_n) = v_n$. In the course of proving [3, Theorem 1.2] it was verified that the mapping f will have a partial recursive extension. It follows from this property that

$$(2.5) \quad u_n \leq^* v_n.$$

Because $d_n = j(a_n, b_n)$ is a (regressive) function that ranges over a set in $\min(A, B)$, and u_n is a function that ranges over the set δ , to establish (b.2) we see that it is sufficient to verify

$$(2.6) \quad u_n \simeq j(a_n, b_n).$$

This will be our approach here. We first note that (2.6) is equivalent to the two relations

$$(2.7) \quad u_n \leq^* j(a_n, b_n),$$

and

$$(2.8) \quad j(a_n, b_n) \leq^* u_n.$$

The first of these follows directly from (2.5) and the definition of the functions u_n and v_n . To verify (2.8), we first recall that for each number n , either $u_n = a_n$ or $u_n = b_n$. It readily follows from our remarks in the proof of (b.1), that given both of the numbers a_n and b_n , we can effectively determine which one is u_n . Since both a_n and b_n can be found from the value of $j(a_n, b_n)$, it follows therefore that the mapping $j(a_n, b_n) \rightarrow u_n$ will have a partial recursive extension. This property is equivalent to the relation of (2.8). This verifies both (2.7) and (2.8); and as we noted earlier, together these establish both (2.6) and (b.2). This completes the proof.

REMARK. Regarding the proof of Theorem 2.1(b), we wish to note here some properties of the two functions u_n and v_n , that are of interest in the next section. These are

- (a) u_n and v_n range over separated sets,
- (b) u_n ranges over a set in $\min(A, B)$, and
- (c) u_n and v_n are each regressive functions.

Because the functions u_n and v_n range over the sets δ and $(\alpha + \beta) - \delta$ respectively, we see that (a) and (b) follow directly from (b.1) and (b.2) respectively. To verify (c), we first note that by combining (2.6) and the fact the function $d_n = j(a_n, b_n)$ is regressive, it follows that u_n will be a regressive function. We now want to show that the function v_n is also regressive, i.e., that the mapping $v_{n+1} \rightarrow v_n$ will have a partial recursive extension. For this purpose, let the value of v_{n+1} be given; we wish to find the value of v_n . We know that for $m \in E$,

$$\begin{aligned} v_m &= a_m, & \text{if } a_m \notin \delta, \\ &= b_m, & \text{if } b_m \notin \delta. \end{aligned}$$

Since α and β are separated sets we can effectively determine whether $v_{n+1} = a_{n+1}$ or $v_{n+1} = b_{n+1}$. Assume that $v_{n+1} = a_{n+1}$. Then, using the regressive property of the function a_x , we can find the value of a_n . In view of (b.1), we can then determine whether or not $a_n \in \delta$. If $a_n \notin \delta$, then $v_n = a_n$ and we are done. In view of (2.5), we see that if $a_n \in \delta$ we can then also find the value of v_n . We had assumed that $v_{n+1} = a_{n+1}$, and then showed that the value of v_n could be found. It is easy to see that we could also have found the value of v_n if it turned out that $v_{n+1} = b_{n+1}$. We can conclude from these remarks that the mapping $v_{n+1} \rightarrow v_n$, will have a partial recursive extension, and therefore also that v_n is a regressive function.

In [8], Dekker introduced the concept of the *degree of unsolvability of a regressive isol*. This is a useful notion and is employed in the proof of Theorem 2.2. We now recall, from [8], how it is defined. Every regressive isol contains at least one retraceable set. By [8, Corollary 2 of Proposition P14], any two retraceable sets that belong to the same isol [i.e., that are recursively equivalent] will have the same (Turing) degree of unsolvability. Let A be a regressive isol. Then its *degree of unsolvability*, denoted by Δ_A , is defined to be the (Turing) degree of unsolvability of any retraceable set belonging to A . In the special case A and B are infinite regressive isols, then the degree has the following properties [8],

- (i) $A + B \in \Lambda_R \Rightarrow \Delta_A = \Delta_B$,
- (ii) $A \leq B \Rightarrow \Delta_A = \Delta_B$,
- (iii) $\Delta_A \leq \Delta_{\min(A, B)}$ and $\Delta_B \leq \Delta_{\min(A, B)}$.

In addition, it was also proved in [8], that there exist infinite regressive isols that have degrees incomparable with respect to the relation \leq .

LEMMA 2.1. *Let A and B be infinite regressive isols. Then*

$$\min(A, B) \leq A + B \Rightarrow \Delta_A \leq \Delta_B \text{ or } \Delta_B \leq \Delta_A.$$

Proof. We note that $\min(A, B)$ is an infinite (regressive) isol, since each of the isols A and B is infinite. Assume that

$$(2.9) \quad \min(A, B) \leq A + B.$$

Combining (2.1) and the refinement property for recursive equivalence types, it follows from (2.9) that there will exist regressive isols M and N such that

$$M + N = \min(A, B), \quad M \leq A \text{ and } N \leq B.$$

Because $\min(A, B)$ is infinite, at least one of the two regressive isols M and N is also infinite. Let us assume that M is infinite. Then $M \leq A$ implies, by (ii), that $\Delta_M = \Delta_A$, and $M \leq \min(A, B)$ implies that $\Delta_M = \Delta_{\min(A, B)}$. Therefore $\Delta_A = \Delta_{\min(A, B)}$. Combining this identity with (iii), gives $\Delta_B \leq \Delta_A$. In a similar manner one can show that if N is infinite then $\Delta_A \leq \Delta_B$. This gives the desired conclusion that Δ_A and Δ_B are comparable degrees with respect to the \leq relation.

Proof of Theorem 2.2. Use Lemma 2.1 and the fact that there exist infinite regressive isols that have incomparable degrees of unsolvability.

REMARK. It is proven in [8] that there are infinite cosimple regressive isols that have incomparable degrees of unsolvability. Also, the minimum of two cosimple regressive isols is again a cosimple regressive isol. Thus, as a consequence of the previous lemma, we see that the following result is also true.

THEOREM. *There are (infinite) cosimple regressive isols A and B , for which the relation $\min(A, B) \leq A + B$ is not true.*

We state the next result without proof. It is a special case of a theorem of Nerode.

LEMMA A. *Let $f: E^n \rightarrow E$ and $g_1, \dots, g_n: E^m \rightarrow E$ be recursive functions. Let $f \circ g$ denote the composition function $f(g_1(X), \dots, g_n(X))$, for $X \in E^m$. Then $f \circ g$ is a recursive function, and*

$$D_{f \circ g}(T) = D_f(D_{g_1}(T), \dots, D_{g_n}(T)), \quad \text{for } T \in \Lambda^m \text{ and } D_{g_1}(T), \dots, D_{g_n}(T) \in \Lambda.$$

THEOREM 2.3. *Let $f: E \rightarrow E$ be an increasing recursive function, and A and B any regressive isols. Then*

- (a) $\min(D_f(A), D_f(B)) = D_f(\min(A, B))$,
- (b) $A \leq^* B \Rightarrow D_f(A) \leq^* D_f(B)$.

Proof. Because f is an increasing recursive function, we know that both $D_f(A)$ and $D_f(B)$ will be regressive isols; the identity of part (a) is therefore meaningful.

Re (a). Let $\min(a, b)$ denote the familiar minimum function from E^2 into E . Then

$$(2.10) \quad \min(f(a), f(b)) = f(\min(a, b)), \quad \text{for } a, b \in E.$$

Application of Lemma A to the identity of (2.10) gives

$$(2.11) \quad D_{\min}(D_f(A), D_f(B)) = D_f(D_{\min}(A, B)).$$

Finally, combining (8) and (2.11) yields (a).

Re (b). Assume that $A \leq^* B$. Then $\min(A, B) = A$ and therefore also

$$(2.12) \quad D_f(\min(A, B)) = D_f(A).$$

Combining part (a) and (2.12) gives

$$\min(D_f(A), D_f(B)) = D_f(A);$$

and this implies that $D_f(A) \leq^* D_f(B)$. This verifies (b) and completes the proof.

REMARK. As an illustration of Theorem 2.3 we mention

$$\min(2^A, 2^B) = 2^{\min(A, B)}$$

and

$$A \leq^* B \Rightarrow 2^A \leq^* 2^B, \text{ for } A, B \in \Lambda_R.$$

The reader is referred to [8, p. 365] for some additional comments that are related to Theorem 2.3. In the special case that f is a strictly increasing recursive function, then the converse of Theorem 2.3(b) is also true, and this will be our next result. We need the following lemma; it is a special case of a theorem of Nerode and we shall state it without proof.

LEMMA B. Let f, g, h and k be recursive functions from E^n into E . If

$$(\forall t \in E^n)[f(t) = g(t) \Rightarrow h(t) = k(t)],$$

then

$$(\forall T \in \Lambda^n)[D_f(T) = D_g(T) \Rightarrow D_h(T) = D_k(T)].$$

THEOREM 2.4. Let $f: E \rightarrow E$ be a strictly increasing recursive function, and A and B any regressive isols. Then $D_f(A) \leq^* D_f(B) \Rightarrow A \leq^* B$.

Proof. Because f is a strictly increasing function, it follows that for any numbers $a, b \in E$, $f(a) \leq f(b) \Rightarrow a \leq b$, and therefore also,

$$(2.13) \quad \min(f(a), f(b)) = f(a) \Rightarrow \min(a, b) = a.$$

In view of Lemmas A and B, it follows from (2.13) that

$$(2.14) \quad \min(D_f(A), D_f(B)) = D_f(A) \Rightarrow \min(A, B) = A.$$

Because f is an increasing recursive function, we know that $D_f(A)$ and $D_f(B)$ are regressive isols. Therefore, by (7), we see that (2.14) is equivalent to $D_f(A) \leq^* D_f(B) \Rightarrow A \leq^* B$, and this is the desired result.

REMARK. Let $f: E \rightarrow E$ be a strictly increasing recursive function and let A and B be any regressive isols. By Theorems 2.3(b) and 2.4 we know that

$$A \leq^* B \Rightarrow D_f(A) \leq^* D_f(B),$$

and

$$D_f(A) \leq^* D_f(B) \Rightarrow A \leq^* B.$$

We wish to consider the following two similar assertions,

$$(I) \quad A \leq B \Rightarrow D_f(A) \leq D_f(B),$$

$$(II) \quad D_f(A) \leq D_f(B) \Rightarrow A \leq B.$$

Concerning (I), let the function \tilde{f} be defined by, $\tilde{f}(n) = f(n+1) - f(n)$ for $n \in E$. Then \tilde{f} will be a recursive function, and it is called the *first difference function* of f .

In [20], Sansone proved that if \tilde{f} is an eventually increasing function then statement (I) is true. He also showed that in the event \tilde{f} is not an eventually increasing function that not only will (I) not be a theorem, but that there will even exist an (infinite) regressive isol T such that $D_f(T) \leq D_f(T+1)$ is not true. It can be readily shown that in the special case that f is a combinatorial function, then its first difference function \tilde{f} is also a combinatorial function; and therefore also an increasing function. In view of these remarks, we see that while statement (I) is not true for all strictly increasing recursive functions f , it is true in the particular case that f is combinatorial.

Concerning statement (II), (II) will be true in the special case that f is a linear function, i.e., $f(x) = ax + b$, where a and b denote some particular constants. This property follows from a result of Nerode (see [12, p. 1]). On the other hand, Ellentuck proved in [12, Theorem 3] that if f is a combinatorial but nonlinear function, then there are isols A and B for which statement (II) is false. We wish to note that such isols A and B can be chosen to be regressive. [To obtain this last property, simply modify the proof of [12, Theorem 3] by selecting U to be a universal regressive isol. The existence of universal isols that are regressive was recently proved by Ellentuck in some notes not yet published.]

3. $\max(A, B)$. Let the ordinary recursive function, maximum (a, b) , be denoted by $\max(a, b)$ for pairs of numbers, and let $\max(A, B)$ denote its canonical extension, $D_{\max}(A, B)$, to Λ^2 . It is easy to show that $\max(a, b)$ is not an almost combinatorial function; and therefore $\max(A, B)$ will not map Λ^2 into Λ . It is also true that $\max(A, B)$ will not map Λ_R^2 into Λ_R ; M. Hassett proved this property, in [13]. However, in the special case that A and B are regressive isols such that their sum $A+B$ is also regressive, then $\max(A, B)$ as well as $\min(A, B)$ will be regressive isols. To prove this fact, first note that the statement

$$\min(a, b) + c = a + b \Rightarrow c = \max(a, b)$$

is valid in the domain E . Hence, by Lemma B,

$$(3.1) \quad \min(A, B) + C = A + B \Rightarrow C = \max(A, B), \quad \text{for } A, B \in \Lambda_R.$$

By Theorem 2.1(b),

$$(3.2) \quad \min(A, B) \leq A + B, \quad \text{when } A, B, A + B \in \Lambda_R.$$

Combining (3.1) and (3.2), we see that in the special case A, B and $A+B$ are regressive isols, then

$$\max(A, B) = (A + B) - \min(A, B), \quad \text{and} \quad \max(A, B) \in \Lambda_R.$$

NOTATION. For the remainder of this section we let A and B denote any regressive isols such that $A+B \in \Lambda_R$. By our previous remarks we then have that both $\min(A, B)$ and $\max(A, B)$ will be regressive isols.

REMARK. In view of (3.1), and properties of the minimum of two regressive isols, it follows that the values of $\min(A, B)$ and $\max(A, B)$ will be closely related. For example, it is readily shown that

$$(\alpha) \quad \max(A, B) = \max(B, A),$$

$$(\beta) \quad \max(A, B) \in (A, B) \Leftrightarrow \min(A, B) \in (A, B),$$

and

$$(\delta) \quad \max(A, B) = A \Leftrightarrow \min(A, B) = B \Leftrightarrow B \leq^* A.$$

In the course of proving Theorem 2.1(b) we assumed that A and B were infinite regressive isols such that $A + B \in \Lambda_R$, and then we proved that $\min(A, B) \leq A + B$. In doing this we defined two (one-to-one) functions u_n and v_n . In the remark following the proof of Theorem 2.1, we observed that both u_n and v_n are regressive functions, and that u_n ranges over a set in $\min(A, B)$. In addition, the ranges of the functions u_n and v_n are separated sets. In view of (3.1), it follows from this fact that the (regressive) function v_n ranges over a set in $\max(A, B)$. If one combines these properties with the definitions of the functions u_n and v_n , then the following properties can be readily established:

$$(\gamma) \quad A, B \in \Lambda_R - E \Rightarrow \min(A, B), \quad \max(A, B) \in \Lambda_R - E,$$

$$(\epsilon) \quad A \leq^* \max(A, B) \quad \text{and} \quad B \leq^* \max(A, B),$$

$$(\zeta) \quad \min(A, B) = \max(A, B) \Leftrightarrow A = B,$$

and

$$(\eta) \quad (\forall T \in \Lambda_R)[A \leq^* T \text{ and } B \leq^* T \Leftrightarrow \max(A, B) \leq^* T].$$

We note that properties (ϵ) and (η) represent analogues for the function $\max(A, B)$ of two well-known properties of $\min(A, B)$, [8, p. 363]. The next theorem we state without proof, because each of its parts can be verified in a manner similar to that employed in the proof of Theorem 2.3(a).

THEOREM 3.1. *Let $f: E \rightarrow E$ be an increasing recursive function. Then*

$$(a) \quad \max(D_f(A), D_f(B)) = D_f(\max(A, B)),$$

$$(b) \quad \min(A, B) \cdot \max(A, B) = A \cdot B,$$

$$(c) \quad \min(D_f(A), D_f(B)) + \max(D_f(A), D_f(B)) = D_f(A) + D_f(B),$$

$$(d) \quad \min(D_f(A), D_f(B)) \cdot \max(D_f(A), D_f(B)) = D_f(A) \cdot D_f(B),$$

$$(e) \quad D_f(\min(A, B)) \cdot D_f(\max(A, B)) = D_f(A) \cdot D_f(B).$$

II. PRIME REGRESSIVE ISOLS

Let U be an isol; we recall, from [10], the following definitions,

$$U \text{ is multiple-free if } 2A \leq U \Rightarrow A \in E,$$

$$U \text{ is prime if } A|U \Rightarrow (A = 1 \text{ or } A = U).$$

Each of the results

(4.1) infinite multiple-free isols exist

and

(4.2) infinite multiple-free isols are prime

is proved in [10].

THEOREM 4.1. *There is an infinite regressive isol T that is multiple-free.*

Proof. This will be a constructive type of proof and we will use a technique introduced in the proof of [10, Theorem 95].

Let $\{p_i\}$ be an enumeration of all partial recursive functions of one variable that are one-to-one [i.e., that are one-to-one on their respective domains]. For each number i , we let p_i^{-1} denote the inverse function of p_i . We note that for each i , the function p_i^{-1} is also partial recursive and one-to-one. If p is any partial function, x any number, and α any set of numbers, then we will write $p(x) \notin \alpha$ to mean either, $p(x)$ is undefined, or $p(x)$ is defined and $p(x) \notin \alpha$. We shall now define a function t_n such that the recursive equivalence type $T = \text{Req } \rho t_n$ will satisfy the condition of the theorem.

Set $t_0 = 1$. Let $n \geq 1$ and assume that the values t_0, \dots, t_{n-1} have already been defined. Let

$$(4.3) \quad \pi_{n-1} = (t_0, \dots, t_{n-1}).$$

We define t_n by setting

$$(4.4) \quad t_n = j(t_{n-1}, u_n),$$

where the number u_n is given by,

$$(4.5) \quad u_n = (\mu n)[(\forall r \leq n)[p_r j(t_{n-1}, u) \notin \pi_{n-1} \text{ and } p_r^{-1} j(t_{n-1}, u) \notin \pi_{n-1}]].$$

In light of the fact that each of the functions p_i and j is one-to-one, it is readily seen that there will be a number u_n satisfying (4.5). This completes the definition of the function t_n . In almost the same fashion as in the proof of [1, Theorem 4], one can easily verify that the function t_n has the following properties,

$$(4.6) \quad t_n \text{ is everywhere defined and one-to-one,}$$

$$(4.7) \quad t_n \text{ is a retraceable function.}$$

Let

$$\tau = \rho t_n \quad \text{and} \quad T = \text{Req } \tau.$$

Then τ is an infinite retraceable set and T is an infinite regressive RET.

We now verify that,

- (a) T is multiple-free,
- (b) τ is an immune set,
- (c) T is an infinite multiple-free regressive isol.

Re (a). Let us assume otherwise, and let A denote an RET such that

$$A \text{ is infinite and } 2A \leq T.$$

Then both A and $2A$ will be (infinite) regressive RET's. In addition, there will exist infinite regressive sets $\alpha, \alpha^* \subseteq \tau$ such that,

$$(4.8) \quad \alpha \cap \alpha^* = \emptyset \quad \text{and} \quad \alpha, \alpha^* \in A.$$

Let u_n and v_n be one-to-one functions such that

$$(4.9) \quad \rho u_n \cap \rho v_n = \emptyset,$$

$$(4.10) \quad \rho t_{u_n} = \alpha \quad \text{and} \quad \rho t_{v_n} = \alpha^*,$$

and

$$(4.11) \quad t_{u_n} \text{ and } t_{v_n} \text{ are each regressive functions.}$$

Because $\alpha, \alpha^* \in A$, it follows that $\alpha \simeq \alpha^*$. Combining this fact with (4.10), (4.11) and [7, Proposition 3], it follows that the two regressive functions t_{u_n} and t_{v_n} are recursively equivalent; this means that there will exist a one-to-one partial recursive function p such that, for each number $n \in E$,

$$(4.12) \quad p(t_{u_n}) \text{ is defined, and } p(t_{u_n}) = t_{v_n}.$$

Such a function p will appear in our enumeration $\{p_i\}$; let i be one of its indices. In view of (4.12), it then follows that, for each number $n \in E$,

$$(4.13) \quad p_i(t_{u_n}) = t_{v_n} \quad \text{and} \quad p_i^{-1}(t_{v_n}) = t_{u_n}.$$

Let m be the smallest number such that $u_m, v_m > i$. By (4.9), we note that $u_m \neq v_m$. We consider two cases:

Case 1. $u_m < v_m$. In light of the definition of the function t_n , parts (4.3), (4.4) and (4.5), it follows then that

$$p_i^{-1}(t_{v_m}) \notin (t_0, \dots, t_{v_m-1}),$$

and therefore also

$$p_i^{-1}(t_{v_m}) \notin (t_{u_0}, \dots, t_{u_m}).$$

Hence $p_i^{-1}(t_{v_m}) \neq t_{u_m}$, which cannot be true in view of (4.13).

Case 2. $u_m > v_m$. Then, as in the previous case, it follows that

$$p_i(t_{u_m}) \notin (t_0, \dots, t_{u_m-1}),$$

and

$$p_i(t_{u_m}) \notin (t_{v_0}, \dots, t_{v_m}).$$

The last relation implies $p_i(t_{u_m}) \neq t_{v_m}$, and this also cannot be true in view of (4.13).

In view of our remarks in each of these cases, we see that there does not exist a one-to-one partial recursive function p that satisfies (4.12). We can therefore conclude that T is a multiple-free RET; and this verifies part (a).

Re (b). We have already noted that τ is an infinite retraceable set. It is well known that retraceable sets are recursive or immune. If τ were a recursive set, then the strictly increasing function that ranges over τ , namely t_n , would be a recursive function. In this event, it would then be an easy consequence that $t_{2n} \simeq t_{2n+1}$, which would imply that T is an even RET. However, this is not possible, since T is infinite and, by part (a), a multiple-free RET. We can conclude therefore that τ will be an immune set.

Re (c). Because $T = \text{Req } \tau$, and τ is an infinite retraceable and immune set, it follows that T is an infinite regressive isol. Combining this fact with part (a), gives that T is an infinite multiple-free regressive isol. This verifies part (c) and completes the proof of the theorem.

COROLLARY 4.1. *There exist infinite regressive isols that are prime.*

Proof. Combine Theorem 4.1 and (4.2).

REMARK. Let T be an infinite multiple-free regressive isol. Let $n \in E$. It can be readily verified that each isol of the form $T+n$ and $T-n$, is also regressive and multiple-free. In view of (4.2), we see then that each isol in the enumeration

$$\dots, T-2, T-1, T, T+1, T+2, \dots,$$

will be infinite, regressive and prime. It is interesting to note here that even though both T and $T+1$ will be infinite prime isols, the number 2 will be a factor of the product $T \cdot (T+1)$ [10, Theorem 73].

NOTATION AND REMARK. For the remainder of the paper we let $w(x)$ be the function given by $w(x) = x$ th prime number, i.e., $w(0) = 2$, $w(1) = 3, \dots$. Because $w(x)$ is a (strictly) increasing recursive function, its canonical extension D_w will map Λ_R into Λ_R . However, its extension will not map Λ into Λ . The reason for this is that $w(x)$ will not be an eventually combinatorial function [this property was mentioned to the author by A. Nerode].

THEOREM 4.2. *Let $A, B \in \Lambda_R$. Then*

- (a) $D_w(A)$ is a prime regressive isol,
- (b) $D_w(A) = D_w(B) \Rightarrow A = B$.

Proof. From the definition of the function $w(x)$ we note that each of the arithmetic statements

$$(4.14) \quad w(a) = (u+2) \cdot v \Rightarrow v = 1,$$

$$(4.15) \quad w(a) = w(b) \Rightarrow a = b,$$

is universally valid in E . Application of Lemmas A and B to (4.14) and (4.15), and substitution gives both

$$(4.16) \quad D_w(A) = (U+2) \cdot V \Rightarrow V = 1, \quad \text{for } U, V \in \Lambda,$$

and

$$(4.17) \quad D_w(A) = D_w(B) \Rightarrow A = B.$$

Part (b) follows directly from (4.17). To verify part (a), we first note that $D_w(A)$ is a regressive isol, because A is a regressive isol. To show that $D_w(A)$ is also prime, assume that

$$(4.18) \quad D_w(A) = T \cdot V$$

for two isols T and V . We wish to show that either $T=1$ or $V=1$. Because $D_w(A)$ will be nonzero it is readily seen that $T \neq 0$. If $T=1$, then we are done. Assume now that $T \neq 1$. Then $T \geq 2$, and we let $U=T-2$. Then (4.18) can be expressed by,

$$(4.19) \quad D_w(A) = (U+2) \cdot V.$$

Finally combining (4.16) and (4.19) implies that $V=1$. It follows therefore that $D_w(A)$ will be a prime regressive isol. This verifies (a) and completes the proof.

REMARK. We note that by combining Theorem 4.2 with the fact that there are c regressive isols, it follows that there will exist c prime regressive isols belonging to $D_w(\Lambda_R)$. We wish also to note that the prime regressive isols obtained from Corollary 4.1 [these were infinite multiple-free regressive isols] will not belong to $D_w(\Lambda_R)$. This property will follow from the next result and the fact that infinite multiple-free isols are neither even nor odd.

THEOREM 4.3. *Let $B \in \Lambda_R$. Then $D_w(B)=2$ if $B=0$, and $D_w(B)$ is an odd (regressive) isol if $B \geq 1$.*

Proof. If $B=0$, then the result is easy.

Assume now that $B \geq 1$. We note that for each number $a \in E$, $w(a+1)$ will be an odd (prime) number. Let the function $u(a)$ be defined by,

$$(4.20) \quad w(a+1) = 2u(a) + 1, \quad \text{for } a \in E.$$

It is readily seen that the function $u(a)$ is recursive and increasing. Therefore, its canonical extension D_u will map Λ_R into Λ_R . Combining Lemma A and (4.20), it follows that

$$(4.21) \quad D_w(A+1) = 2D_u(A) + 1, \quad \text{for } A \in \Lambda_R.$$

Because $B \in \Lambda_R$ and $B \geq 1$, it follows that $B-1 \in \Lambda_R$ and hence also that $D_u(B-1) \in \Lambda_R$. If we substitute $B-1$ for A in (4.21), we obtain $D_w(B)=2D_u(B-1)+1$, from which it follows that $D_w(B)$ will be an odd regressive isol.

LEMMA C. Let $f: E \rightarrow E$ be an increasing recursive function, and A and B any regressive isols. Then

$$A + B \in \Lambda_R \Rightarrow D_f(A) + D_f(B) \in \Lambda_R.$$

Proof. Assume that $A + B$ is a regressive isol. Let $g: E \rightarrow E$ be an increasing recursive function chosen such that the recursive function $h(x) = f(x) + g(x)$, will be combinatorial. By Lemma A it follows that

$$(4.22) \quad D_h(A) = D_f(A) + D_g(A), \quad D_h(B) = D_f(B) + D_g(B).$$

Because each of the recursive functions f , g and h is increasing, it follows that each of the isols appearing in (4.22) is regressive. In addition, $D_h(A + B)$ will also be a regressive isol, since $A + B \in \Lambda_R$. Consider the following implications,

$$\begin{aligned} A \leq A + B &\Rightarrow D_h(A) \leq D_h(A + B) \\ &\Rightarrow D_f(A) + D_g(A) \leq D_h(A + B) \\ &\Rightarrow D_f(A) \leq D_h(A + B). \end{aligned}$$

The first implication follows from [20, Theorem 2] and the fact that since h is a combinatorial function its first difference function will also be combinatorial and hence increasing. The second implication follows from (4.22), and the third one is clear. Clearly $A \leq A + B$, and therefore,

$$(4.23) \quad D_f(A) \leq D_h(A + B).$$

In a similar manner it can be shown that

$$(4.24) \quad D_f(B) \leq D_h(A + B).$$

Since $D_h(A + B) \in \Lambda_R$, it follows from (2.1), (4.23) and (4.24) that

$$D_f(A) + D_f(B) \in \Lambda_R;$$

and this is the desired result.

THEOREM 4.4. There exist distinct prime regressive isols U_1 , U_2 , V_1 and V_2 such that

- (a) $U_1, U_2, \hat{V}_1, V_2 \in D_w(\Lambda_R)$,
- (b) $U_1 + U_2 = V_1 + V_2$,
- (c) $U_1 \cdot U_2 = V_1 \cdot V_2$,
- (d) $U_1 + U_2, U_1 \cdot U_2 \in \Lambda_R$.

Proof. By [3, Theorem 2.2], there will exist two (infinite) regressive isols that are not \leq^* comparable and yet whose sum is a regressive isol. Let A and B denote two regressive isols of this kind. We now verify that the regressive isols

$$\begin{aligned} U_1 &= D_w(A), & V_1 &= D_w(\min(A, B)), \\ U_2 &= D_w(B), & V_2 &= D_w(\max(A, B)), \end{aligned}$$

will satisfy the conditions of the theorem. It follows from (6), (7), (α), (β) and (ζ), and the fact that A and B are not \leq^* comparable that the four regressive isols A , B , $\min(A, B)$ and $\max(A, B)$ are distinct. By Theorem 4.2(b), this also means that the four regressive isols U_1 , U_2 , V_1 and V_2 are also distinct. In light of the definitions of U_1 , U_2 , V_1 and V_2 , we see that part (a) follows directly. Part (b) follows from Theorems 2.3(a) and 3.1(c), and part (c) follows from parts (a) and (d) of Theorem 3.1. Lastly, combining Lemma C and (2.2) gives part (d).

REMARK. We wish to give an analogue for regressive isols of the following well-known theorem from number theory, called Bertrand's conjecture and proved by Chebyshev in 1850; see [14, p. 108]:

THEOREM A. *For each number $a \in E$, there is a prime p such that, $a < p \leq 2a$.*

We will need two lemmas; the second of these is proved in [6].

LEMMA 4.1. *Let $f, g: E \rightarrow E$ be increasing recursive functions. If*

$$(\forall t \in E)[f(t) \leq g(t)],$$

then

$$(\forall T \in \Lambda_R)[D_f(T) \leq^* D_g(T)].$$

Proof. Assume that $f(t) \leq g(t)$, for each number t . Then

$$\min(f(t), g(t)) = f(t), \quad \text{for } t \in E.$$

Therefore, by Lemma A, (7) and the fact that the canonical extensions of increasing recursive functions map Λ_R into Λ_R , it follows that

$$\min(D_f(T), D_g(T)) = D_f(T), \quad \text{for } T \in \Lambda_R,$$

and

$$D_f(T) \leq^* D_g(T), \quad \text{for } T \in \Lambda_R.$$

LEMMA 4.2 [6, THEOREM 1(a)]. *Let $f, g: E \rightarrow E$ be increasing recursive functions. Then*

$$f(E) \subseteq g(E) \Rightarrow D_f(\Lambda_R) \subseteq D_g(\Lambda_R).$$

THEOREM 4.5. *For each regressive isol A , there is a prime regressive isol P such that $P \in D_w(\Lambda_R)$ and $A \leq^* P \leq^* 2A$.*

Proof. In view of Theorem A, it is easily seen that there will exist an increasing recursive function $p(a)$ such that, for each number $a \in E$, $a \leq p(a) \leq 2a$, and $p(a)$ is a prime number.

Therefore, by Lemmas 4.1 and 4.2, it follows that for each regressive isol A ,

$$A \leq^* D_p(A) \leq^* 2A \quad \text{and} \quad D_p(A) \in D_w(\Lambda_R).$$

Since $D_w(\Lambda_R)$ consists entirely of prime regressive isols, the desired result follows.

REMARK. Regarding Theorem 4.5, we do not know whether or not it is the case

that for each regressive isol A , there is a prime regressive isol P such that $P \in D_w(\Lambda_R)$ and $A \leq P \leq 2A$.

REFERENCES

1. J. Barback, *Recursive functions and regressive isols*, Math. Scand. **15** (1964), 29–42.
2. ———, *A note on regressive isols*, Notre Dame J. Formal Logic **7** (1966), 203–205.
3. ———, *Two notes on regressive isols*, Pacific J. Math. **16** (1966), 407–420.
4. ———, *Double series of isols*, Canad. J. Math. **19** (1967), 1–15.
5. ———, *Regressive upper bounds*, Seminario Matematico **39** (1967), 248–272.
6. ———, *On recursive sets and regressive isols*, Michigan Math. J. **15** (1968), 27–32.
7. J. C. E. Dekker, “Infinite series of isols” in *Recursive function theory*, Proc. Sympos. Pure Math., Vol. 5, Amer. Math. Soc., Providence, R. I., 1962, pp. 77–96.
8. ———, *The minimum of two regressive isols*, Math. Z. **83** (1964), 345–366.
9. ———, “Regressive isols” in *Sets, models and recursion theory*, North-Holland, Amsterdam, 1967, pp. 272–296.
10. J. C. E. Dekker and J. Myhill, *Recursive equivalence types*, Univ. Calif. Publ. Math. (N.S.) **3** (1960), 67–213.
11. E. Ellentuck, Review of “*Extensions to isols*”, by A. Nerode (see [16]), Math. Reviews **24** (1962), #A1215.
12. ———, *Universal isols*, Math. Z. **98** (1967), 1–8.
13. M. J. Hassett, *Some theorems on regressive isols and isolic groups*, Doctoral Thesis, Rutgers, The State University, New Brunswick, N. J., 1966.
14. W. J. LeVeque, *Topics in number theory*, Vol. 1, Addison-Wesley, Reading, Mass., 1956, pp. 1–198.
15. J. Myhill, *Recursive equivalence types and combinatorial functions*, Proc. 1960 Internat. Congress in Logic, Methodology and Philosophy of Science, Stanford, Stanford Univ. Press, Stanford, Calif., 1962, pp. 46–55.
16. A. Nerode, *Extensions to isols*, Ann. of Math. **73** (1961), 362–403.
17. ———, *Extensions to isolic integers*, Ann. of Math. **75** (1962), 419–448.
18. I. Niven and H. S. Zuckerman, *An introduction to the theory of numbers*, Wiley, New York, 1962, pp. 1–250.
19. F. J. Sansone, *Combinatorial functions and regressive isols*, Pacific J. Math. **13** (1963), 703–707.
20. ———, *On order-preserving extensions to regressive isols*, Michigan Math. J. **13** (1966), 353–355.

STANFORD UNIVERSITY,
 STANFORD, CALIFORNIA
 STATE UNIVERSITY OF NEW YORK,
 BUFFALO, NEW YORK