

BIFUNCTORS AND ADJOINT PAIRS

BY

J. FISHER PALMQUIST AND DAVID C. NEWELL

Abstract. We use a definition of tensor products of functors to generalize some theorems of homological algebra. We show that adjoint pairs of functors between additive functor categories correspond to bifunctors and that composition of such adjoint pairs corresponds to the tensor product of the bifunctors. We also generalize some homological characterizations of finitely generated projective modules to characterizations of small projectives in a functor category. We apply our results to adjoint pairs arising from satellites and from a functor on the domain categories.

0. Introduction. Let R and S be rings, and let Mod_R and Mod_S denote the categories of right R -modules and right S -modules respectively. Given an S - R bimodule E , we obtain a functor $\text{Hom}_R(E, _): \text{Mod}_R \rightarrow \text{Mod}_S$ with an adjoint $_ \otimes_S E: \text{Mod}_S \rightarrow \text{Mod}_R$, which we shall consider as an adjoint pair

$$(\text{Hom}_R(E, _), _ \otimes_S E): \text{Mod}_R \rightarrow \text{Mod}_S.$$

Furthermore, given any adjoint pair $(u_1, u_2): \text{Mod}_R \rightarrow \text{Mod}_S$ (i.e., a functor $u_1: \text{Mod}_R \rightarrow \text{Mod}_S$ having $u_2: \text{Mod}_S \rightarrow \text{Mod}_R$ as an adjoint) it is well known that there is a bimodule E which represents this adjoint pair, that is, one has equivalences $u_1 \cong \text{Hom}_R(E, _)$ and $u_2 \cong _ \otimes_S E$ (see [3] or [6]). In fact, let $E = u_2(S)$, then using the fact that u_2 is right continuous (i.e., u_2 is right exact and preserves direct sums) one obtains $N \otimes_S E = u_2(N)$ for every S -module N by examining the effect of u_2 on an S -free presentation of N . Furthermore, if T is a ring and $(v_1, v_2): \text{Mod}_S \rightarrow \text{Mod}_T$ is an adjoint pair represented by a T - S bimodule E' , then $(v_1 \circ u_1, u_2 \circ v_2): \text{Mod}_R \rightarrow \text{Mod}_T$ is an adjoint pair represented by the T - R bimodule $E' \otimes_S E$.

Although it is well known that functor categories are generalizations of categories of modules, many of the techniques of homological algebra could not be extended because of a lack of a proper definition of a tensor product. However, in [4], a suitable definition has been given, and in this paper we demonstrate how this definition can be exploited. In particular, we show that the above theorems for modules can be easily extended to functor categories.

Before giving a specific summary of the contents of this paper, we introduce some notation and conventions.

Received by the editors March 29, 1970.

AMS 1969 subject classifications. Primary 1810, 1820.

Key words and phrases. Functor categories, tensor product of functors, bifunctors, adjoint pairs on functor categories, satellites, duality on functors, small projective functors, adjoints of induced functors.

Copyright © 1971, American Mathematical Society

We shall assume throughout this paper that all of our categories are at least preadditive and that all of our functors are covariant and additive. If \mathcal{C} and \mathcal{D} are two categories, we shall let $[\mathcal{C}, \mathcal{D}]$ denote the category whose objects are functors from \mathcal{C} to \mathcal{D} and whose morphisms are natural transformations between these functors. Similarly, we let $[\mathcal{C}^\circ, \mathcal{D}]$ denote the category whose objects are contravariant functors from \mathcal{C} to \mathcal{D} (that is, covariant functors from the dual category of \mathcal{C} , \mathcal{C}° , to \mathcal{D}). If F and G are objects of $[\mathcal{C}, \mathcal{D}]$, we let $[F, G]$ denote the abelian group of natural transformations from F to G . (We shall assume throughout that we are in a proper set-theoretic setting so that all of our categories are well defined.)

Let Ab denote the category of abelian groups. If \mathcal{C} is a category, then for objects A and B of \mathcal{C} we denote the abelian group of morphisms in \mathcal{C} from A to B by $\mathcal{C}(A, B)$, we let $h_A^\mathcal{C} = \mathcal{C}(A, _)$ (or h^A if \mathcal{C} is unambiguous) denote the representable functor from \mathcal{C} to Ab , and similarly we let $h_B^\mathcal{C} = \mathcal{C}(_, B)$, or h_B , denote the contravariant representable functor from \mathcal{C}° to Ab .

In §1, we define, for functors $F: \mathcal{C} \rightarrow \text{Ab}$ and $G: \mathcal{C}^\circ \rightarrow \text{Ab}$, an abelian group $F \otimes_{\mathcal{C}} G$, called the tensor product of F and G over \mathcal{C} . This definition and properties (which generalize the properties of the tensor product of modules) have been given in [4], and we present here in a way suitable for applications to this paper a summary of these properties.

If \mathcal{C}_1 and \mathcal{C}_2 are categories and $U: \mathcal{C}_1^\circ \times \mathcal{C}_2 \rightarrow \text{Ab}$ is a bifunctor, we have a functor $\otimes_{\mathcal{C}_1} U: [\mathcal{C}_1, \text{Ab}] \rightarrow [\mathcal{C}_2, \text{Ab}]$ defined as follows: if $F: \mathcal{C}_1 \rightarrow \text{Ab}$ is a functor and B is an object of \mathcal{C}_2 , then $(F \otimes_{\mathcal{C}_1} U)(B) = F \otimes_{\mathcal{C}_1} U(_, B)$. We also have a functor $[U, _]_{\mathcal{C}_2}: [\mathcal{C}_2, \text{Ab}] \rightarrow [\mathcal{C}_1, \text{Ab}]$ defined as follows: for $G: \mathcal{C}_2 \rightarrow \text{Ab}$ a functor and A an object of \mathcal{C}_1 , we let $[U, G]_{\mathcal{C}_2}(A) = [U(A, _), G]$. We complete §1 by showing that $\otimes_{\mathcal{C}_1} U$ is adjoint to $[U, _]_{\mathcal{C}_2}$.

It follows easily from the additive case of a theorem of André (see [1, Proposition 9.1]) that one has an equivalence of categories $[\mathcal{C}_1^\circ \times \mathcal{C}_2, \text{Ab}] \cong \text{Adj}([\mathcal{C}_2, \text{Ab}], [\mathcal{C}_1, \text{Ab}])$, where the objects of the second category consist of pairs of functors (u_1, u_2) , $u_1: [\mathcal{C}_2, \text{Ab}] \rightarrow [\mathcal{C}_1, \text{Ab}]$ and $u_2: [\mathcal{C}_1, \text{Ab}] \rightarrow [\mathcal{C}_2, \text{Ab}]$ its adjoint. In other words, every adjoint pair (u_1, u_2) is represented by a bifunctor U . We give a simpler proof of this theorem which generalizes the proof mentioned above for the case of modules. Furthermore, if $V: \mathcal{C}_2^\circ \times \mathcal{C}_3 \rightarrow \text{Ab}$ is a bifunctor representing an adjoint pair $(v_1, v_2): [\mathcal{C}_3, \text{Ab}] \rightarrow [\mathcal{C}_2, \text{Ab}]$, then there is a natural way to define a bifunctor $U \otimes_{\mathcal{C}_2} V: \mathcal{C}_1^\circ \times \mathcal{C}_3 \rightarrow \text{Ab}$ so that $U \otimes_{\mathcal{C}_2} V$ represents the adjoint pair $(u_1 \circ v_1, v_2 \circ u_2)$. These results are given in §2.

As an example, in §3 we show that if $F: \mathcal{C} \rightarrow \mathcal{D}$ is a functor between abelian categories and $S^n F$ and $S_n F$ denote the n th right satellite of F and the n th left satellite of F respectively, then $S^n: [\mathcal{C}, \mathcal{D}] \rightarrow [\mathcal{C}, \mathcal{D}]$ is adjoint to $S_n: [\mathcal{C}, \mathcal{D}] \rightarrow [\mathcal{C}, \mathcal{D}]$ (if they exist), and if $\mathcal{D} = \text{Ab}$, the corresponding bifunctor representing this adjoint pair (S_n, S^n) is $\text{Ext}^n(_, _): \mathcal{C}^\circ \times \mathcal{C} \rightarrow \text{Ab}$.

In §4, we define for a functor $F: \mathcal{C} \rightarrow \text{Ab}$ its "dual" functor $F^*: \mathcal{C}^\circ \rightarrow \text{Ab}$,

natural transformations $\otimes_{\mathcal{C}} F^* \rightarrow [F, \]$ and $F \rightarrow F^{**}$, and we are able to prove the usual theorems about these natural transformations when F is a small projective. We make the easy step of generalizing these results to bifunctors, and as an example, we give a simple description of the adjoint to a functor $u: [\mathcal{C}_2, \text{Ab}] \rightarrow [\mathcal{C}_1, \text{Ab}]$ when u is induced by a functor $f: \mathcal{C}_1 \rightarrow \mathcal{C}_2$.

We note that the results of this paper still hold when Ab is replaced by Ens , the category of sets, and all of our preadditive categories are replaced by ordinary ones. We do the additive case as the technique involved is more clearly an extension of that of homological algebra.

1. Tensor products of functors. In this section we summarize the properties of the tensor product of functors which are found in [4] in a way convenient for our use in this paper. This summarization is given in the following:

PROPOSITION 1.1. *Let \mathcal{C} be a category and let $F: \mathcal{C} \rightarrow \text{Ab}$ and $G: \mathcal{C}^\circ \rightarrow \text{Ab}$. Then there is a unique abelian group $F \otimes_{\mathcal{C}} G$ so that the assignment $(F, G) \rightarrow F \otimes_{\mathcal{C}} G$ defines a functor of two variables having the following properties:*

- (a) *For any object A of \mathcal{C} , $h^A \otimes_{\mathcal{C}} G \cong G(A)$ and $F \otimes_{\mathcal{C}} h_A \cong F(A)$,*
- (b) *$\otimes_{\mathcal{C}} G$, considered as a functor from $[\mathcal{C}, \text{Ab}]$ to Ab , is right continuous, and $F \otimes_{\mathcal{C}} \ ,$ considered as a functor from $[\mathcal{C}^\circ, \text{Ab}]$ to Ab , is right continuous.*

For the proof we refer to [4]. As a consequence, we have the following:

PROPOSITION 1.2. *Let \mathcal{C} be a category, $G: \mathcal{C}^\circ \rightarrow \text{Ab}$ and $\nu: [\mathcal{C}, \text{Ab}] \rightarrow \text{Ab}$ be functors, and suppose that for each object A of \mathcal{C} there is a homomorphism $\varphi'_A: G(A) \rightarrow \nu(h^A)$ which is natural in A . Then there is a unique natural transformation $\varphi: \otimes_{\mathcal{C}} G \rightarrow \nu$ such that $\varphi(h^A)$ is the composition of the isomorphism $h^A \otimes_{\mathcal{C}} G \cong G(A)$ followed by φ'_A . Furthermore, if ν is right continuous and φ'_A is an isomorphism of abelian groups for all objects A of \mathcal{C} , then φ is an isomorphism of functors.*

The proof follows from 1.1 and the observation that for any functor $F: \mathcal{C} \rightarrow \text{Ab}$, there are collections of objects $(A_i)_i$ and $(A_j)_j$ of \mathcal{C} and an exact sequence

$$(*) \quad \sum_i h^{A_i} \rightarrow \sum_j h^{A_j} \rightarrow F \rightarrow 0$$

in $[\mathcal{C}, \text{Ab}]$.

Recall that, for \mathcal{A} and \mathcal{B} categories, a functor $u_2: \mathcal{B} \rightarrow \mathcal{A}$ is said to be adjoint to a functor $u_1: \mathcal{A} \rightarrow \mathcal{B}$ (or equivalently u_1 is co-adjoint to u_2) if for any objects F of \mathcal{A} and G of \mathcal{B} there is an isomorphism

$$\mathcal{A}(u_2G, F) \simeq \mathcal{B}(G, u_1F)$$

which is natural in F and G .

PROPOSITION 1.3. *For each bifunctor $U: \mathcal{C}_1 \times \mathcal{C}_2 \rightarrow \text{Ab}$ the functor*

$$\otimes_{\mathcal{C}_1} U: [\mathcal{C}_1, \text{Ab}] \rightarrow [\mathcal{C}_2, \text{Ab}]$$

is adjoint to $[U, \]_{\mathcal{C}_2}$.

Proof. For each $G: \mathcal{C}_2 \rightarrow \text{Ab}$ the functors $[\otimes_{\mathcal{C}_1} U, G]$ and $[\ , [U, G]_{\mathcal{C}_2}] : [\mathcal{C}_1, \text{Ab}]^\circ \rightarrow \text{Ab}$, given by $[\otimes_{\mathcal{C}_1} U, G](F) = [F \otimes_{\mathcal{C}_1} U, G]$ and $[\ , [U, G]_{\mathcal{C}_2}](F) = [F, [U, G]_{\mathcal{C}_2}]$ for $F: \mathcal{C}_1 \rightarrow \text{Ab}$, are left continuous and have isomorphic values on representable functors, i.e., for A in \mathcal{C}_1 , $[h^A \otimes_{\mathcal{C}_1} U, G] \simeq [U(A, \), G] \simeq [h^A, [U, G]_{\mathcal{C}_2}]$ by the Yoneda lemma. Thus using the exact sequence (*) we see that for each $F: \mathcal{C}_1 \rightarrow \text{Ab}$, $[F \otimes_{\mathcal{C}_1} U, G] \simeq [F, [U, G]_{\mathcal{C}_2}]$ and this isomorphism is natural in F and G .

2. Representing adjoint pairs. Let \mathcal{A} and \mathcal{B} be categories and let $u_1: \mathcal{A} \rightarrow \mathcal{B}$ and $u_2: \mathcal{B} \rightarrow \mathcal{A}$ be functors. We call (u_1, u_2) an *adjoint pair from \mathcal{A} to \mathcal{B}* if u_2 is adjoint to u_1 .

Given two adjoint pairs (u_1, u_2) and (v_1, v_2) from \mathcal{A} to \mathcal{B} , we define a *morphism of adjoint pairs* $\alpha: (u_1, u_2) \rightarrow (v_1, v_2)$ by $\alpha = (\alpha_1, \alpha_2)$, where α_1 and α_2 are natural transformations $\alpha_1: v_1 \rightarrow u_1$ and $\alpha_2: u_2 \rightarrow v_2$ so that for all objects F of \mathcal{A} and G of \mathcal{B} , the diagram

$$\begin{array}{ccc} \mathcal{A}(v_2 G, F) \cong \mathcal{B}(G, v_1 F) & & \\ \downarrow (\alpha_2 G)^* & & \downarrow (\alpha_1 F)_* \\ \mathcal{A}(u_2 G, F) \cong \mathcal{B}(G, u_1 F) & & \end{array}$$

commutes.

Given two morphisms of adjoint pairs from \mathcal{A} to \mathcal{B} , $\alpha = (\alpha_1, \alpha_2): (u_1, u_2) \rightarrow (v_1, v_2)$ and $\beta = (\beta_1, \beta_2): (v_1, v_2) \rightarrow (w_1, w_2)$, we define their composition by $\beta \circ \alpha = (\alpha_1 \circ \beta_1, \beta_2 \circ \alpha_2): (u_1, u_2) \rightarrow (w_1, w_2)$, which is easily seen to be a morphism of adjoint pairs. We may therefore define a category $\text{Adj}(\mathcal{A}, \mathcal{B})$ whose objects are adjoint pairs from \mathcal{A} to \mathcal{B} and whose morphisms and composition are defined above.

It is well known that

- (1) Given an object (u_1, u_2) of $\text{Adj}(\mathcal{A}, \mathcal{B})$, u_2 is determined up to an isomorphism of functors by u_1 and conversely, and given a morphism $\alpha = (\alpha_1, \alpha_2)$ of $\text{Adj}(\mathcal{A}, \mathcal{B})$, α_2 is determined by α_1 and conversely.
- (2) For objects (u_1, u_2) of $\text{Adj}(\mathcal{A}, \mathcal{B})$ and (u'_1, u'_2) of $\text{Adj}(\mathcal{B}, \mathcal{B}')$, $(u'_1 \circ u_1, u_2 \circ u'_2)$ is an object of $\text{Adj}(\mathcal{A}, \mathcal{B}')$ which is called the composition of (u_1, u_2) and (u'_1, u'_2) .
- (3) Given an object (u_1, u_2) of $\text{Adj}(\mathcal{A}, \mathcal{B})$, u_1 is left continuous and u_2 is right continuous.

It follows from 1.3 that for any two categories \mathcal{C}_1 and \mathcal{C}_2 we have a functor

$$\Phi: [\mathcal{C}_1^\circ \times \mathcal{C}_2, \text{Ab}] \rightarrow \text{Adj}([\mathcal{C}_2, \text{Ab}], [\mathcal{C}_1, \text{Ab}])$$

given by $\Phi(U) = ([U, \]_{\mathcal{C}_2}, \otimes_{\mathcal{C}_1} U)$ for any bifunctor $U: \mathcal{C}_1^\circ \times \mathcal{C}_2 \rightarrow \text{Ab}$.

The following theorem and corollary are easy consequences of Proposition 9.1 of [1]. We give a simple proof that generalizes the proof for the case of modules.

THEOREM 2.1. *If $u: [\mathcal{C}_1, \text{Ab}] \rightarrow [\mathcal{C}_2, \text{Ab}]$ is right continuous, then there is a bifunctor $U: \mathcal{C}_1^\circ \times \mathcal{C}_2 \rightarrow \text{Ab}$ such that $u \cong \otimes_{\mathcal{C}_1} U$.*

Proof. Let $h: \mathcal{C}_1 \rightarrow [\mathcal{C}_1, \text{Ab}]$ denote the functor such that $h(A) = h_{\mathcal{C}_1}^A (= h^A)$ for any object A of \mathcal{C}_1 . Then $u \circ h$ is an object of $[\mathcal{C}_1^\circ, [\mathcal{C}_2, \text{Ab}]]$. There is a canonical isomorphism

$$\tau: [\mathcal{C}_1^\circ, [\mathcal{C}_2, \text{Ab}]] \cong [\mathcal{C}_1^\circ \times \mathcal{C}_2, \text{Ab}]$$

given by $\tau(t)(A, B) = t(A)(B)$ for all functors $t: \mathcal{C}_1^\circ \rightarrow [\mathcal{C}_2, \text{Ab}]$ and for all objects A of \mathcal{C}_1 and B of \mathcal{C}_2 . Let $U = \tau(u \circ h)$. Since, for each object A of \mathcal{C}_1 , we have an isomorphism $\varphi'_A: U(A, \) \rightarrow u(h^A)$ defined by the equalities

$$U(A, \) = \tau(u \circ h)(A, \) = (u \circ h)(A) = u(h^A),$$

and u is right continuous, it follows from 1.2 that there is an isomorphism $\varphi: \otimes_{\mathcal{C}_1} U \cong u$.

It follows from this theorem and our remarks about adjoint functors that

COROLLARY 2.2. $\Phi: [\mathcal{C}_1^\circ \times \mathcal{C}_2, \text{Ab}] \rightarrow \text{Adj}([\mathcal{C}_2, \text{Ab}], [\mathcal{C}_1, \text{Ab}])$ is an equivalence of categories.

Therefore, if (u_1, u_2) is an adjoint pair in $\text{Adj}([\mathcal{C}_2, \text{Ab}], [\mathcal{C}_1, \text{Ab}])$, and $U: \mathcal{C}_1^\circ \times \mathcal{C}_2 \rightarrow \text{Ab}$ is a bifunctor such that $\Phi(U) \cong (u_1, u_2)$, then U is unique up to a canonical isomorphism. We shall say that U represents the adjoint pair (u_1, u_2) , and that U represents u_1 on the left and U represents u_2 on the right.

As a trivial example, consider a category \mathcal{C} and the adjoint pair (I, I) from $[\mathcal{C}, \text{Ab}]$ to itself, where I is the identity functor on $[\mathcal{C}, \text{Ab}]$. Then, if we let $h_{\mathcal{C}}: \mathcal{C}^\circ \times \mathcal{C} \rightarrow \text{Ab}$ denote the bifunctor $h_{\mathcal{C}}(A_1, A_2) = \mathcal{C}(A_1, A_2)$ for all objects A_1 and A_2 of \mathcal{C} , $h_{\mathcal{C}}$ represents (I, I) . That $h_{\mathcal{C}}$ represents I on the left is a restatement of the "Yoneda lemma" and that $h_{\mathcal{C}}$ represents I on the right is a restatement of 1.1(a).

Suppose we have an adjoint pair (u_1, u_2) from $[\mathcal{C}_2, \text{Ab}]$ to $[\mathcal{C}_1, \text{Ab}]$ represented by a bifunctor $U: \mathcal{C}_1^\circ \times \mathcal{C}_2 \rightarrow \text{Ab}$, and an adjoint pair (v_1, v_2) from $[\mathcal{C}_3, \text{Ab}]$ to $[\mathcal{C}_2, \text{Ab}]$ represented by a bifunctor $V: \mathcal{C}_2^\circ \times \mathcal{C}_3 \rightarrow \text{Ab}$. The composition of these adjoint pairs is an adjoint pair $(u_1 \circ v_1, v_2 \circ u_2)$ from $[\mathcal{C}_3, \text{Ab}]$ to $[\mathcal{C}_1, \text{Ab}]$.

We may define a bifunctor $U \otimes_{\mathcal{C}_2} V: \mathcal{C}_1^\circ \times \mathcal{C}_3 \rightarrow \text{Ab}$ by $U \otimes_{\mathcal{C}_2} V(A, C) = U(A, \) \otimes_{\mathcal{C}_2} V(\ , C)$ for all objects A of \mathcal{C}_1 and C of \mathcal{C}_3 . Furthermore, if $F: \mathcal{C}_1 \rightarrow \text{Ab}$ is a functor, it follows from the associativity of the tensor product (see [4]) that $(F \otimes_{\mathcal{C}_1} U) \otimes_{\mathcal{C}_2} V \cong F \otimes_{\mathcal{C}_1} (U \otimes_{\mathcal{C}_2} V)$. As an easy consequence, we have

PROPOSITION 2.3. *Let (u_1, u_2) and (v_1, v_2) be adjoint pairs represented by $U: \mathcal{C}_1^\circ \times \mathcal{C}_2 \rightarrow \text{Ab}$ and $V: \mathcal{C}_2^\circ \times \mathcal{C}_3 \rightarrow \text{Ab}$, respectively. Then the adjoint pair $(u_1 \circ v_1, v_2 \circ u_2)$ is represented by $U \otimes_{\mathcal{C}_2} V$.*

Since for any functor $F: \mathcal{C}_3 \rightarrow \text{Ab}$, $u_1 \circ v_1(F) \cong [U, [V, F]_{\mathcal{C}_3}]_{\mathcal{C}_2}$, and left adjoints of a functor are canonically isomorphic, we have

COROLLARY 2.4. *For any bifunctors $U: \mathcal{C}_1^\circ \times \mathcal{C}_2 \rightarrow \text{Ab}$ and $V: \mathcal{C}_2^\circ \times \mathcal{C}_3 \rightarrow \text{Ab}$ and for any functor $F: \mathcal{C}_3 \rightarrow \text{Ab}$, there is a canonical isomorphism*

$$[U \otimes_{\mathcal{C}_2} V, F]_{\mathcal{C}_3} \cong [U, [V, F]_{\mathcal{C}_3}]_{\mathcal{C}_2}$$

natural in U, V , and F .

3. Satellites. In this section, \mathcal{C} and \mathcal{D} will be abelian categories.

For a functor $F: \mathcal{C} \rightarrow \text{Ab}$, there are functors $S^n F$ and $S_n F$ for each integer $n \geq 0$, the n th right and left satellites of F respectively (which are defined below). We may consider S^n and S_n as functors from $[\mathcal{C}, \text{Ab}]$ to itself. The result of this section is that (S_n, S^n) is an adjoint pair which is represented by the bifunctor $\text{Ext}_{\mathcal{C}}^n(,) : \mathcal{C}^\circ \times \mathcal{C} \rightarrow \text{Ab}$.

Let F_1 and F_2 be functors from a category \mathcal{C} to a category \mathcal{D} . Then F_1 and F_2 are *connected* if for every short exact sequence $E: 0 \rightarrow A' \rightarrow A \rightarrow A'' \rightarrow 0$ in \mathcal{C} there is a map $\Delta_E: F_1(A'') \rightarrow F_2(A')$ natural in E so that the complex $F_1(A') \rightarrow F_1(A) \rightarrow F_1(A'') \rightarrow F_2(A') \rightarrow F_2(A) \rightarrow F_2(A'')$ is of order 2 (that is, composition of two successive morphisms is the zero morphism). We shall indicate that F_1 and F_2 are connected by Δ , by the triple (F_1, Δ, F_2) , which we call a *connected triple in $[\mathcal{C}, \mathcal{D}]$* .

Given two connected triples (F_1, Δ, F_2) and (G_1, Δ', G_2) , we may define a *morphism of connected triples* to be a pair $(\alpha_1, \alpha_2): (F_1, \Delta, F_2) \rightarrow (G_1, \Delta', G_2)$, where $\alpha_1: F_1 \rightarrow G_1$ and $\alpha_2: F_2 \rightarrow G_2$ are natural transformations such that, for every exact sequence $E: 0 \rightarrow A' \rightarrow A \rightarrow A'' \rightarrow 0$, we have $\Delta'_E \circ \alpha_1(A'') = \alpha_2(A') \circ \Delta_E$.

If $(\alpha_1, \alpha_2): (F_1, \Delta, F_2) \rightarrow (G_1, \Delta', G_2)$ and $(\beta_1, \beta_2): (G_1, \Delta', G_2) \rightarrow (H_1, \Delta'', H_2)$ are morphisms of connected triples, then the composition $(\beta_1, \beta_2) \circ (\alpha_1, \alpha_2) = (\beta_1 \circ \alpha_1, \beta_2 \circ \alpha_2)$ is also. Consequently, we may define a category $\Delta[\mathcal{C}, \mathcal{D}]$ whose objects are connected triples in $[\mathcal{C}, \mathcal{D}]$ and whose morphisms and composition are defined above. We have two functors $\pi_i: \Delta[\mathcal{C}, \mathcal{D}] \rightarrow [\mathcal{C}, \mathcal{D}]$ defined by $\pi_i(F_1, \Delta, F_2) = F_i$, $i = 1, 2$, for any connected triple (F_1, Δ, F_2) in $[\mathcal{C}, \mathcal{D}]$.

Let $F: \mathcal{C} \rightarrow \mathcal{D}$ be a functor. Then F has a *right satellite* if there is a functor $S^1 F: \mathcal{C} \rightarrow \mathcal{D}$ so that F and $S^1 F$ are connected (via Δ , let us say), and given any connected triple (G_1, Δ', G_2) in $[\mathcal{C}, \mathcal{D}]$ and any natural transformation $\alpha: F \rightarrow G_1$, there is a unique natural transformation $\alpha': S^1 F \rightarrow G_2$ so that $(\alpha, \alpha'): (F, \Delta, S^1 F) \rightarrow (G_1, \Delta', G_2)$ is a morphism of connected triples. Similarly, F has a *left satellite* if there is a functor $S_1 F: \mathcal{C} \rightarrow \mathcal{D}$ so that $S_1 F$ and F are connected (by Δ), and given any connected triple (G_1, Δ', G_2) and a natural transformation $\alpha: G_2 \rightarrow F$, there is a unique natural transformation $\alpha': G_1 \rightarrow S_1 F$ so that $(\alpha', \alpha): (G_1, \Delta', G_2) \rightarrow (S_1 F, \Delta, F)$ is a morphism of connected pairs.

We shall denote the connected triples $(F, \Delta, S^1 F)$ and $(S_1 F, \Delta, F)$ by $\Delta S^1(F)$ and $\Delta S_1(F)$ respectively. If $[\mathcal{C}, \mathcal{D}]$ has right satellites (that is, all functors $F: \mathcal{C} \rightarrow \mathcal{D}$ have right satellites), then ΔS^1 defines a functor $\Delta S^1: [\mathcal{C}, \mathcal{D}] \rightarrow \Delta[\mathcal{C}, \mathcal{D}]$, and similarly if $[\mathcal{C}, \mathcal{D}]$ has left satellites, ΔS_1 defines a functor $\Delta S_1: [\mathcal{C}, \mathcal{D}] \rightarrow \Delta[\mathcal{C}, \mathcal{D}]$. In fact, we may translate the definition of ΔS^1 and ΔS_1 into the following:

PROPOSITION 3.1. *If $[\mathcal{C}, \mathcal{D}]$ has right satellites, then ΔS^1 is the adjoint of π_1 . If $[\mathcal{C}, \mathcal{D}]$ has left satellites, then ΔS_1 is the co-adjoint of π_2 .*

Suppose $[\mathcal{C}, \mathcal{D}]$ has satellites (both right and left). Then we have adjoint pairs $(\Delta S_1, \pi_2)$ from $[\mathcal{C}, \mathcal{D}]$ to $\Delta[\mathcal{C}, \mathcal{D}]$ and $(\pi_1, \Delta S^1)$ from $\Delta[\mathcal{C}, \mathcal{D}]$ to $[\mathcal{C}, \mathcal{D}]$, and composing gives us an adjoint pair $(\pi_1 \circ \Delta S_1, \pi_2 \circ \Delta S^1)$ from $[\mathcal{C}, \mathcal{D}]$ to itself. But $\pi_1 \circ \Delta S_1(F) = S_1F$ and $\pi_2 \circ \Delta S^1(F) = S^1F$, which proves the following:

COROLLARY 3.2⁽¹⁾. *If $[\mathcal{C}, \mathcal{D}]$ has satellites, then $S^1: [\mathcal{C}, \mathcal{D}] \rightarrow [\mathcal{C}, \mathcal{D}]$ is adjoint to S_1 .*

If $F: \mathcal{C} \rightarrow \mathcal{D}$ is a functor, we define, in the usual way, the higher satellites of F as follows:

$$\begin{aligned} S^0F &= F, & S^{n+1}F &= S^1(S^nF) \quad \text{for all } n \geq 0, \\ S_0F &= F, & S_{n+1}F &= S_1(S_nF) \quad \text{for all } n \geq 0. \end{aligned}$$

COROLLARY 3.3. *If $[\mathcal{C}, \mathcal{D}]$ has satellites, then S^n is adjoint to S_n for all $n \geq 0$.*

The proof is by induction, using the fact that composition of adjoint pairs is an adjoint pair.

If $\mathcal{D} = \text{Ab}$, then $[\mathcal{C}, \text{Ab}]$ has satellites for any abelian category \mathcal{C} (see [2]). To compute the bifunctor representing the adjoint pair (S_n, S^n) , we may use the proof of Theorem 2.1. Since for any object A of \mathcal{C} , it is known that $S^n(h_A^A) \cong \text{Ext}_{\mathcal{C}}^n(A, _)$, we have the following:

THEOREM 3.4. *For all integers $n \geq 0$, the functors S_n and S^n from $[\mathcal{C}, \text{Ab}]$ to itself form an adjoint pair (S_n, S^n) which is represented by the bifunctor $\text{Ext}_{\mathcal{C}}^n(_, _)$.*

We remark that the fact that S_n is represented on the left by $\text{Ext}_{\mathcal{C}}^n(_, _)$ is a result of Yoneda [7], while the fact that S^n is represented on the right by $\text{Ext}_{\mathcal{C}}^n(_, _)$ is a result in [4].

4. Duality and adjoints of induced functors. In this section we define, for a function $F: \mathcal{C} \rightarrow \text{Ab}$, its dual $F^*: \mathcal{C}^\circ \rightarrow \text{Ab}$ and prove some theorems concerning F^* . Using these results, we are able to prove some results about a bifunctor U when U represents an adjoint pair (u_1, u_2) from $[\mathcal{C}_2, \text{Ab}]$ to $[\mathcal{C}_1, \text{Ab}]$ with u_1 exact and continuous. In particular, we deal with the case where u_1 is induced by a functor $\mathcal{C}_1 \rightarrow \mathcal{C}_2$.

Let $V: \mathcal{C}_1^\circ \times \mathcal{C}_2 \rightarrow \text{Ab}$ be a bifunctor. Then we define a functor $H_{\mathcal{C}_2}[_, V]: [\mathcal{C}_2, \text{Ab}]^\circ \rightarrow [\mathcal{C}_1^\circ, \text{Ab}]$ as follows: for a functor $F: \mathcal{C}_2 \rightarrow \text{Ab}$ and an object A of \mathcal{C}_1 , we let $H_{\mathcal{C}_2}[F, V](A) = [F, V(A, _)]$. Similarly, we define a functor

$$H^{\mathcal{C}_1}[_, V]: [\mathcal{C}_1^\circ, \text{Ab}]^\circ \rightarrow [\mathcal{C}_2, \text{Ab}]$$

⁽¹⁾ We thank our colleague, Paul Palmquist, for suggesting using universal properties of satellites to prove this theorem.

as follows: for a functor $G: \mathcal{C}_1 \rightarrow \text{Ab}$ and an object B of \mathcal{C}_2 , $H^{\mathcal{C}_1}[G, V](B) = [G, V(\cdot, B)]$.

As a direct consequence of 1.2 we have

PROPOSITION 4.1. *Let $F: \mathcal{C} \rightarrow \text{Ab}$ be a functor and let $F^* = H_{\mathcal{C}}[F, h_{\mathcal{C}}]$. Then there exists a unique natural transformation*

$$\beta(\cdot, F): \otimes_{\mathcal{C}} F^* \rightarrow [F, \cdot]$$

of functors from $[\mathcal{C}, \text{Ab}]$ to Ab , natural in F , such that $\beta(h^A, F)$ is the composition of isomorphisms

$$h^A \otimes F^* \simeq F^*(A) = H_{\mathcal{C}}[F, h_{\mathcal{C}}](A) = [F, h^A].$$

Recall that, for an object P of an abelian category \mathcal{A} , P is a small projective if and only if $h^P_{\mathcal{A}}$ is right continuous. In particular, it follows that an object F in $[\mathcal{C}, \text{Ab}]$ is a small projective if and only if there is a finite collection of objects $(B_i)_I$ of \mathcal{C} such that F is a direct summand of $\sum_I h^B_i$.

PROPOSITION 4.2. *For $F: \mathcal{C} \rightarrow \text{Ab}$ the following are equivalent:*

- (a) $\beta(\cdot, F): \otimes_{\mathcal{C}} F^* \rightarrow [F, \cdot]$ is a natural equivalence.
- (b) $\beta(G, F)$ is an epimorphism for each $G: \mathcal{C} \rightarrow \text{Ab}$.
- (c) $\beta(F, F)$ is an epimorphism.
- (d) F is a small projective in $[\mathcal{C}, \text{Ab}]$.

Proof. $a \Rightarrow b \Rightarrow c$ is trivial.

$d \Rightarrow a$ is a direct consequence of the definition of small projective and 1.2.

$c \Rightarrow d$. We know that there exists a collection $(A_j)_J$ of objects in \mathcal{C} and transformations $\{s_j: h^A_j \rightarrow F\}$ such that $\sum s_j: \sum_{j \in J} h^A_j \rightarrow F$ is an epimorphism. Since $\otimes F^*$ is right continuous we have the commutative diagram

$$\begin{array}{ccc} \sum_J (h^A_j \otimes F^*) \simeq \left(\sum_J h^A_j \right) \otimes F^* & \xrightarrow{(\sum s_j) \otimes F^*} & F \otimes F^* \\ \downarrow \sum \beta(h^A_j, F) \cong & & \downarrow \beta(F, F) \\ \sum_J [F, h^A_j] & \xrightarrow{\sum_J [F, s_j]} & [F, F] \end{array}$$

with the top map an epimorphism. Thus $\beta(F, F)$ is an epimorphism implies $\sum_J [F, s_j]$ is an epimorphism and hence the existence of a finite subset I of J and elements $t_i \in [F, h^A_i]$, $i \in I$, such that

$$\left(F \xrightarrow{\sum t_i} \sum_I h^A_i \xrightarrow{\sum s_i} F \right) = 1_F.$$

Thus by the remark preceding this proposition F is a small projective.

Let $F: \mathcal{C} \rightarrow \text{Ab}$, and let F^{**} denote the functor $H^{\mathcal{C}}[H_{\mathcal{C}}[F, h], h]$. Let A and B be objects of \mathcal{C} . Then we have a map

$$\nu_F(A, B): F(B) \rightarrow \text{Hom}(\text{Hom}(F(B), \mathcal{C}(A, B)), \mathcal{C}(A, B))$$

given by $\nu_F(A, B)(x)(\lambda) = \lambda(x)$ for all x in $F(B)$ and $\lambda \in \text{Hom}(F(B), \mathcal{C}(A, B))$. It is easy to show that this defines a natural transformation $\nu_F: F \rightarrow F^{**}$ which is natural in F .

PROPOSITION 4.3. *If F is a small projective in $[\mathcal{C}, \text{Ab}]$, then $\nu_F: F \rightarrow F^{**}$ is an isomorphism.*

Proof. It is easy to check that for each object A of \mathcal{C} , $h^{A*} \cong h_A$ and $h^{A**} \cong h^A$, and that ν_{h^A} is this isomorphism. Therefore, by a direct sum argument, ν_F is an isomorphism for any small projective F .

We now generalize these results to bifunctors. Let $\mathcal{C}_1, \mathcal{C}_2$, and \mathcal{C}_3 be categories. Given bifunctors $U: \mathcal{C}_3^{\circ} \times \mathcal{C}_2 \rightarrow \text{Ab}$ and $V: \mathcal{C}_1^{\circ} \times \mathcal{C}_2 \rightarrow \text{Ab}$, we define a bifunctor $H_{\mathcal{C}_2}[U, V](A, B) = [U(B, _), V(A, _)]$ for any objects A of \mathcal{C}_1 and B of \mathcal{C}_3 . If $W: \mathcal{C}_1^{\circ} \times \mathcal{C}_3 \rightarrow \text{Ab}$ is a bifunctor and V is as above, we define a bifunctor

$$H^{\mathcal{C}_1}[W, V]: \mathcal{C}_3^{\circ} \times \mathcal{C}_2 \rightarrow \text{Ab}$$

by $H^{\mathcal{C}_1}[W, V](C, B) = [W(_, C), V(_, B)]$ for any objects B of \mathcal{C}_2 and C of \mathcal{C}_3 .

PROPOSITION 4.4. *Let $U: \mathcal{C}_1^{\circ} \times \mathcal{C}_2 \rightarrow \text{Ab}$ be a bifunctor. Then*

(a) *there is a natural transformation*

$$\beta_U: \otimes_{\mathcal{C}_2} H_{\mathcal{C}_2}[U, h_{\mathcal{C}_2}] \rightarrow [U, _]_{\mathcal{C}_2}$$

of functors from $[\mathcal{C}_2, \text{Ab}]$ to $[\mathcal{C}_1, \text{Ab}]$ such that β_U is an equivalence of functors if and only if $U(A, _)$ is a small projective of $[\mathcal{C}_2, \text{Ab}]$ for every A in \mathcal{C}_1 , and

(b) *there is a natural transformation of bifunctors $\nu_U: U \rightarrow H^{\mathcal{C}_2}[H_{\mathcal{C}_2}[U, h_{\mathcal{C}_2}], h_{\mathcal{C}_2}]$ such that if $U(A, _)$ is a small projective of $[\mathcal{C}_2, \text{Ab}]$ for every A in \mathcal{C}_1 , then ν_U is an isomorphism.*

Proof. (a) For each $G: \mathcal{C}_2 \rightarrow \text{Ab}$ and A in \mathcal{C}_1 , we let $\beta_U(G)(A)$ equal $\beta(G, U(A, _))$ defined in 4.1. The remainder of (a) follows from 4.2.

(b) For each A in \mathcal{C}_1 , we let $\nu_U(A, _)$ be the natural transformation

$$\nu_{U(A, _)}: U(A, _) \rightarrow U(A, _)^{**} = H^{\mathcal{C}_2}[H_{\mathcal{C}_2}[U, h_{\mathcal{C}_2}], h_{\mathcal{C}_2}](A, _)$$

defined above in 4.3. By 4.3 if $U(A, _)$ is a small projective for every A in \mathcal{C}_1 then ν_U is an equivalence of functors.

Let \mathcal{C}_1 and \mathcal{C}_2 be two categories, (u_1, u_2) an adjoint pair from $[\mathcal{C}_2, \text{Ab}]$ to $[\mathcal{C}_1, \text{Ab}]$, and $U: \mathcal{C}_1^{\circ} \times \mathcal{C}_2 \rightarrow \text{Ab}$ the bifunctor representing (u_1, u_2) .

u_1 is an exact continuous functor if and only if $u_1 \cong [U, _]_{\mathcal{C}_2}$ is right continuous. Since exactness and limits in $[\mathcal{C}_1, \text{Ab}]$ are computed "pointwise," we see that $[U, _]_{\mathcal{C}_2}$ is right continuous if and only if for each A in \mathcal{C}_1 , $U(A, _)$ is a small projective in $[\mathcal{C}_2, \text{Ab}]$. These remarks, and 4.4(a) give us the following:

PROPOSITION 4.5. *Let (u_1, u_2) be an adjoint pair from $[\mathcal{C}_2, \text{Ab}]$ to $[\mathcal{C}_1, \text{Ab}]$ represented by $U: \mathcal{C}_1^\circ \times \mathcal{C}_2 \rightarrow \text{Ab}$. Then the following statements are equivalent:*

- (a) u_1 is an exact continuous functor.
- (b) $U(A, \)$ is a small projective for each object A of \mathcal{C}_1 .
- (c) u_1 is represented on the right by $H_{\mathcal{C}_2}[U, h_{\mathcal{C}_2}]$, i.e., $u_1 \simeq \otimes_{\mathcal{C}_2} H_{\mathcal{C}_2}[U, h_{\mathcal{C}_2}]$.

For example, suppose $u_1 = f^*: [\mathcal{C}_2, \text{Ab}] \rightarrow [\mathcal{C}_1, \text{Ab}]$, where $f: \mathcal{C}_1 \rightarrow \mathcal{C}_2$ is a functor. Then, since f^* is exact and continuous, f^* is represented on the left by a bifunctor $U: \mathcal{C}_1^\circ \times \mathcal{C}_2 \rightarrow \text{Ab}$ such that for all objects A of \mathcal{C}_1 , $U(A, \)$ is a small projective in $[\mathcal{C}_2, \text{Ab}]$. Since f^* is represented on the right by $H_{\mathcal{C}_2}[U, h_{\mathcal{C}_2}]$, we can compute $H_{\mathcal{C}_2}[U, h_{\mathcal{C}_2}]$ by the method used in the proof of 2.1 with the result that

$$H_{\mathcal{C}_2}[U, h_{\mathcal{C}_2}] \simeq \mathcal{C}_2(\ , f).$$

Since, by 4.4(b), $U \simeq H^{\mathcal{C}_2}[H_{\mathcal{C}_2}[U, h_{\mathcal{C}_2}], h_{\mathcal{C}_2}]$, we can compute U , with the result that $U \simeq \mathcal{C}_2(f, \)$. Therefore, it follows that

THEOREM 4.6. *Let $f: \mathcal{C}_1 \rightarrow \mathcal{C}_2$ be a functor, let $f^*: [\mathcal{C}_2, \text{Ab}] \rightarrow [\mathcal{C}_1, \text{Ab}]$ be the induced functor, and let f_* denote its adjoint. Then the adjoint pair (f^*, f_*) is represented by the bifunctor $\mathcal{C}_2(f, \): \mathcal{C}_1^\circ \times \mathcal{C}_2 \rightarrow \text{Ab}$.*

We can give a trivial proof of the following known result:

COROLLARY 4.7. *If f, f^* , and f_* are as in 4.6, and $f: \mathcal{C}_1 \rightarrow \mathcal{C}_2$ is a full and faithful functor (for example, \mathcal{C}_1 a full subcategory of \mathcal{C}_2 , and f the inclusion functor), then $f^* \circ f_* \simeq I$, the identity on $[\mathcal{C}_1, \text{Ab}]$.*

Proof. Since f is full and faithful, we have $\mathcal{C}_1(\ , \) \simeq \mathcal{C}_2(f, f)$. Since f_* is represented on the right by $\mathcal{C}_2(f, \)$ and f^* is represented on the right by $\mathcal{C}_2(\ , f)$, we have that $f^* \circ f_*$ is represented on the right by $\mathcal{C}_2(f, \) \otimes_{\mathcal{C}_2} \mathcal{C}_2(\ , f)$ by Proposition 2.3.

Now, for all objects A_1 and A_2 of \mathcal{C}_1 ,

$$\begin{aligned} \mathcal{C}_2(f, \) \otimes_{\mathcal{C}_2} \mathcal{C}_2(\ , f)(A_1, A_2) &= \mathcal{C}_2(f(A_1), \) \otimes_{\mathcal{C}_2} \mathcal{C}_2(\ , f(A_2)) \\ &= h_{\mathcal{C}_2}^{f(A_1)} \otimes_{\mathcal{C}_2} \mathcal{C}_2(\ , f(A_2)) \simeq \mathcal{C}_2(f(A_1), f(A_2)) \\ &\simeq \mathcal{C}_1(A_1, A_2); \end{aligned}$$

so that $f^* \circ f_*$ is represented on the right by the bifunctor $\mathcal{C}_1(\ , \) = h_{\mathcal{C}_1}$, which also represents the identity functor on the right. The equivalence of categories given in 2.2 completes the proof.

REFERENCES

1. M. André, *Categories of functors and adjoint functors*, Amer. J. Math. **88** (1966), 529–543. MR **33** #5693.
2. D. A. Buchsbaum, *Satellites and universal functors*, Ann. of Math. (2) **71** (1960), 199–209. MR **22** #3751.

3. S. Eilenberg, *Abstract description of some basic functors*, J. Indian Math. Soc. **24** (1960), 231–234. MR **23** #A2454.
4. J. L. Fisher, *The tensor product of functors; satellites; and derived functors*, J. Algebra **8** (1968), 277–294. MR **38** #5874.
5. D. M. Kan, *Adjoint functors*, Trans. Amer. Math. Soc. **87** (1958), 294–329. MR **24** #A1301.
6. C. E. Watts, *Intrinsic characterizations of some additive functors*, Proc. Amer. Math. Soc. **11** (1960), 5–8. MR **22** #9528.
7. N. Yoneda, *On Ext and exact sequences*, J. Fac. Sci. Univ. Tokyo Sect. I **8** (1960), 507–576. MR **37** #1445.

UNIVERSITY OF CALIFORNIA,
IRVINE, CALIFORNIA 92664