

THE ORIENTED BORDISM OF Z_{2^k} ACTIONS

BY

E. R. WHEELER

ABSTRACT. Let R_2 be the subring of the rationals given by $R_2 = Z[1/2]$. It is shown that for $G = Z_{2^k}$ the bordism group of orientation preserving G actions on oriented manifolds tensored with R_2 is a free $\Omega_* \otimes R_2$ module on even dimensional generators (where Ω_* is the oriented bordism ring).

1. Introduction. Let G be a group. Let Ω_*^G denote the bordism group of differentiable orientation preserving G actions on closed oriented manifolds. In R. E. Stong's paper [10] Ω_*^G is understood for G a p -group and p an odd prime. In [9] H. L. Rosenzweig shows that $\Omega_*^{Z^2} \otimes Q = 0$ if $* \neq 4k$. In this paper the module structure of Ω_*^G is determined up to 2-torsion for $G = Z_{2^k}$.

§§ 2, 3, and 4 are largely preliminary material. In §5 it is shown that $\Omega_*^{Z_{2^k}} \otimes R_2$ is a free $\Omega_* \otimes R_2$ module on even dimensional generators (where $R_n = \{a/b \mid a \text{ is an integer and } b \text{ is a power of } n\}$ is a subring of the rationals).

This paper discusses part of the research undertaken while I was a Ph. D. candidate at the University of Virginia. I would like to express my appreciation to my advisor, R. E. Stong, who directed this research in a most generous way.

2. Equivariant bordism. For a finite abelian group G a family F'' of subgroups of G is a collection of subgroups of G such that if $H \in F''$ and $K < H$, then $K \in F''$. If (M, σ) is a manifold with G action, then (M, σ) is F'' -free if for each $x \in M$, the isotropy subgroup of x is an element of F'' .

Let $F' \subset F''$ be families of subgroups of G . Let (X, A) be a space pair with G action. Consider 5-tuples (M, M_0, M_1, σ, f) where

(1) M, M_0, M_1 are compact differentiable oriented manifolds with n the dimension of M .

(2) $\partial M = M_0 \cup M_1, \partial M_0 = \partial M_1 = M_0 \cap M_1$.

(3) $\sigma: G \times M \rightarrow M$ is a differentiable G -action which preserves M_0 and M_1 and which preserves the orientation on M .

(4) (M, σ) is F'' -free while $(M_0, \sigma|_{G \times M_0})$ is F' -free.

(5) $f: (M, M_1) \rightarrow (X, A)$ is an equivariant map.

Received by the editors June 27, 1973 and, in revised form, November 5, 1973.

AMS (MOS) subject classifications (1970). Primary 57D85.

Key words and phrases. Equivariant bordism, orientation preserving group action.

Copyright © 1974, American Mathematical Society

Under the usual equivariant bordism relation (see [10, §2]) one forms a set of equivalence classes of such 5-tuples, denoted $\Omega_n^G(F'', F')(X, A)$, with an abelian group structure induced by disjoint union. The graded sum of these groups has an Ω_* module structure induced by cartesian product and is denoted by $\Omega_*^G(F'', F')(X, A)$.

Now if $h: (X, A) \rightarrow (Y, B)$ is an equivariant map between spaces with G action, one has an induced homomorphism $h_*: \Omega_*^G(F'', F')(X, A) \rightarrow \Omega_*^G(F'', F')(Y, B)$ sending $[M, M_0, M_1, \sigma, f]$ into $[M, M_0, M_1, \sigma, h \circ f]$. Let \emptyset denote the empty set. Then there is a degree -1 boundary map $\partial_*: \Omega_*^G(F'', F')(X, A) \rightarrow \Omega_*^G(F'', F')(A, \emptyset) \equiv \Omega_*^G(F'', F')(A)$ sending $[M, M_0, M_1, \sigma, f]$ into $[M_1, \partial M_1, \emptyset, \sigma/G \times M_1, f/M_1]$. From [10, Proposition 2.1], $\Omega_*^G(F'', F')(-)$ and ∂_* define an equivariant homology theory on the category of G -pairs to the category of Ω_* modules. Specifically this theory satisfies equivariant homotopy, excision, and exactness axioms.

From [10, Proposition 2.2], one knows that for families of subgroups $F' \subset F''$ there is an exact triangle

$$\begin{array}{ccc} \Omega_*^G(F')(X, A) & \xrightarrow{\alpha_*} & \Omega_*^G(F'')(X, A) \\ \partial'_* \swarrow & & \swarrow \beta_* \\ & \Omega_*^G(F'', F')(X, A) & \end{array}$$

in which α_* and β_* , respectively, forget F' and F'' -freeness while ∂'_* sends $[M, M_0, M_1, \sigma, f]$ into $[M_0, \emptyset, \partial M_0, \sigma/G \times M_0, f/M_0]$.

Note. If G is an abelian group and $H < G$, the collection of all subgroups of H is a family of subgroups of G . If $H \not\leq G$, this family is denoted by F_H . In particular F_e denotes the family consisting of the identity subgroup. Let F denote the family of all subgroups of G .

3. Classifying spaces for bundles with G action. Let G be a finite abelian group with exactly r distinct irreducible complex representations. Let $C^\infty = C_1^\infty \oplus C_2^\infty \oplus \dots \oplus C_r^\infty$. Define a G action on C^∞ by considering C_i^∞ as a countable direct sum of the i th irreducible representation. Now let BU_s be the Grassmannian of complex s -planes in C^∞ and γ_s be the universal complex s -plane bundle over BU_s . Since the elements of G act on C^∞ via complex linear transformations, there is an induced G action on BU_s and γ_s (see [10, §3]). One learns from Atiyah [2, §1.6] that $\gamma_s \rightarrow BU_s$ is the universal complex n -plane bundle in the category of G -spaces.

One can perform essentially the same construction in the real case by taking the Grassmannian of real n -planes in C^∞ . In this way one gets BO_s together with its canonical bundle γ_s , the universal real s -plane bundle in the category of G -spaces. Note that in what follows these G -spaces are called BU_s and BO_s .

except in cases where the context does not make the meaning clear. In these cases the notation (BU_s, G) and (BO_s, G) is used.

In the process of defining BU_s and BO_s together with their canonical bundles one may place a metric on the γ_s such that the G action is orthogonal with respect to this metric. Further, for any G -bundle $E \rightarrow X$ of dimension s over a compact Hausdorff space X , one may assume there is a metric on E such that

- (a) the G action on E is orthogonal with respect to this metric,
- (b) the bundle map covering the classifying map takes

$$(D(E), S(E)) \rightarrow (D(\gamma_s), S(\gamma_s))$$

where $D(-)$ denotes the unit disc bundle and $S(-)$ denotes the unit sphere bundle.

Now consider the G -spaces BO_s and BU_s and the fixed sets of subgroups of G acting on BO_s and BU_s . Let $H < G$ and X be a compact Hausdorff G -space. The isomorphism classes of G -bundles over X of real dimension s , $\text{vect}_s^G(X)$, are in 1-1 correspondence with the G -homotopy classes of equivariant maps from X into $BO_s, [X, BO_s]_G$. Now if $H < G$ fixes X , any equivariant map $X \rightarrow BO_s$ goes into the fixed set of H acting on $BO_s, F_H(BO_s)$. Hence if H fixes $X, \text{vect}_s^G(X) \leftrightarrow [X, F_H(BO_s)]_G$. It follows that $F_H(BO_s)$ is the classifying space of G bundles of dimension s over base spaces X such that H fixes X . Exactly the same analysis is true for complex s -bundles over X and $F_H(BU_s)$.

Further, if $E \rightarrow X$ is a complex G bundle and $H < G$ fixes X, E splits into G subbundles according to the nontrivial irreducible complex representations of H [2, §1.6]. The classifying space for G -bundles over a base which H fixes can be understood in terms of these subbundles. Using this information one can compute explicitly the fixed sets $F_H(BU_s)$. Using similar techniques one can understand $F_H(BO_s)$. In particular, for the purposes of this paper one records the following computations.

PROPOSITION 3.1. *If $H < G$ with $d =$ the order of H , then $F_H(BU_s, G)$ is G homotopy equivalent to $\bigcup BU_{t_1} \times \cdots \times BU_{t_d}$ where $\sum t_i = s$. \square*

Since the real irreducible representations of Z_2 are multiplication by $+1$ and by -1 on one-dimensional vector spaces, a Z_{2^k} bundle E over a Z_{2^k} space which is fixed by Z_2 decomposes into $E_1 \oplus E_{-1}$ where Z_2 acts in the fibers of E_i by multiplication by i . Thus the classifying space for s -dimensional real vector bundles over Z_{2^k} spaces fixed by Z_2 is $\bigcup BO_{t_1} \times BO_{t_{-1}}$ where $t_{-1} + t_1 = s$. Thus

PROPOSITION 3.2. $F_{Z_2}(BO_s, Z_{2k})$ is Z_{2k} homotopy equivalent to $\bigcup BO_{t-1} \times BO_{t1}$. \square

It is evident that the component of $F_{Z_2}(BO_s, Z_{2k})$ above which Z_2 acts as -1 in the fibers of the canonical bundle is a BO_s . Denote this component by $F_{Z_2}^-(BO_s, Z_{2k})$. The Z_{2k} action restricted to $F_{Z_2}(BO_s, Z_{2k})$ can be considered a Z_{2k-1} action. If $k > 1$ it is necessary to know the fixed set of $Z_{2j} < Z_{2k-1}$ acting on $F_{Z_2}^-(BO_s, Z_{2k})$. $F_{Z_{2j}}[F_{Z_2}^-(BO_s, Z_{2k})]$ is the classifying space for Z_{2k} bundles $E \rightarrow X$ which have the properties

(a) $Z_{2j+1} < Z_{2k}$ fixes X .

(b) $Z_2 < Z_{2j+1} < Z_{2k}$ acts on the fibers of E as multiplication by -1 .

For such a bundle E splits into subbundles with respect to the irreducible representations of Z_{2j+1} . Since each irreducible representation of Z_{2j+1} which satisfies (b) is the realification of an irreducible complex representation, each of the subbundles of E has a complex structure. Thus if there are r irreducible real representations of Z_{2j+1} satisfying (b) one has

PROPOSITION 3.3. $F_{Z_{2j}}[F_{Z_2}^-(BO_s, Z_{2k})]$ is Z_{2k-1} homotopy equivalent to $\bigcup BU_{t_1} \times \dots \times BU_{t_r}$ with $\sum t_i = s$. \square

4. A special case of equivariant transverse regularity. Let γ_{2s} represent the canonical $2s$ plane bundle over $F_{Z_2}^-(BO_{2s}, Z_{2k})$. (Note that (BO_{2s}, Z_{2k-1}) is Z_{2k-1} homotopy equivalent to $F_{Z_2}^-(BO_{2s}, Z_{2k})$.) Since $Z_2 < Z_{2k}$ acts as multiplication by -1 in the fibers of γ_{2s} and since the determinant of -1 acting on an even dimensional vector space is $+1$, the Z_2 action dies when one takes the determinant bundle of γ_{2s} together with its induced action. In other words, $\det \gamma_{2s} \rightarrow F_{Z_2}^-(BO_{2s}, Z_{2k})$ is a Z_{2k-1} bundle.

PROPOSITION 4.1. If

$$f: (M, \partial M, Z_{2k-1} \text{ action}) \rightarrow (D(\det \gamma_{2s}), S(\det \gamma_{2s}), \det(Z_{2k} \text{ action}))$$

is an equivariant map, then f may be equivariantly homotoped to be transverse regular on the zero section of $\det \gamma_{2s}$. Further, if $A \subset M$ is a closed subspace and if $f|_A$ is already transverse regular, the homotopy can be chosen to fix A .

PROOF. One needs only to check that the hypotheses for Lemma 4.2 in [10] are satisfied. Therefore one looks at the fixed set of $Z_{2j} < Z_{2k-1}$ acting on $F_{Z_2}^-(BO_{2s}, Z_{2k})$ for all $1 \leq j \leq k-1$, and one checks that if $x \in BO_{2s}$ is fixed by Z_{2j} , then v is fixed by Z_{2j} for all $v \in \det \gamma_{2s}/x$.

If T is the generator of Z_{2j+1} acting on γ_{2s} , T acts as a real linear transformation on γ_{2s}/x such that T^{2j} acts as multiplication by -1 . Further, the minimum polynomial of T , m_T , must divide $y^{2^{j+1}} - 1 = (y - 1) \cdot (y + 1) \cdot q_1(y) \cdot \dots \cdot q_{j-1}(y)$ where $q_i(y)$ is an irreducible quadratic of the form

$y^2 + ay + 1$. $y - 1$ does not divide m_T since this would imply that T is multiplication by 1 on some one-dimensional subspace of γ_{2s}/x . Elementary linear algebra then yields that $\det T = +1$ which implies that $\det T$ fixes pointwise the fiber $\det \gamma_{2s}/x$. \square

5. The oriented bordism of Z_{2k} . For a group G , denote by Ω_*^G the equivariant bordism module $\Omega_*^G(F)(pt)$. In this section $\Omega_*^{Z_{2k}} \otimes R_2$ is computed. Let (X, A) be a c.w. pair with Z_{2k} action having the property that $F_{Z_{2^j}}(X, A)$ is a c.w. pair for $0 \leq j \leq k$ where $F_{Z_{2^j}}(X, A)$ is the fixed set of Z_{2^j} , acting on (X, A) . For a bundle E with unit disc, $D(E)$, and unit sphere, $S(E)$, one denotes by $T(E)$ the space $D(E)/S(E)$, the Thom space of E . The primary tool of this paper is the following theorem.

THEOREM 5.1. $\Omega_*^{Z_{2k}}(F, F_e)(X, A)$ is isomorphic to

$$\bigoplus_{s=0}^{[* / 2]} \tilde{\Omega}_{*-2s+1}^{Z_{2^{k-1}}}(F)(F_{Z_2}(X)/F_{Z_2}(A) \wedge T(\det \gamma_{2s}))$$

where γ_{2s} is the canonical $2s$ plane bundle over $F_{Z_2}^-(BO_{2s}, Z_{2k})$.

PROOF. Let $[M, M_0, M_1, T, f] \in \Omega_*^{Z_{2k}}(F, F_e)(X, A)$ where T generates the Z_{2k} action on M . Let F_2 be the $(n - s)$ -dimensional component of the fixed set of $Z_2 < Z_{2k}$ acting on M . Then F_2 is a submanifold of M with an induced action of $Z_{2^{k-1}}$ which is covered in the normal bundle to F_2 in M , ν , by an action of Z_{2k} . Further, $\partial F_2 = F_2 \cap M_1$. Since one may identify the disc of the normal bundle equivariantly with a small tubular neighborhood of F_2 , one knows that no elements of the disc of the normal bundle - {zero section} can be fixed by Z_{2^j} for $1 \leq j \leq k$. Since each fiber of ν is a representation space for Z_2 , ν is a Z_{2k} bundle over F_2 such that Z_2 acts as -1 in the fibers. One then knows that $\nu \rightarrow F_2$ is classified equivariantly into $F_{Z_2}^-(BO_{2s}, Z_{2k})$ yielding a Z_{2k} bundle map

$$\begin{array}{ccc} \nu & \xrightarrow{g'} & \gamma_{2s} \\ \pi \downarrow & & \downarrow \\ F_2 & \xrightarrow{g} & BO_{2s} \end{array}$$

By taking the determinant bundles of ν and γ_{2s} one gets a similar diagram of $Z_{2^{k-1}}$ bundle maps.

One may assume that $\det g'$ maps the (D, S) pair of $\det \nu$ into the (D, S) pair of $\det \gamma_{2s}$. Letting $\tilde{\pi}: \det \nu \rightarrow F_2$ be the projection, and crossing $\det g'$ with $f \circ \tilde{\pi}$, one gets a map from the pair

$$(D(\det \nu), D(\det \nu / \partial F_2) \cup S(\det \nu))$$

into

$$(F_{Z_2}(X, A) \times (D(\det \gamma_{2s}), S(\det \gamma_{2s}))).$$

Since the first Stiefel-Whitney classes of ν and the tangent bundle of F_2 , $\tau(F_2)$, are equal, $D(\det \nu)$ is an oriented manifold. Let T' generate the $Z_{2^{k-1}}$ action on $D(\det \nu)$. One notes that T' is orientation preserving if $\det(dT') = T' \times 1$ on $\det \tau(D(\det \nu))$ [6, Lemma 3]. Since $\det dT' = \tilde{\pi}^*(\det dT)$ on $\tilde{\pi}^*(\det \tau(M)/F_2) \cong \det \tau(D(\det \nu))$ it follows that T' is orientation preserving since T is orientation preserving. Thus by summing over the discs of the determinant bundles of all possible components of the fixed set of Z_2 , one may define a map

$$F: \Omega_*^{Z_2^k}(F, F_e)(X, A) \rightarrow \bigoplus_{s=0}^{[* / 2]} \Omega_*^{Z_2^{k-1}}(F)(F_{Z_2}(X, A) \times (D(\det \gamma_{2s}), S(\det \gamma_{2s}))).$$

In order to define an inverse to F consider

$$[N, \partial N, S, h] \in \Omega_*^{Z_2^{k-1}}(F)(F_{Z_2}(X, A) \times (D(\det \gamma_{2s}), S(\det \gamma_{2s}))).$$

One has $N \xrightarrow{p_2 \circ h} D(\det \gamma_{2s})$ and $p_2 \circ h$ is an equivariant map which, by Proposition 4.1, may be considered to be transverse regular on the zero section, BO_{2s} , of $\det(\gamma_{2s})$. Let $N' = (p_2 \circ h)^{-1}(BO_{2s})$. Since γ_{2s} has a Z_{2^k} action covering the $Z_{2^{k-1}}$ action on BO_{2s} , $(p_2 \circ h)^*(\gamma_{2s}) \xrightarrow{\pi'} N'$ is a bundle with an induced Z_{2^k} action such that $Z_2 < Z_{2^k}$ acts as -1 in the fibers. Let S' generate the Z_{2^k} action on $(p_2 \circ h)^*(\gamma_{2s})$. $D((p_2 \circ h)^*(\gamma_{2s}))$ is oriented and one checks that $\det dS'$ acts as $S' \times 1$ on the determinant of the tangent bundle. Hence S' is orientation preserving by [6, Lemma 3]. Hence by mapping $[N, \partial N, S, h]$ into

$$[D((p_2 \circ h)^*(\gamma_{2s})), S(p_2 \circ h)^*(\gamma_{2s}), D((p_2 \circ h)^*(\gamma_{2s})/\partial N'), S', p_1 \circ h \circ \pi']$$

one defines a map K from

$$\Omega_*^{Z_2^{k-1}}(F)(F_{Z_2}(X, A) \times (D(\det \gamma_{2s}), S(\det \gamma_{2s})))$$

into $\Omega_*^{Z_2^k}(F, F_e)(X, A)$.

To see that $F \circ K = \text{id}$ one notes that $D[(p_2 \circ h)^*(\gamma_{2s})]$ may be regarded as a tubular neighborhood of N' in N . By a deformation one may assume that $p_2 \circ h$ maps

$$\{N - [D(\det(p_2 \circ h)^*(\gamma_{2s})) - S(\det(p_2 \circ h)^*(\gamma_{2s}))]\}$$

into $S(\det \gamma_{2s})$. Let π'' be the bundle projection, $\pi'': (p_2 \circ h)^*(\det \gamma_{2s}) \rightarrow N'$. Since N' is a strong equivariant homotopy retract of its tubular neighborhood, there is an equivariant homotopy $J: N \times I \rightarrow F_{Z_2}(X)$ giving a homotopy

between $p_1 \circ h$ and J_1 where J_1 has the property that J_1 on $D((p_2 \circ h)^*(\det \gamma_{2s}))$ is given by $(p_1 \circ h/N') \circ \pi''$. It follows that

$$\{N \times I, \partial N \times I \cup N \times 1 - \text{int } D((p_2 \circ h)^*(\det \gamma_{2s})), S \times 1, (J \times (p_2 \circ h)) \times 1\}$$

gives a bordism between $[N, \partial N, S, h]$ and $F \circ K([N, \partial N, S, h])$.

To see that $K \circ F = \text{id}$ it suffices to observe that F_2 is a strong equivariant retract of its tubular neighborhood, $D(\nu)$, and hence one may suppose f is homotopic to a map H such that $H/D(\nu) = f/F_2 \circ \pi$. Now $K \circ F$ is obtained by restricting to $D(\nu)$. Since Z_{2^k} acts freely in the complement of F_2 , $[M, M_0, M_1, T, f] = K \circ F([M, M_0, M_1, T, f])$. \square

Now suppose (X, A) is a c.w. pair acted on by $G = Z_{2^k}$. Let q denote the quotient map onto the space pair $(X/G, A/G)$ obtained by identifying the orbits of the G action. It is a well-known fact that $q^*: H^*(X/G, A/G; R_2) \rightarrow H^*(X, A; R_2)$ is a monomorphism onto the elements of $H^*(X, A; R_2)$ which are invariant under the G action (see [5, Corollary 2.3]). This fact together with the appropriate universal coefficient theorem indicates that if $H_*(X, A; R_2)$ is a free R_2 module on even [odd] dimensional generators, then $H_*(X/G, A/G; R_2)$ is a free R_2 module on even [odd] dimensional generators.

In light of this fact one defines a space pair (X, A) to be (2-even) [(2-odd)] if and only if $H_*(X, A; R_2)$ is a free R_2 module on even [odd] dimensional generators.

LEMMA 5.2. *Let $G = Z_{2^k}$. If (X, A) is a G pair and if (X, A) is (2-even) [(2-odd)], then $\Omega_*^G(F_e)(X, A) \otimes R_2$ is a free $\Omega_* \otimes R_2$ module on even [odd] dimensional generators.*

PROOF. From Proposition 2.3 in [10] one learns that $\Omega_*^G(F_e)(X, A) \cong \Omega_*(X \times_G EG, A \times_G EG)$ where EG is the total space of the universal principal G bundle. From the discussion preceding this lemma one learns that $(X \times_G EG, A \times_G EG)$ is (2-even) [(2-odd)]. As in [11, p. 145] one can show that if (X, A) is a c.w. pair such that $H_*(X, A; R_2)$ is a torsion free R_2 module, then

$$\Omega_*(X, A) \otimes R_2 \cong (\Omega_* \otimes R_2) \otimes_{R_2} H_*(X, A; R_2).$$

This yields the desired result. \square

Thus it is of interest to examine the homology of the spaces introduced in Theorem 5.1. From the homology exact sequence of the cofibration $S(\det \gamma_{2s}) \rightarrow D(\det \gamma_{2s}) \rightarrow T(\det \gamma_{2s})$ in which BSO_{2s} is homotopy equivalent to $S(\det \gamma_{2s})$ and BO_{2s} is homotopy equivalent to $D(\det \gamma_{2s})$ one learns that $T(\det \gamma_{2s})$ is (2-odd). From the proof of Proposition 4.1 one knows that for $Z_{2^j} < Z_{2^{k-1}}$,

$$F_{Z_2^j}(T(\det \gamma_{2s})) = T(\det \gamma_{2s}/F_{Z_2^j}[F_{Z_2}^-(BO_{2s}, Z_2^k)]).$$

If $E \rightarrow X$ is an oriented bundle, $\det E$ is a trivial line bundle and thus $T(\det E) = \Sigma X^+$ where Σ denotes reduced suspension. Now by Proposition 3.3, $F_{Z_2^j}[F_{Z_2}^-(BO_{2s}, Z_2^k)]$ is homotopic to $\bigcup BU_{(t)}$ where (t) is a q -tuple of nonnegative integers (t_1, t_2, \dots, t_q) and $BU_{(t)} = BU_{t_1} \times BU_{t_2} \times \dots \times BU_{t_q}$. Since $\gamma_{2s}/BU_{(t)}$ is complex, $\det \gamma_{2s}/BU_{(t)}$ is trivial and $T(\det \gamma_{2s}/\bigcup BU_{(t)}) = \bigvee \Sigma BU_{(t)}^+$. It follows that $F_{Z_2^j}(T(\det \gamma_{2s}))$ is (2-odd) for $0 \leq j \leq k-1$. Now from the appropriate cofibrations $X \vee Y \rightarrow X \times Y \rightarrow X \wedge Y$ one reads off the result:

LEMMA 5.3. *If $F_{Z_2^j}(X, A)$ is (2-even) [(2-odd)] for $0 \leq j \leq k$ and if $Y = F_{Z_2}(X)/F_{Z_2}(A) \wedge T(\det \gamma_{2s})$, then Y is a space with $Z_{2^{k-1}}$ action such that $F_{Z_2^j}(Y)$ is (2-odd) [(2-even)] for $0 \leq j \leq k-1$. \square*

This brings one finally to the computations.

THEOREM 5.4. *If $F_{Z_2^j}(X, A)$ is (2-even) [(2-odd)] for $0 \leq j \leq k$, then $\Omega_*^{Z_2^k}(F)(X, A) \otimes R_2$ is a free $\Omega_* \otimes R_2$ module on even [odd] dimensional generators.*

PROOF. If $k = 0$ then both the even and the [odd] case follow from Lemma 5.2. Assume that the theorem is true for $k' < k$. Let (X, A) have a Z_{2^k} action satisfying the hypotheses. By Lemma 5.2, $\Omega_*^{Z_2^k}(F_e)(X, A) \otimes R_2$ is a free $\Omega_* \otimes R_2$ module on even [odd] dimensional generators. Theorem 5.1 yields that

$$\Omega_*^{Z_2^k}(F, F_e)(X, A) \cong \bigoplus \tilde{\Omega}_{*-2s+1}^{Z_2^{k-1}}(F)(F_{Z_2}(X)/F_{Z_2}(A) \wedge T(\det \gamma_{2s})).$$

By Lemma 5.3 and induction hypothesis, this implies that $\Omega_*^{Z_2^k}(F, F_e)(X, A) \otimes R_2$ is a free $\Omega_* \otimes R_2$ module on even [odd] dimensional generators.

Now consider the exact triangle

$$\Omega_*^{Z_2^k}(F_e)(X, A) \otimes R_2 \rightarrow \Omega_*^{Z_2^k}(F)(X, A) \otimes R_2 \rightarrow \Omega_*^{Z_2^k}(F, F_e)(X, A) \otimes R_2.$$

$$\begin{array}{c} \uparrow \hspace{15em} \downarrow \\ \hline \partial'_* \text{ of degree } -1 \end{array}$$

Note that it is in fact a split short exact sequence. This gives the induction step. \square

Note. If $(X, A) = (pt, \emptyset)$, Theorem 5.4 says that $\Omega_*^{Z_2^k} \otimes R_2$ is a free $\Omega_* \otimes R_2$ module on even dimensional generators.

Note. This is the best possible result in the following sense. In [3, p. 105] P. E. Conner computes the torsion of $\Omega_*^{Z_2}$. There is too much torsion for $\Omega_*^{Z_2}$ to be a free Ω_* module.

Note. In the paper as originally submitted the author asserted that $\Omega_*^G \otimes R_2$ is a free $\Omega_* \otimes R_2$ module for G any finite cyclic group. However, the

referee kindly noted a logical error in the author's proof of this statement. Nonetheless it is still a very reasonable conjecture, and in fact seems to be true in certain special cases (e.g. $Z_2 \times Z_p$). Along this line it should also be noted that in the author's dissertation [13] he proves via a somewhat arduous and noninstructive argument that for G a finite cyclic group the torison of Ω_*^G is all 2-torsion.

BIBLIOGRAPHY

1. M. F. Atiyah, *Bordism and cobordism*, Proc. Cambridge Philos. Soc. 57 (1961), 200–208. MR 23 #A4150.
2. ———, *K-theory*, Lecture notes by D. W. Anderson, Benjamin, New York, 1967. MR 36 #7130.
3. P. E. Conner, *Lectures on the action of a finite group*, Lecture Notes in Math., no. 73, Springer-Verlag, Berlin and New York, 1968. MR 41 #2670.
4. P. E. Conner and E. E. Floyd, *Differentiable periodic maps*, Ergebnisse der Math. und ihrer Grenzgebiete, Band 33, Academic Press, New York; Springer-Verlag, Berlin, 1964. MR 31 #750.
5. E. E. Floyd, *Periodic maps via Smith theory*, Seminar on Transformation Groups, Ann. of Math. Studies, no. 46, Princeton Univ. Press, Princeton, N. J., 1960, Chap. III.
6. K. Komiya, *Oriented bordism and involutions*, Osaka J. Math. 9 (1972), 165–181. MR 46 #6341.
7. P. S. Landweber, *Equivariant bordism and cyclic groups*, Proc. Amer. Math. Soc. 31 (1972), 564–570. MR 45 #6028.
8. E. Ossa, *Unitary bordism of abelian groups*, Proc. Amer. Math. Soc. 33 (1972), 568–571. MR 45 #2743.
9. H. L. Rosenzweig, *Bordism of involutions on manifolds*, Illinois J. Math. 16 (1972), 1–10. MR 44 #7568.
10. R. E. Stong, *Complex and oriented equivariant bordism*, Topology of Manifolds (Proc. Inst., Univ. of Georgia, Athens, Ga., 1969), Markham, Chicago, Ill., 1970, pp. 291–316. MR 42 #8521.
11. ———, *Notes on cobordism theory*, Mathematical Notes, Princeton Univ. Press, Princeton, N. J.; Univ. of Tokyo Press, Tokyo, 1968. MR 40 #2108.
12. ———, *Unoriented bordism and actions of finite groups*, Mem. Amer. Math. Soc. No. 103 (1970). MR 42 #8522.
13. E. R. Wheeler, *The oriented bordism of cyclic, group actions*, Dissertation, University of Virginia, Charlottesville, Va., 1973.

DEPARTMENT OF MATHEMATICS, NORTHERN KENTUCKY STATE COLLEGE,
HIGHLAND HEIGHTS, KENTUCKY 41076