

ω -COHESIVE SETS

BY

BARBARA F. RYAN⁽¹⁾

ABSTRACT. We define and investigate ω -cohesiveness, a strong notion of indecomposability for subsets of the integers and their isols. This notion says, for example, that if X is the isol of an ω -cohesive set then, for any integer n , $Y + Z = \binom{X}{n}$ implies that, for some integer k , $\binom{X-k}{n} \in Y$ or Z . From this it follows that if $f(x) \in T_1$, the collection of almost recursive combinatorial polynomials, then the predecessors of $f_\Lambda(X)$ are limited to isols $g_\Lambda(X)$ where $g(x) \in T_1$. We show existence of ω -cohesive sets. And we show that the isol of an ω -cohesive set is an n -order indecomposable isol as defined by Manaster. This gives an alternate proof to one half of Ellentuck's theorem showing a simple algebraic difference between the isols and cosimple isols. In the last section we study functions of several variables when applied to isols of ω -cohesive sets.

1. Introduction. Let E denote the nonnegative integers, $P(E)$ all subsets of E , Λ the isols, and $\langle \omega \rangle$ the recursive equivalence type of $\alpha \subseteq E$. Let J be a fully effective map from $\bigcup E^n$ ($n \in E$), one-one onto E . For $\alpha \in P(E)$, $n > 0$, define

$$\alpha^{(n)} = \{(a_1, \dots, a_n) \in \alpha^n \mid a_1 > \dots > a_n\} \quad \text{and} \quad \binom{\alpha}{n} = J(\alpha^{(n)}).$$

For $X \in \Lambda$, let $\binom{X}{n}$ denote $\langle J(\binom{\alpha}{n}) \rangle$ where $\langle \omega \rangle = X$. An isolated set α is called cohesive if for all r.e. sets ω , there is a finite set $\beta \subseteq \alpha$ such that either $(\alpha - \beta) \subseteq \omega$ or $(\alpha - \beta) \subseteq E - \omega$. In this paper, we investigate the following stronger notions of cohesiveness.

DEFINITION. For $n > 0$, an infinite set $\alpha \subseteq E$ is n -cohesive if for all r.e. sets ω , there is a finite set $\beta \subseteq \alpha$ such that either $\binom{\alpha - \beta}{n} \subseteq \omega$ or $\binom{\alpha - \beta}{n} \subseteq E - \omega$.

DEFINITION. A set $\alpha \subseteq E$ is ω -cohesive if α is n -cohesive for every integer $n \in E$.

Received by the editors November 13, 1972 and, in revised form, January 16, 1974.

AMS (MOS) subject classifications (1970). Primary 02F40.

Key words and phrases. Isols, ω -cohesive sets, almost recursive combinatorial functions, predecessors of isols, higher-order indecomposable isols, universal isols.

⁽¹⁾ This paper is essentially the author's doctoral dissertation at Cornell University. The author wishes to thank Professor Anil Nerode for his patience, encouragement and many stimulating hours of conversation.

It is clear that if $\alpha \subseteq E$ is n -cohesive (ω -cohesive) and β is recursively isomorphic to α then β is also n -cohesive (ω -cohesive). Let Λ_c be the collection of all isols of ω -cohesive sets.

In §2, we show the existence of ω -cohesive sets, and in particular show that any infinite set of integers has an ω -cohesive subset. We also show that if α is n -cohesive then $\binom{\alpha}{n} \in P_n - S_n$, the n -order indecomposables (Manaster's sets P_n, S_n are defined in §2). Thus we have a very simple method of constructing n -order indecomposable isols, where n is a finite ordinal. Unfortunately our techniques do not seem to extend to the transfinite case. Now if we take $X \in \Lambda_c$ then $\binom{X}{n} \in P_n - S_n$ for all $n > 0$. So we have a "uniform" procedure for constructing finite indecomposable isols. Using a more general technique than ours, Ellentuck in [3] proves the following restatement of the last result: There is an $X \in \Lambda$ such that $\binom{X}{n} \in P_n - S_n$ for all $n > 0$. By showing that the same statement is false in the cosimple isols, he has a simple algebraic difference between the two theories.

We also investigate how a function of an isol in Λ_c can be decomposed. Let $T_1 =$ the collection of all almost recursive combinatorial polynomials of one variable, i.e. a function $f: E \rightarrow E$ is in T_1 iff there is an integer $k \geq 0$ such that, for some finite string of nonnegative integers $c_0, c_1, \dots, c_n, f(x + k) = \sum c_i \binom{x}{i}$. Let f_Λ be the canonical extension of f to the isols. (For a discussion of combinatorial functions, almost recursive combinatorial functions and extension procedures see [5] and [6].) For $f, g \in T_1$, define $f \leq g$ if there is $h \in T_1$ such that $f + h = g$. And define an equivalence relation on T_1 by $f \sim g$ if there is an integer $k \geq 0$ such that $f(x + k) = g(x + k)$. Then Theorem 3 says that the only predecessors of $f_\Lambda(X)$, where $f \in T_1$ and $X \in \Lambda_c$, are isols of the form $g_\Lambda(X)$ for some $g \in T_1$ with $g \leq f$. Since $g \leq f$ implies that $g_\Lambda(X) \leq f_\Lambda(X)$ for all $X \in \Lambda$, it is always the case that an isol of the form $g_\Lambda(X)$ is a predecessor of $f_\Lambda(X)$. Theorem 3 says these are the only predecessors of $f_\Lambda(X)$ if $X \in \Lambda_c$.

It is easy to show that any ω -cohesive set has a subset whose isol is universal. This allows us (in Theorem 4) to define a map $\theta_1: T_1 \rightarrow \Lambda$ which preserves addition and composition, and such that $f \sim g$ iff $\theta_1(f) = \theta_1(g)$, and $\theta_1(T_1)$ is an ideal in Λ .

Let $T_i =$ the collection of all almost recursive combinatorial polynomials $f(x_i)$ of the variable x_i . Let $T_\infty =$ the collection of all finite sums of functions in $\bigcup T_i$ ($i \in E$). T_∞ is closed under addition, composition and predecessor. In §3, we extend Theorems 3 and 4 of §2 to functions in T_∞ .

Theorem 3 does not hold if we allow product terms in our functions. For example if $f(x_1, x_2) = x_1 \cdot x_2$ and $X_1, X_2 \in \Lambda_c$ then in general the predecessors of $f_\Lambda(X_1, X_2)$ are not restricted to isols of the form $g_\Lambda(X_1, X_2)$ for

some $g \leq f$. §4 gives a characterization of the predecessors of $f_{\wedge}(X_1, \dots, X_n)$ where f is an arbitrary almost recursive combinatorial polynomial and X_1, X_2, X_3, \dots is a "universal sequence" of isols (see §§3 and 4 for definitions).

2. Basic results.

LEMMA 1. *For any n and any infinite subset α of E , α has an n -cohesive subset β .*

PROOF. Let $\omega_1, \omega_2, \dots$ be a list of all r.e. sets. We will construct a sequence $\alpha_0 \supseteq \alpha_1 \supseteq \alpha_2 \supseteq \dots$ of infinite subsets of α such that, for any $i > 0$,

$$\binom{\alpha_i}{n} \subseteq \omega_i \quad \text{or} \quad \binom{\alpha_i}{n} \subseteq E - \omega_i.$$

Let $\alpha_0 = \alpha$. Suppose we have α_i . Then

$$\mathcal{J}^{-1}\left[\binom{\alpha_i}{n} \cap \omega_{i+1}\right] \quad \text{and} \quad \mathcal{J}^{-1}\left[\binom{\alpha_i}{n} \cap (E - \omega_{i+1})\right]$$

are partitions of $\alpha_i^{(n)}$ into two disjoint sets. By Ramsey's theorem [7] α_i has an infinite subset γ such that either

$$\gamma^{(n)} \subseteq \mathcal{J}^{-1}\left[\binom{\alpha_i}{n} \cap \omega_{i+1}\right] \quad \text{or} \quad \gamma^{(n)} \subseteq \mathcal{J}^{-1}\left[\binom{\alpha_i}{n} \cap (E - \omega_{i+1})\right].$$

Therefore $\binom{\gamma}{n} \subseteq \omega_{i+1}$ or $\binom{\gamma}{n} \subseteq (E - \omega_{i+1})$. Let $\alpha_{i+1} = \gamma$. Now, for each $i \in E$, choose $x_i \in \alpha_i$ different from x_0, \dots, x_{i-1} (since each α_i is infinite this is always possible). Let $\beta = \{x_i | i \in E\}$. Then β is an infinite subset of α .

We claim β is n -cohesive. Suppose ω is any r.e. set. Then there is an i such that $\omega = \omega_i$. Let $\beta' = \{x_j | j \geq i\}$. Then $\beta' \subseteq \alpha_i$. We chose α_i such that either

$$\binom{\alpha_i}{n} \subseteq \omega_i \quad \text{or} \quad \binom{\alpha_i}{n} \subseteq E - \omega_i.$$

Therefore, $\binom{\beta'}{n} \subseteq \omega_i$ or $\binom{\beta'}{n} \subseteq E - \omega_i$. Since $\beta' = \beta - \{x_1, \dots, x_{i-1}\}$, β is n -cohesive.

LEMMA 2. *Any infinite subset of an n -cohesive set is n -cohesive.*

PROOF. Follows directly from the definition.

LEMMA 3. *If α is n -cohesive and $m \leq n$, then α is m -cohesive.*

PROOF. We need to show only that α n -cohesive implies α $(n - 1)$ -cohesive. Suppose α is n -cohesive but not $(n - 1)$ -cohesive. Then there is an r.e. set ω such that, for all finite subsets β of α , $\binom{\alpha - \beta}{n-1} \not\subseteq \omega$ and $\binom{\alpha - \beta}{n-1} \not\subseteq E - \omega$. Let $A =$

$J^{-1}(\omega) \times E$. Let $\omega_1 = J(A)$. Then ω_1 is r.e. Suppose there is a finite subset β of α such that $(\alpha - \beta) \subseteq \omega_1$. Then $(\alpha - \beta)^{(n)} \subseteq A$. Let x_0 be the smallest element of $(\alpha - \beta)$. Then it follows that $[\alpha - (\beta \cup \{x_0\})]^{(n-1)} \subseteq J^{-1}(\omega)$. Therefore,

$$\binom{\alpha - (\beta - \{x_0\})}{n - 1} \subseteq \omega.$$

Similarly if $(\alpha - \beta) \subseteq E - \omega_1$.

COROLLARY 1. *If α is n -cohesive then $\langle \alpha \rangle \in \Lambda$.*

PROOF. If α is n -cohesive then α is 1-cohesive or cohesive and hence isolated.

LEMMA 4. *If X is the isol of an n -cohesive set then $Y + Z = \binom{X}{n}$ implies that there is a $k \in E$ such that $\binom{X-n-k}{n} \leq Y$ or $\binom{X-n-k}{n} \leq Z$.*

PROOF. Clear from the definition of n -cohesive.

In [4] Manaster defines a sequence of triples $(P_\alpha, S_\alpha, I_\alpha)$ as follows: $I_0 = \{X | X \text{ is finite}\}$. For ordinals $\alpha > 0$,

$$P_\alpha = \left\{ X | X = Y + Z \Rightarrow Y \in \bigcup_{\alpha' < \alpha} I_{\alpha'}, \forall Z \in \bigcup_{\alpha' < \alpha} I_{\alpha'} \right\}$$

$$S_\alpha = \left\{ X | X = Y + Z \wedge Z \notin \bigcup_{\alpha' < \alpha} I_{\alpha'} \right.$$

$$\left. \Rightarrow (\exists V)(\exists W) \left[Z = V + W \wedge V \notin \bigcup_{\alpha' < \alpha} I_{\alpha'}, \wedge W \notin \bigcup_{\alpha' < \alpha} I_{\alpha'} \right] \right\}.$$

I_α is the ideal generated by $P_\alpha \cup S_\alpha$. It can be shown that $\alpha < \beta \Rightarrow P_\alpha \subseteq P_\beta$, $S_\alpha \subseteq S_\beta$, $I_\alpha \subseteq I_\beta$; $\bigcup_{\alpha' < \alpha+1} I_{\alpha'} = I_\alpha$; $P_\alpha \cap S_\alpha = \bigcup_{\alpha' < \alpha} I_{\alpha'}$.

In the following theorem we consider only $(P_\alpha, S_\alpha, I_\alpha)$ where α is finite. Elements of $P_n - S_n$ are called n -order indecomposables.

THEOREM 1. *If X is the isol of an n -cohesive set then $\binom{X}{n} \in P_n - S_n$ and $X^n \in I_n - S_n$.*

PROOF. We prove the first part of the theorem by induction on n . If $n = 1$ then X is the isol of a cohesive set and therefore $X \in P_1 - S_1$. Assume the theorem is true for $m \leq n - 1$. Suppose $\binom{X}{n} = Y + Z$. Then by Lemma 4, there is a $k \in E$ such that $\binom{X-n-k}{n} \leq Y$ or $\binom{X-n-k}{n} \leq Z$. Since

$$\binom{X}{n} = \binom{X-k}{n} + k \binom{X-k}{n-1} + \binom{k}{2} \binom{X-k}{n-2} + \dots + \binom{k}{n-1} \binom{X-k}{n-k} + \binom{k}{n},$$

either

$$Z \leq \binom{k}{1} \binom{X-k}{n-1} + \binom{k}{2} \binom{X-k}{n-2} + \dots + \binom{k}{n-1} \binom{X-k}{1} + \binom{k}{n}$$

or

$$Y \leq \binom{k}{1} \binom{X-k}{n-1} + \binom{k}{2} \binom{X-k}{n-2} + \dots + \binom{k}{n-1} \binom{X-k}{1} + \binom{k}{n}$$

Since X is the isol of an n -cohesive set, $X - k$ is the isol of an n -cohesive set. Then, by Lemma 3, $X - k$ is the isol of an m -cohesive set for $1 \leq m \leq n - 1$. By the inductive hypothesis, for $1 \leq m \leq n - 1$, $\binom{X-k}{m} \in P_m - S_m \subseteq P_m \subseteq P_{n-1} \subseteq I_{n-1}$. Since I_{n-1} is an ideal

$$k \binom{X-k}{n-1} + \dots + \binom{k}{n-1} \binom{X-k}{1} + \binom{k}{n} \in I_{n-1}.$$

Therefore, $Y \in I_{n-1}$ or $Z \in I_{n-1}$. Therefore, by definition of P_n , $\binom{X}{n} \in P_n$. By inductive hypothesis $\binom{X}{n-1} \in P_{n-1} - I_{n-1}$. And, for any integer r , $r \binom{X}{n-1} \leq \binom{X}{n}$. It is shown in Theorem 8.1 of [4] that for any $Z \in I_\alpha$, there is a finite m such that any linear decomposition of Z includes at most m isols of $P_\alpha - S_\alpha$. Hence $\binom{X}{n} \notin I_{n-1}$. Since

$$P_n - S_n = P_n - \bigcup_{\alpha < n} I_\alpha = P_n - I_{n-1}, \quad \binom{X}{n} \in P_n - S_n.$$

It is clear that the function x^n is an almost recursive combinatorial polynomial of the form, $x^n = c_n \binom{x}{n} + c_{n-1} \binom{x}{n-1} + \dots + c_1 x + c_0$. By the first part of the theorem, for $1 \leq i \leq n$, $\binom{x}{i} \in P_i - S_i \subseteq I_n$. Therefore, $X^n \in I_n - S_n$.

THEOREM 2. *If α is an infinite subset of E , then α has a subset β which is ω -cohesive.*

PROOF. By Lemma 1, there is a sequence $\alpha \supseteq \alpha_1 \supseteq \dots$ of infinite sets such that α_n is n -cohesive. For each n , choose $x_n \in \alpha_n$ different from x_1, \dots, x_{n-1} . Let $\beta = \{x_n | n > 0\}$. Claim β is ω -cohesive. For any n , $\beta = \{x_1, \dots, x_{n-1}\} \cup \beta'$ where $\beta' \subseteq \alpha_n$. Thus β' is n -cohesive. Therefore β is n -cohesive.

COROLLARY 2. *There is an isol X such that $\binom{X}{n} \in P_n - S_n$ for all $n > 0$.*

LEMMA 5. *Any infinite subset of an ω -cohesive set is ω -cohesive.*

PROOF. Follows from the definition of ω -cohesive and Lemma 2.

THEOREM 3. *If $f(x) \in T_1$, $X \in \Lambda_c$ and $Y \leq f_\Lambda(X)$, then there is a $g(x) \in T_1$ such that $g_\Lambda(X) = Y$ and $g \leq f$.*

PROOF. First suppose $f(x) = \binom{x}{i}$. Induct on i . If $i = 0$, then $f(x) = 1$ and the theorem is true. Assume the theorem is true for $i - 1$. Suppose $f(x) = \binom{x}{i}$. Suppose $Y \leq f_\Lambda(X) = \binom{X}{i}$. Then by Lemma 4, there is a $k \in E$ such that either $\binom{X-k}{i} \leq Y$ or $\binom{X-k}{i} \leq \binom{X}{i} - Y$. In the first case, $Y = \binom{X-k}{i} + Z$ where $Z \leq \binom{X}{i} - \binom{X-k}{i} \leq r \binom{X-1}{i-1}$ for some integer r . Then $Z = Z_1 + \dots + Z_r$ where $Z_j \leq \binom{X}{i-1}$ for each $1 \leq j \leq r$. By the inductive hypothesis, for each $1 \leq j \leq r$ there is a $g_j(x) \leq \binom{x-1}{i-1}$ such that $g_{j\Lambda}(X) = Z_j$. Let $g_0(x) = \binom{x-k}{i}$ and $g = g_0 + g_1 + \dots + g_r$. Then $g(x) \leq f(x)$ and $g(X) = Y$. In the second case, $\binom{X-k}{i} \leq \binom{X}{i} - Y$ implies that $Y \leq \binom{X}{i} - \binom{X-k}{i}$ and then the rest follows as above.

Suppose $f(x) \in T_1$ and $Y \leq f_\Lambda(X)$ then there exists $k \in E$ such that $f(x+k) = \sum_{i=0}^n c_i \binom{x}{i}$ for some $n, c_0, \dots, c_n \in E$ and $Y \leq f_\Lambda(X) \leq f_\Lambda(X+k) = \sum_{i=0}^n c_i \binom{X}{i}$. Then

$$Y = \sum_i \sum_j Y_{ij} \quad (0 \leq i \leq n, 1 \leq j \leq c_i)$$

where $Y_{ij} \leq \binom{X}{i}$ for $0 \leq i \leq n$ and $1 \leq j \leq c_i$. Therefore, for $0 \leq i \leq n$ and $1 \leq j \leq c_i$, there exists $g_{ij}(x) \in T_1$ such that $g_{ij\Lambda}(X) = Y_{ij}$. Let

$$g = \sum_i \sum_j g_{ij} \quad (0 \leq i \leq n, 1 \leq j \leq c_i).$$

Then $g_\Lambda(X) = Y$.

In [2], Ellentuck defines the notion of a universal isol. It is easy to see that his definition is equivalent to the following: an isol X is universal if for any pair of almost recursive combinatorial functions f and g , $f_\Lambda(X) = g_\Lambda(X)$ implies that there is an integer k such that for any $x \geq k$, $f(x) = g(x)$. We want to show that any ω -cohesive set has a subset whose isol is universal. Modify Ellentuck's method of showing the existence of universal isols [2] by replacing the set of integers E with an arbitrary infinite set $\alpha \subseteq E$ and topologize $P(\alpha)$ as he topologizes $P(E)$. Then Lemmas 1, 2 and Theorem 1 of [2] go through routinely to give the following lemma.

LEMMA 6. *Any infinite set of integers has a subset whose isol is universal.*

THEOREM 4. *There is a map $\theta_1: T_1 \rightarrow \Lambda$ such that for any $f, g \in T_1$*

- (1) $f \sim g$ iff $\theta_1(f) = \theta_1(g)$,
- (2) $\theta_1(f(g)) = f_\Lambda(\theta_1(g))$,

- (3) $\theta_1(f + g) = \theta_1(f) + \theta_1(g)$,
- (4) $\theta_1(T_1)$ is an ideal in Λ .

PROOF. Choose a set α which is ω -cohesive. By Lemma 6, α has a subset β such that $X = \langle \beta \rangle$ is universal. Define $\theta_1: T_1 \rightarrow \Lambda$ by $\theta_1(f) = f_\Lambda(X)$. If $f \sim g$, then $f_\Lambda(X) = g_\Lambda(X)$ since X is infinite. Hence $\theta_1(f) = \theta_1(g)$. If $\theta_1(f) = \theta_1(g)$ then $f \sim g$ since X is universal. Parts (2) and (3) follow directly from the fact that the extension procedure preserves both composition and addition of functions. Part (4) follows from part (3) and Theorem 3.

Let T_1^* be the set of all equivalence classes in T_1 . Since " \sim " preserves addition in T_1 , we can define addition in T_1^* by $[f] + [g] = [f + g]$, and $[f] \leq [g]$ if there is $[h] \in T_1^*$ such that $[f] + [h] = [g]$. Then T_1^* is an ideal.

COROLLARY 3. T_1^* under addition is isomorphic to an ideal in Λ .

3. Functions in T_∞ . We want to extend Theorem 3 to functions of more than one variable. Let S be the collection of all almost recursive combinatorial polynomials, i.e. $f(x_1, \dots, x_n) \in S$ iff f is an almost recursive combinatorial function such that for some integer k ,

$$f(x_1 + k, \dots, x_n + k) = \sum c_{i_1, \dots, i_n} \binom{x_1}{i_1} \cdots \binom{x_n}{i_n}$$

with all $c_{i_1, \dots, i_n} \geq 0$ and only a finite number of them nonzero and, for all $j < k$,

$$f(j, x_2, \dots, x_n), \dots, f(x_1, \dots, x_{n-1}, j) \in S.$$

Then T_∞ is the subset of S consisting of all those functions which do not involve product terms. Theorem 3 easily generalizes to functions in T_∞ .

THEOREM 5. If $f \in T_\infty$ and $X_1, \dots, X_n \in \Lambda_c$, and $Y \leq f_\Lambda(X_1, \dots, X_n)$ then there is a function $g \in T_\infty$ such that $Y = f_\Lambda(X_1, \dots, X_n)$ and $g \leq f$.

PROOF. Since $f \in T_\infty$, f is of the form $f(x_1, \dots, x_n) = f_1(x_1) + \dots + f_n(x_n)$ where $f_1, \dots, f_n \in T_1$. $Y \leq f_\Lambda(X_1, \dots, X_n)$ implies $Y = Y_1 + \dots + Y_n$ where each $Y_i \leq f_{i\Lambda}(X_i)$. By Theorem 3, $Y_i = g_i(X_i)$ for some $g_i \in T_1$ and $g_i \leq f_i$. Hence $g = g_1 + \dots + g_n$.

We can define an equivalence relation on T_∞ (or S) just as we did on T_1 . That is, for $f, g \in S$, $f \sim g$ iff $f(x_1 + k, \dots, x_n + k) = g(x_1 + k, \dots, x_n + k)$ for some integer k . Let T_∞^* be the set of equivalence classes in T_∞ and define $+$ and \leq as on T_1^* . In order to extend Theorem 4, we need an analogue of Lemma 6, to give us a "universal sequence" of isols.

Let $X = (P(\mu))^\omega$ where μ is an arbitrary infinite subset of E . Denote a vector $(\alpha_1, \alpha_2, \dots) \in X$ by $\underline{\alpha}$. Let $Q =$ collection of all vectors $\underline{\alpha} = (\alpha_1, \alpha_2, \dots) \in X$ such that (1) all the sets α_i are finite and (2) except for a finite number of coordinates, the sets $\alpha_i = \emptyset$. For $\delta \subseteq E$, let $\|\delta\|$ denote the cardinality of δ . If $\underline{\alpha}, \underline{\beta} \in Q$, define $N(\underline{\alpha}, \underline{\beta}) = \{\underline{\xi}: \underline{\alpha} \subseteq \underline{\xi}, \underline{\xi} \cap \underline{\beta} = \emptyset\}$. The collection $N = \{N(\underline{\alpha}, \underline{\beta}): \underline{\alpha}, \underline{\beta} \in Q\}$ forms a basis, and X with the topology induced by this basis is easily shown to be a complete metric space. Then by the Baire Category Theorem, if a set $A \subseteq X$ is of the first category (i.e. the countable union of nowhere dense sets) then $X - A$ is nonempty. We will use a straightforward generalization of Ellentuck's techniques [2] to prove the following lemma.

LEMMA 7. *Any infinite set of integers μ has a sequence of subsets $\delta_1, \delta_2, \dots$ such that for any $f, g \in S$, if $f_\Lambda(\langle \delta_1 \rangle, \dots, \langle \delta_n \rangle) = g_\Lambda(\langle \delta_1 \rangle, \dots, \langle \delta_n \rangle)$ then $f \sim g$.*

PROOF. Let (f_i, g_i) be a list of all pairs of recursive combinatorial functions (of any number of variables) such that $f_i \not\sim g_i$. For each i , let φ_i, ψ_i be recursive operators inducing f_i, g_i respectively. Let p_i be a list of all one-one partial recursive functions from E to E . Define $H(\varphi, \psi, p) = \{\underline{\xi} \in X: \varphi(\underline{\xi}) \subseteq \text{domain of } p, \text{ and } p[\varphi(\underline{\xi})] = \psi(\underline{\xi})\}$. By a straightforward generalization of Lemma 2 of [2] to vectors we can show that each set $H(\varphi_i, \psi_i, p_i)$ is nowhere dense.

Let $\underline{\alpha}_k$ be a list of all vectors in Q . Define $H(\varphi, \psi, p, \underline{\alpha}_k) = \{\underline{\xi} \cup \underline{\alpha}_k: \underline{\xi} \in H(\varphi, \psi, p)\}$. Since $H(\varphi, \psi, p, \underline{\alpha}_k)$ is just a translation of $H(\varphi, \psi, p)$, it is also a nowhere dense set. Let $A = \bigcup H(\varphi_i, \psi_i, p_i, \underline{\alpha}_k)$ union over all integers i, j, k . Then A is the countable union of nowhere dense sets. Hence $X - A \neq \emptyset$. Let $\underline{\delta} \in X - A$. First notice that all coordinates of $\underline{\delta}$ are infinite. For suppose not, i.e. suppose δ_i is finite for some i . Let $f(x_i) = x_i$ and $g(x_i) = \|\delta_i\|$. Then $f \not\sim g$, but $\varphi(\underline{\delta})$ is recursively isomorphic to $\psi(\underline{\delta})$ where φ, ψ induce f, g respectively.

Now we show that $\underline{\delta}$ satisfies our lemma. Suppose $f, g \in S, f \not\sim g$, but $f_\Lambda(\langle \delta_1 \rangle, \dots, \langle \delta_n \rangle) = g_\Lambda(\langle \delta_1 \rangle, \dots, \langle \delta_n \rangle)$. Since $f, g \in S$, there is an integer k such that $f(x_1 + k, \dots, x_n + k) = f_i(x_1, \dots, x_n)$ and $g(x_1 + k, \dots, x_n + k) = g_i(x_1, \dots, x_n)$ for some pair of recursive combinatorial functions (f_i, g_i) . Since $f \not\sim g, f_i \not\sim g_i$. For $i = 1, \dots, n$, let α_i be a subset of δ_i with $\|\alpha_i\| = k$. Then

$$f_{i\Lambda}(\langle \delta_1 - \alpha_1 \rangle, \dots, \langle \delta_n - \alpha_n \rangle) = g_{i\Lambda}(\langle \delta_1 - \alpha_1 \rangle, \dots, \langle \delta_n - \alpha_n \rangle).$$

Hence there is a p_j such that $(\delta_1 - \alpha_1, \dots, \delta_n - \alpha_n, \delta_{n+1}, \dots) \in H(\varphi_j, \psi_j, p_j)$. But then $\underline{\delta} \in A$, which contradicts the fact that $\underline{\delta} \in X - A$.

THEOREM 6. *There is a map $\theta: T_\infty \rightarrow \Lambda$ such that*

- (1) $f \sim g$ iff $\theta(f) = \theta(g)$,
- (2) $\theta(f(g_1, \dots, g_n)) = f_\Lambda(\theta(g_1), \theta(g_2), \dots, \theta(g_n))$,
- (3) $\theta(f + g) = \theta(f) + \theta(g)$
- (4) $\theta(T_\infty)$ is an ideal in Λ .

PROOF. Let μ be ω -cohesive. Let $\delta_1, \delta_2, \dots$ be subsets of μ , such that $X_i = \langle \delta_i \rangle, i = 1, 2, \dots$ form a “universal sequence” of isols. For $f(x_1, \dots, x_n) \in T_\infty$ define $\theta(f) = f_\Lambda(X_1, \dots, X_n)$. Then properties (1)–(4) follow as in the proof of Theorem 4.

COROLLARY 4. T_∞^* under addition is isomorphic to an ideal in Λ .

4. Functions with product terms. In Theorem 5, we defined a map, θ , such that $\theta(T_\infty)$ is closed under predecessors. In this section we want to investigate the predecessors of isols in $\theta(S)$. That is, let μ be a fixed, ω -cohesive set. By Lemma 7, μ has a sequence of subsets, $\delta_1, \delta_2, \delta_3, \dots$ such that $(X_1, X_2, X_3, \dots) = (\langle \delta_1 \rangle, \langle \delta_2 \rangle, \langle \delta_3 \rangle, \dots)$ is a “universal” sequence of isols. Use this sequence to define a map $\theta: S \rightarrow \Lambda$ by $\theta(f(x_1, \dots, x_n)) = f_\Lambda(X_1, \dots, X_n)$. The following lemma shows that $\theta(S)$ is not closed under predecessors.

LEMMA 8. *If $f \in S - T_\infty$ then $\theta(f)$ has a predecessor $U \notin \theta(S)$.*

PROOF. Since $f(x_1, \dots, x_n) \in S$, there is an integer k such that $f(x_1 + k, \dots, x_n + k)$ is a recursive combinatorial function. Since $f(x_1, \dots, x_n) \notin T_\infty$, $f(x_1 + k, \dots, x_n + k)$ contains a term of the form

$$\binom{x_1}{i_1} \dots \binom{x_n}{i_n}$$

where for some $r \neq s, i_r \geq 1$ and $i_s \geq 1$. Then the function

$$x_r x_s \leq \binom{x_1}{i_1} \dots \binom{x_n}{i_n} \leq f(x_1 + k, \dots, x_n + k)$$

and $(x_r - k)(x_s - k) \leq f(x_1, \dots, x_n)$. So we can restrict ourselves to predecessors of $(X_r - k)(X_s - k)$. Choose $\alpha_r \subseteq \delta_r$ and $\alpha_s \subseteq \delta_s$ such that $\|\alpha_r\| = \|\alpha_s\| = k$. Let

$$\beta_1 = J[(\delta_r - \alpha_r) \times (\delta_s - \alpha_s)] \cap \{(x, y) \in E^2 | x < y\}$$

and

$$\beta_2 = J[(\delta_r - \alpha_r) \times (\delta_s - \alpha_s)] \cap \{(x, y) \in E^2 | x \geq y\}.$$

Let $U = \langle \beta_1 \rangle$, $V = \langle \beta_2 \rangle$. Then $U + V = (X_r - k)(X_s - k)$. Suppose $U, V \in \theta(S)$. Then there exist $g, h \in S$ such that $\theta(g) = U$, $\theta(h) = V$. Since $\theta((x_r - k)(x_s - k)) = (X_r - k)(X_s - k) = U + V = \theta(g) + \theta(h) = \theta(g + h)$, $(x_r - k)(x_s - k) \sim g + h$. Thus there is a k' such that

$$(x_r + k')(x_s + k') = g(x_r + k', x_s + k') + h(x_r + k', x_s + k').$$

It is easy to check that if $g(x_r + k', x_s + k') + h(x_r + k', x_s + k') = (x_r + k')(x_s + k')$ then either g or h is of the form $ax_r + bx_s \pm c$, $a, b, c \in E$. Suppose g is. Then $U = aX_r + bX_s \pm c$. From the definition of the set β_1 , it is easily seen that $mX_s \leq U$ for all $m \in E$. Thus, in particular, $(a + b)X_s \leq U = aX_r + bX_s \pm c$. Therefore, $X_s \leq X_r + c$. Since X_s is infinite and X_r is ω -cohesive, $X_s = X_r \pm d$ for some $d \in E$. However, $x_s \not\sim x_r \pm d$. Thus $U \notin \theta(S)$.

Essentially what we did to construct the predecessor in Lemma 8 was to divide the "rectangle" $X_r \cdot X_s$ into two "triangles", U and V . This is really the only way we can get a predecessor of $X_r \cdot X_s$ which is not an isol in $\theta(S)$. Consider an arbitrary $f \in S$. Then there is a k such that $f_\Delta(X_1, \dots, X_m) \leq (X_1 \cdots X_m)^k$. So we can restrict ourselves to predecessors of the form $X_{i_1} \cdots X_{i_n}$ where X_{i_1}, \dots, X_{i_n} are from the fixed "universal" sequence of isols X_1, X_2, \dots and need not be distinct. We will show that essentially the only predecessors of $X_{i_1} \cdots X_{i_n}$ which are not isols in $\theta(S)$ are obtained by taking the " n -dimensional rectangle" $X_{i_1} \cdots X_{i_n}$ and dividing it into $n!$ " n -simplexes".

Let $p_1, p_2, \dots, p_n!$ be the permutations on $\{1, \dots, n\}$, with p_1 the identity permutation. Let $E_{n,k} = \{(x_1, \dots, x_n) \mid x_{p_k(1)} > \dots > x_{p_k(n)}\}$ for $k = 1, \dots, n!$ and $E_{n,0} = E^n - \bigcup_{k=1}^{n!} E_{n,k}$. Let $\beta_k = (\delta_{i_1} \times \dots \times \delta_{i_n}) \cap E_{n,k}$ and $Y_k = \langle \beta_k \rangle$. Then

$$X_{i_1} \cdots X_{i_k} = Y_0 + Y_1 + \dots + Y_{n!}.$$

Suppose $Z \leq X_{i_1} \cdots X_{i_n}$. Then $Z = Y'_0 + Y'_1 + \dots + Y'_{n!}$, where each $Y'_k \leq Y_k$. In the following lemma, we show that the only predecessors of Y_k are isols U such that either U or $Y_k - U$ is of "lower degree" than n .

LEMMA 9. *Suppose $U + V = Y_k$. Then either U or V is of the form $c_1 Z_1 + \dots + c_n Z_n$, where $c_1, \dots, c_n \in E$ and, for each $j = 1, \dots, n$, $Z_j \leq X_{i_1} \cdots X_{i_{j-1}} \cdot X_{i_{j+1}} \cdots X_{i_n}$.*

PROOF. First consider $Y_1 = \langle \beta_1 \rangle$. Notice $\beta_1 \subseteq \mu^{(n)}$. If $U + V = Y_1$, then there exist disjoint sets $\omega_1, \omega_2 \subseteq E^n$ such that $\langle J(\omega_1 \cap \beta_1) \rangle = U$, $\langle J(\omega_2 \cap \beta_1) \rangle = V$ and $J(\omega_1), J(\omega_2)$ are r.e. Since μ is ω -cohesive, it easily

follows that there is a finite subset ν of μ such that either $(\mu - \nu)^{(n)} \subseteq \omega_1$ or $(\mu - \nu)^{(n)} \subseteq \omega_2$. Assume the first case holds. Then $\mu^{(n)} - (\mu - \nu)^{(n)} \supseteq \mu^{(n)} - \omega_1 \supseteq \beta_1 \cap \omega_2$. Therefore

$$\begin{aligned} \beta_1 \cap \omega_2 &\subseteq \beta_1 \cap [\mu^{(n)} - (\mu - \nu)^{(n)}] \\ &= \beta_1 - (\delta_{i_1} - \nu) \times \cdots \times (\delta_{i_n} - \nu) \\ &\subseteq \delta_{i_1} \times \cdots \times \delta_{i_n} - (\delta_{i_1} - \nu) \times \cdots \times (\delta_{i_n} - \nu), \end{aligned}$$

and clearly $\beta_1 \cap \omega_2$ is recursively separated from its complement in $\delta_{i_1} \times \cdots \times \delta_{i_n} - (\delta_{i_1} - \nu) \times \cdots \times (\delta_{i_n} - \nu)$. Hence

$$V = \langle \beta_1 \cap \omega_2 \rangle \leq X_{i_1} \cdots X_{i_n} - (X_{i_1} - k_1) \cdots (X_{i_n} - k_n).$$

where $k_j = \|\delta_{i_k} \cap \nu\|$. Then clearly $X_{i_1} \cdots X_{i_n} - (X_{i_1} - k_1) \cdots (X_{i_n} - k_n)$ and hence V is of the desired form $c_1 Z_1 + \cdots + c_n Z_n$.

Now consider Y_k , for $k > 1$. Define $q: E^n \rightarrow E^n$ by $q((x_1, \dots, x_n)) = (x_{p_k(1)}, \dots, x_{p_k(n)})$. Then $q(\beta_k) \subseteq \mu^{(n)}$ and predecessors of Y_k can be mapped into predecessors of Y_1 and the above argument for Y used.

REFERENCES

1. J. C. E. Dekker and J. Myhill, *Recursive equivalence types*, Univ. California Publ. Math. **3** (1960), 67–213. MR 22 #7938.
2. E. Ellentuck, *Universal isols*, Math. Z. **98** (1967), 1–8. MR 35 #5315.
3. ———, *An algebraic difference between isols and cosimple isols*, J. Symbolic Logic **37** (1972), 557–561.
4. A. B. Manaster, *Higher-order indecomposable isols*, Trans. Amer. Math. Soc. **125** (1966), 363–383. MR 37 #67.
5. J. Myhill, *Recursive equivalence types and combinatorial functions*, Bull. Amer. Math. Soc. **64** (1958), 373–376. MR 21 #7.
6. A. Nerode, *Extensions to isols*, Ann. of Math. (2) **73** (1961), 362–403. MR 24 #A1215.
7. F. P. Ramsey, *On a problem of formal logic*, Proc. London Math. Soc. **30** (1930), 264–286.

DEPARTMENT OF MATHEMATICS, PENNSYLVANIA STATE UNIVERSITY, UNIVERSITY PARK, PENNSYLVANIA 16802