

## COMPARISON THEOREMS FOR BOUNDED SOLUTIONS OF $\Delta u = Pu$

BY

MOSES GLASNER

**ABSTRACT.** Let  $P$  and  $Q$  be  $C^1$  densities on a hyperbolic Riemann surface  $R$ . A characterization of isomorphisms between the spaces of bounded solutions of  $\Delta u = Pu$  and  $\Delta u = Qu$  on  $R$  in terms of the Wiener harmonic boundary is given.

In 1959, H. Royden [6] proved the following comparison theorem: *If  $P$  and  $Q$  are nonnegative  $C^1$  densities on a Riemann surface  $R$  such that there is a constant  $c$  with  $c^{-1}P \leq Q \leq cP$  outside some compact subset of  $R$ , then the spaces of bounded solutions of  $\Delta u = Pu$  and  $\Delta u = Qu$  on  $R$  are isomorphic.* In response to a question posed by Royden, M. Nakai [4] in 1960 showed that the same conclusion follows under the assumption that  $\int_R |P - Q| < +\infty$ . The conditions  $\int_R |P - Q| < +\infty$  and  $c^{-1}P \leq Q \leq cP$  outside a compact subset are independent and neither is a necessary condition for the conclusion. Recently A. Lahtinen [2] gave a necessary and sufficient condition for the bounded solutions of  $\Delta u = Pu$  on  $R$  to be isomorphic to the harmonic bounded functions. This result actually is implicit in the paper of Loeb and Walsh [3] in the axiomatic setting.

In this paper a necessary and sufficient condition is given for the existence of an isomorphism  $S$  between the spaces of bounded solutions of  $\Delta u = Pu$  and  $\Delta u = Qu$  on  $R$  with  $|u - Su|$  bounded by a potential. This result contains the ones mentioned above as special cases.

1. Let  $R$  be a Riemann surface and  $P$  a nonnegative  $C^1$  density on  $R$ . To avoid trivial considerations we assume  $R$  is hyperbolic. We denote by  $P(U)$  the space of solutions of  $\Delta u = Pu$  on an open subset  $U$  of  $R$ . The subspace of bounded solutions of  $P(U)$  will be denoted by  $PB(U)$ . The superscript  $c$  in the notation  $P^c(U)$  and  $P^cB(U)$  denotes the subspaces with continuous extensions to  $\partial U$ . It is conventional to use the symbol  $H$  in case  $P \equiv 0$ .

---

Received by the editors January 18, 1974.

AMS (MOS) subject classifications (1970). Primary 31A35; Secondary 30A50, 35J05.

Key words and phrases. Solution of  $\Delta u = Pu$ , Riemann surfaces, Green's function, maximum principle, Wiener harmonic boundary.

The Wiener ideal boundary will play a fundamental role and therefore we briefly mention some of its properties here (for more details, cf. [7]). The Wiener algebra  $N$  associated with  $R$  is the set of bounded continuous harmonizable functions on  $R$ . The bounded continuous superharmonic functions on  $R$ , for example, are contained in  $N$ . It can be seen that  $N$  is also a vector lattice. The potential subalgebra  $N_\delta$  consists of the functions in  $N$  with 0 harmonizations.

The Wiener compactification  $R^*$  of  $R$  is a compact Hausdorff space which contains  $R$  as an open dense subset and such that the functions in  $N$  extend continuously to  $R^*$  and separate points there. The space  $R^*$  is unique up to a homeomorphism fixing  $R$  pointwise. In general we shall use the same symbol for a function in  $N$  and its continuous extension to  $R^*$ . We use  $\bar{A}$  to denote the closure of  $A$  in  $R^*$  and  $\partial A$  for the boundary of  $A$  with respect to  $R$ . If we use  $A^*$  to denote a subset of  $R^*$ , then by  $A$  we shall mean  $A^* \cap R$ .

The Wiener harmonic boundary  $\delta$  is the set of common zeros of functions in  $N_\delta$  and hence is a compact subset of  $R^* \setminus R$ . The main properties of  $\delta$  that we shall need are the following: If  $U$  is an open subset of  $R$  with  $\partial U$  piecewise analytic and  $f \in N$ , then there is a function  $h \in N$  such that  $h \in H^c(U)$  and  $h - f|_{(R \setminus U) \cup (\bar{U} \cap \delta)} = 0$ . The function  $h$  is called the harmonic projection of  $f$  with respect to  $U$ . If  $s$  is bounded and subharmonic on  $U$  with continuous extension to  $\partial U$  then  $s \leq \max_{\partial U \cup (\bar{U} \cap \delta)} s$ . Therefore a superharmonic function on  $U$  with continuous extension to  $\partial U$  is a potential if and only if it vanishes on  $\partial U \cup (\bar{U} \cap \delta)$ .

We shall use the notation  $\|\varphi\|_A$  for the supremum of the function  $|\varphi|$  on  $A$ .

2. We consider solutions of  $\Delta u = Pu$  on subregions  $G$  with  $\partial G$  piecewise analytic ( $\partial G$  may be  $\emptyset$ ). The fact that nonnegative solutions are subharmonic gives the following (cf. [3]).

LEMMA. *If  $u \in P^c B(G)$ , then  $\|u\|_G \leq \|u\|_{\partial G \cup (\bar{G} \cap \delta)}$ . Moreover, if  $u|_{\partial G \cup (\bar{G} \cap \delta)} \geq 0$ , then  $u \geq 0$ .*

We now describe the integral operator  $T_G$  which is the basic tool here. Let  $\Omega$  be a relatively compact region in  $R$  with  $\partial\Omega$  piecewise analytic. Define  $\tau_{G \cap \Omega}$  on the bounded  $C^1$  functions on  $\Omega$  by setting  $\tau_{G \cap \Omega} \varphi = \int_\Omega g_{G \cap \Omega}(\cdot, z) \varphi P$ , where  $g_{G \cap \Omega}(\cdot, z)$  denotes the Green's function of  $G \cap \Omega$  with pole at  $z$ . Then  $\tau_{G \cap \Omega} \varphi$  is a  $C^2$  function on  $G \cap \Omega$  vanishing continuously on  $\partial(G \cap \Omega)$  and satisfies  $\Delta \tau_{G \cap \Omega} \varphi = -\varphi P$ .

Therefore setting  $T_{G \cap \Omega} u = u + \tau_{G \cap \Omega} u$  gives an operator  $T_{G \cap \Omega}$ :

$P^c(G \cap \Omega) \rightarrow H^c(G \cap \Omega)$  such that  $u - T_{G \cap \Omega} u | \partial(G \cap \Omega) = 0$ . For  $u \in P^cB(G)$  with  $u \geq 0$ ,  $T_G u = \lim_{\Omega \rightarrow R} T_{G \cap \Omega} u$  exists and is given by  $T_G u = u + \tau_G u$ , where  $\tau_G u = \int_G g_G(\cdot, z) u P$ . If  $u \in P^cB(G)$  is arbitrary, then  $T_G u$  is defined to be  $T_G u_1 - T_G u_2$  where  $u = u_1 - u_2$ ,  $u_i \in P^cB(G)$ ,  $u_i \geq 0$ . We collect here some of the known properties of  $T_G$  that will be needed in later arguments (cf. [4], [5]).

**THEOREM.** *The operator  $T_G: P^cB(G) \rightarrow H^cB(G)$  gives an isometric isomorphism such that  $u - T_G u | \partial G \cup (\bar{G} \cap \delta) = 0$ . If  $\int_G g_G(\cdot, z) P < +\infty$ , for every  $z \in G$ , then  $T_G$  is onto.*

3. We introduce the set  $\delta^P = \{p \in \delta \mid p \text{ has a nbd } U^* \text{ in } R^* \text{ with } \int_U g_U(\cdot, z) P < +\infty \text{ for each } z \in U\}$ . Here  $g_U(\cdot, z)$  for an arbitrary open set  $U$  is defined as follows: Let the component of  $U$  containing  $z$  be denoted by  $U_z$ . Then  $g_U(\cdot, z)$  is the Green's function of  $U_z$  on  $U_z$  and zero on  $U \setminus U_z$ .

Clearly  $\delta^P$  is an open subset of  $\delta$ . The significance of  $\delta^P$  stems from the following

**THEOREM.** *The functions in  $PB(R)$  restricted to  $\delta$  vanish on  $\delta \setminus \delta^P$ .*

It is sufficient to prove the assertion for  $u \in PB(R)$  with  $u \geq 0$ . By Theorem 2,  $\tau_R u(z) < +\infty$  for each  $z \in R$ . Suppose  $p \in \delta$  and  $u(p) \neq 0$ . By the continuity of  $u$  on  $R^*$  there is a neighborhood  $U^*$  of  $p$  and an  $\epsilon > 0$  such that  $u | U^* \geq \epsilon$ . Then

$$+\infty > \tau_R u(z) > \int_U g_R(\cdot, z) u P \geq \epsilon \int_U g_R(\cdot, z) P \geq \epsilon \int_U g_U(\cdot, z) P,$$

for each  $z \in U$ . Thus  $p \in \delta^P$ .

Combining this with Lemma 2 gives the

**COROLLARY.** *If  $u \in PB(R)$ , then  $\|u\|_R \leq \|u\|_{\delta^P}$ . If in addition  $u | \delta^P \geq 0$ , then  $u \geq 0$ .*

4. Denote by  $C_C(A)$  the continuous functions with compact support in  $A$  and by  $C_0(A)$  the closure of  $C_C(A)$  with respect to  $\|\cdot\|_A$ .

**THEOREM.** *The spaces  $PB(R)$  and  $C_0(\delta^P)$  are isomorphic vector lattices. The isomorphism is obtained by restriction to  $\delta^P$ , i.e.  $PB(R) | \delta^P = C_0(\delta^P)$ .*

We begin the proof by observing that for any  $f \in N$  with  $K = \text{supp}(f | \delta)$  a compact subset of  $\delta^P$ , there exists a  $u \in PB(R)$  with  $u - f | \delta = 0$ . In fact we may assume  $f \geq 0$ . Cover  $K$  by a finite number of sets  $U_i^*$ ,  $i = 1, \dots, m$  such that  $\int_{U_i} g_{U_i}(\cdot, z) P < +\infty$  for each  $z \in U_i$ . By taking  $U_i^*$  slightly smaller if necessary we may assume that  $\partial U_i$  is piecewise analytic.

By the Urysohn property for  $N$  we can find  $\varphi_i \in N$  such that  $\text{supp } \varphi_i \subset U_i^*$ ,  $\varphi_i \geq 0$  and  $\sum_1^m \varphi_i|K = 1$ . Let  $f_i = \varphi_i f \in N$  and note that  $f = \sum_1^m f_i$  on  $K$ . Denote by  $h_i$  the harmonic projection of  $f_i$  with respect to  $U_i$ , i.e.  $h_i \in H^c B(U_i)$  and  $0 = f_i|_{\partial U_i} = h_i|_{\partial U_i}$ ,  $f_i|_{U_i^* \cap \delta} = h_i|_{U_i^* \cap \delta}$ . Let  $G$  be any component of  $U_i$ . Then  $\int_G g_G(\cdot, z) P < +\infty$  for every  $z \in G$ . Thus by Theorem 2 there is a function  $u \in P^c B(G)$  such that  $u|_{\partial G} = 0$ ,  $u|_{\bar{G} \cap \delta} = h_i|_{\bar{G} \cap \delta}$ . Repeating this in each component of  $U_i$  we obtain  $v_i \in P^c B(U_i)$  with  $v_i \geq 0$ ,  $v_i|_{\partial U_i} = 0$  and  $v_i - f|_{U_i^* \cap \delta} = 0$ . Setting  $v_i = 0$  on  $R \setminus U_i$  gives a subsolution on  $R$ .

Let  $k_i$  be the least harmonic majorant of  $v_i$ . Take an exhaustion  $\{\Omega_n\}$  of  $R$  by regular regions. Denote by  $u_{in}$  the function in  $P^c(\Omega_n)$  such that  $u_{in} - v_i|_{\partial \Omega_n} = 0$ . Then we have  $0 \leq v_i \leq u_{in} \leq u_{i,n+1} \leq k_i$ . Thus  $u_i = \lim_n u_{in}$  is in  $PB(R)$  and  $0 \leq v_i \leq u_i \leq k_i$ . Since  $k_i - v_i$  is a potential on  $R$  it vanishes on  $\delta$  and consequently  $k_i - u_i$  also vanishes there. That is,  $u_i - v_i|_{\delta} = 0$ . The function  $u = \sum_1^m u_i \in PB(R)$  has the property that  $u|_{\delta} = f$ .

For an arbitrary  $f \in C_0(\delta^P)$  there is a sequence  $\{f_k\}$  of functions of the sort considered above such that  $\|f - f_k\|_{\delta^P} \rightarrow 0$ . Then the corresponding sequence of solutions  $\{u_k\}$  is a Cauchy sequence with respect to  $\|\cdot\|_R$  by Corollary 3. Thus there is a  $u \in PB(R)$  such that  $\|u - u_k\|_R \rightarrow 0$ . Let  $u|_{\delta^P} = g$  and for a given  $\epsilon > 0$  take  $k$  such that  $\|u - u_k\|_R < \epsilon$  and  $\|f - f_k\|_{\delta^P} < \epsilon$ . Then the denseness of  $R$  in  $R^*$  gives  $\|g - f_k\|_{\delta^P} < \epsilon$ . This means  $g = f$  on  $\delta^P$ .

Thus far we have shown  $C_0(\delta^P) \subset PB(R)|_{\delta^P}$ . For the proof of the reverse inclusion take  $u \in PB(R)$  which without loss of generality can be assumed to be nonnegative. By Theorem 3,  $u|_{\delta \setminus \delta^P} = 0$ . Set  $K_n = \{p \in \delta \mid u(p) \geq 1/n\}$ .  $K_n$  is a closed and hence compact subset of  $\delta$ . Since  $K_n \subset \delta^P$  it is compact in  $\delta^P$ . Now choose  $\varphi_n \in N$  such that  $\varphi_n|_{K_n} = 1$ ,  $\varphi_n|_{\delta \setminus K_{n+1}} = 0$  and  $0 \leq \varphi_n \leq 1$ . Then  $\varphi_n u \in C_c(\delta^P)$  and  $\|u - \varphi_n u\|_{\delta^P} \leq 1/n$ .

Corollary 3 implies that the mapping of  $PB(R)$  onto  $C_0(\delta^P)$  obtained by restriction to  $\delta^P$  preserves order and sup norm.

A slightly more tractable description of  $\delta^P$  can be derived from this theorem.

**COROLLARY.**  $\delta^P = \{p \in \delta \mid p \text{ has a nbd } U^* \text{ in } R^* \text{ such that } \int_U g_R(\cdot, z) P < +\infty \text{ for each } z \in U\}$ .

Since  $g_R(\cdot, z) \geq g_U(\cdot, z)$  for  $z \in U$  we need only show that for each  $p \in \delta^P$  there is a neighborhood  $U^*$  of  $p$  such that  $\int_U g_R(\cdot, z) P < +\infty$ . But by the theorem there is a function  $u \in PB(R)$  such that  $u(p) = 2$ . Set  $U^* = \{q \in R^* \mid u(q) > 1\}$ . By Theorem 2,  $\tau_R u$  exists and, in particular,  $+\infty > \tau_R u(z) \geq \int_U g_R(\cdot, z) P$ , for  $z \in U$ .

5. The main result is as follows:

**THEOREM.** *Suppose  $P$  and  $Q$  are nonnegative  $C^1$  densities on a hyperbolic Riemann surface  $R$ . There is an isomorphism  $S$  between  $PB(R)$  and  $QB(R)$  such that  $|u - Su|$  is bounded by a potential on  $R$  if and only if  $\delta^P = \delta^Q$ .*

If  $\delta^P = \delta^Q$ , then both  $PB(R)$  and  $QB(R)$  are isomorphic to  $C_0(\delta^P)$  by restriction to  $\delta^P$ . Thus define  $S: PB(R) \rightarrow QB(R)$  by  $u - Su|_{\delta^P} = 0$ . In order to show that  $|u - Su|$  is bounded by a potential, express  $u$  as  $u = u_1 - u_2$  with  $u_i \in PB(R)$  and  $u_i \geq 0$ . Note that  $u|_{\delta^P} = u_1|_{\delta^P} - u_2|_{\delta^P}$  which implies that  $Su = Su_1 - Su_2$ ,  $Su_i \geq 0$ . Take  $h_i \in HB(R)$  with  $h_i|_{\delta} = u_i|_{\delta}$ ,  $i = 1, 2$ . Since  $u_i - h_i$  and  $Su_i - h_i$  are bounded subharmonic functions on  $R$  which vanish on  $\delta$  we have  $u_i - h_i \leq 0, Su_i - h_i \leq 0$ . Thus  $|u - Su|$  is bounded by the potential  $(h_1 - u_1) + (h_1 - Su_1) + (h_2 - u_2) + (h_2 - Su_2)$ .

Conversely, suppose an isomorphism  $S$  as described in the theorem exists. Then  $|u - Su|_{\delta} = 0$  for each  $u \in PB(R)$ . If  $p \in \delta^P$ , then by Theorem 4 we can find  $u \in PB(R)$  with  $u(p) = 1$ . This implies that  $Su(p) = 1$  and in view of Theorem 3 we conclude that  $p \in \delta$ , i.e.  $\delta^P \subset \delta^Q$ . By symmetry we obtain  $\delta^P = \delta^Q$ .

The assumption on  $|u - Su|$  implies that  $S$  preserves the behavior of functions on  $\delta^P$ . In view of Theorem 4 this means that  $S$  commutes the lattice operations. If the assumption on  $|u - Su|$  is replaced by the assumption that  $S$  is a vector lattice isomorphism, then by the Kakutani theorem we see that  $\delta^P$  and  $\delta^Q$  are homeomorphic.

The results of Royden [6] and Nakai [4] are immediate consequences.

**COROLLARY.** *If  $P$  and  $Q$  are  $C^1$  densities on a hyperbolic Riemann surface  $R$  such that  $c^{-1}P \leq Q \leq cP$  outside some compact subset and for some constant  $c$ , then  $PB(R)$  and  $QB(R)$  are isomorphic.*

**COROLLARY.** *If  $P$  and  $Q$  are  $C^1$  densities on a hyperbolic Riemann surface  $R$  and  $\int_R |P - Q| < +\infty$ , then  $PB(R)$  and  $QB(R)$  are isomorphic.*

In the first case it is clear that the hypothesis implies that  $\delta^P = \delta^Q$ . In the second case note that  $g_R(\cdot, z)|_{\delta} = 0$  and hence there is a neighborhood  $V^*$  of  $\delta$  in  $R^*$  with  $g_R(\cdot, z)|_{V^*} \leq 1$ . Thus  $\int_V g_R(\cdot, z)|P - Q| < +\infty$ . By the Harnack inequality this is also valid if  $z$  is allowed to vary and the conclusion  $\delta^P = \delta^Q$  now follows.

Actually the corollaries followed from the slightly weaker hypotheses  $c^{-1}P \leq Q \leq cP$  in  $V$ ,  $\int_V |P - Q| < +\infty$ , where  $V^*$  is a neighborhood of  $\delta$  in  $R^*$ .

Denote by  $w$  the greatest solution of  $\Delta u = Pu$  on  $R$  which is less than 1 on  $R$ . The following result is due to Lahtinen [2] and Loeb and Walsh [3].

COROLLARY.  $HB(R)$  and  $PB(R)$  are isomorphic vector lattices if and only if 1 is the least harmonic majorant of  $w$  on  $R$ .

Let  $h$  be the least harmonic majorant of  $w$ . Then  $h - w$  is a potential and hence vanishes on  $\delta$ . Therefore,  $h$  is the constant 1 if and only if  $w|_{\delta} = 1$ . This in turn is equivalent to  $\delta^P = \delta$  which is equivalent to  $PB(R)$  being isomorphic to  $HB(R)$ .

6. Denote by  $PBE(R)$  (resp.  $PBD(R)$ ) the subspace of  $PB(R)$  such that  $E(u) = \int_R du \wedge *du + u^2 P < +\infty$  (resp.  $D(u) = \int_R du \wedge *du < +\infty$ ). Denote by  $\Delta$  the Royden harmonic boundary of  $R$ ,  $R^*$  the corresponding compactification and define

$$\Delta^P = \left\{ p \in \Delta \mid p \text{ has a nbd } U^* \text{ in } R^* \text{ with } \int_U P < +\infty \right\},$$

$$\Delta_P = \left\{ p \in \Delta \mid p \text{ has a nbd } U^* \text{ in } R^* \text{ with } \iint_{U \times U} g_R(x, y) P(x) P(y) < +\infty \right\}.$$

These definitions lead to criteria for isomorphisms between the closures with respect to the sup norm of the bounded energy finite or bounded Dirichlet finite solutions.

THEOREM. Suppose  $P$  and  $Q$  are nonnegative  $C^1$  densities on  $R$ . There is an isomorphism  $S$  between  $\overline{PBE(R)}$  and  $\overline{QBE(R)}$  (resp.  $\overline{PBD(R)}$  and  $\overline{QBD(R)}$ ) such that  $|u - Su|$  is bounded by a potential on  $R$  if and only if  $\Delta^P = \Delta^Q$  (resp.  $\Delta_P = \Delta_Q$ ).

The proof is analogous to that of Theorem 5 and therefore we only mention some differences. The operator  $T_G$  defined in §2 also maps the spaces  $P^c BE(G)$  and  $P^c BD(G)$  into  $H^c BD(G)$ . If  $\int_G P < +\infty$ , then

$$T_G(P^c BE(G)) = H^c BD(G)$$

(cf. [1]) and if  $\iint_{G \times G} g_G(x, y) P(x) P(y) < +\infty$ , then

$$T_G(P^c BD(G)) = H^c BD(G)$$

(cf. [5]). This is the motivation for the choice of  $\Delta^P$  and  $\Delta_P$ . The proper analogue of Theorem 4 is that the closure of  $PBE(R)$  (resp.  $PBD(R)$ ) with respect to the sup norm restricted to  $\Delta^P$  (resp.  $\Delta_P$ ) is the space  $C_0(\Delta^P)$  (resp.  $C_0(\Delta_P)$ ) but this causes no complications. In the proof some complications do occur because of the need to establish the convergence of sequences of functions in the  $D$  or  $E$  norm.

ADDED IN PROOF. M. Nakai (*Banach spaces of bounded solutions of  $\Delta u = Pu$  on hyperbolic Riemann surfaces*, Nagoya Math. J. 53 (1974), 141–155) has

simultaneously discovered Theorem 5 and also has given a more detailed analysis of its consequences.

## REFERENCES

1. M. Glasner and R. Katz, *On the behavior of solutions of  $\Delta u = Pu$  at the Royden boundary*, J. Analyse Math. 22 (1969), 343–354. MR 41 #1995.
2. A. Lahtinen, *On the solutions of  $\Delta u = Pu$  for acceptable densities*, Ann. Acad. Sci. Fenn. Ser. A I No. 515 (1972).
3. P. A. Loeb and B. Walsh, *A maximal regular boundary for solutions of elliptic differential equations*, Ann. Inst. Fourier (Grenoble) 18 (1968), fasc. 1, 283–308. MR 39 #4423.
4. M. Nakai, *The space of bounded solutions of  $\Delta u = Pu$  on a Riemann surface*, Proc. Japan Acad. 36 (1960), 267–272. MR 22 #12216.
5. ———, *Dirichlet finite solutions of  $\Delta u = Pu$  and classification of Riemann surfaces*, Bull. Amer. Math. Soc. 77 (1971), 381–385. MR 45 #2162.
6. H. L. Royden, *The equation  $\Delta u = Pu$  and the classification of open Riemann surfaces*, Ann. Acad. Sci. Fenn. Ser. A I No. 271 (1959). MR 22 #12215.
7. L. Sario and M. Nakai, *Classification theory of Riemann surfaces*, Die Grundlehren der math. Wissenschaften, Band 164, Springer-Verlag, Berlin and New York, 1970. MR 41 #8660.

DEPARTMENT OF MATHEMATICS, PENNSYLVANIA STATE UNIVERSITY, UNIVERSITY PARK, PENNSYLVANIA 16802