

ON THE DIMENSION OF VARIETIES OF SPECIAL DIVISORS

BY

R. F. LAX

ABSTRACT. Let T_g denote the Teichmüller space and let V denote the universal family of Teichmüller surfaces of genus g . Let $V_{T_g}^{(n)}$ denote the n th symmetric product of V over T_g and let J denote the family of Jacobians over T_g . Let $f: V_{T_g}^{(n)} \rightarrow J$ be the natural relativization over T_g of the classical map defined by integrating holomorphic differentials. Let

$$u: f^* \Omega_{J/T_g}^1 \rightarrow \Omega_{V_{T_g}^{(n)}/T_g}^1$$

be the map induced by f . We define G_n^r to be the analytic subspace of $V_{T_g}^{(n)}$ defined by the vanishing of $\wedge^{n-r+1} u$.

Put $\tau = (r + 1)(n - r) - rg$. We show that $G_n^1 - G_n^2$, if nonempty, is smooth of pure dimension $3g - 3 + \tau + 1$. From this result, we may conclude that, for a generic curve X , the fiber of $G_n^1 - G_n^2$ over the module point of X , if nonempty, is smooth of pure dimension $\tau + 1$, a classical assertion.

Variational formulas due to Schiffer and Spencer and Rauch are employed in the study of G_n^r .

0. Introduction. Let X be a complete, nonsingular curve of genus g over an algebraically closed field K . Let $X^{(n)}$ denote the n th symmetric product of X . Let $G_n^r(X)$ denote the subvariety of $X^{(n)}$ of all divisors D of degree n such that $\dim |D| \geq r$. (In the literature, e.g. [12], $G_n^r(X)$ is often used to denote the subvariety of the Jacobian of X consisting of all linear systems of degree n and projective dimension at least r .)

Put τ equal to $(r + 1)(n - r) - rg$. Brill and Noether [2] asserted that if τ were nonnegative and X were a generic curve, then $G_n^r(X)$ would have dimension $\tau + r$. The recent work of Kleiman and Laksov ([10], [11]) and Kempf [8] shows that for X any curve, if $\tau \geq 0$, then $G_n^r(X)$ has dimension *at least* $\tau + r$. We will show, in the case $K = \mathbb{C}$, that if X is a generic curve, then $G_n^1(X) - G_n^2(X)$, if nonempty, has dimension $\tau + 1$.

Received by the editors December 19, 1973.

AMS (MOS) subject classifications (1970). Primary 14H15, 32G15; Secondary 14H40, 14C20.

Key words and phrases. Analytic space, special divisor, Teichmüller surface, moduli.

Copyright © 1975, American Mathematical Society

We work in the category of analytic spaces over \mathbb{C} . We do this because we want to consider the Teichmüller space, an analytic, but not algebraic, variety [5]. We take the Séminaire Cartan, 1960–61, as our foundational reference. In particular, we allow the structure sheaf of an analytic space to contain nilpotents.

Let Y be an analytic space over \mathbb{C} and let E and F be locally free \mathcal{O}_Y -modules of ranks g and n respectively. Suppose we are given a map $u: E \rightarrow F$. In §1, we define the analytic space $Z'(u)$ to be given by the vanishing of the map $\bigwedge^{n-r+1} u$. We then study the infinitesimal structure of $Z'(u)$.

Let S be an analytic space over \mathbb{C} and let X be a family of nonsingular curves of genus g over S . Let $X_S^{(n)}$ denote the n th symmetric product of X over S and let J_S denote the family of Jacobians over S (cf. [7], [15]). Suppose we are given a map $f: X_S^{(n)} \rightarrow J_S$. Let

$$u: f^* \Omega_{J_S/S}^1 \rightarrow \Omega_{X_S^{(n)}/S}^1$$

be the map induced by f . We study the analytic space $Z'(u) \subseteq X_S^{(n)}$ in the following situation: $S = T_g$, the Teichmüller space, X is the universal family of Teichmüller surfaces of genus g , and f is the natural relativization over T_g of the classical map from the n th symmetric product of a curve into its Jacobian defined by integrating a basis of homomorphic differentials (cf. §2). We let G_n^r denote $Z'(u)$ in this situation.

In order to understand explicitly the above map f , we must use certain variational formulas which are similar to those derived by Schiffer and Spencer [19], but much closer in form to those appearing in Rauch [18]. We also need a theorem due to Patt [17] concerning local coordinates at a point of T_g .

Our main result is:

THEOREM. *Suppose $y \in G_n^1 - G_n^2$. Then the dimension of the tangent space to G_n^1 at y is $3g - 3 + \tau + 1$.*

From this result, we can conclude that if X is a generic compact Riemann surface, then $G_n^1(X) - G_n^2(X)$, if nonempty, is smooth of pure dimension $\tau + 1$.

As an application, we show that the subvariety of T_g , for $g \geq 4$, of curves with nonempty G_3^1 (so-called "trigonal" curves), is of dimension $2g + 1$, a result which was known to Severi [22] and B. Segre [20].

In a sequel to this paper, we will show that if $\tau \geq 0$ then G_n^2 (resp. G_n^3) has a component of dimension $3g - 3 + \tau + 2$ (resp. $3g - 3 + \tau + 3$). The proof involves computations using the examples of Riemann surfaces given by Meis [16].

The author would like to thank D. Laksov, M. Artin, A. Mattuck and, most of all, his thesis advisor S. L. Kleiman for their generous help.

1. $Z^r(u)$ and its infinitesimal structure. Let S be an analytic space over C . Denote by $((An/S))$ the category of analytic spaces over S . Let Y be an analytic space over S and let E and F be locally free \mathcal{O}_Y -modules of ranks g and n respectively. Suppose we are given a map $u: E \rightarrow F$. Define the functor $Z^r(u): ((An/S))^0 \rightarrow ((Sets))$ by

$$Z^r(u)(T) = \left\{ g \in \text{Hom}(T, Y) \mid \bigwedge^{n-r+1} g^*u = 0 \right\}.$$

We wish to show that this functor is represented by an analytic subspace of Y .

DEFINITION 1 [5]. Let S be an analytic space and let

$$G: ((An/S))^0 \rightarrow ((Sets))$$

be a functor. We say that G is of a local nature if for every T the presheaf $U \mapsto G(U)$, where U runs through the open sets of T , is a sheaf.

REMARK. This is the analog to the notion of a Zariski sheaf in the category of contravariant functors from $((Schemes))$ to $((Sets))$.

LEMMA 1. Let (S_i) be a covering of an analytic space S by open sets. Let $G: ((An/S))^0 \rightarrow ((Sets))$ be a functor. Then G is representable iff G is of a local nature and for every i , the functor $G/S_i: ((An/S_i))^0 \rightarrow ((Sets))$ is representable.

PROOF. [5, Corollary 5.7 of Exposé 7].

Our functor $Z^r(u)$ is clearly of a local nature. Hence, by the lemma, its representability is a local question.

Let y be a point of Y . Since E and F are locally free of ranks g and n respectively, the map u is given locally at y by an $n \times g$ matrix $[f_{jk}]$ of functions regular at y . The functor $Z^r(u)$ is then locally represented by the closed analytic subspace defined by the vanishing of the minors of order $n - r + 1$ of the matrix $[f_{jk}]$. Thus we have

PROPOSITION 1. $Z^r(u)$ is represented by a closed analytic subspace of Y .

We will also use $Z^r(u)$ to denote this analytic subspace.

Put $\rho = \text{rank}(u \otimes \kappa(y))$. Locally at y , both E and F split off a direct summand of rank ρ , and u maps one summand isomorphically onto the other. The map that u induces on the other two summands is given by an $(n - \rho) \times (g - \rho)$ matrix $[e_{jk}]$ of functions regular at y . The analytic space $Z^r(u)$ is also defined locally at y by the vanishing of the minors of order $(n - r + 1 - \rho)$ of the matrix $[e_{jk}]$ (cf. [10]).

PROPOSITION 2. Assume $r > 0$. Then the points of $Z^{r+1}(u)$ are singular points of $Z^r(u)$.

PROOF. Suppose $y \in Z^{r+1}(u)$. Then we have $\rho < n - r$. By construction, the e_{jk} above vanish at y , hence are in the maximal ideal m of $\mathcal{O}_{Y,y}$. The analytic space $Z^r(u)$ is defined locally at y by the vanishing of the minors of order $(n - r + 1 - \rho)$ of the matrix $[e_{jk}]$ and, since $\rho < n - r$, all these minors are of order at least 2, hence are in m^2 . Thus y cannot be a smooth point of $Z^r(u)$.

We want now to study the infinitesimal structure of $Z^r(u)$. Let ξ denote a tangent vector to Y at y . We will also use ξ to denote the comorphism, which is a \mathbb{C} -homomorphism of local rings $\xi: \mathcal{O}_{Y,y} \rightarrow \mathbb{C}[\epsilon]/(\epsilon^2)$.

We are interested in seeing when ξ is a tangent vector to $Z^r(u)$ at y . By definition, this will be true if $\bigwedge^{n-r+1} \xi^*u = 0$.

PROPOSITION 3. ξ is a tangent vector to $Z^r(u)$ at y iff the minors of order $n - r + 1$ of the matrix $[\xi(f_{jk})]$ are all zero.

PROOF. It is easy to see that the map ξ^*u is given by the matrix $[\xi(f_{jk})]$. Thus we have $\bigwedge^{n-r+1} \xi^*u = 0$ iff the minors of order $n - r + 1$ of $[\xi(f_{jk})]$ all vanish.

We now assume that Y is smooth of dimension m over \mathbb{C} . Let $y \in Y$ and let $\sigma_1, \dots, \sigma_m$ be local parameters on Y at y . Let s_l in \mathbb{C} be given by

$$\xi(\sigma_l) = s_l \epsilon, \quad l = 1, 2, \dots, m.$$

Then, by Taylor's Theorem, we have

$$\xi(f_{jk}) = f_{jk}(y) + \epsilon \sum_{l=1}^m s_l \frac{\partial f_{jk}}{\partial \sigma_l}(y).$$

The vanishing of the minors of order $n - r + 1$ of the matrix $[\xi(f_{jk})]$ gives rise to linear equations in the s_l . These equations must be satisfied for ξ to be a tangent vector to $Z^r(u)$ at y . If we view s_1, \dots, s_m as being unknowns, then the dimension of the solution space of this system of equations is the dimension of the tangent space to $Z^r(u)$ at y .

If $y \in Z^r(u) - Z^{r+1}(u)$, we will want to use the following lemma.

LEMMA 2. Let A be a commutative ring (with unit). Let $M = [a_{jk}]$ be an $m \times n$ matrix over A . Suppose that a minor μ of order r is a unit, and that every minor of order $r + 1$ containing μ vanishes. Then every minor of order $r + 1$ vanishes.

PROOF. Without loss of generality, we may assume that μ is the leading (i.e., upper left) minor of order r . Since μ is a unit, we may perform column operations using the first r columns to change M to the matrix

$$M' = \begin{bmatrix} \mu & \cdots & \cdots & \cdots & \cdots & 0 \\ a_{r+1,1} & \cdots & \cdots & \cdots & a_{r+1,r} & \vdots \\ \vdots & \cdots & \cdots & \cdots & \vdots & \vdots \\ a_{m,1} & \cdots & \cdots & \cdots & a_{m,r} & N \end{bmatrix}$$

where N is an $(m - r) \times (n - r)$ matrix.

Then by row operations, using the first r rows, we may change M' to the matrix

$$M'' = \begin{bmatrix} \mu & \vdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \vdots & N \end{bmatrix}$$

where N is the same matrix as before.

Now, no minor containing μ is affected by performing these row and column operations. Hence, the minors of order $r + 1$ of M'' which contain μ are all zero. Thus N is the zero matrix.

But this implies that every column of M is a linear combination of the first r columns of M . Hence, every minor of order $r + 1$ of M is zero. \square

Suppose now that $y \in Z^r(u) - Z^{r+1}(u)$. Then the matrix $[f_{jk}]$ has rank $n - r$. We may thus assume that the leading minor of order $n - r$ of $[f_{jk}]$, call it μ , is nonzero. Let μ' denote the leading minor of order $n - r$ of $[\xi(f_{jk})]$. Then $\mu' = \mu + c\epsilon$, for some $c \in \mathbb{C}$. Since μ is nonzero, μ' does not lie in the maximal ideal of $\mathbb{C}[\epsilon]/(\epsilon^2)$, hence is a unit. We then have, by Proposition 3 and Lemma 2, that ξ is a tangent vector to $Z^r(u)$ at y iff the minors of order $n - r + 1$ of $[\xi(f_{jk})]$ which contain μ' all vanish. Obviously, there are $r(g - n + r)$ such minors. If the equations in the s_i given by the vanishing of these minors are linearly independent (over \mathbb{C}), then the dimension of the tangent space to $Z^r(u)$ at y is $m - r(g - n + r)$. We could then conclude that y is a smooth point of $Z^r(u)$ by virtue of the following proposition.

PROPOSITION 4. *Either $Z^r(u)$ is empty, or each component has codimension at most $r(g - n + r)$ in Y .*

PROOF. This is proved in [9] for Y a scheme. With the obvious modifications, the proof is valid for Y an analytic space.

2. The universal family of Teichmüller surfaces. In [5], Grothendieck proved the following

THEOREM 1. *There exist an analytic space T_g and a family V of Teichmüller surfaces of genus g over T_g which is universal in the following sense: for every family X of Teichmüller surfaces of genus g over an analytic space S , there exists a unique map $\Phi: S \rightarrow T_g$, such that X is isomorphic (as a family of Teichmüller surfaces) to the pullback via Φ of V/T_g .*

T_g is called the *Teichmüller space* (for Teichmüller surfaces of genus g). The Teichmüller space is a smooth, irreducible, and simply connected analytic space [5].

Let $h: V \rightarrow T_g$ denote the structural morphism. By well-known topological facts, since T_g is simply connected, the fiber bundle $R^1 h_* \mathbf{Z}$ is trivial. Thus, there are sections of this bundle which give rise to cycles $\gamma_i(s), \delta_i(s), i = 1, \dots, g$, which form a canonical homology basis for $H_1(V_s, \mathbf{Z}), s \in T_g$ [15].

Consider the sheaf Ω^1_{V/T_g} . For all $s \in T_g$, we have

$$\dim H^0(V_s, \Omega^1_{V/T_g} \otimes \kappa(s)) = \dim H^0(V_s, \Omega^1_{V_s}) = g.$$

Hence, $h_* \Omega^1_{V/T_g}$ is a vector bundle of rank g over T_g and we have

$$h_* \Omega^1_{V/T_g} \otimes \kappa(s) \cong H^0(V_s, \Omega^1_{V_s})$$

by [4].

Choose holomorphic sections $d\zeta_i^*$, $i = 1, \dots, g$, of $h_* \Omega^1_{V/T_g}$ such that $\{d\zeta_i^*(s)\}_{i=1}^g$ is a basis for $H^0(V_s, \Omega^1_{V_s}), s \in T_g$ (cf. [15]). Put

$$a_{ij}(s) = \int_{\gamma_i(s)} d\zeta_j^*(s), \quad b_{ij}(s) = \int_{\delta_i(s)} d\zeta_j^*(s), \quad i, j = 1, \dots, g.$$

For each $s \in T_g$, the matrix $[a_{ij}(s), b_{ij}(s)]$ is the period matrix of V_s . Recall that the columns of this matrix generate a maximal lattice subgroup of \mathbf{C}^g . Let J be the quotient of $T_g \times \mathbf{C}^g$ by this family of lattices. The induced projection $J \rightarrow T_g$ gives a complex analytic family of complex tori, the fiber J_s being the Jacobian variety of the Teichmüller surface V_s [15].

Since our concern will only be local, we assume that there exist sections of $V \rightarrow T_g$. Let $P_0^*(s)$ be such a section. As in [15], define a map $\psi: V \rightarrow J$ by

$$\psi(s, P) = \left(s, \int_{P_0^*(s)}^P d\zeta_1^*(s), \dots, \int_{P_0^*(s)}^P d\zeta_g^*(s) \right) \text{ mod periods}$$

for $P \in V_s$.

Denote by $V_T^{(n)}$ the n th symmetric product of V over T_g (cf. [7]). Extend ψ to a map $f: V_T^{(n)} \rightarrow J$ as follows. If $s \in T_g$ and $D \in (V_T^{(n)})_s$ is the divisor $\sum_{i=1}^n P_i$ on V_s , then

$$f(s, D) = \left(s, \sum_{i=1}^n \int_{P_0^*(s)}^{P_i} d\zeta_1^*(s), \dots, \sum_{i=1}^n \int_{P_0^*(s)}^{P_i} d\zeta_g^*(s) \right) \text{ mod periods.}$$

Let

$$u: f^*\Omega_{J/T_g}^1 \longrightarrow \Omega_{V_{T_g}^{(n)}/T_g}^1$$

be the map induced by f . Since J and $V_{T_g}^{(n)}$ are smooth over T_g of relative dimensions g and n respectively, the sheaves

$$f^*\Omega_{J/T_g}^1 \quad \text{and} \quad \Omega_{V_{T_g}^{(n)}/T_g}^1$$

are locally free of ranks g and n respectively. Thus, we may consider the analytic subspace $Z^r(u) \subseteq V_{T_g}^{(n)}$ of §1. We will denote by G_n^r the analytic space $Z^r(u)$ which arises in this situation. We will see in §4 that $(G_n^r)_s$ is what was denoted by $G_n^r(V_s)$ in §0.

We wish to study the infinitesimal structure of G_n^r . To do this, we need explicit knowledge of the above map f . And to obtain this knowledge, we need certain variational formulas which are contained in the next section.

3. The variational formula. For a detailed treatment of the material in this section, the reader is referred to Rauch [18] or Patt [17].

Let X be a compact Riemann surface of genus $g > 0$. Let $\Gamma = (\gamma_1, \dots, \gamma_g)$ and $\Delta = (\delta_1, \dots, \delta_g)$ be a canonical homotopy basis and let Π be the simply connected surface obtained by the canonical dissection of X determined by Γ and Δ (cf. [23]).

Let w be a point in the interior of Π and let $\tau_{w,\nu}(z)$ denote the (normalized) elementary integral of the second kind with pole of order $\nu + 1$ at w and zero Γ -periods.

Let ζ be an Abelian integral of the first kind. Let $a_i, i = 1, \dots, g$, denote the Γ -periods of $d\zeta$; that is,

$$a_i = \int_{\gamma_i} d\zeta, \quad i = 1, \dots, g.$$

The value of the derivatives of a determination of ζ at w and the periods of the differentials $d\tau_{w,\nu}$ are related by

$$\zeta^{(\nu+1)}(w) = \frac{\nu!}{2\pi i} \sum_{j=1}^g a_j \int_{\delta_j} d\tau_{w,\nu}(z)$$

which follows from the bilinear relation for differentials of the first and second kinds [23, p. 260].

Let Q_1, \dots, Q_n be distinct points in the interior of Π such that all the Q_j are different from w and none of the Q_j is a zero of $d\zeta$. Let $t_j, j = 1, \dots, n$, be a local parameter at Q_j . Let D_1, \dots, D_n be disjoint disks about Q_1, \dots, Q_n respectively, such that D_j lies in the domain of t_j , is completely

contained in the interior of Π and such that no D_j contains either w or any zero of $d\xi$.

Inside D_j , we can vary the local parameter t_j to a new parameter t_j^* given by

$$t_j^* = t_j + c_j/t_j, \quad j = 1, \dots, n,$$

where c_j is sufficiently small. This defines a new Riemann surface X^* , having the same canonical homotopy basis as X has (since all variations take place in the interior of Π).

Let ζ^* be the Abelian integral of first kind on X^* with the same Γ -periods as ζ . We wish to compute

$$\Delta \zeta^{(\nu+1)}(w) = \zeta^{*(\nu+1)}(w) - \zeta^{(\nu+1)}(w).$$

Notation. $d\tau_{w,\nu}d\xi$ is a (not necessarily finite) quadratic differential on X . Locally at Q_j , we may write $d\tau_{w,\nu}d\xi = h(t_j)dt_j^2$. We now introduce the notation $\tau'_{w,\nu}(Q_j)\xi'(Q_j)$ for $h(0)$.

Utilizing the techniques and formulas in [18] and [17], one can obtain the following proposition:

PROPOSITION 5.

$$\Delta \zeta^{(\nu+1)}(w) = \nu! \sum_{m=1}^n c_m \tau'_{w,\nu}(Q_m) \xi'(Q_m) + O(c^2)$$

where $c = \max_{1 \leq m \leq n} |c_m|$.

We will also want to use the following theorem, due to Patt [17]:

THEOREM 2. *One may choose $3g - 3$ points Q_1, \dots, Q_{3g-3} on X such that, if c_m is the variation parameter at Q_m , then a neighborhood of the origin in the c_1, \dots, c_{3g-3} space describes a complex-analytic structure for a neighborhood of X in the Teichmüller space. Moreover, the set of collections of $3g - 3$ points with this property is open in X^{3g-3} .*

PROOF. The first assertion follows from Theorems 2 and 4 of [17]. Although Patt does not state the second assertion, his proofs demonstrate it, as was noted by Farkas [3, p. 885].

4. The equations which define the tangent space. Let X be a compact Riemann surface of genus $g > 1$. Let $\{\gamma_j, \delta_j\}_{j=1}^g$ be a canonical homotopy basis and let $\{d\xi_k\}_{k=1}^g$ be a basis of the holomorphic differentials. Put

$$A_{jk} = \int_{\gamma_j} d\xi_k, \quad j, k = 1, \dots, g.$$

Let P be a point of X and let t be a local parameter on X at P . Write

$$d\xi_k = \sum_{l=0}^{\infty} a_{k,l} t^l dt.$$

Fix a point P_0 different from P . Choose a point (Q_1, \dots, Q_{3g-3}) from the open subset of X^{3g-3} in Theorem 2 such that all the Q_m are different from P and P_0 and such that none of the Q_m is a zero of any $d\xi_k$. Perform the variation described in §3, taking the disk about each Q_m sufficiently small so that no two disks intersect and no disk contains P, P_0 , or any zero of any $d\xi_k$. Let c_m denote the variation parameter at Q_m , $m = 1, \dots, 3g-3$, as in §3.

Let $s_0 \in T_g$ be the module point of X (i.e., $V_{s_0} = X$). By definition of the variation, there exists a complex-analytic neighborhood U of s_0 in T_g such that, for all $s' \in U$, the curves $\{\gamma_j, \delta_j\}_{j=1}^g$ are a canonical homotopy basis on $V_{s'}$, the points P_0 and P are on $V_{s'}$, and t is a local parameter on $V_{s'}$ at P . Choose holomorphic sections $d\xi_k^*$, $k = 1, \dots, g$, of $h_*\Omega^1_{V/T_g}$ such that

$$\int_{\gamma_j} d\xi_k^*(s') = A_{jk}, \quad s' \in U, \quad j, k = 1, \dots, g$$

(cf. [15, §3]).

PROPOSITION 6. *With notation as in §3 and above, if we define $a_{k,l}^*$ by $d\xi_k^* = \sum_{l=0}^{\infty} a_{k,l}^* t^l dt$, then we have*

$$a_{k,l}^* = a_{k,l} + \sum_{m=1}^{3g-3} c_m \tau'_{P,l}(Q_m) \xi'_k(Q_m) + O(c^2).$$

PROOF. The variational formula (Proposition 5) shows that this equality holds in a complex-analytic neighborhood of (s_0, P) on V . This is the main import of the variational formula.

In order to study the map

$$u: f^*\Omega^1_{J/T_g} \rightarrow \Omega^1_{V_{T_g}^{(n)}/T_g}$$

of §2, we first consider the divisor nP on X . Let t_1, \dots, t_n be n copies of t , and let $\sigma_1, \dots, \sigma_n$ denote the n elementary symmetric functions in t_1, \dots, t_n .

PROPOSITION 7. *Local parameters on $V_{T_g}^{(n)}$ at (s_0, nP) are given by $c_1, \dots, c_{3g-3}, \sigma_1, \dots, \sigma_n$.*

PROOF. By Theorem 2, local parameters on T_g at s_0 are given by c_1, \dots, c_{3g-3} . By [1], local parameters on $X^{(n)}$ at nP are given by $\sigma_1, \dots, \sigma_n$. By the definition of the variation in §3, local parameters on $(V_{T_g}^{(n)})_{s'}$ at nP , for $s' \in U$, are also given by $\sigma_1, \dots, \sigma_n$. Thus, local parameters on $V_{T_g}^{(n)}$ at (s_0, nP) are given by $c_1, \dots, c_{3g-3}, \sigma_1, \dots, \sigma_n$.

Put

$$\tau_j = t_1^j dt_1 + \dots + t_n^j dt_n, \quad j = 0, 1, 2, \dots$$

We have

PROPOSITION 8. *The space of holomorphic 1-forms on X is naturally isomorphic to the space of holomorphic 1-forms on $X^{(n)}$. Both these spaces are isomorphic to the space of symmetric holomorphic 1-forms on the Cartesian product X^n . If $d\zeta = \sum_{i=1}^{\infty} a_i t^i dt$ is a holomorphic 1-form on X and $d\tilde{\zeta}$ is the corresponding symmetric holomorphic 1-form on X^n , then $d\tilde{\zeta} = \sum_{i=0}^{\infty} a_i \tau_i$.*

PROOF. [14, pp. 226–227].

This result is easily seen to relativize to the following proposition.

PROPOSITION 9. *The space of relative holomorphic 1-forms on $V_{T_g}^{(n)}$ over T_g and the space of relative holomorphic 1-forms on V over T_g are naturally isomorphic. Both spaces are isomorphic to the space of relative symmetric holomorphic 1-forms on $V_{T_g}^n$, the product over T_g of n copies of V , over T_g . If $d\tilde{\zeta}_k^*$ is the relative symmetric holomorphic 1-form on $V_{T_g}^n$ over T_g corresponding to $d\zeta_k^*$ (cf. Proposition 6), then*

$$d\tilde{\zeta}_k^* = \sum_{l=0}^{\infty} a_{k,l}^* \tau_l.$$

We will identify relative symmetric holomorphic 1-forms on $V_{T_g}^n$ over T_g and relative holomorphic 1-forms on $V_{T_g}^{(n)}$ over T_g .

Now, we can express $d\tilde{\zeta}_k^*$ in terms of $d\sigma_1, \dots, d\sigma_n$ by using the following identities [14]:

$$\sigma_k \tau_0 - \sigma_{k-1} \tau_1 + \dots + (-1)^k \tau_k = d\sigma_{k+1}.$$

(By convention, $\sigma_k = 0$ and $d\sigma_k = 0$ if $k > n$.) Inverting these identities, and writing out only the linear terms, we obtain

$$\tau_k = (-1)^k (d\sigma_{k+1} - \sigma_1 d\sigma_k - \dots - \sigma_k d\sigma_1)$$

+ higher order terms.

Thus we may write

$$\begin{aligned}
 \text{(\#)} \quad d\tilde{\xi}_k^* &= \sum_{l=0}^{\infty} (-1)^l \left[\left(a_{k,l} + \sum_{m=1}^{3g-3} c_m \tau'_{P,l}(Q_m) \xi'_k(Q_m) \right) \right. \\
 &\quad \left. (d\sigma_{l+1} - \sigma_1 d\sigma_l - \cdots - \sigma_l d\sigma_1) \right] + O(\sigma^2, c^2)
 \end{aligned}$$

where $O(\sigma^2, c^2)$ denotes higher order terms in the σ_j and the c_m .

By definition of the map $f: V_{T_g}^{(n)} \rightarrow J$ in §2, it is easy to see that f is given at (s_0, nP) by

$$f(s_0, nP) = \left(s_0, \int_{P_0}^P d\tilde{\xi}_1^*(s_0), \dots, \int_{P_0}^P d\tilde{\xi}_g^*(s_0) \right) \text{ mod periods}$$

where the integrals $\int_{P_0}^P d\tilde{\xi}_k^*(s_0)$ are evaluated by recalling that t_1, \dots, t_n are just copies of t . Let $\partial\tilde{\xi}_k^*/\partial\sigma_j$ be given by

$$d\tilde{\xi}_k^* = \sum_{j=1}^n \frac{\partial\tilde{\xi}_k^*}{\partial\sigma_j} d\sigma_j.$$

Then we have

PROPOSITION 10. *The map*

$$u: f^* \Omega_{J/T_g}^1 \rightarrow \Omega_{V_{T_g}^{(n)}/T_g}^1$$

is given locally at (s_0, nP) by the matrix

$$[\partial\tilde{\xi}_k^*/\partial\sigma_j], \quad j = 1, \dots, n, k = 1, \dots, g.$$

PROOF. This follows easily from the definitions of f and $\partial\tilde{\xi}_k^*/\partial\sigma_j$. (Compare with [3] and [6].)

REMARK. Let J denote the Jacobian variety of X and let $f_0: X^{(n)} \rightarrow J$ be the classical map (i.e., the map $f \otimes \kappa(s_0)$). Then the matrix $M = [(\partial\tilde{\xi}_k^*/\partial\sigma_j)(s_0, nP)]$ is the matrix of the map $u_0: f_0^* \Omega_J^1 \rightarrow \Omega_{X^{(n)}}^1$ at nP (cf. [3], [6]). It is then easy to see that $(G_n^r)_{s_0}$ is what was denoted by $G_n^r(X)$ in §0.

Now let ξ be a tangent vector to $V_{T_g}^{(n)}$ at (s_0, nP) . Let s_j and b_m in \mathbb{C} be given by

$$\begin{aligned}
 \xi(\sigma_j) &= s_j \epsilon, \quad j = 1, \dots, n, \\
 \xi(c_m) &= b_m \epsilon, \quad m = 1, \dots, 3g - 3.
 \end{aligned}$$

Then, using Taylor's Theorem as in §1, we have

$$(1) \quad \xi \left(\frac{\partial \tilde{\zeta}_k^*}{\partial \sigma_j} \right) = \frac{\partial \tilde{\zeta}_k^*}{\partial \sigma_j} (s_0, nP) + \epsilon \sum_{i=1}^n s_i \frac{\partial^2 \tilde{\zeta}_k^*}{\partial \sigma_i \partial \sigma_j} (s_0, nP) + \epsilon \sum_{m=1}^{3g-3} b_m \frac{\partial^2 \tilde{\zeta}_k^*}{\partial c_m \partial \sigma_j} (s_0, nP).$$

We will now use (#) to compute the partial derivatives of $\partial \tilde{\zeta}_k^* / \partial \sigma_j$ with respect to σ_i and with respect to c_m . (We remind the reader that the functions σ_i and c_m vanish at (s_0, nP) .) We obtain

$$\frac{\partial^2 \tilde{\zeta}_k^*}{\partial \sigma_i \partial \sigma_j} (s_0, nP) = (-1)^{j+l} a_{k,j+l-1}$$

and

$$\frac{\partial^2 \tilde{\zeta}_k^*}{\partial \sigma_m \partial \sigma_j} (s_0, nP) = \tau'_{P,j-1}(Q_m) \zeta'_k(Q_m).$$

Substituting these expressions for the partial derivatives evaluated at (s_0, nP) into (1) gives us

PROPOSITION 11.

$$\xi \left(\frac{\partial \tilde{\zeta}_k^*}{\partial \sigma_j} \right) = \frac{\partial \tilde{\zeta}_k^*}{\partial \sigma_j} (s_0, nP) + \epsilon \sum_{i=1}^n (-1)^j s_i a_{k,j+l-1} + \epsilon \sum_{m=1}^{3g-3} b_m \tau'_{P,j-1}(Q_m) \zeta'_k(Q_m).$$

Now on to the general case. Consider a divisor D on X of the form $D = m_1 P_1 + \dots + m_d P_d$. Assume D is in $G_n^r(X)$ and choose a basis $\{d\zeta_k\}_{k=1}^g$ of the holomorphic differentials on X such that the last $i = \dim H^1(X, \mathcal{O}_X(D))$ of them vanish on D .

In performing the variation in §3, choose a point (Q_1, \dots, Q_{3g-3}) from the open set in X^{3g-3} in Theorem 2 so that each Q_m is different from P_0, P_1, \dots, P_d and any other zero of any $d\zeta_k$. (The choice of this point will be further modified later.) Take the disk about each Q_m sufficiently small so that no two disks intersect and such that no disk contains P_0, P_1, \dots, P_d or any other zero of any $d\zeta_k$.

Let $f_j: V_{T_g}^{(m_j)} \rightarrow J$ be the map defined in §2 and let

$$u_j: f_j^* \Omega_{J/T_g}^1 \rightarrow \Omega_{V_{T_g}/T_g}^1(m_j)$$

be the map induced by f_j . The obvious map

$$V_{T_g}^{(m_1)} \times_{T_g} V_{T_g}^{(m_2)} \times_{T_g} \cdots \times_{T_g} V_{T_g}^{(m_d)} \rightarrow V_{T_g}^{(n)}$$

is a local analytic isomorphism by an argument analogous to that given in [14] in the case of a curve over a field. Locally, the map f is the one induced by the f_j and the map u is the one induced by the u_j . Thus, the matrix of u locally at (s_0, D) is obtained by “stacking” the matrices of the u_j locally at $(s_0, m_j P_j)$.

Let ξ be a tangent vector to $V_{T_g}^{(n)}$ at (s_0, D) and let ξ_j be the tangent vector to $V_{T_g}^{(m_j)}$ at $(s_0, m_j P_j)$ induced by ξ , for $j = 1, \dots, d$. Then the matrix of $\xi^* u$ is obtained by “stacking” the matrices of the $\xi_j^* u_j$, for $j = 1, \dots, d$.

Let M' denote the matrix of $\xi^* u$. Let μ denote the leading minor of order $n - r$ of M , the matrix $[(\partial \tilde{\xi}_k^* / \partial \sigma_j)(s_0, D)]$, and let μ' denote the leading minor of order $n - r$ of M' . Then we have $\mu' = \mu + c\epsilon$ for some c in \mathbb{C} . Now, by our choice of a basis of the holomorphic differentials on X , the last i columns of M are identically zero, hence the last i columns of M' contain “pure” ϵ terms (i.e., members of the maximal ideal of $\mathbb{C}[\epsilon]/(\epsilon^2)$). Thus, in computing a minor of order $n - r + 1$ containing μ' , any ϵ 's in the first $n - r$ columns will be “killed” by the ϵ in the last column of the minor of order $n - r + 1$. Hence, we have established

LEMMA 3. *For purposes of computing the minors of order $n - r + 1$ of M' , we may replace the first $n - r$ columns of M' by the first $n - r$ columns of M .*

Let M denote the resulting matrix.

M has a particularly nice form in the case that $D = P_1 + P_2 + \cdots + P_n$, with all points distinct. Let t_j be a local parameter at P_j and write $d\xi_k = \varphi_{j,k} dt_j$. Then we have

$$M = \begin{bmatrix} \varphi_{j,k}(P_j) & \vdots & \left(s_j \varphi'_{j,k}(P_j) + \sum_{m=1}^{3g-3} b_m \tau'_{P_j,0}(Q_m) \xi'_k(Q_m) \right) \\ j = 1, \dots, n & & j = 1, \dots, n \\ k = 1, \dots, g - i & & k = g - i + 1, \dots, g \end{bmatrix}$$

Going back to the general case, recall that, by Proposition 1, ξ will be a tangent vector to G_n^r at (s_0, D) iff the minors of order $n - r + 1$ of the matrix M all vanish. Assume $D = m_1 P_1 + \cdots + m_d P_d$ is in $G_n^r(X) - G_n^{r+1}(X)$. Then the matrix M has rank precisely $n - r$. Hence, by permuting the rows of

M , if necessary, we end up with a matrix whose leading minor of order $n - r$, which we will denote by μ , is nonzero. We will continue to denote this matrix by M , although its form may differ slightly from that specified earlier.

Perform the same row permutations as above on the matrix M and denote the resulting matrix also by M . Then μ is also the leading minor of order $n - r$ of M , so we may apply Lemma 2. Thus, for all the minors of order $n - r + 1$ of M to vanish, it is sufficient that every minor of order $n - r + 1$ which contains μ vanishes. The vanishing of each of these minors gives rise to a linear equation in the s_j and the b_m .

Let $\mu_{j,k}$ denote the minor of order $n - r + 1$ of M obtained by adjoining to μ the first $n - r$ elements of the $(n - r + j)$ th row of M and the first $n - r$ elements and the $(n - r + j)$ th element of the $(n - r + k)$ th column of M (thus j runs from 1 through r and k runs from 1 through i). The equation $\mu_{j,k} = 0$ is of the form $\epsilon E_{j,k} = 0$ where $E_{j,k}$ is a linear equation in the s_j and the b_m with coefficients in C .

We will now view the s_j and the b_m as being unknowns (as in §1). Thus, $E_{j,k}$ is an equation in $3g - 3 + n$ unknowns. By the discussion after Proposition 1, the dimension of the tangent space to G'_n at (s_0, D) is

$$3g - 3 + n - (\text{the number of } E_{j,k} \text{ which are linearly independent}).$$

Consider the coefficient of b_m in $E_{j,k}$. This coefficient will be a linear combination of certain of the $\tau'_{P_j, \nu}(Q_m) \zeta'_k(Q_m)$. That is, the coefficient of b_m will be a certain quadratic differential (the above linear combination of certain of the $d\tau_{P_j, \nu} d\zeta_k$) evaluated at the point Q_m . It should be noted that, by the symmetry of the matrix M in the b_m , this quadratic differential does not depend on m , but only on j and k . The coefficient of b_1 in $E_{j,k}$ is the value of this quadratic differential at Q_1 , the coefficient of b_2 in $E_{j,k}$ the value at Q_2 , etc. Put $\alpha_{j,k}$ equal to the above linear combination of certain of the $d\tau_{P_j, \nu} d\zeta_k$. Then $\alpha_{j,k}$ is a (not necessarily finite) quadratic differential.

Notation. Choose a local parameter u_m on X at Q_m and write $\alpha_{j,k} = g(u_m) du_m^2$. Then we will write $\alpha_{j,k}(Q_m)$ for $g(0)$. Hence, by the above discussion, $\alpha_{j,k}(Q_m)$ is the coefficient of b_m in $E_{j,k}$.

Our aim now is to show that, in certain situations, by suitably choosing the point (Q_1, \dots, Q_{3g-3}) , we may conclude that the $E_{j,k}$ are linearly independent. Assume that $ri \leq 3g - 3$. By elementary linear algebra, to conclude that the $E_{j,k}$ are linearly independent, it is sufficient to show that the matrix of coefficients

$$A = [\alpha_{j,k}(Q_m)], \quad j = 1, \dots, r; k = 1, \dots, i; m = 1, \dots, ri,$$

is nonsingular.

LEMMA 4. Assume that the $\alpha_{j,k}$ for $j = 1, \dots, r$ and $k = 1, \dots, i$, are linearly independent and that $ri \leq 3g - 3$. Then we may choose a point (Q_1, \dots, Q_{3g-3}) from the open set in X^{3g-3} in Theorem 2 such that each Q_m is different from P_0 and no Q_m is a zero of $d\xi_1, \dots, d\xi_g$ and such that the above matrix A is nonsingular.

PROOF. The lemma will follow readily from the following

SUBLEMMA. Let β_1, \dots, β_n be n linearly independent quadratic differentials on X . Let U be an open set contained in X^n . Then we may choose a point $(P_1, \dots, P_n) \in U$ such that each P_m is different from a finite set of points of X and such that the matrix $[\beta_j(P_k)]$ ($j = 1, \dots, n, k = 1, \dots, n$) is nonsingular.

PROOF. By induction on n . If $n = 1$, then β_1 is a nontrivial quadratic differential. Hence, β_1 is nonzero and finite on a dense open set of X . So, given any open set in X , there exists a point in that set satisfying the requirements of the Sublemma.

Now suppose U is an open set contained in X^n . Let V be the projection of U onto X^{n-1} . Then V is open and, by induction, we may choose a point $(P_0, \dots, P_{n-1}) \in V$ such that each P_m is different from a finite set of points of X and such that the leading subdeterminant of order $n - 1$ of the determinant

$$\begin{vmatrix} \beta_1(P_1) \cdots \beta_1(P_{n-1}) & \beta_1 \\ \cdot & \cdot \\ \cdot & \cdot \\ \cdot & \cdot \\ \beta_n(P_1) \cdots \beta_n(P_{n-1}) & \beta_n \end{vmatrix}$$

is nonzero. Expanding the full determinant by the last column, we obtain a nontrivial linear combination of β_1, \dots, β_n . By the linear independence of these quadratic differentials, this linear combination is a nontrivial quadratic differential, hence is nonzero and finite on an open dense set W contained in X . Since U is open in X^n and W is dense in X , we may choose a point in the intersection of U and $\{(P_0, \dots, P_{n-1})\} \times W$ which satisfies the requirements of our Sublemma.

Now, since the set of points in X^{3g-3} in Theorem 2 is open, it is easy to see that we may choose a point (Q_1, \dots, Q_{3g-3}) in this set such that each Q_m is different from P_0 and the zeros of $d\xi_1, \dots, d\xi_g$ and so that Q_1, \dots, Q_{ri} make the matrix A nonsingular. This completes the proof of the lemma.

We then have

PROPOSITION 12. Suppose D is in $G_n^r(X) - G_n^{r+1}(X)$ and that $ri \leq 3g - 3$. Then if all the $\alpha_{j,k}$ are linearly independent, the dimension of the tangent space to G_n^r at (s_0, D) is $3g - 3 + \tau + r$.

PROOF. By Lemma 4, we may choose a point (Q_1, \dots, Q_{3g-3}) from the open set in Theorem 2 such that each of the Q_m is different from P_0 and the zeros of $d\zeta_1, \dots, d\zeta_g$ (note that this latter set includes the points of D), and such that the equations $E_{j,k}$ are linearly independent. Thus the dimension of the tangent space to G_n^r at (s_0, D) is $3g - 3 + n - ir = 3g - 3 + \tau + r$.

In the next section, we show that if D is in $G_n^1(X) - G_n^2(X)$, then the $\alpha_{j,k}$ are linearly independent. (Note that we have $i < 3g - 3$ if $g > 1$.)

5. The dimension of $G_n^1 - G_n^2$. For simplicity, we will first treat a divisor consisting of n distinct points. So assume $D = P_1 + P_2 + \dots + P_n$, all points distinct, is in $G_n^1(X) - G_n^2(X)$. Recall that the matrix M is

$$M = \begin{bmatrix} \varphi_{j,k}(P_j) & \vdots & \epsilon \left(s_j \varphi'_{j,k}(P_j) + \sum_{m=1}^{3g-3} b_m \tau'_{P_j,0}(Q_m) \zeta'_k(Q_m) \right) \\ j = 1, \dots, n & \vdots & j = 1, \dots, n \\ k = 1, \dots, g - i & \vdots & k = g - i + 1, \dots, g \end{bmatrix}.$$

Let $|\hat{j}|$ denote the minor of order $n - 1$ obtained by omitting the j th row from the matrix $[\varphi_{j,k}(P_j)]$ ($j = 1, \dots, n, k = 1, \dots, g - i$). Then we have

$$\alpha_{1,k}(Q) = \sum_{j=1}^n (-1)^{j-1} |\hat{j}| \tau'_{P_j,0}(Q) \zeta'_{n+k-1}(Q)$$

for $k = 1, 2, \dots, i$. Suppose we had a linear relation of the form $\sum_{k=1}^i a_k \alpha_{1,k} = 0$ with some a_i nonzero. Then this would imply that

$$(*) \quad \left(\sum_{j=1}^n (-1)^{j-1} |\hat{j}| \tau'_{P_j,0}(Q) \right) \left(\sum_{k=1}^i a_k \zeta'_{n+k-1}(Q) \right) = 0.$$

But the $d\tau_{P_j,0}, j = 1, \dots, n$, are linearly independent, since they have poles at different points. This, together with the fact that $|\hat{n}| \neq 0$, implies that there is a dense open set of points of X where the expression $\sum_{j=1}^n (-1)^{j-1} |\hat{j}| \tau'_{P_j,0}(Q)$ is nonzero.

And the linear independence of $d\zeta_n, \dots, d\zeta_g$, together with the fact that some a_i is nonzero, implies that the expression $\sum_{k=1}^i a_k \zeta'_{n+k-1}(Q)$ is nonzero on a dense open set of points of X . Hence, we may choose a point Q such that $(*)$ is nonzero, contradicting the assumption that $\alpha_{1,1}, \dots, \alpha_{1,i}$ are linearly dependent.

Now suppose $D = m_1P_1 + \dots + m_dP_d$ is in $G_n^1(X) - G_n^2(X)$. Then we have

$$\alpha_{1,k}(Q) = \zeta'_{n+k-1}(Q)(|\hat{1}|_{\tau'_{P_1,0}}(Q) + \dots + (-1)^{n-1}|\hat{n}|_{\tau'_{P_d,m_d-1}}(Q)).$$

Hence, if there existed a linear relation $\sum_{k=1}^i a_k \alpha_{1,k} = 0$, we would have

$$\left(\sum_{k=1}^i a_k \zeta'_{n+k-1}(Q)\right) \left(|\hat{1}|_{\tau'_{P_1,0}}(Q) + \dots + (-1)^{n-1}|\hat{n}|_{\tau'_{P_d,m_d-1}}(Q)\right) = 0.$$

The same reasoning as in the case of simple points applies, since $d\tau_{P_1,0}, \dots, d\tau_{P_d,m_d-1}$ are easily seen to be linearly independent (they either have poles at different points or have poles of differing orders at the same point).

REMARK. The above reasoning shows that if $D \in G_n^r - G_n^{r+1}$, then the $\alpha_{j,k}$ for a fixed j are linearly independent.

THEOREM 3. $G_n^1 - G_n^2$, if nonempty, is smooth of pure dimension $3g - 3 + \tau + 1$.

PROOF. Let (s_0, D) be any point of $G_n^1 - G_n^2$. By Proposition 12 and the work of this section, we may conclude that the dimension of the tangent space to G_n^1 at (s_0, D) is $3g - 3 + \tau + 1$. By Proposition 4, the dimension of G_n^1 at (s_0, D) is at least $3g - 3 + \tau + 1$, hence G_n^1 is smooth at (s_0, D) and has dimension precisely $3g - 3 + \tau + 1$.

REMARK. Theorem 3 does not depend upon τ being nonnegative.

THEOREM 4. Suppose that $G_n^1(X) - G_n^2(X)$ is nonempty for a generic curve X . Then $G_n^1(X) - G_n^2(X)$, for a generic X , is smooth of pure dimension $\tau + 1$.

PROOF. Under our assumption, the image of $G_n^1 - G_n^2$ in T_g would be a dense open subspace U . By Sard's Theorem, since $G_n^1 - G_n^2$ is smooth, the generic fiber of the map $G_n^1 - G_n^2 \rightarrow U$ is smooth. And since U has dimension $3g - 3$ and $G_n^1 - G_n^2$ has dimension $3g - 3 + \tau + 1$, the generic fiber has dimension $\tau + 1$. Thus, for a generic curve, $G_n^1(X) - G_n^2(X)$ is smooth of dimension $\tau + 1$.

REMARK. If $\tau \geq 0$, then by [10] we know that $G_n^r(X)$ is nonempty. If we knew that $G_n^r(X)$ were reduced for a generic X , then, since the points of G_n^{r+1} are singular points of G_n^r , we could conclude that $G_n^r(X) - G_n^{r+1}(X)$ is nonempty for generic X if $\tau \geq 0$.

6. Moduli of trigonal curves. A trigonal curve is a curve X such that $G_3^1(X)$ is nonempty. We can use Theorem 3 to compute the moduli of trigonal curves. By Clifford's Theorem, G_3^2 is empty hence, by Theorem 3, G_3^1 , if

nonempty, is smooth of pure dimension $3g - 3 + \tau + 1$. Now $\tau = 2(3 - 1) - g = 4 - g$, so G_3^1 , if nonempty, has dimension $2g + 2$.

By Theorem 1 of [12], we have that, for $g \geq 4$, if $G_3^1(X)$ is nonempty, then every component has dimension at least $5 - g$ and at most 2, with the upper bound occurring if and only if X is hyperelliptic. So, if there exists a nonhyperelliptic trigonal curve of genus g , then we must have that the dimension of the generic fiber of the map $G_3^1 \rightarrow T_g$ is 1. Examples of such curves (for every $g \geq 3$) are given in [1a, p. 196].⁽¹⁾ Hence, the dimension of the subvariety of T_g , for $g \geq 4$, of trigonal curves is $2g + 2 - 1 = 2g + 1$. This agrees with the number which appears in Segre [20] and Severi [22].

BIBLIOGRAPHY

1. A. Andreotti, *On a theorem of Torelli*, Amer. J. Math. **80** (1958), 801–828. MR 21 #1309.
- 1a. A. Andreotti and A. L. Mayer, *On period relations for abelian integrals on algebraic curves*, Ann. Scuola Norm. Sup. Pisa (3) **21** (1967), 189–238. MR 36 #3792.
2. A. Brill and M. Noether, *Über die algebraischen Funktionen und ihre Anwendung in der Geometrie*, Math. Ann. **7** (1874).
3. H. M. Farkas, *Special divisors and analytic subloci of the Teichmüller space*, Amer. J. Math. **88** (1966), 881–901. MR 35 #4406.
4. G. Grauert, *Ein Theorem der analytischen Garbentheorie und die Modulräume komplexer Strukturen*, Inst. Hautes Études Sci. Publ. Math. No. 5 (1960). MR 22 #12544.
5. A. Grothendieck, *Exposés in Séminaire Cartan*, 1960/61, Secrétariat mathématique, Paris.
6. R. C. Gunning, *Jacobi varieties*, Princeton Univ. Press, Princeton, N. J., 1972.
7. B. Iversen, *Linear determinants with applications to the Picard scheme of a family of algebraic curves*, Lecture Notes in Math., vol. 174, Springer-Verlag, Berlin, 1970. MR 45 #1917.
8. G. Kempf, *Schubert methods with an application to algebraic curves*, Stichting Mathematisch Centrum, Amsterdam, 1971.
9. G. Kempf and D. Laksov, *The determinantal formula of Schubert calculus*, Acta Math. **132** (1974), 153–162.
10. S. L. Kleiman and D. Laksov, *On the existence of special divisions*, Amer. J. Math. **94** (1972), 431–436.
11. ———, *Another proof of the existence of special divisors*, Acta Math. **132** (1974), 163–176.
12. H. H. Martens, *On the varieties of special divisors on a curve. I*, J. Reine Angew. Math. **227** (1967), 111–120. MR 35 #6682.
13. ———, *Varieties of special divisors on a curve. II*, J. Reine Angew. Math. **233** (1968), 89–100. MR 39 #2760.
14. A. Mattuck and A. Mayer, *The Riemann-Roch theorem for algebraic curves*, Ann. Scuola Norm. Sup. Pisa (3) **17** (1963), 223–237. MR 29 #102.
15. A. Mayer, *Rauch's variational formula and the heat equation*, Math. Ann. **181** (1969), 53–59. MR 40 #1588.
16. T. Meis, *Die minimale Blätterzahl der Konkretisierung einer kompakten Riemannschen Fläche*, Schr. Math. Inst. Univ. Münster No. 16 (1960). MR 26 #5158.
17. C. Patt, *Variations of Teichmüller and Torelli surfaces*, J. Analyse Math. **11** (1963), 221–247. MR 28 #4103.
18. H. E. Rauch, *Weierstrass points, branch points and moduli of Riemann surfaces*, Comm. Pure Appl. Math. **12** (1959), 543–560. MR 22 #1666.

⁽¹⁾ The author wishes to thank the referee for pointing out this reference.

19. M. Schiffer and D. C. Spencer, *Functionals of finite Riemann surfaces*, Princeton Univ. Press, Princeton, N. J., 1964.
20. B. Segre, *Sui moduli delle curve poligonali* · · · , Math. Ann. **100** (1928), 537–551.
21. F. Severi and E. Löffler, *Vorlesungen über algebraischen Geometrie*, Teubner, Leipzig, 1921.
22. ———, *Sul teorema di esistenza di Riemann*, Rend. Circ. Mat. Palermo **46** (1922).
23. G. Springer, *Introduction to Riemann surfaces*, Addison-Wesley, Reading, Mass., 1957. MR **19**, 1169.

DEPARTMENT OF MATHEMATICS, LOUISIANA STATE UNIVERSITY, BATON
ROUGE, LOUISIANA 70803