

ERGODIC TRANSFORMATIONS FROM AN INTERVAL INTO ITSELF¹

BY

TIEN-YIEN LI AND JAMES A. YORKE

ABSTRACT. A class of piecewise continuous, piecewise C^1 transformations on the interval $J \subset R$ with finitely many discontinuities n are shown to have at most n invariant measures.

1. The way phenomena or processes evolve or change in time is often described by differential equations or difference equations. One of the simplest mathematical situations occurs when the phenomenon can be described by a single number and when this number can be estimated purely as a function of the previous number. That is, when the number x_{n+1} can be written as $x_{n+1} = \tau(x_n)$ where τ maps an interval $J \subset R$ into itself. For $x \in J$, let $\tau^0(x)$ denote x and $\tau^{n+1}(x)$ denote $\tau(\tau^n(x))$ for $n = 0, 1, \dots$. We will say $p \in J$ is a *periodic point with period n* if $p = \tau^n(p)$ and $p \neq \tau^k(p)$ for $1 < k < n$. We say p is a *periodic point* if p is periodic with some period $n \geq 1$. In this paper we assume τ is piecewise continuous and piecewise twice continuous differentiable. We also assume that

$$(1.1) \quad \inf_{x \in J_1} \left| \frac{d}{dx} \tau(x) \right| > 1 \quad \text{where } J_1 = \left\{ x \in J, \frac{d}{dx} \tau(x) \text{ exists} \right\}.$$

We will refer to the points of $J - J_1$ as the *points of discontinuity*. For such a transformation all periodic points of τ are unstable. See, for example, [4]. For $x \in J$, let $\Lambda(x)$ be the set of limit points of $\tau^n(x)$, that is,

$$\Lambda(x) = \bigcap_{N=1}^{\infty} \overline{\{\tau^n(x)\}_{n=N}^{\infty}}.$$

Notice that $\tau(\Lambda(x)) = \Lambda(x)$. We show that $\Lambda(x)$ is the union of (one or more) intervals of positive length for almost all $x \in J$. Furthermore, there is a finite collection of sets L_1, \dots, L_n , where each L_i ($i = 1, \dots, n$) is a union of disjoint intervals, such that for almost all $x \in J$, $\Lambda(x)$ is one of the sets L_i .

Received by the editors February 25, 1975 and, in revised form, March 9, 1976.

AMS (MOS) subject classifications (1970). Primary 28A65.

Key words and phrases. Frobenius-Perron operator, invariant measures, function of bounded variation, invariant sets.

¹Research was partially supported by National Science Foundation under grant MSC 76-24432.

© American Mathematical Society 1978

The intersection $L_i \cap L_j$ contains at most a finite number of points when $i \neq j$ and each L_i contains in its interior a point of discontinuities of τ and/or $d\tau/dx$. Hence as a corollary we have that if τ has at most one point of discontinuity then $n = 1$ and $\Lambda(x) = \Lambda(y)$ for almost all x and y in J .

A measure μ is said to be *invariant (under τ)* if for all measurable $S \subset J$, we have $\mu(S) = \mu(\tau^{-1}(S))$ where $\tau^{-1}(S) = \{x \in J: \tau(x) \in S\}$. In [3], Lasota and Yorke showed for the transformation τ under study here there exists an absolutely continuous invariant measure. This result generalizes some previous results of Gelfond [1], Lasota [2], Parry [5] and Rényi [6]. We show in §2 that for each L_i there exists a unique absolutely continuous invariant measure μ_i which is invariant for τ such that

$$\mu_i(L_i) = 1 \quad \text{and} \quad \mu_i(L_j) = 0 \quad \text{for } i \neq j.$$

Furthermore, any absolutely continuously invariant measure μ can be written as $\sum a_i \mu_i$ for appropriately chosen constants a_i .

If τ and τ' have at most one point of discontinuity then there is a unique absolutely continuous invariant measure for τ . Hence τ is ergodic on J . Other properties of these transformations are discussed in [4].

The referee has pointed out that in [8] a similar result is proved using different techniques. They do not estimate the number of invariant measures and so do not discuss conditions under which τ is ergodic.

2. Let L^1 denote the space of all integrable functions defined on the interval $[0,1]$ and let $\|\cdot\|$ be the L^1 -norm. Lebesgue measure on $[0,1]$ will be denoted by m . We say $f \in L^1$ is a *function of bounded variation in L^1* if f equals almost everywhere some function of bounded variation. Let $\tau: [0,1] \rightarrow [0,1]$ be a piecewise continuous and piecewise c^2 -function with $\{x_1, \dots, x_k\} = J - J_1$, the points of discontinuity of τ and τ' . We assume (1.1) holds for τ . The Frobenius-Perron operator $P_\tau: L^1 \rightarrow L^1$ is defined as a linear operator such that for $f \in L^1$, $P_\tau f$ is the function with

$$(2.1) \quad \int_E P_\tau f = \int_{\tau^{-1}(E)} f$$

for all measurable sets E . We say f is *invariant (under τ)* or is an *invariant function* (of P_τ) if

$$(2.2) \quad \int_E f = \int_{\tau^{-1}(E)} f$$

for every measurable set $E \subset [0,1]$. Notice that f is invariant if and only if $P_\tau f = f$ almost everywhere. It is well known that f is invariant under τ if and only if the measure $d\mu = f dm$ is invariant under τ . In [3], Lasota and Yorke prove that invariant functions of P_τ exist and every invariant function of P_τ is a function of bounded variation in L^1 .

DEFINITION 2.1. Let f be a function defined on $[0,1]$. We call the set, on which the function f is nonzero, the *support* of f and denote it, $\text{spt } f$. Notice that $\text{spt } f$ need not be closed in our definition.

The following property concerning the structure of the support of a bounded variation function is essential for this paper.

PROPOSITION 2.1. *If f is a function of bounded variation then*

$$\text{spt } f = \bigcup_{n=0}^P K_n \cup M, \quad 0 \leq P < \infty,$$

where K_n are disjoint intervals, M is a countable set and

$$\bigcup_{n=0}^P K_n \cap M = \emptyset.$$

PROOF. We only need to show that M is at most countable. Suppose this is not the case. Then there exists $m > 0$ such that the set $S = \{x: f(x) \geq 1/m\} \cap M$ is uncountable. For, otherwise, the set M would be a union of countable sets. Since M contains no interior points the variation of f over any interval containing S would be unbounded. Hence M is at most countable.

□

Let F be the set $f \in L^1$ which is invariant under τ (so F is a subspace). As mentioned earlier, by [3], each f in F represents a class of functions which is equal almost everywhere to a function f_0 of bounded variation. By Proposition 2.1, we can write $\text{spt } f_0$ as a disjoint union of countable intervals $\{K_n\}$ and a set M which is at most countable. Let $f_1 = f_0$ on $\bigcup K_n$ and $f_1 = 0$ elsewhere. Then, f_1 equals f_0 almost everywhere. Hence, we may assume, without loss of generality, that every f in F is a function of bounded variation, and its support consists of closed intervals.

THEOREM 1. *There is a finite collection of sets L_1, \dots, L_n and a set of functions $\{f_1, \dots, f_n\} \subset F$ such that*

- (1) each L_i ($1 \leq i \leq n$) is a finite union of closed intervals;
- (2) $L_i \cap L_j$ contains at most a finite number of points when $i \neq j$;
- (3) each L_i ($1 \leq i \leq n$) contains at least one point of discontinuity x_j ($j \in \{1, \dots, k\}$) in its interior; hence $n \leq k$;
- (4) $f_i(x) = 0$ for $x \notin L_i$, $1 \leq i \leq n$, and $f_i(x) > 0$ for almost all x in L_i ;
- (5) $\int_{L_i} f_i(x) dx = 1$ for $1 \leq i \leq n$;
- (6) if $g \in F$ satisfies (4) and (5) for some $1 \leq i \leq n$, then $g = f_i$ almost everywhere;
- (7) every $f \in F$ can be written as $f = \sum_{i=1}^n a_i f_i$ with suitable chosen $\{a_i\}$.

REMARK. Consider the simple transformation of the form

$$\tau(x) = 2x, \quad 0 \leq x < 1/2,$$

$$\tau(x) = (2 - a) + 2(a - 1), \quad 1/2 \leq x < 1,$$

where $0 < a < 1/2$ (see Figure 1). In [7], S. Ulam pointed out that it was not known even in this simple case whether there was a function invariant under τ . The complete answer to this problem is now clear. In fact, the existence of an invariant function is guaranteed by Theorem 1 of Lasota-Yorke [3] and the uniqueness (up to constant multiples) follows immediately from Theorem 1 stated above.

Before proving Theorem 1, we give the following definitions and a series of lemmas.

DEFINITION 2.2. We write " $A \subset B$ a.e." if $A, B \subset [0,1]$ and $x \in B$ for almost all x in A . We write " $A = B$ a.e." if both $A \subset B$ a.e. and $B \subset A$ a.e. are satisfied. We say a set A is *invariant* (or is *invariant under τ*) if A is a measurable subset of $[0,1]$ and $\tau(A) = A$ a.e. (Notice this does not imply $\tau^{-1}(A) = A$ a.e.)




FIGURE 1

We now list some obvious properties of invariant sets which are for later reference. If A is an invariant set, then

$$(2.3) \quad m(\tau(A)) = m(A),$$

$$(2.4) \quad \tau(A) \subset A \text{ a.e.},$$

(2.5) for any invariant measure μ with $d\mu = f dm$ (where $f \in L^2$ is invariant), we have

$$\mu(\tau^{-1}(A) - A) = 0.$$

If A_1 and A_2 are invariant sets, then

- (a) $\tau(A_1 \cap A_2) = A_1 \cap A_2$ a.e.,
- (b) $\tau(A_1 \cup A_2) = A_1 \cup A_2$ a.e.,
- (c) $\tau(A_1 - A_2) = A_1 - A_2$ a.e.

These are all quite simple except possibly (2.5) which we obtain as Corollary 2.1.

LEMMA 2.1. *Let A be a measurable set satisfying $\tau(A) \subset A$ a.e. Then, for any invariant function f , we have*

$$(2.7) \quad \int_{\tau^{-1}(A)-A} f = 0.$$

PROOF. Since $\tau(A) \subset A$ a.e., so $A \subset \tau^{-1}(A)$ a.e. Then,

$$\int_{\tau^{-1}(A)-A} f = \int_{\tau^{-1}(A)} f - \int_A f = 0.$$

COROLLARY 2.1. *If A is invariant, then (2.7) is satisfied.*

Write X_S for the characteristic function of a set $S \subset [0,1]$.

LEMMA 2.2. *Let A be an invariant set. For any invariant function f , the product (pointwise multiplication) $f \cdot X_A$ is invariant.*

PROOF. Let S be any measurable set. Then,

$$\begin{aligned} \int_S f \cdot X_A &= \int_{S \cap A} f = \int_{\tau^{-1}(S \cap A)} f = \int_{\tau^{-1}(S) \cap \tau^{-1}(A)} f \\ &= \int_{\tau^{-1}(S) \cap (\tau^{-1}(A)-A)} f + \int_{\tau^{-1}(S) \cap A} f = \int_{\tau^{-1}(S) \cap A} f. \end{aligned}$$

The last equality follows by Corollary 2.1. \square

For $f \in L^2$, we write $P(f) = \{x: f(x) > 0\}$ and $N(f) = \{x: f(x) < 0\}$. We will often write P, N for $P(f), N(f)$ when no clarification is needed.

LEMMA 2.3. *If f is invariant, the sets P and N are invariant.*

PROOF. Write

$$\tau^{-1}(P) = (\tau^{-1}(P) \cap P) \cup (\tau^{-1}(P) \cap N) \cup (\tau^{-1}(P) \cap Z)$$

where $Z = \{x: f(x) = 0\}$. Then,

$$\int_P f = \int_{\tau^{-1}(P)} f = \int_{\tau^{-1}(P) \cap N} f + \int_{\tau^{-1}(P) \cap P} f + \int_{\tau^{-1}(P) \cap Z} f \geq \int_P f$$

where the equal sign holds only if

$$m(\tau^{-1}(P) \cap P) = m(P) \quad \text{and} \quad m(\tau^{-1}(P) \cap N) = 0.$$

That is,

$$\tau^{-1}(P) \supset P \text{ a.e. and } \tau^{-1}(P) \cap N = \emptyset \text{ a.e.}$$

Hence, $\tau(P) \subset P$ a.e. Let $A = \tau(P)$ then $\tau(A) \subset A$ a.e. By Lemma 2.1

$$\int_{\tau^{-1}(A)-A} f = 0 \quad \text{and} \quad \int_{\tau^{-1}(P)-P} f = 0.$$

But $P \subset \tau^{-1}(A) \subset \tau^{-1}(P)$ a.e., it follows that

$$\int_{\tau^{-1}(A)-A} f = \int_{P-\tau(P)} f + \int_{\tau^{-1}(A)-P} f = \int_{P-\tau(P)} f = 0.$$

Since $f > 0$ in $P - \tau(P)$, hence $m(P - \tau(P)) = 0$. Therefore $\tau(P) = P$ a.e. Similarly $\tau(N) = N$ a.e. \square

LEMMA 2.4. *If f is invariant then $f \cdot X_P$ and $f \cdot X_N$ are invariant.*

PROOF. By Lemma 2.2 and Lemma 2.3 these functions are invariant. \square

Lemmas 2.5 through 2.8 are for demonstrating how to “transform” a set of linearly independent invariant functions into a set of invariant functions with disjoint supports.

LEMMA 2.5. *Let f_1 and f_2 be invariant and let $S_1 = \text{spt } f_1$ and $S_2 = \text{spt } f_2$. Then,*

(2.8) S_1 and S_2 are invariant, and

(2.9) $S_1 - S_1 \cap S_2$ and $S_2 - S_1 \cap S_2$ are invariant.

PROOF. Since $\text{spt } f_i = P(f_i) \cup N(f_i)$ for $i = 1, 2$, by Lemma 2.3 and (2.6) we easily see that both S_1 and S_2 are invariant. (2.9) is a direct result of repeated application of (2.6). \square

LEMMA 2.6. *If f_1 and f_2 are linearly independent functions in F with $\|f_1\| = \|f_2\| = 1$ then there exist f_1^* and f_2^* such that*

(a) $f_1^* \geq 0, f_2^* \geq 0$ and $\|f_1^*\| = \|f_2^*\| = 1$;

(b) $\text{spt } f_1^*$ and $\text{spt } f_2^*$ are disjoint;

(c) for each $i = 1, 2$, $\text{spt } f_i^*$ is a union of disjoint intervals contained in $\text{spt } f_1 \cup \text{spt } f_2$.

PROOF. If $i = 1$ or 2 , we have $P(f_i)$ and $N(f_i)$ are both nonempty, then we may let

$$f_1^* = (f_i \wedge 0) / \|f_i \wedge 0\| \quad \text{and} \quad f_2^* = (f_i \vee 0) / \|f_i \vee 0\|$$

where $f_i \vee 0 = \max\{f_i, 0\}$ and $f_i \wedge 0 = \max\{-f_i, 0\}$. We have the remaining cases when $f_i \geq 0$ or $f_i \leq 0$ for each i . In the following construction we assume $f_i \geq 0$ for each i , replacing f_i by $-f_i$ if necessary. In this case, neither $f_1 \geq f_2$ a.e. nor $f_2 \geq f_1$ a.e. is true, otherwise, since $\|f_1\| = \|f_2\|, f_1 = f_2$ a.e. Hence, neither $(f_1 - f_2) \vee 0$ nor $(f_1 - f_2) \wedge 0$ is equal to zero almost everywhere. The lemma is proved by letting $f_1^* = ((f_1 - f_2) \vee 0) / \|(f_1 - f_2) \vee 0\|$ and $f_2^* = ((f_1 - f_2) \wedge 0) / \|(f_1 - f_2) \wedge 0\|$. \square

LEMMA 2.7. *Let $\{f_1, \dots, f_m\}$ be a subset of F with disjoint supports and $\|f_i\| = 1, f_i \geq u$ for all $1 \leq i \leq m$. If f_{m+1} is linearly independent of*

$\{f_1, \dots, f_m\}$ then there exists a set of nonnegative functions $\{f_1^*, \dots, f_{m+1}^*\} \subset F$ with disjoint supports and $\|f_i^*\| = 1$ for $1 \leq i \leq m + 1$.

PROOF. Without loss of generality we may suppose $f_{m+1} \geq 0$ a.e. For if both $P(f_{m+1})$ and $N(f_{m+1})$ are nonempty, then either $f^+ = f_{m+1} \vee 0$ or $f^- = f_{m+1} \wedge 0$ is linearly independent of $\{f_1, \dots, f_m\}$. Otherwise,

$$f^+ = \sum_{i=1}^m a_i f_i, \quad f^- = \sum_{i=1}^m b_i f_i$$

implies $f_{m+1} = f^+ - f^- = \sum_{i=1}^m (a_i - b_i) f_i$ and f_{m+1} is linearly dependent of $\{f_1, \dots, f_m\}$.

Let $S_1 = \cup_{i=1}^m \text{spt } f_i$ and $S_2 = \text{spt } f_{m+1}$. Consider the following cases:

(1) S_1 and S_2 are disjoint. In this case the lemma is obvious. f_1^* may be chosen as f_i for $1 \leq i \leq m$ and $f_{m+1}^* = f_{m+1} / \|f_{m+1}\|$.

(2) $S_1 \subset S_2$ and $S_1 \neq S_2$. By repeated application of (2.6) and Lemma 2.5, $S_2 - S_1$ is invariant. Hence, by Lemma 2.1, $f_{m+1} \cdot X_{(S_2 - S_1)}$ is invariant and the lemma holds for $f_i^* = f_i$ for all $1 \leq i \leq m$ and

$$f_{m+1}^* = (f_{m+1} \cdot X_{(S_2 - S_1)}) / \|f_{m+1} \cdot X_{(S_2 - S_1)}\|.$$

(3) $S_2 - S_1 = \emptyset$ and $S_2 \subsetneq S_1$. There exists a function, say f_1 , such that $A = \text{spt } f_1 \cap S_2 \neq \emptyset$, and $f_1 \neq f_{n+1}$ in A . Let $f_1' = (f_1 \cdot X_A) / \|f_1 \cdot X_A\|$. By applying Lemma 2.6, we have f_1^* and f_{n+1}^* with $\|f_1^*\| = \|f_{n+1}^*\| = 1$ and disjoint supports. The results of the lemma follow by letting $f_i^* = f_i$ for all $2 \leq i \leq n$.

(4) $S_1 = S_2$. Suppose for each $1 \leq i \leq m$, there exists $\alpha_i \neq 0$ such that $f_{n+1} \cdot X_{\text{spt } f_i} = \alpha_i f_i$. Then,

$$f_{m+1} = \sum_{i=1}^m f_{m+1} \cdot X_{\text{spt } f_i} = \sum_{i=1}^m \alpha_i f_i.$$

This is impossible since f_{m+1} is independent of $\{f_1, \dots, f_m\}$. Hence there exists a $j \in \{1, \dots, m\}$ such that $f_{m+1} \cdot X_{\text{spt } f_j} / \|f_{m+1} \cdot X_{\text{spt } f_j}\|$ and f_j are linearly independent functions with disjoint supports. By Lemma 2.6 there exists $f_{m+1}^* \geq 0$ and $f_j^* \geq 0$ in F with disjoint supports and $\|f_{m+1}^*\| = \|f_j^*\| = 1$. Hence, the lemma is proved by choosing $f_i^* = f_i$ for $i \neq j$. \square

LEMMA 2.8. If $f \in F$ and $\text{spt } f = \cup_{k=0}^P I_k$, $1 \leq P \leq \infty$, where all I_k are closed intervals, then

(a) there exists $k_0 \geq 0$ such that I_{k_0} contains at least one discontinuity x_j , $j \in \{1, \dots, k\}$ in its interior;

(b) $p < \infty$.

PROOF. (a) Suppose for each I_k , I_k does not contain any discontinuity x_j in its interior. Choose any k_1 with $0 \leq k_1 \leq p$. Then, since I_{k_1} contains no x_j in its interior, τ is strictly monotonic and continuous in I_{k_1} with $|\tau'| > 1$. So,

$m(\tau(I_{k_1})) > m(I_{k_1})$ and $\tau(I_{k_1}) \neq I_{k_1}$. By Lemma 2.4,

$$\bigcup_{k=0}^P \tau(I_k) = \tau\left(\bigcup_{k=0}^P I_k\right) = \bigcup_{k=0}^P I_k \text{ a.e.}$$

Hence, $\tau(I_{k_1}) \subset \bigcup_{k=1}^P I_k$ a.e. On the other hand, $\tau(I_{k_1})$ is an interval and the I_k 's are disjoint. So, if $\tau(I_{k_1}) \cap I_{k_2} \neq \emptyset$ for some $k_2 \neq k_1$ then $\tau(I_{k_1}) = I_{k_2}$ and $m(I_{k_2}) > m(I_{k_1})$. By repeating the same argument, we may construct a sequence of I_{k_j} 's with strictly increasing measures which are bounded below by $m(I_{k_1})$. This is impossible since all I_k 's are disjoint and all of them are contained in finite interval $[0,1]$.

(b) By (a), there exists $k_0 \geq 0$ such that I_{k_0} contains at least one x_j in its interior. Let

$$D = \{k \in \{1, \dots, P\} : I_k \text{ contains } x_j \text{ for some } j\}.$$

Then D is finite, since there are only finitely many points of discontinuities.

Let r be such that I_r is the shortest interval in the collection of intervals:

$$\{\tau(I_k)\}_{k \in D} \cup \{I_k\}_{k \in D}.$$

Notice that $\{\tau(I_k)\}_{k \in D}$ consists of finite pieces of intervals. Let S be the union of those intervals I_k for which $m(I_k) \geq m(I_r)$. Then S contains finitely many intervals and it is obvious that $\tau(S) \subset S$ a.e. since if $m(I_k) \geq m(I_r)$ and I_k does not contain any discontinuity x_j , then $m(\tau(I_k)) > m(I_k) \geq m(I_r)$. By Lemma 2.1,

$$(2.10) \quad \int_{\tau^{-1}(S)-S} f = 0.$$

Suppose there exists interval K with $K \subset \text{spt } f - S$. Choose I_1 to be the largest such interval. Since K contains no discontinuity x_j , we have $\tau(K)$ is an interval of length greater than $m(K)$. Hence $\tau(K) \subset S$ and $K \subset \tau^{-1}(S)$. By (2.10), $\int_K f = 0$. This is a contradiction since K is in the support of f . Therefore $S = \text{spt } f$ and $P < \infty$. \square

PROOF OF THEOREM 1. By Lemmas 2.7 and 2.8, there exist at most n functions in F with disjoint supports and the support of each function contains at least one discontinuity x_j in its interior. Let $H = \{f_1, \dots, f_n\}$ be the set in F , which has the maximum number of functions having the above properties and $\|f_i\| = 1$ for $f_i \in H$. Let $L_i = \text{spt } f_i$. Then (1), (2), (3), (5) of Theorem 1 are satisfied. Suppose for some $i \in \{1, \dots, n\}$, f_i assumes both positive and negative values on subsets of L_i with positive measures. Then, by letting $f_i^1 = f_i \vee 0$ and $f_i^2 = f_i \wedge 0$, we may increase the number of functions in H by 1. This is a contradiction. Hence, by replacing f_i by $-f_i$ if necessary, we may assume $f_i > 0$ on L_i for all $1 < i \leq n$. In order to prove (6), let $g \in F$ satisfy (4), (5) for some $i \in \{1, \dots, n\}$. If $g \neq f_i$ almost everywhere, then

both $(g - f) \vee 0$ and $(g - f) \wedge 0$ are not equal to zero almost everywhere. This is impossible since H has the maximum number of functions in F having disjoint supports. Hence (6) is proved. If $f \in F$ and f is linearly independent of H , then, by Lemma 2.7, we may again construct $n + 1$ functions in F with disjoint support. Hence $f = \sum_{i=1}^n a_i f_i$ with suitable choice of a_i 's. \square

3. Let $\{L_i\}_{i=1}^n$ be the collection of sets stated in Theorem 1. In this section we relate the limit set

$$\Lambda(x) = \bigcap_{N=1}^{\infty} \overline{\{\tau^n(x)\}_{n=N}^{\infty}}$$

to the sets $\{L_i\}$.

THEOREM 2. For almost every $x \in J$, $\Lambda(x) = L_i$ for some $i \in \{1, \dots, n\}$.

PROOF. Let

$$\mathcal{L}_i = \bigcup_{k=0}^{\infty} \tau^{-k}(L_i)$$

where $\tau^{-0}(L_i) \equiv L_i$. We first prove that $\bigcup_{i=1}^n \mathcal{L}_i = J$ almost everywhere. Suppose this is not the case. Then, there exists an interval $[a, b] \subset J - \bigcup_{i=1}^n \mathcal{L}_i$. Let $f = X_{[a, b]}$. By a theorem of Lasota and Yorke [3],

$$\frac{1}{m} \sum_{k=0}^{m-1} P_{\tau^k} f$$

converges to a function $g \neq 0$ in the L^1 norm and g is invariant under τ . Let $L_0 = \text{spt } g$. Without loss of generality we may suppose $g > 0$ in L_0 . Then, $m(L_i \cap L_0) = 0$ for $i = 1, \dots, n$. For, if $A \subset L_i$ for some $i \in \{1, \dots, n\}$, then $\tau^{-k}(A) \subset \mathcal{L}_i$ for all $k = 0, 1, \dots$. Hence,

$$\int_A P_{\tau^k} f = \int_A P_{\tau^k} f = \int_{\tau^{-k}(A)} f = 0$$

for all $k = 0, 1, \dots$. Therefore $\int_A g = 0$, and $m(A \cap L_i) = 0$. This is a contradiction to condition (7) of Theorem 1. Thus, we have $J = \bigcup_{i=1}^n \mathcal{L}_i$ a.e. Now, for almost every x in L_i by applying the Birkhoff Ergodic Theorem, we have

$$\lim_{m \rightarrow \infty} \frac{1}{m} \sum_{k=0}^{m-1} X_{L_i}(\tau^k(x)) = \int_{L_i} f_i = 1.$$

Hence, $\Lambda(x) = \bigcap_{N=1}^{\infty} \overline{\{\tau^n(x)\}_{n=N}^{\infty}} \subset L_i$. Since $\Lambda(x)$ is invariant under τ , therefore $\Lambda(x) = L_i$. For, if not, f_i restricted on $\Lambda(x)$ would be an invariant function which cannot be written as a linear combination of $\{f_1 \dots f_n\}$. \square

REFERENCES

1. A. O. Gelfond, *A common property of number systems*, *Izv. Akad. Nauk SSSR Ser. Mat.* **23** (1959), 809–814. (Russian) MR **22** #702.
2. A. Lasota, *Invariant measures and functional equations*, *Aequationes Math.* **9** (1973), 193–200. MR **48** #6368.
3. A. Lasota and J. A. Yorke, *On the existence of invariant measures for piecewise monotonic transformations*, *Trans. Amer. Math. Soc.* **186** (1973), 481–488. (1974) MR **49** #538.
4. T. Y. Li and J. A. Yorke, *Period three implies chaos*, *Amer. Math. Monthly* **82** (1975), 985–992.
5. W. Parry, *On the β -expansions of real numbers*, *Acta Math. Acad. Sci. Hungar.* **11** (1960), 401–416. MR **26** #288.
6. A. Rényi, *Representation for real numbers and their ergodic properties*, *Acta Math. Acad. Sci. Hungar.* **8** (1957), 477–493. MR **20** #3843.
7. S. M. Ulam, *A collection of mathematical problems*, Interscience Tracts in Pure and Appl. Math., no. 8, Interscience, New York, 1960. MR **22** #10884.
8. A. A. Kosjakin and E. A. Sandler, *Ergodic properties of a certain class of piecewise smooth transformations of a segment*, *Izv. Vysš. Učebn. Zaved. Matematika* **1972**, no. 3(118), 32–40. MR **45** #8802.

DEPARTMENT OF MATHEMATICS, MICHIGAN STATE UNIVERSITY, EAST LANSING, MICHIGAN 48823

INSTITUTE FOR FLUID DYNAMICS AND APPLIED MATHEMATICS, UNIVERSITY OF MARYLAND, COLLEGE PARK, MARYLAND 20742