

COMPLETE CHARACTERIZATION OF FUNCTIONS WHICH ACT, VIA SUPERPOSITION, ON SOBOLEV SPACES

BY

MOSHE MARCUS¹ AND VICTOR J. MIZEL²

ABSTRACT. Given a domain $\Omega \subset R_N$ and a Borel function $h: R_m \rightarrow R$, conditions on h are sought ensuring that for every m -tuple of functions u_i belonging to the first order Sobolev space $W^{1,p}(\Omega)$, the function $h(u_1(\cdot), \dots, u_m(\cdot))$ will belong to a first order Sobolev space $W^{1,r}(\Omega)$, $1 < r < p < \infty$. In this paper conditions are found which are both necessary and sufficient in order that h have the above property. This result is based on a characterization obtained here for those Borel functions $g: R_m \times (R_N)_m \rightarrow R$ satisfying the requirement that for every m -tuple of functions $u_i \in W^{1,p}(\Omega)$ the function $g(u_1(\cdot), \dots, u_m(\cdot), \nabla u_1(\cdot), \dots, \nabla u_m(\cdot))$ belongs to $L^r(\Omega)$. A needed result on the measurability of the set of R_N -Lebesgue points of a function on R_N is presented in an appendix.

1. Introduction. Let Ω be a domain in R_N and let $\mathfrak{N}(\Omega)$ denote the space of real measurable functions in Ω . Given a Borel function $h: R_m \rightarrow R$ we define the superposition mapping $T_h: \mathfrak{N}(\Omega) \rightarrow \mathfrak{N}(\Omega)$ by

$$T_h \mathbf{u} = h \circ \mathbf{u}, \quad \mathbf{u} = (u_1, \dots, u_m) \in \mathfrak{N}(\Omega)^m. \quad (1.1)$$

Let $W_{1,p}(\Omega)$, $1 < p < \infty$, denote the space of functions in $L_p(\Omega)$ whose first order distribution derivatives belong to $L_p(\Omega)$. The standard norm in this space is

$$\|u\|_{W_{1,p}(\Omega)} = \|u\|_{L_p(\Omega)} + \sum_{q=1}^N \|D_q u\|_{L_p(\Omega)}, \quad u \in W_{1,p}(\Omega), \quad (1.2)$$

where $D_1 u, \dots, D_N u$ denote the first order distribution derivatives of u with respect to some coordinate system in R_N .

Suppose that Ω is bounded and satisfies the cone condition. If h is locally Lipschitz and if the first order partial derivatives of h satisfy a.e. in R_m polynomial growth conditions, then T_h maps $W_{1,p}(\Omega)^m$ into $W_{1,r}(\Omega)$, where $1 < r < p < \infty$ and the relation between p and r depends on the order of

Received by the editors April 5, 1978.

AMS (MOS) subject classifications (1970). Primary 46E35, 47H15; Secondary 46J15.

Key words and phrases. Locally Lipschitz function, cone condition, k -dimensional Hausdorff measure, regular Lebesgue point.

¹Research supported by a grant from the Israeli National Academy of Sciences.

²Research supported by National Science Foundation Grants MPS71-03776-A05 and MCS-77-03643.

the polynomials in the growth conditions. Furthermore, this mapping is bounded. This result, in a somewhat different form, was obtained in [5]. For $m = 1$ the result was previously obtained by Stampacchia [9] when h is uniformly Lipschitz and by Serrin in the general case. Serrin's result also shows that $D_i(T_h u)$ can be computed by the standard chain rule. (For details see [4].)

The question whether the above mentioned conditions are also necessary in order that T_h should map $W_{1,p}(\Omega)^m$ into $W_{1,r}(\Omega)$ proved to be considerably more difficult and it remained open for several years. The main purpose of the present paper is to provide an affirmative answer to this question. To achieve this result we first consider a related problem for superposition operators which map Sobolev spaces into Lebesgue spaces.

Let $g: R_m \times R_{mN} \rightarrow R$ be a Borel function and let $S_g: W_{1,1}^{loc}(\Omega)^m \rightarrow \mathcal{N}(\Omega)$ be the mapping defined by

$$S_g \mathbf{u} = g(u_1, \dots, u_m, D_1 u_1, \dots, D_N u_1, \dots, D_1 u_m, \dots, D_N u_m) \quad (1.3)$$

where $\mathbf{u} = (u_1, \dots, u_m)$. We show that if S_g maps $W_{1,p}(\Omega)^m$ into $L_r(\Omega)$, $1 < r < p < \infty$, then g must satisfy certain polynomial growth conditions depending on p and r . For instance, if $p < N$ the condition is

$$|g(\xi, \eta)| < \text{const}(1 + |\xi|^{q/r} + |\eta|^{p/r}), \quad \text{with } q = Np / (N - p),$$

for every η in R_{mN} , for a.e. ξ in R_m . This result leads almost directly to the necessity of the conditions on h when $m = 1$. This is due to the fact that, when $m = 1$, the chain rule for $D_i(T_h u)$ is available. When $m > 1$ the chain rule for $D_i(T_h \mathbf{u})$ does not hold in general. In this case the necessity of the conditions on h is established by combining the result already proved for $m = 1$ with the result concerning the mapping S_g for $m > 1$.

Results similar to those described above hold also in the case where Ω is unbounded. In this case the conditions are stronger than in the case of bounded domains.

The plan of the paper is as follows. In §2 we state the main results in the case of bounded domains. §3 and §4 (together with an appendix) are devoted to the development of auxiliary results. The main result for the mapping S_g is proved in §5 and for the mapping T_h in §6 and §7. The parallel results in the case of unbounded domains are presented in §8. Finally in §9 and §10 we discuss some additional results and open questions related to the mappings S_g and T_h .

2. Statement of main results. In this section Ω denotes a domain in R_N satisfying the cone condition. The generic points in R_N , R_m and R_{mN} will be denoted by $x = (x_1, \dots, x_n)$, $\xi = (\xi_1, \dots, \xi_m)$ and $\eta = (\eta_{1,1}, \dots, \eta_{1,N}, \dots, \eta_{m,1}, \dots, \eta_{m,N})$ respectively. The norm $|\cdot|$ of such a point will always mean Euclidean norm.

The first result concerns superposition mappings of the type (1.1).

THEOREM 1. *Suppose that Ω is bounded. Let $h: R_m \rightarrow R$ be a Borel function and let p, r be two numbers such that $1 < r < p < N$. Then T_h maps $W_{1,p}(\Omega)^m$ into $W_{1,r}(\Omega)$ if and only if the following conditions hold:*

- (i) *h is locally Lipschitz in R_m .*
- (ii) *The first order partial derivatives of h satisfy the inequality*

$$\left| \frac{\partial h}{\partial \xi_i}(\xi) \right| < a_0(1 + |\xi|^v) \quad \text{a.e. in } R_m, i = 1, \dots, m, \quad (2.1)$$

where a_0 is a constant and $v = N(p - r)/(r(N - p))$.

If $N < p$ (or $N = 1$ and $1 < p$) and $1 < r < p$ then T_h maps $W_{1,p}(\Omega)^m$ into $W_{1,r}(\Omega)$ if and only if condition (i) holds.

In either case, if T_h maps $W_{1,p}(\Omega)^m$ into $W_{1,r}(\Omega)$ then the mapping is bounded and the following inequalities hold:

$$\begin{aligned} \|T_h \mathbf{u}\|_{W_{1,r}(\Omega)} &< b_0 \left(1 + \sum_{i=1}^m \|u_i\|_{W_{1,p}(\Omega)}^{p+1} \right), & \text{if } p < N, \\ \|T_h \mathbf{u}\|_{W_{1,r}(\Omega)} &< b(M) \left(1 + \sum_{i=1}^m \|u_i\|_{W_{1,p}(\Omega)} \right), & \text{if } p > N \text{ (or } N = 1 \text{ and } p > 1) \\ & & \text{and } \sum_{i=1}^m \|u_i\|_{W_{1,p}(\Omega)} < M. \end{aligned} \quad (2.2)$$

The constants b_0 and $b(M)$ depend on $m, a_0, h(0)$ and Ω . (Of course, $b(M)$ depends also on M .)

The second result concerns superposition mappings of the type (1.3).

THEOREM 2. *Suppose that Ω is bounded. Let $g: R_m \times R_{mN} \rightarrow R$ be a Borel function and let p, r be two numbers such that $1 < r < p < \infty$. Suppose that S_g maps $W_{1,p}(\Omega)^m$ into $L_r(\Omega)$. Then g satisfies the following conditions.*

If $p < N$, there exists a constant c_0 such that

$$|g(\xi, \eta)| < c_0(1 + |\xi|^{q/r} + |\eta|^{p/r}), \quad \text{with } q = Np/(N - p), \quad (2.3)$$

for every η in R_{mN} and a.e. ξ in R_m . The exceptional null set in R_m may vary with η .

If $p > N$ (or $N = 1$ and $p > 1$), then for every $M > 0$ there exists a constant $c(M)$ such that

$$|g(\xi, \eta)| < c(M)(1 + |\eta|^{p/r}) \quad (2.4)$$

for every η in R_{mN} and a.e. ξ in R_m such that $|\xi| < M$. Again, the exceptional null set in R_m may vary with η .

If a function g satisfies (2.3) or (2.4) *everywhere* in $R_m \times R_{mN}$, then (in view of the Sobolev imbedding theorem) S_g maps $W_{1,p}(\Omega)^m$ into $L_r(\Omega)$ for the appropriate values of p and r . Therefore if g is continuous the conditions described in Theorem 2 are necessary and sufficient for S_g to map $W_{1,p}(\Omega)^m$ into $L_r(\Omega)$. However the assumption of continuity can be replaced by various weaker assumptions. Results of this type will be discussed in §9.

Results similar to those stated above hold also in the case where Ω is unbounded. These will be presented in §8.

3. Construction of a class of elementary functions. The proof of Theorems 1 and 2 calls for the construction of functions in $W_{1,p}(\Omega)$ according to a given set of specifications. One of the requirements is that the gradient of the function should attain a prescribed sequence of values on sets of prescribed measure and at the same time the values of the function should lie within a given range in each of these sets. In the construction of these functions we use certain piecewise linear functions which we describe below.

LEMMA 1. *Let $z \in R_N$, $z \neq 0$, and let σ be a positive number. Then there exists a piecewise linear function w in the cube $E_\sigma = \{x \in R_N: |x_i| < \sigma/2, i = 1, \dots, N\}$ such that,*

(i) $w = 0$ on ∂E_σ ,

(ii) $|\nabla w|$ is either zero or $|z|$ in E_σ ,

(iii) $\nabla w = z$ in a cube F contained in E_σ , whose sides are parallel to the axes and which satisfies

$$\mu_N(F)/\sigma^N \geq a_N > 0. \quad (3.1)$$

Here μ_N denotes N -dimensional Lebesgue measure and a_N is a constant which depends only on N .

PROOF. Let $b = \sigma/2\sqrt{N}$ and let T be a cube in R_N of side length b such that one of its vertices (say P) lies in the half space $z \cdot x > 0$, the N vertices nearest to P lie in the hyperplane $z \cdot x = 0$ and one of these vertices is at the origin. Let $T' = T \cap \{z \cdot x > 0\}$ and define a function w_0 in T as follows:

$$w(x) = z \cdot x \quad \text{in } T', \quad w(x) = 0 \quad \text{in } T \setminus T'.$$

Note that $\mu_N(T')/\mu_N(T)$ depends only on N .

Let P' be that vertex of T for which $\overline{PP'}$ is one of the main diagonals of T . Let U be a closed cube of side $2b$ such that $U \supset T$, the sides of U are parallel to those of T and P' is a vertex of U . We extend w_0 to U by N successive reflexions. In order to describe these reflexions we use a set of coordinates $(\bar{x}_1, \dots, \bar{x}_N)$ whose origin is at P' and with respect to which $T = \{\bar{x}: 0 < \bar{x}_i < b, i = 1, \dots, N\}$. Let T^* be the image of T by reflexion with respect to $\bar{x}_1 = b$ and set $T_1 = T \cup T^*$. We extend w_0 to T_1 by reflexion with

respect to $\bar{x}_1 = b$. Let k be an integer, $1 \leq k \leq N$, and set $T_k = \{\bar{x}: 0 \leq \bar{x}_i \leq 2b \text{ for } i = 1, \dots, k \text{ and } 0 \leq \bar{x}_i \leq b \text{ for } i = k + 1, \dots, N\}$.

If $k < N$, denote by T_k^* the image of T_k by reflexion with respect to $\bar{x}_{k+1} = b$. Then $T_{k+1} = T_k \cup T_k^*$. Assuming that w_0 is defined in T_k we extend it to T_{k+1} by reflexion with respect to $\bar{x}_{k+1} = b$. Thus after N steps we obtain an extension of w_0 to the entire cube U .

Next let V be a cube whose sides are parallel to the axes (x_1, \dots, x_N) , whose center is at the center of U and whose side length is $2\sqrt{N}b$. We extend w_0 to V by setting $w_0 = 0$ in $V \setminus U$. Let B be the largest ball contained in T' and let F' be the cube inscribed in B whose sides are parallel to the axes. Then $\mu_N(F')/b^N$ depends only on N . Finally, let x^0 be the center of V and set $w(x) = w_0(x + x^0)$ for every x in $E_\sigma = -x_0 + V$. Clearly the function w satisfies all the assumptions of the lemma. In particular (iii) holds with respect to the cube $F = -x_0 + F'$.

We shall often use a variant of Lemma 1 which, for purposes of reference, we shall state separately.

LEMMA 2. Let $z^1, \dots, z^m \in R_N, z^i \neq 0 (i = 1, \dots, m)$, and let σ be a positive number. Then there exist piecewise linear functions w_1, \dots, w_m in E_σ such that w_i satisfies the conditions of Lemma 1 with $z = z^i$ but with F independent of i such that

$$\mu_N(F)/\sigma^N \geq a_N^m. \tag{3.2}$$

PROOF. Let w_1 be as in Lemma 1 with $z = z^1$. Let F_1 be the cube mentioned in (iii) where $\nabla w_1 = z^1$. Let x^1 be the center of F_1 and σ_1 its side length. Denote by \tilde{w}_2 the function described in Lemma 1 when $\sigma = \sigma_1$ and $z = z^2$, and define w_2 in E_σ as follows:

$$w_2(x) = \tilde{w}_2(x - x^1) \quad \forall x \in F_1, \quad w_2(x) = 0 \quad \forall x \in E_\sigma \setminus F_1.$$

Proceeding in this manner we obtain functions w_1, \dots, w_m and cubes F_1, \dots, F_m such that w_i satisfies conditions (i)–(iii) of Lemma 1 with z and F replaced by z^i and F_i . Moreover, $F_1 \supset F_2 \supset \dots \supset F_m$, and $\mu_N(F_1)/\sigma^N > a_N$ and $\mu_N(F_{i+1})/\mu_N(F_i) \geq a_N$ for $i = 1, \dots, m - 1$. Thus the lemma holds with $F = F_m$.

REMARK. In the set F the functions $w_i, i = 1, \dots, m$, are of the form $z^i \cdot (x - y^i)$ where $|y^i| < 2\sqrt{N} \sigma$. Thus

$$\begin{cases} w_i(x) = \sum_{n=1}^N z_n^i x_n + b_i & \forall x \in F, \text{ and} \\ |b_i| \leq 2\sqrt{N} \sigma |z^i|. \end{cases} \tag{3.3}$$

4. Auxiliary results. Let $g: R_m \times R_{mN} \rightarrow [0, \infty)$ be a Borel function. As before, we denote by S_g the mapping from $W_{1,p}(\Omega)^m$ into $\mathcal{N}(\Omega)$ given by

$$(u_1, \dots, u_m) \rightarrow g(u_1, \dots, u_m, \nabla u_1, \dots, \nabla u_m).$$

Here, as before, Ω is a domain in R_N .

With every point $\eta \in R_{mN}$, $\eta = (\eta_{1,1}, \dots, \eta_{1,N}, \dots, \eta_{m,1}, \dots, \eta_{m,N})$, we associate the $m \times N$ matrix

$$A(\eta) = (\eta_{i,j})_{i=1, \dots, m; j=1, \dots, N}.$$

The rank of this matrix will be denoted by $k(\eta)$. The image of R_N by the linear transformation $x \rightarrow A(\eta)x$ will be denoted by M_η . Thus M_η is a subspace of R_m of dimension $k(\eta)$.

Suppose that S_g maps $W_{1,p}(\Omega)^m$ into $L_1^{loc}(\Omega)$. Then, for every η in R_{mN} and every ξ^0 in R_m , the function $f(\cdot; \xi^0, \eta) : R_N \rightarrow R$ given by

$$f(x; \xi^0, \eta) = g(A(\eta)x + \xi^0, \eta) \quad \forall x \in R_N \tag{4.1}$$

is locally integrable in R_N . Using this fact we shall show that for every η in R_{mN} , the function $\xi \rightarrow g(\xi, \eta)$ is locally integrable in $\xi^0 + M_\eta$ (with respect to $k(\eta)$ -dimensional Hausdorff measure) for every fixed ξ^0 in R_m .

Let us assume (as we may) that the first $k(\eta)$ columns of $A(\eta)$ are linearly independent. We assume of course that $\eta \neq 0$ so that $k(\eta) > 0$. Set

$$B(\eta) = (\eta_{i,j})_{i=1, \dots, m; j=1, \dots, k}$$

where $k = k(\eta)$. If $k < N$ and $\zeta \in R_{N-k}$ set

$$\pi_k(\zeta) = \{x \in R_N : x_{k+1} = \zeta_1, \dots, x_N = \zeta_{N-k}\}.$$

If $k = N$ set $\pi_k = \pi_k(0) = R_N$. Clearly, the transformation $x \rightarrow A(\eta)x$ maps $\pi_k(\zeta)$ onto M_η in a (1-1) manner.

Now, let ξ^0 be a point in R_m and denote by $U_0(\cdot) = U(\cdot; \xi^0, \eta)$ the mapping given by $U_0(x) = A(\eta)x + \xi^0$, $x \in R_N$. Let ζ be a point in R_{N-k} such that $f(\cdot; \xi^0, \eta)$ restricted to $\pi_k(\zeta)$ is locally integrable with respect to μ_k (= k -dimensional Lebesgue measure). Let E be a μ_k -measurable set in $\pi_k(\zeta)$. Then by the change of variables formula for integration (see e.g. [1, p. 244]) we have

$$\int_{U_0(E)} g(\xi, \eta) d\mathcal{H}_k(\xi) = \int_E g(A(\eta)x + \xi^0, \eta) J_k(B(\eta)) d\mu_k(x), \tag{4.2}$$

where \mathcal{H}_k denotes k -dimensional Hausdorff measure and $J_k(B(\eta))$ is the square root of the sum of the squares of the $k \times k$ minors of $B(\eta)$. By our choice of ζ the right-hand side of (4.2) is finite for every bounded E . Therefore $g(\cdot, \eta)$ is locally integrable with respect to \mathcal{H}_k in $\xi^0 + M_\eta$.

From the last assertion we deduce (again using (4.2)) that $f(\cdot; \xi^0, \eta)$ restricted to $\pi_k(\zeta)$ is locally integrable for every ζ in R_{N-k} .

Given $\eta \neq 0$ in R_{mN} we denote by $P(\eta)$ the set of points $\{\xi\}$ in R_m such that ξ is a (regular) Lebesgue point of the function $g(\cdot, \eta)$ restricted to $\xi + M_\eta$, with respect to the measure $\mathcal{H}_{k(\eta)}$. Some properties of $P(\eta)$ will be needed in the next section. These properties are the subject of the following two lemmas.

LEMMA 3. $P(\eta)$ is a set of total measure in R_m , i.e. its complement is a Lebesgue null set.

LEMMA 4. Let $\xi' \in P(\eta)$. Let ξ^0 and x' be points in R_m and R_N respectively such that

$$\xi' = A(\eta)x' + \xi^0. \tag{4.3}$$

Then x' is a Lebesgue point of $f(\cdot; \xi^0, \eta)$ with respect to μ_N .

We defer the proof of Lemma 3 to the appendix where it will be presented in a slightly more general form. Here we turn to

PROOF OF LEMMA 4. As before we may assume that $\text{rank } A(\eta) = \text{rank } B(\eta) = k$. In this proof we shall use the notations previously introduced in this section.

First we note that by (4.2), if E is a μ_k -measurable subset of $\pi_k(\zeta)$ (for some ζ in R_{N-k}) then

$$\mathcal{H}_k(U_0(E)) = J_k(B(\eta))\mu_k(E). \tag{4.4}$$

Let c be a positive constant. We denote by $\mathcal{E}_c(x')$ the family of cubes in R_N whose sides are parallel to the axes, such that

$$E \in \mathcal{E}_c(x') \Rightarrow \exists r > 0 \text{ such that } E \subset \{x \in R_n : |x - x'| < r\} \\ \text{and } \mu_N(E)/r^N > c.$$

Further we denote by $\mathcal{F}_c(\xi')$ the family of \mathcal{H}_k -measurable subsets of $\xi' + M_\eta$ such that

$$F \in \mathcal{F}_c(\xi') \Rightarrow \exists r > 0 \text{ such that } F \subset \{\xi \in \xi' + M_\eta : |\xi - \xi'| < r\} \\ \text{and } \mathcal{H}_k(F)/r^k > c.$$

Now let us consider the case $k = N$. If $E \in \mathcal{E}_c(x')$ then $E^* = U_0(E) \in \mathcal{F}_c(\xi')$, where c' is a positive number depending only on c and η . Indeed, if E is contained in a ball of radius r around x' then (in view of (4.3)) the set E^* is contained in a ball of radius $|\eta|r$ around ξ' . From this and (4.4) it follows that $E^* \in \mathcal{F}_c(\xi')$ if we choose, for instance, $c' = J_N(B(\eta))c|\eta|^{-N}$. Furthermore, by (4.2) and (4.4),

$$\frac{1}{\mu_N(E)} \int_E f(x; \xi^0, \eta) d\mu_N = \frac{1}{\mathcal{H}_N(E^*)} \int_{E^*} g(\xi, \eta) d\mathcal{H}_N(\xi). \tag{4.5}$$

Since $\xi' \in P(\eta)$, the right-hand side of (4.5) tends to $g(\xi', \eta)$ when $\mathcal{I}_{\mathcal{C}_N}(E^*) \rightarrow 0$. This implies the assertion of the lemma when $k = N$.

Next, assume that $k < N$. If E is a set in R_N we denote by $E_k(\zeta)$ the set $E \cap \pi_k(\zeta)$, $\zeta \in R_{N-k}$, and by E_{N-k} the set

$$\{\zeta \in R_{N-k}: (x_1, \dots, x_k, \zeta_1, \dots, \zeta_{N-k}) \in E \text{ for some } x_1, \dots, x_k\}.$$

If $E \in \mathcal{E}_c(x')$ then $U_0(E_k(\zeta)) \in \mathcal{F}_c(\xi')$ for every ζ in E_{N-k} , where $c' = J_k(B(\eta))c^{k/N}|\eta|^{-k}$. This can be verified in the same way as before. Now let E be a cube in $\mathcal{E}_c(x')$ of side length δ . Then, by (4.2) and (4.4),

$$\begin{aligned} &\delta^{-N} \int_E f(x; \xi^0, \eta) d\mu_N \\ &= \delta^{-N} \int_{E_{N-k}} \left(\int_{E_k(\zeta)} f(x; \xi^0, \eta) d\mu_k(x_1, \dots, x_k) \right) d\mu_{N-k}(\zeta) \\ &= \delta^{-N+k} \int_{E_{N-k}} \left(\mathcal{I}_{\mathcal{C}_k}(F(\zeta))^{-1} \int_{F(\zeta)} g(\xi, \eta) d\mathcal{I}_{\mathcal{C}_k} \right) d\mu_{N-k}(\zeta) \end{aligned} \tag{4.6}$$

where $F(\zeta) = U_0(E_k(\zeta))$. Since $\xi' \in P(\eta)$, $F(\zeta) \in \mathcal{F}_c(\xi')$ and $\mathcal{I}_{\mathcal{C}_k}(F(\zeta)) = \text{const } \delta^k$ for every ζ in E_{N-k} it follows that

$$\mathcal{I}_{\mathcal{C}_k}(F(\zeta))^{-1} \int_{F(\zeta)} g(\xi, \eta) d\mathcal{I}_{\mathcal{C}_k} \rightarrow g(\xi', \eta)$$

when $\delta \rightarrow 0$ uniformly with respect to ζ in E_{N-k} . Hence, by (4.6),

$$\mu_N(E)^{-1} \int_E f(x; \xi^0, \eta) d\mu_N(x) \rightarrow f(x'; \xi^0, \eta)$$

when $E \in \mathcal{E}_c(x')$ and $\mu_N(E) \rightarrow 0$. This completes the proof of the lemma.

5. Proof of Theorem 2. First we note that it is sufficient to prove the theorem in the case where $g \geq 0$ and $r = 1$. The result in the general case will follow immediately from this by considering the function $\tilde{g} = |g|^{1/r}$. Secondly, we may assume that Ω is the unit cube $\{x \in R_N: 0 < x_i < 1\}$. Indeed, if S_g maps $W_{1,p}(\Omega)$ into $L_r(\Omega)$ for some domain Ω in R_N , then

(i) S_g maps $W_{1,p}(\Omega')^m \rightarrow L_r(\Omega')$ for any bounded domain Ω' contained in Ω , whose boundary is Lipschitz;

(ii) S_g maps $W_{1,p}(\tilde{\Omega})^m \rightarrow L_r(\tilde{\Omega})$, where $\tilde{\Omega}$ is the image of Ω by a translation.

The first of these statements follows from the fact that any function in $W_{1,p}(\Omega')$ can be extended to a function in $W_{1,p}(R_N)$. The second statement is obvious.

Therefore in this section we shall consider Theorem 2 when $g \geq 0$, $r = 1$ and Ω is the unit cube mentioned before.

Let $P(\eta)$ be defined as in §4 and let M be a positive number. Set

$$\begin{cases} Q = \{(\xi, \eta) \in R_m \times R_{mN}: \xi \in P(\eta)\}, \\ Q_M = \{(\xi, \eta) \in Q: |\xi| < M\}. \end{cases} \tag{5.1}$$

Further denote by G_p the real function on $R_m \times R_{mN}$ given by

$$G_p(\xi, \eta) = 1 + |\xi|^q + |\eta|^p, \tag{5.2}$$

where $q = Np/(N - p)$ if $p < N$ and $q = 0$ if $p > N$.

We shall prove the following assertions:

(5.3)₁ If $1 < p < N$ condition (2.3) holds everywhere in Q .

(5.3)₂ If $p > N$ or $p \geq 1$ and $N = 1$ condition (2.4) holds everywhere in Q_M .

Since $P(\eta)$ is a set of total measure in R_m (Lemma 3) these assertions imply the statement of the theorem.

Suppose that assertion (5.3)₁ (respectively (5.3)₂) does not hold. Then there exists a sequence $\{(\xi^n, \eta^n)\}$ in Q (resp. Q_M) such that

$$g(\xi^n, \eta^n)/G_p(\xi^n, \eta^n) = \alpha_n \rightarrow \infty \quad \text{when } n \rightarrow \infty. \tag{5.4}$$

We wish to show that this leads to a contradiction. To this purpose we shall consider several cases, in each of which, using (5.4), we shall construct functions w_1, \dots, w_m in $W_{1,p}(\Omega)$ such that $S_g(w_1, \dots, w_m)$ is not in $L_1(\Omega)$.

Case I. Suppose that $1 < p < N$ and that there exists a positive number β such that

$$\beta < |\xi^n|^q / (1 + |\eta^n|^p), \quad \text{where } q = Np / (N - p). \tag{5.5}$$

Let $\theta_n = 1/\alpha_n$. By considering a subsequence if necessary, we may assume that

$$\sum_{n=1}^{\infty} \theta_n^{1/N} < \frac{1}{2} \beta^{1/N} \quad \text{and} \quad \sum_{n=1}^{\infty} \theta_n^{(N-p)/N} < \infty. \tag{5.6}$$

Set

$$\delta_n^N = \theta_n |\xi^n|^{-q}, \quad n = 1, 2, \dots \tag{5.7}$$

Note that, in view of (5.5) and (5.6), $\sum_1^{\infty} \delta_n < \frac{1}{2}$.

By assumption $\xi^n \in P(\eta^n)$. Therefore, by Lemma 4, $x = 0$ is a Lebesgue point of the function $f(\cdot; \xi^n, \eta^n)$ defined in (4.1). Hence, for every $c > 0$ there exists a positive ρ_n such that

$$\mu_N(B)^{-1} \int_B f(\cdot; \xi^n, \eta^n) d\mu_N \geq g(\xi^n, \eta^n)/2, \tag{5.8}$$

for every cube B in R_N such that $\mu_N(B) \leq \rho_n^N$ and the distance between B and the origin is not larger than $c\mu_N(B)^{1/N}$. In the sequel ρ_n will correspond to $c = \sqrt{N} a_N^{-m/N}$, where a_N is the constant appearing in (3.1) and (3.2).

Let σ_n be the largest number of the form $\delta_n/2j$, $j = 1, 2, \dots$, such that $\sigma_n \leq \rho_n$. Let $\{v_{i,n}\}_{n=1}^m$ be a set of functions in $E_n^* = \{x: |x_i| < \sigma_n/2, i = 1, \dots, N\}$, constructed as in Lemma 2 with $\sigma = \sigma_n$ and $z^i = (\eta_{i,1}^n, \dots, \eta_{i,N}^n)$,

$i = 1, \dots, m$. The cube F mentioned in Lemma 2 will be denoted here by F_n^* . By (3.3) we have

$$v_{i,n}(x) = \sum_{s=1}^N \eta_{i,s}^n x_s + b_{i,n}, \quad x \in F_n^*,$$

and

$$|b_{i,n}| < 2\sqrt{N} \sigma_n |(\eta_{i,1}, \dots, \eta_{i,N})|.$$

Let

$$\begin{cases} \gamma_{i,n} = \xi_i^n - b_{i,n}, & i = 1, \dots, m, \\ \tilde{v}_{i,n}(x) = v_{i,n}(x) + \gamma_{i,n} & \forall x \in E_n^*, i = 1, \dots, m. \end{cases} \quad (5.9)$$

Then,

$$\begin{cases} \tilde{v}_{i,n}(x) = \sum_{s=1}^N \eta_{i,s}^n x_s + \xi_i^n & \forall x \in F_n^*, i = 1, \dots, m, \\ \tilde{v}_{i,n}(x) = \gamma_{i,n} & \forall x \in \partial E_n^*, \text{ and} \\ |\gamma_{i,n}| < |\xi_i^n| + 2\sqrt{N} \sigma_n |(\eta_{i,1}^n, \dots, \eta_{i,N}^n)|. \end{cases} \quad (5.10)$$

Next let $E_n = \{x: |x_i| < \delta_n/2, i = 1, \dots, N\}$ and let $\{E_{n,\nu}\}_{\nu=1}^k$ be a family of cubes whose interiors are disjoint such that each cube $E_{n,\nu}$ is a translate of E_n^* and $\cup_{\nu=1}^k E_{n,\nu} = E_n$. (These conditions uniquely determine the family $\{E_{n,\nu}\}$.) Denote by x^ν the center of the cube $E_{n,\nu}$. Then we define functions $u_{i,n}, i = 1, \dots, m$, in E_n as follows

$$u_{i,n}(x) = \tilde{v}_{i,n}(x - x^\nu), \quad \forall x \in E_{n,\nu}, \nu = 1, \dots, k. \quad (5.11)$$

Let

$$F_n = \bigcup_{\nu=1}^k (x^\nu + F_n^*).$$

Then by (5.8), (5.10), (5.11) and (3.2) we have

$$\begin{aligned} \int_{F_n} S_g(u_{1,n}, \dots, u_{m,n}) d\mu_N &= k \int_{F_n^*} S_g(\tilde{v}_{1,n}, \dots, \tilde{v}_{m,n}) d\mu_N \\ &> k \mu_N(F_n^*) g(\xi^n, \eta^n)/2 \\ &> k \mu_N(E_n^*) a_N^m g(\xi^n, \eta^n)/2 = b_1 \delta_n^N g(\xi^n, \eta^n) \end{aligned} \quad (5.12)$$

where $b_1 = a_N^m/2$. (Note that the distance between F_n^* and the origin is not larger than $\sqrt{N} (\mu_N(F_n^*)/a_N^m)^{1/N}$.)

Finally, set $D_n = \{x: |x_i| < \delta_n, i = 1, \dots, N\}$ and

$$D_{n,\nu} = \{x: \delta_n/2 < |x_\nu| < \delta_n, |x_\nu| \geq |x_k|, k = 1, \dots, N\}, \quad \nu = 1, \dots, N.$$

Thus, $D_n = (\cup_{\nu=1}^N D_{n,\nu}) \cup E_n$. In D_n we define m functions $w_{1,n}, \dots, w_{m,n}$ as follows:

$$\begin{cases} w_{i,n}(x) = u_{i,n}(x) & \forall x \in E_n, \\ w_{i,n}(x) = 2(\gamma_{i,n}/\delta_n) \cdot (\delta_n - |x_\nu|) & \forall x \in D_{n,\nu}, \nu = 1, \dots, N. \end{cases} \quad (5.13)$$

The functions $w_{1,n}, \dots, w_{m,n}$ are continuous, piecewise linear in D_n , they vanish on ∂D_n and their gradients satisfy the inequalities

$$\begin{cases} |\nabla w_{i,n}| < |\eta^n| & \text{in } E_n; \\ |\nabla w_{i,n}| < 2|\gamma_{i,n}|\delta_n^{-1} < 2\sqrt{N}(|\xi^n|\delta_n^{-1} + |\eta^n|) & \text{in } D_n \setminus E_n, i = 1, \dots, m. \end{cases} \quad (5.14)$$

For the second inequality we used (5.10).

Let y^n be the point $(\tau_n, \frac{1}{2}, \dots, \frac{1}{2}) \in R_N$, where $\tau_n = 2 \sum_1^{n-1} \delta_k + \delta_n$, $n = 1, 2, \dots$, and set $\tilde{D}_n = y^n + D_n$. Clearly, the interiors of the cubes \tilde{D}_n are disjoint and (in view of the fact that $2 \sum_1^\infty \delta_n < 1$) $\tilde{D}_n \subset \bar{\Omega}$, $n = 1, 2, \dots$. Now, we define m functions w_1, \dots, w_m in $\bar{\Omega}$ as follows:

$$\begin{cases} w_i(x) = w_{i,n}(x - y^n) & \text{in } \tilde{D}_n, n = 1, 2, \dots; \\ w_i(x) = 0 & \text{in } \bar{\Omega} \setminus \bigcup_1^\infty \tilde{D}_n. \end{cases} \quad (5.15)$$

We claim that

$$w_i \in W_{1,p}(\Omega), \quad i = 1, \dots, m, \quad (5.16)_1$$

and

$$\int_{\bar{\Omega}} S_g(w_1, \dots, w_m) d\mu_N = \infty. \quad (5.16)_2$$

By construction the functions w_1, \dots, w_m are continuous in $\bar{\Omega}$, except at the point $(2 \sum_1^\infty \delta_n, \frac{1}{2}, \dots, \frac{1}{2})$, and vanish on $\partial \Omega$. Furthermore they are Lipschitz and, in fact, piecewise linear in every subset of $\bar{\Omega}$ which is bounded away from the point mentioned above. Therefore, in order to prove (5.16)₁ it is sufficient to show that $\nabla w_i \in L_p(\Omega)^N$, $i = 1, \dots, m$. By (5.14) and (5.15) we have

$$\begin{aligned} \int_{\tilde{D}_n} |\nabla w_i|^p d\mu_N &= \int_{D_n \setminus E_n} |\nabla w_{i,n}|^p d\mu_N + \int_{E_n} |\nabla w_{i,n}|^p d\mu_N \\ &< (2^N - 1)\delta_n^N (2\sqrt{N})^p (|\xi^n|\delta_n^{-1} + |\eta^n|)^p + \delta_n^N |\eta^n|^p \\ &< \text{const}(\delta_n^{N-p} |\xi^n|^p + \delta_n^N |\eta^n|^p), \quad i = 1, \dots, m; n = 1, 2, \dots \end{aligned} \quad (5.17)$$

By (5.5) and (5.7),

$$\delta_n^N |\eta^n|^p < \delta_n^N |\xi^n|^q / \beta = \theta_n / \beta \quad \text{and} \quad \delta_n^{N-p} |\xi^n|^p = \theta_n^{(N-p)/N}.$$

Therefore, by (5.6) and (5.17),

$$\int_{\Omega} |\nabla w_i|^p d\mu_N < \text{const} \left(\sum_{n=1}^{\infty} \theta_n^{(N-p)/N} + \beta^{-1} \sum_1^{\infty} \theta_n \right) < \infty \quad (i = 1, \dots, m).$$

This proves (5.16)₁.

Now, by (5.15), (5.13) and (5.12) we have

$$\begin{aligned} \int_{\bar{D}_n} S_g(w_1, \dots, w_m) d\mu_N &= \int_{D_n} S_g(w_{1,n}, \dots, w_{m,n}) d\mu_N \\ &> \int_{F_n} S_g(u_{1,n}, \dots, u_{m,n}) d\mu_N > b_1 g(\xi^n, \eta^n) \delta_n^N. \end{aligned} \quad (5.18)$$

Hence, using (5.4) and (5.7), we obtain

$$\begin{aligned} \int_{\Omega} S_g(w_1, \dots, w_m) d\mu_N &> b_1 \sum_{n=1}^{\infty} \delta_n^N g(\xi^n, \eta^n) = b_1 \sum_1^{\infty} \delta_n^N \alpha_n G_p(\xi^n, \eta^n) \\ &= b_1 \sum_1^{\infty} \delta_n^N \theta_n^{-1} (1 + |\xi^n|^q + |\eta^n|^p) > b_1 \sum_1^{\infty} \delta_n^N |\xi^n|^q \theta_n^{-1} = +\infty. \end{aligned}$$

This proves (5.16)₂. Therefore we have proved that (5.4) and (5.5) lead to a contradiction.

Case II. Suppose that $1 < p < N$ and that

$$|\xi^n|^q / (1 + |\eta^n|^p) < \gamma \quad \text{and} \quad \beta^{1/p} < |\eta^n| \quad (5.19)$$

for some positive numbers β, γ . Let $\theta_n = 1/\alpha_n$. By considering a subsequence if necessary, we may assume that (5.6) holds. Set

$$\delta_n^N = \theta_n |\eta^n|^{-p}, \quad n = 1, 2, \dots \quad (5.20)$$

In view of (5.19) and (5.6) we have $\sum_1^{\infty} \delta_n < \frac{1}{2}$.

Let w_1, \dots, w_m be defined exactly as in Case I. As before we shall show that (5.16) holds.

By (5.19) and (5.20)

$$\begin{aligned} \delta_n^{N-p} |\xi^n|^p &= (\delta_n^N |\xi^n|^q)^{(N-p)/N} < \text{const} (\delta_n^N (1 + |\eta^n|^p))^{(N-p)/N} \\ &= \text{const} (\delta_n^N + \theta_n)^{(N-p)/N} < \text{const} \theta_n^{(N-p)/N}, \quad n = 1, 2, \dots \end{aligned}$$

From this inequality together with (5.17) and (5.20) we obtain

$$\int_{\bar{D}_n} |\nabla w_i|^p d\mu_N < \text{const} (\theta_n^{(N-p)/N} + \theta_n) \quad (i = 1, \dots, m; n = 1, 2, \dots).$$

Hence, by (5.6)

$$\int_{\Omega} |\nabla w_i|^p d\mu_N < \text{const} \left(\sum_1^{\infty} \theta_n^{(N-p)/N} + \sum_1^{\infty} \theta_n \right) < \infty, \quad i = 1, \dots, m.$$

This implies (5.16)₁.

Next, by (5.18) and (5.20),

$$\begin{aligned} \int_{\Omega} S_g(w_1, \dots, w_m) d\mu_N &> b_1 \sum_{n=1}^{\infty} \delta_n^N g(\xi^n, \eta^n) = b_1 \sum_{n=1}^{\infty} \delta_n^N \alpha_n G_p(\xi^n, \eta^n) \\ &\geq b_1 \sum_{n=1}^{\infty} \delta_n^N \alpha_n |\eta^n|^p = b_1 \sum_1^{\infty} \alpha_n \theta_n = \infty. \end{aligned}$$

Thus we have shown that (5.16) holds. Therefore (5.4) and (5.19) lead to a contradiction.

Case III. Suppose that $1 < p < \infty$ and that

$$|\xi^n| + |\eta^n| < C, \quad n = 1, 2, \dots, \tag{5.21}$$

for some constant C .

By considering a subsequence if necessary, we may assume that

$$\xi^n \rightarrow \xi^0 \quad \text{and} \quad \sum_1^{\infty} |\xi^n - \xi^0| < \infty, \tag{5.22}$$

and also that

$$\sum_1^{\infty} \alpha_n^{-1} < \frac{1}{2}. \tag{5.23}$$

Let ρ_n, σ_n and $\tilde{v}_{i,n}$ be as in Case I with $\delta_n = \alpha_n^{-1}$. Set

$$\begin{aligned} J_n &= \{x: 0 < x_1 < \delta_n, 0 < x_i < 1, i = 2, \dots, N\}, \\ \tilde{J}_n &= \{x: 0 < x_1 < \delta_n, 0 < x_i < \kappa_n, i = 2, \dots, N\}, \end{aligned}$$

where κ_n is the largest integral multiple of σ_n such that $\kappa_n < 1$. Note that, since $\sigma_n < \frac{1}{2}$, we have $\kappa_n > \frac{1}{2}$.

Let $\{J_{n,\nu}\}_{\nu=1}^s$ be a family of cubes whose interiors are disjoint such that the union of the cubes equals \tilde{J}_n and each cube $J_{n,\nu}$ is a translate of E_n^* . Denote by x^ν the center of the cube $J_{n,\nu}$. Now we define in J_n functions $u_{1,n}, \dots, u_{m,n}$ as follows:

$$\begin{cases} u_{i,n}(x) = \tilde{v}_{i,n}(x - x^\nu), & \forall x \in J_{n,\nu}, \nu = 1, \dots, s; \\ u_{i,n}(x) = \gamma_{i,n}, & \forall x \in J_n \setminus \tilde{J}_n, \end{cases} \tag{5.24}$$

with $\gamma_{i,n}$ as in (5.9). Let $F_n = \cup_{\nu=1}^s (x^\nu + F_n^*)$ where F_n^* is as in Case I. Then, by (5.24), (5.10), (5.8), (5.4) and (3.2) we have

$$\begin{aligned} \int S_g(u_{1,n}, \dots, u_{m,n}) d\mu_N &= s \int_{F_n^*} S_g(\tilde{v}_{1,n}, \dots, \tilde{v}_{m,n}) d\mu_N \\ &> s\mu_N(E_n^*) a_N^m g(\xi^n, \eta^n) / 2 = \delta_n \kappa_n a_N^m g(\xi^n, \eta^n) / 2 \\ &\geq (a_N^m / 4) \alpha_n^{-1} g(\xi^n, \eta^n) > a_N^m / 4. \end{aligned} \tag{5.25}$$

Next let $\beta = 2 \sum_1^\infty |\Gamma_{n+1} - \Gamma_n|$, where $\Gamma_n = (\gamma_{1,n}, \dots, \gamma_{m,n})$. Note that by (5.21)–(5.23) and (5.9) β is finite. Indeed,

$$|\Gamma_n - \xi^n| < 2\sqrt{N} \sigma_n |\eta^n| < \sqrt{N} \delta_n |\eta^n| \quad (5.26)$$

and hence,

$$|\Gamma_{n+1} - \Gamma_n| < |\xi^{n+1} - \xi^n| + \sqrt{N} (\delta_{n+1} |\eta^{n+1}| + \delta_n |\eta^n|) < |\xi^{n+1} - \xi^n| + \sqrt{N} C.$$

Let $J_n^* = \{x: 0 < x_1 < \delta_n^*, 0 < x_i < 1, i = 2, \dots, N\}$, where

$$\delta_n^* = \begin{cases} |\Gamma_{n+1} - \Gamma_n| \beta^{-1} & \text{if } \beta \neq 0, \\ 0 & \text{if } \beta = 0. \end{cases}$$

In J_n^* we define functions $u_{i,n}^*$ ($i = 1, \dots, m$) by

$$\begin{cases} u_{i,n}^*(x) = ((\gamma_{i,n+1} - \gamma_{i,n})/\delta_n^*)x_1 + \gamma_{i,n} & \text{if } \delta_n^* > 0, \\ u_{i,n}^*(x) = \gamma_{i,n} & \text{if } \delta_n^* = 0, \forall x \in J_n^*. \end{cases} \quad (5.27)$$

We are now ready to define functions w_1, \dots, w_m in Ω which will be shown to satisfy (5.16). Let $e^1 = (1, 0, \dots, 0)$ and set $I_n = \beta_n e^1 + J_n$, $I_n^* = \beta_n^* e^1 + J_n^*$ where

$$\beta_1 = 0, \quad \beta_n = \sum_1^{n-1} (\delta_k + \delta_k^*), \quad n = 2, 3, \dots;$$

$$\beta_n^* = \beta_{n+1} - \delta_n^*, \quad n = 1, 2, \dots$$

Let

$$\begin{cases} w_i(x) = u_{i,n}(x - \beta_n e^1), & \forall x \in I_n, \quad n = 1, 2, \dots, \\ w_i(x) = u_{i,n}^*(x - \beta_n^* e^1), & \forall x \in I_n^*, \quad n = 1, 2, \dots, \\ w_i(x) = \xi^0, & \forall x \in \bar{\Omega} \setminus \cup_i^\infty (I_n + I_n^*), \quad i = 1, \dots, m, \end{cases} \quad (5.28)$$

with ξ^0 as in (5.22).

The function w_i is bounded and continuous in $\bar{\Omega}$ and it is piecewise linear in every subdomain bounded away from the hyperplane $x_1 = \sum_1^\infty (\delta_n + \delta_n^*)$. Furthermore, by (5.21), (5.24), (5.27) and (5.28),

$$|\nabla w_i| < C \text{ in } I_n \quad \text{and} \quad |\nabla w_i| < \beta \text{ in } I_n^*.$$

Hence $w_i \in W_{1,\infty}(\Omega)$.

It remains to show that w_1, \dots, w_m satisfy (5.16)₂. By (5.28) and (5.25),

$$\begin{aligned} \int_{\Omega} S_g(w_1, \dots, w_m) d\mu_N &> \sum_{n=1}^{\infty} \int_{I_n} S_g(w_1, \dots, w_m) d\mu_N \\ &= \sum_{n=1}^{\infty} \int_{J_n} S_g(u_{1,m}, \dots, u_{m,n}) d\mu_N > \sum_{n=1}^{\infty} \int_{F_n} S_g(u_{1,n}, \dots, u_{m,n}) d\mu_N = \infty. \end{aligned}$$

Thus (5.4) and (5.21) lead to a contradiction.

Case IV. Suppose that $1 < p < \infty$ and that

$$|\xi^n| \leq C_0, \quad C_1 \leq |\eta^n|^p \tag{5.29}$$

for some positive C_0, C_1 . By considering a subsequence if necessary we may assume $\sum_1^{\infty} \alpha_n^{-1} \leq C_1/2$ and that (5.22) holds. Set

$$\delta_n = (\alpha_n |\eta^n|^p)^{-1}. \tag{5.30}$$

Note that $\sum_1^{\infty} \delta_n \leq \sum_1^{\infty} (\alpha_n C_1)^{-1} \leq \frac{1}{2}$. Let β be as in Case III. By (5.22), (5.26), (5.29) and (5.30), β is finite. Indeed,

$$\begin{aligned} |\Gamma_{n+1} - \Gamma_n| &\leq |\xi^{n+1} - \xi^n| + \sqrt{N} (\delta_{n+1} |\eta^{n+1}| + \delta_n |\eta^n|) \\ &\leq |\xi^{n+1} - \xi^n| + \sqrt{N} C_1^{(p-1)/p} (\alpha_{n+1}^{-1} + \alpha_n^{-1}). \end{aligned}$$

Let δ_n^* be as in Case III. Define $u_{i,n}, u_{i,n}^*$ and w_i as in the previous case.

The function w_i is piecewise linear in I_n and

$$w_i = \gamma_{i,n} \text{ on } \partial I_n, \quad |\nabla w_i| \leq |\eta^n| \text{ in } I_n \quad (n = 1, 2, \dots).$$

Thus $|w_i - \gamma_{i,n}| \leq |\eta^n| \delta_n \leq C_1^{(p-1)/p} \alpha_n^{-1}$ everywhere in I_n . Furthermore, by (5.26), $|\xi^n - \gamma_{i,n}| \leq \sqrt{N} \delta_n |\eta^n| \leq \text{const } \alpha_n^{-1}$ ($n = 1, 2, \dots$). Finally, in I_n^* ($n = 1, 2, \dots$), w_i is linear and its range lies between $\gamma_{i,n}$ and $\gamma_{i,n+1}$. Therefore (in view of (5.22)) w_i is bounded and continuous in Ω and w_i is piecewise linear in every subdomain bounded away from the hyperplane $x_1 = \sum_1^{\infty} (\delta_n + \delta_n^*)$. In addition, by (5.24), (5.27), (5.28) and (5.30)

$$\begin{aligned} \int_{\Omega} |\nabla w_i|^p d\mu_N &= \sum_1^{\infty} \left(\int_{I_n} |\nabla w_i|^p d\mu_N + \int_{I_n^*} |\nabla w_i|^p d\mu_N \right) \\ &\leq \sum_1^{\infty} \left[|\eta^n|^p \delta_n + \left(\frac{|\Gamma_{n+1} - \Gamma_n|}{\delta_n^*} \right)^p \delta_n^* \right] \\ &= \sum_1^{\infty} \alpha_n^{-1} + \beta^{p-1} \sum_1^{\infty} |\Gamma_{n+1} - \Gamma_n| < \infty. \end{aligned}$$

Thus (5.16)₁ holds.

By (5.4), (5.25), (5.28) and (5.30),

$$\int_{\Omega} S_g(w_1, \dots, w_m) d\mu_N \geq \sum_1^{\infty} \int_{F_n} S_g(u_{1,n}, \dots, u_{m,n}) d\mu_N$$

$$> \left(\frac{a_N^m}{4}\right) \sum_1^{\infty} \delta_n g(\xi^n, \eta^n) \geq \left(\frac{a_N^m}{4}\right) \sum_1^{\infty} (\alpha_n |\eta^n|^p)^{-1} \alpha_n (1 + |\eta^n|^p) = \infty.$$

Thus (5.16)₂ holds. This shows that (5.4) and (5.29) lead to a contradiction.

We claim that the contradictions obtained in Cases I-IV prove (5.3)₁ and (5.3)₂. Clearly, if one of the assumptions (5.5), (5.19), (5.21) or (5.29) holds for a subsequence of $\{(\xi^n, \eta^n)\}$ then (5.4) is impossible. Thus, Case III shows that if $\{(\xi^n, \eta^n)\}$ contains a bounded subsequence then we cannot have (5.4). Now suppose that $\{\xi^n\}$ is bounded. Then we must have $|\eta^n| \rightarrow \infty$ and Case IV show that (5.4) is impossible. This already proves (5.3)₂.

Next, if $1 \leq p < N$, Case I shows that (5.4) cannot hold unless

$$\lim |\xi^n|^q / (1 + |\eta^n|^p) = 0. \tag{5.31}$$

If (5.31) holds and if $\{\eta^n\}$ contains a subsequence that is bounded away from zero then Case II shows that (5.4) is impossible. On the other hand if (5.31) holds and $\eta^n \rightarrow 0$ then the sequence $\{(\xi^n, \eta^n)\}$ is bounded and this again contradicts (5.4). This proves (5.3)₁ and completes the proof of the theorem.

6. Proof of Theorem 1; the necessity of the conditions. We start with the case $m = 1$. Thus $h: R \rightarrow R$ is a Borel function such that T_h maps $W_{1,p}(\Omega)$ into $W_{1,r}(\Omega)$. Without loss of generality we may assume that Ω is the unit cube $\{x: 0 < x_i < 1, i = 1, \dots, N\}$.

First we claim that h is l.a.c. (i.e. locally absolutely continuous) in R . Let $u(x) = x_1 - c$ where c is an arbitrary constant. Since $u \in W_{1,p}(\Omega)$, we have $h \circ u \in W_{1,r}(\Omega)$. This implies that there exists an l.a.c. function $h_0: R \rightarrow R$ such that $h = h_0$ a.e. in R . Suppose that there exists a point ξ_0 such that $h(\xi_0) \neq h_0(\xi_0)$. Consider the function u given by

$$u(x) = \begin{cases} (x_1 - \frac{1}{2}) + \xi_0, & \frac{1}{2} \leq x_1 < 1, \\ \xi_0, & 0 < x_1 < \frac{1}{2}, \forall x \in \Omega. \end{cases}$$

Let $v = h \circ u$. Thus $v(x) = h(x_1 - \frac{1}{2} + \xi_0) = h_0(x_1 - \frac{1}{2} + \xi_0)$ a.e. in the set $\{x \in \Omega: \frac{1}{2} < x_1\}$ and $v(x) = h(\xi_0) \neq h_0(\xi_0)$ in $\{x \in \Omega: x_1 \leq \frac{1}{2}\}$. Therefore $\partial v / \partial x_1$ (in the distribution sense) is not a function. On the other hand, by assumption, $v \in W_{1,r}(\Omega)$ because $u \in W_{1,p}(\Omega)$. This contradiction shows that h is identical with h_0 .

Let $u \in W_{1,1}^{loc}(\Omega)$. Since h is l.a.c., $h \circ u \in W_{1,1}^{loc}(\Omega)$ if and only if $(h^* \circ u) \nabla u \in L_1^{loc}(\Omega)^N$, where $h^*: R \rightarrow R$ is any Borel function which is equivalent to h' . Moreover $\nabla(h \circ u) = (h^* \circ u) \nabla u$ a.e. in Ω . For a proof of this result see [4].

Therefore, if $g: R \times R_N \rightarrow R$ is defined by

$$g(\xi, \eta) = |h^*(\xi)\eta|, \quad (\xi, \eta) \in R \times R_N, \tag{6.1}$$

then S_g maps $W_{1,p}(\Omega)$ into $L_r(\Omega)$. Hence, by Theorem 2, if $1 < r < p < N$, we have

$$g(\xi, \eta) \leq \text{const}(1 + |\xi|^{q/r} + |\eta|^{p/r}), \quad q = Np / (N - p), \tag{6.2}$$

for every η in R_N and every ξ in $R \setminus C_\eta$ where C_η is a null subset of R which may depend on η . In the present case, g is continuous with respect to η . Therefore the above statement implies that there exists a null subset of R , say A , such that (6.2) holds for every (ξ, η) in $(R \setminus A) \times R_N$. In fact if D is a countable dense subset of R_N we could set $A = \bigcup_{\eta \in D} C_\eta$.

Similarly, if $p > N$ (or $p > 1$ and $N = 1$) and $M > 0$, we have

$$g(\xi, \eta) \leq c(M)(1 + |\eta|^{p/r}), \tag{6.3}$$

for every (ξ, η) in $(R \setminus A) \times R_N$ such that $|\xi| \leq M$, with A (as before) a null subset of R .

From (6.1), (6.3) we deduce that h^* is locally essentially bounded and hence that h is locally Lipschitz in R .

If $p < N$, (6.2) implies that

$$|h^*(\xi)| \leq \text{const} \inf_{0 < \gamma < \infty} H(|\xi|, \gamma), \quad \xi \in R \setminus A, \tag{6.4}$$

where

$$H(\beta, \gamma) = (1 + \beta^{q/r} + \gamma^{p/r})/\gamma \quad (0 < \beta, \gamma). \tag{6.5}$$

A simple computation show that (6.4) and (6.5) imply that

$$|h^*(\xi)| \leq \text{const}(1 + |\xi|^\nu) \quad \forall \xi \in R \setminus A, \text{ where } \nu = N(p - r) / (r(N - p)). \tag{6.6}$$

This completes the proof in the case $m = 1$.

Next we consider the case $m > 1$. Thus $h: R_m \rightarrow R$ is a Borel function such that T_h maps $W_{1,p}(\Omega)^m$ into $W_{1,r}(\Omega)$. Again we may assume that Ω is the unit cube.

Let $f_i: R \rightarrow R, i = 1, \dots, m$, be uniformly Lipschitz functions. Let $\tilde{h}: R \rightarrow R$ be the function given by

$$\tilde{h}(t) = h(f_1(t), \dots, f_m(t)), \quad \forall t \in R. \tag{6.7}$$

If $u \in W_{1,p}(\Omega)$ then $f_i \circ u \in W_{1,p}(\Omega), i = 1, \dots, m$. Therefore $T_{\tilde{h}}$ maps $W_{1,p}(\Omega)$ into $W_{1,r}(\Omega)$. Hence, by the first part of the proof, \tilde{h} is locally Lipschitz in R . This implies in particular that h is locally Lipschitz on every line parallel to one of the axes.

Now we claim that h is continuous and moreover

$$\limsup_{\xi \rightarrow \xi^0} |h(\xi) - h(\xi^0)| / |\xi - \xi^0| = M(\xi^0) < \infty, \quad \forall \xi^0 \in R_m. \quad (6.8)$$

Suppose that (6.8) fails at some point ξ^0 . Then there exists a sequence $\{\xi^n\}$ in R_m such that $\xi^n \rightarrow \xi^0$ and

$$|h(\xi^n) - h(\xi^0)| > n|\xi^n - \xi^0|, \quad n = 1, 2, \dots \quad (6.9)$$

By considering a subsequence if necessary, we may assume that

$$|\xi^n - \xi^0| > 2|\xi^{n+1} - \xi^0|, \quad n = 1, 2, \dots \quad (6.10)$$

Let

$$\alpha = \sum_{k=1}^{\infty} |\xi^{k+1} - \xi^k|, \quad \alpha_{n+1} = \sum_{k=1}^n |\xi^{k+1} - \xi^k| \quad (n = 1, 2, \dots) \text{ and } \alpha_1 = 0.$$

In view of (6.10) α is finite. We define a curve $\xi = \xi(t)$, $0 < t < \alpha$, as follows.

$$\begin{aligned} \xi(t) &= \xi^n + ((t - \alpha_n) / (\alpha_{n+1} - \alpha_n))(\xi^{n+1} - \xi^n), \\ & \qquad \qquad \qquad t \in [\alpha_n, \alpha_{n+1}], \quad n = 1, 2, \\ \xi(\alpha) &= \xi^0. \end{aligned}$$

Clearly, $|\xi(t) - \xi(t')| < |t - t'|$ for every t, t' in $[0, \alpha]$. Therefore the function \tilde{h} given by $\tilde{h}(t) = h(\xi(t))$ is Lipschitz in $[0, \alpha]$. Thus, using (6.10),

$$|h(\xi^n) - h(\xi^0)| = |\tilde{h}(\alpha_n) - \tilde{h}(\alpha)| < \text{const}|\xi^n - \xi^0|, \quad n = 1, 2, \dots$$

But this contradicts (6.9).

By Stepanov's theorem [1, p. 218], (6.8) implies that h possesses a differential a.e. in R_m . Furthermore, since h is continuous, the set B where its differential is defined is a Borel set and the partial derivatives $\partial h / \partial \xi_i$, $i = 1, \dots, m$, are Borel functions on B [1, p. 211].

Let $g: R_m \times R_{mN} \rightarrow [0, \infty)$ be the function given by

$$g(\xi, \eta) = \begin{cases} \left| \sum_{i=1}^m \frac{\partial h}{\partial \xi_i}(\xi)(\eta_{i,1}, \dots, \eta_{i,N}) \right|, & \text{if } \xi \in B, \\ 0, & \text{if } \xi \in R_m \setminus B. \end{cases} \quad (6.11)$$

(As usual we write $\eta = (\eta_{1,1}, \dots, \eta_{1,N}, \dots, \eta_{m,1}, \dots, \eta_{m,N})$.) Clearly g is a Borel function. We claim that

$$\mathbf{u} = (u_1, \dots, u_m) \in W_{1,p}(\Omega)^m \Rightarrow S_g(\mathbf{u}) \in L_r(\Omega). \quad (6.12)$$

Since \mathbf{u} belongs to $W_{1,p}(\Omega)^m$, it is equivalent to a function \mathbf{u}^* such that, for a.e. line τ parallel to one of the axes in R_N , $\mathbf{u}^*|_{\tau \cap \Omega}$ is l.a.c. Furthermore $\partial \mathbf{u}^* / \partial x_j$ (the partial derivative in the classical sense) exists a.e. in Ω and is equivalent to the corresponding distribution derivative of \mathbf{u} , $j = 1, \dots, m$

(see [7, §3.1]). Without loss of generality we may assume that \mathbf{u} is identical with \mathbf{u}^* . Then the function $h \circ \mathbf{u}$, which by assumption belongs to $W_{1,r}(\Omega)$, is continuous in $\tau \cap \Omega$ for a.e. line τ parallel to one of the axes. (Recall that h is continuous.) It follows from this that $h \circ \mathbf{u}$ is l.a.c. in $\tau \cap \Omega$ for a.e. τ as above. Now, if x^0 is a point in Ω such that $\mathbf{u}(x^0) \in B$ and such that $\partial \mathbf{u} / \partial x_j$ exists at x^0 then $\partial (h \circ \mathbf{u}) / \partial x_j$ exists at x^0 and

$$\left(\frac{\partial (h \circ \mathbf{u})}{\partial x_j} \right) (x^0) = \sum_{i=1}^m \left(\frac{\partial h}{\partial \xi_i} \right) (\mathbf{u}(x^0)) \left(\frac{\partial u_i}{\partial x_j} \right) (x^0).$$

Hence, by (6.11), $S_g(\mathbf{u}) = |\nabla(h \circ \mathbf{u})|$ a.e. in $B^* = \{x \in \Omega: \mathbf{u}(x) \in B\}$ and $S_g(\mathbf{u}) = 0$ in $\Omega \setminus B^*$. Since $h \circ \mathbf{u} \in W_{1,r}(\Omega)$, this implies that $S_g(\mathbf{u}) \in L_r(\Omega)$.

Suppose that $1 < r < p < N$. By Theorem 2, (6.12) implies that g satisfies inequality (6.2) for every η in R_N and every ξ in $R_m \setminus C_\eta$, where C_η is a null set. From this we deduce, by the same argument as in the case $m = 1$, that the partial derivatives of h satisfy the inequality

$$\left| \frac{\partial h}{\partial \xi_i} (\xi) \right| \leq \text{const}(1 + |\xi|^r) \quad \text{a.e. in } R_m, i = 1, \dots, m, \quad (6.13)$$

with ν as in (2.2).

Similarly when $p > N$ (or $p \geq 1$ if $N = 1$) we deduce that for every $M > 0$ there exists a constant κ_M such that

$$\left| \frac{\partial h}{\partial \xi_i} (\xi) \right| \leq \kappa_M \quad \text{a.e. in the sphere } |\xi| < M, i = 1, \dots, m. \quad (6.14)$$

By (6.13) and (6.14) the partial derivatives $\partial h / \partial \xi_i$ ($i = 1, \dots, m$) are locally essentially bounded in R_m . This, together with the fact that h is continuous in R_m and locally Lipschitz on every line parallel to one of the axes, implies that h is locally Lipschitz in R_m . This completes the proof of the necessity part of Theorem 1.

7. Proof of Theorem 1; the sufficiency of the conditions. The sufficiency of the conditions of Theorem 1 could be derived from [5, Theorems 2.1 and 3.3] in which a much more general situation was considered. However in order to apply the results of [5] directly to our case we would have to assume that the inequality in (2.1) holds at every point where $\partial h / \partial \xi_i$ exists. An examination of the arguments in [5] shows that, in the present case, this stronger condition can be replaced by (2.1). However, for the convenience of the reader we shall present here a simple, independent proof of the sufficiency part of Theorem 1.

First some general remarks. Let $A(\Omega)$ denote the space of real measurable functions in Ω such that $v \in A(\Omega)$ iff for almost every line τ parallel to one of the axes $v|_{\tau \cap \Omega}$ is l.a.c. (i.e. locally absolutely continuous). A function u

belongs to $W_{1,p}(\Omega)$ if and only if u is equivalent to a function u^* belonging to $A(\Omega)$ such that u^* and $\partial u^*/\partial x_i$ ($i = 1, \dots, N$) belong to $L_p(\Omega)$. Here $\partial u^*/\partial x_i$ denotes the classical partial derivative of u^* . (It is known that $\partial u^*/\partial x_i$ is defined a.e. in Ω and is measurable.) Furthermore $\partial u^*/\partial x_i$ is equivalent to the corresponding distribution derivative of u . (See [7], [2].)

If $h: R_m \rightarrow R$ is locally Lipschitz and $\mathbf{u} = (u_1, \dots, u_m) \in A(\Omega)^m$ then, clearly $h \circ \mathbf{u} \in A(\Omega)$.

We turn now to the proof of the sufficiency part of Theorem 1 and we start with the case where $p > N$ or $N = 1$ and $p \geq 1$. In this case every function $u \in W_{1,p}(\Omega)$ is equivalent to a function in $C(\Omega)$ and

$$\|u\|_{L_\infty(\Omega)} \leq c_1 \|u\|_{W_{1,p}(\Omega)} \quad (7.1)$$

where c_1 is a constant which depends on Ω .

Let $\mathbf{u} \in W_{1,p}(\Omega)^m$. Without loss of generality we may assume that $\mathbf{u} \in C(\Omega)^m$. This implies that $\mathbf{u} \in A(\Omega)^m$ and hence $h \circ \mathbf{u} \in A(\Omega)$. Let $\|\mathbf{u}\|_{W_{1,p}(\Omega)} \leq M$ and let b_M be the Lipschitz constant of h in the sphere $|\xi| \leq c_1 M$. Then, by (7.1)

$$|h \circ \mathbf{u}(x) - h(\mathbf{0})| \leq b_M |\mathbf{u}(x)|$$

and therefore

$$\|h \circ \mathbf{u}\|_{L_p(\Omega)} \leq |h(\mathbf{0})| \mu_N(\Omega)^{1/p} + b_M \|\mathbf{u}\|_{L_p(\Omega)}. \quad (7.2)$$

Let τ be a line parallel to the x_i -axis such that $\mathbf{u}|_{\tau \cap \Omega}$ is l.a.c. and let x, x' be two points in $\tau \cap \Omega$. Since

$$|h \circ \mathbf{u}(x) - h \circ \mathbf{u}(x')| \leq b_M |\mathbf{u}(x) - \mathbf{u}(x')|,$$

dividing by $|x - x'|$ and letting x' tend to x we obtain

$$|\partial(h \circ \mathbf{u})/\partial x_i| \leq b_M |\partial \mathbf{u}/\partial x_i| \quad \mu_1\text{-a.e. on } \tau \cap \Omega. \quad (7.3)$$

In view of the fact that $h \circ \mathbf{u} \in A(\Omega)$, (7.2) and (7.3) imply that $h \circ \mathbf{u} \in W_{1,p}(\Omega)$ and

$$\|h \circ \mathbf{u}\|_{W_{1,p}(\Omega)} \leq |h(\mathbf{0})| \mu_N(\Omega)^{1/p} + b_M \|\mathbf{u}\|_{W_{1,p}(\Omega)^m}. \quad (7.4)$$

Next we consider the case where $1 < r < p < N$. Let Z_k ($k = 1, \dots, m$) be the family of lines $\{\tau\}$ in R_m , parallel to the ξ_k axis, such that (2.1) holds μ_1 -a.e. on τ . The projection of Z_k on the hyperplane $\xi_k = 0$ is a set of total measure (with respect to μ_{m-1}) in this hyperplane.

Given two points ξ, ξ' in R_m we shall say that they are connected through Z_1, \dots, Z_m if there exists a continuous piecewise linear curve C connecting ξ with ξ' , which consists of j linear segments C_1, \dots, C_j such that C_i is a segment of a line belonging to Z_{k_i} ($i = 1, \dots, j$) and k_1, \dots, k_j are distinct. Suppose that ξ, ξ' have this property. Then, by (2.1),

$$|h(\xi) - h(\xi')| < \sum_{i=1}^j \int_{C_i} \left| \frac{\partial h}{\partial \xi_{k_i}} \right| d\mu_1 < ma_0(1 + |\xi|^r + |\xi'|^r)|\xi - \xi'|.$$

Now, if ξ, ξ' are two arbitrary points in R_m there exist sequences of points in R_m , say $\{\xi^n\}$ and $\{\xi'^n\}$, such that $\xi^n \rightarrow \xi, \xi'^n \rightarrow \xi'$ and each pair ξ^n, ξ'^n is connected through Z_1, \dots, Z_m . Therefore

$$|h(\xi^n) - h(\xi'^n)| < ma_0(1 + |\xi^n|^r + |\xi'^n|^r)|\xi^n - \xi'^n|$$

for every n and by continuity

$$|h(\xi) - h(\xi')| < ma_0(1 + |\xi|^r + |\xi'|^r)|\xi - \xi'|. \tag{7.5}$$

Now let $u \in W_{1,p}(\Omega)^m$. Without loss of generality we may assume that u belongs also to $A(\Omega)^m$ and so $h \circ u \in A(\Omega)$. By (7.5),

$$|h \circ u(x)| < |h(0)| + ma_0(1 + |u(x)|^r)|u(x)|.$$

Hence, by Sobolev's imbedding theorem

$$\begin{aligned} \|h \circ u\|_{L_r(\Omega)} &< |h(0)| \mu_N(\Omega)^{1/r} + ma_0(\|u\|_{L_r(\Omega)} + \|u\|_{L_q(\Omega)}^r \|u\|_{L_r(\Omega)}) \\ &< |h(0)| \mu_N(\Omega)^{1/r} + \text{const} \|u\|_{W_{1,r}^+(\Omega)} \end{aligned} \tag{7.6}$$

where the constant depends on Ω and $q = Np/(N - p)$.

Let τ be a line parallel to the x_i -axis such that $u|_{\tau \cap \Omega}$ is l.a.c. and let $x, x' \in \tau \cap \Omega$. Then by (7.5),

$$|h \circ u(x) - h \circ u(x')| < ma_0(1 + |u(x)|^r + |u(x')|^r)|u(x) - u(x')|.$$

Hence

$$|\partial(h \circ u)/\partial x_i| < ma_0(1 + 2|u(x)|^r)|\partial u/\partial x_i|$$

μ_1 -a.e. in $\tau \cap \Omega$. This inequality implies

$$\begin{aligned} \|\partial(h \circ u)/\partial x_i\|_{L_r(\Omega)} &< 2ma_0(\|\partial u/\partial x_i\|_{L_r(\Omega)} + \|u\|_{L_q(\Omega)}^r \|\partial u/\partial x_i\|_{L_r(\Omega)}) \\ &< \text{const} \|u\|_{W_{1,r}^+(\Omega)} \end{aligned} \tag{7.7}$$

where the constant depends on Ω . In view of the fact that $h \circ u \in A(\Omega)$, (7.6) and (7.7) imply that $h \circ u \in W_{1,r}(\Omega)$ and

$$\|h \circ u\|_{W_{1,r}(\Omega)} < \text{const}(1 + \|u\|_{W_{1,r}^+(\Omega)}). \tag{7.8}$$

This completes the proof of Theorem 1.

8. The case of unbounded domains. Let Ω be an unbounded domain in R_N satisfying the cone condition. Then we have the following result, parallel to Theorems 1 and 2.

THEOREM 3. *Let $h: R_m \rightarrow R$ be a Borel function and let r, p be two numbers such that $1 < r \leq p < N$. Then T_h maps $W_{1,p}(\Omega)^m$ into $W_{1,r}(\Omega)$ if and only if the following conditions hold.*

- (i) $h(0) = 0$ and h is locally Lipschitz in R_m .
- (ii) The first order partial derivatives of h satisfy the inequality

$$\left| \frac{\partial h}{\partial \xi_i}(\xi) \right| \leq a_0(|\xi|^v + |\xi|^{v'}) \quad \text{a.e. in } R_m, i = 1, \dots, m, \tag{8.1}$$

where a_0 is a constant, and $v = N(p - r)/(r(N - p))$ and $v' = (p - r)/r$.

If $p > N$ (or $N = 1$ and $p \geq 1$) and $r = p$ then T_h maps $W_{1,p}(\Omega)^m$ into $W_{1,r}(\Omega)$ if and only if (i) holds.

In either case, if T_h maps $W_{1,p}(\Omega)^m$ into $W_{1,r}(\Omega)$ then the mapping is bounded and the following inequalities hold:

$$\begin{aligned} \|T_h \mathbf{u}\|_{W_{1,r}(\Omega)} &\leq b_0 \sum_{i=1}^m (\|u_i\|_{W_{1,p}(\Omega)}^{v+1} + \|u_i\|_{W_{1,p}(\Omega)}^{v'+1}), \quad \text{if } p < N; \\ \|T_h \mathbf{u}\|_{W_{1,p}(\Omega)} &\leq b(M) \sum_{i=1}^m \|u_i\|_{W_{1,p}(\Omega)}, \quad \text{if } p > N \text{ (or } p \geq 1 \text{ when } N = 1) \\ &\text{and } \sum_{i=1}^m \|u_i\|_{W_{1,p}(\Omega)} \leq M. \end{aligned} \tag{8.2}$$

The constants b_0 and $b(M)$ depend on m, a_0 and Ω .

THEOREM 4. *Let g, p, r be as in Theorem 2. Suppose that S_g maps $W_{1,p}(\Omega)^m$ into $L_r(\Omega)$ (where Ω is unbounded). Then g satisfies the conditions described in Theorem 2 with (2.3) and (2.4) replaced by the first and second of the inequalities below:*

$$|g(\xi, \eta)| \leq C_0(|\xi|^{q/r} + |\xi|^{p/r} + |\eta|^{p/r}), \tag{8.3}$$

$$|g(\xi, \eta)| \leq C(M)(|\xi|^{p/r} + |\eta|^{p/r}). \tag{8.4}$$

In addition we have $g(0, 0) = 0$.

We shall start with the proof of Theorem 4, in which we use the result of Theorem 2. Then, using Theorems 1 and 4, we shall prove Theorem 3.

PROOF OF THEOREM 4. Without loss of generality we shall assume that $g \geq 0$ and $r = 1$. As in the proof of Theorem 2 we shall show that:

(8.5)₁ If $1 < p < N$ then (8.3) holds everywhere in Q .

(8.5)₂ If $p > N$ or $p \geq 1$ and $N = 1$ then (8.4) holds everywhere in Q_M .

If Ω' is a cube contained in Ω then S_g maps $W_{1,p}(\Omega')^m$ into $L_1(\Omega')$. Therefore g satisfies inequality (2.3) in Q when $p < N$ and inequality (2.4) in Q_M when $p > N$ or $p \geq 1$ and $N = 1$. But inequality (8.3) (resp. (8.4)) is equivalent to inequality (2.3) (resp. (2.4)) in every domain of the form $\{(\xi, \eta): |\xi| + |\eta| > a > 0\}$. Therefore, if either (8.5)₁ or (8.5)₂ fails then there must exist a sequence

$\{(\xi^n, \eta^n)\}$ in Q such that

$$|\xi^n| + |\eta^n| \rightarrow 0 \tag{8.6}$$

and

$$g(\xi^n, \eta^n) / (|\xi^n|^p + |\eta^n|^p) = \alpha_n \rightarrow \infty \quad \text{as } n \rightarrow \infty. \tag{8.7}$$

We shall show that (8.6) and (8.7) lead to a contradiction. But first some general remarks.

By considering a subsequence of $\{(\xi^n, \eta^n)\}$ if necessary, we may assume that

$$\sum_1^\infty \alpha_n^{-1} < \infty \quad \text{and} \quad \sum_1^\infty (|\xi^n|^p + |\eta^n|^p) < \infty. \tag{8.8}$$

Since Ω is unbounded and satisfies the cone condition, it follows that there exists an infinite family \mathcal{Q} of disjoint closed cubes of equal volume such that each cube is contained in Ω and its sides are parallel to the axes. We shall denote by 2λ the side length of these cubes.

Now we shall consider two cases in each of which we shall show that (8.6) and (8.7) lead to a contradiction to the assumptions of the theorem. The conclusion of the theorem will follow directly from these results.

Case (a). Suppose that there exists a positive β such that

$$g(\xi^n, \eta^n) = \alpha_n (|\xi^n|^p + |\eta^n|^p) < \beta, \quad n = 1, 2, \dots \tag{8.9}$$

In view of (8.7) it is clear that $g(\xi^n, \eta^n) \neq 0$ for sufficiently large n ; we may assume that $g(\xi^n, \eta^n) \neq 0$ for all n . Let

$$K_n = [\beta g(\xi^n, \eta^n)^{-1}] + 1 \tag{8.10}$$

where (for $b > 0$) $[b]$ denotes the largest integer that is not larger than b . Note that, in view of (8.9),

$$K_n < 2\beta g(\xi^n, \eta^n)^{-1}. \tag{8.11}$$

Let $\{\mathcal{Q}_n\}^\infty$ be a partition of \mathcal{Q} such that \mathcal{Q}_n consists of exactly K_n cubes, $A_{n,j}$ ($j = 1, \dots, K_n$). Let $w_{i,n}$ be defined as in the proof of Theorem 2 (see (5.13)) with D_n, E_n replaced by

$$D = \{x: |x_i| < \lambda, i = 1, \dots, R\}, \quad E = \{x: |x_i| < \lambda/2, i = 1, \dots, N\}.$$

Then, by (5.9)–(5.11), (5.13) and (5.14) we have

$$\begin{cases} |w_{i,n}(x)| < |\gamma_{i,n}| + \lambda|\eta^n| < \text{const}(|\xi^n| + |\eta^n|), & \forall x \in D, \\ |\nabla w_{i,n}(x)| < |\eta^n|, & \text{a.e. in } E, \\ |\nabla w_{i,n}(x)| < 2|\gamma_{i,n}|\lambda^{-1} < \text{const}(|\xi^n| + |\eta^n|), & \text{a.e. in } D \setminus E, \end{cases} \tag{8.12}$$

$n = 1, 2, \dots$. Now we define m functions w_1, \dots, w_m in $\bar{\Omega}$ as follows:

$$\begin{cases} w_i(x) = w_{i,n}(x - x^{n,j}), & \forall x \in A_{n,j} \ (n = 1, 2, \dots; j = 1, \dots, K_n), \\ w_i(x) = 0, & x \in \bar{\Omega} \setminus \Omega_0, \end{cases} \quad (8.13)$$

where $x^{n,j}$ is the center of the cube $A_{n,j}$ and $\Omega_0 = \cup \{A: A \in \mathcal{Q}\}$.

Let $\tilde{D}_n = \cup_{j=1}^{K_n} A_{n,j}$. Then by (8.9)–(8.13),

$$\begin{aligned} \int_{\tilde{D}_n} |w_i|^p d\mu_N &< \text{const } K_n \lambda^N (|\xi^n| + |\eta^n|)^p < \text{const } K_n (|\xi^n|^p + |\eta^n|^p) \\ &< \text{const } \alpha_n^{-1}, \quad n = 1, 2, \dots \end{aligned}$$

and

$$\int_{\tilde{D}_n} |\nabla w_i|^p d\mu_N < \text{const } K_n \lambda^N (|\xi^n| + |\eta^n|)^p < \text{const } \alpha_n^{-1}, \quad n = 1, 2, \dots$$

Therefore, by (8.8), w_i and $|\nabla w_i|$ belong to $L_p(\Omega)$, $i = 1, \dots, m$. In addition, w_i is locally Lipschitz in Ω ; hence, $w_i \in W_{1,p}(\Omega)$, $i = 1, \dots, m$.

On the other hand, by (8.10), (8.11), (8.13) and (5.12), (5.13) we have

$$\begin{aligned} \int_{\tilde{D}_n} S_g(w_1, \dots, w_m) d\mu_N &= K_n \int_D S_g(w_{1,n}, \dots, w_{m,n}) d\mu_N \\ &> K_n \int_E S_g(u_{1,n}, \dots, u_{m,n}) d\mu_N > K_n b_1 \lambda^N g(\xi^n, \eta^n) > b_1 \lambda^N \beta. \end{aligned}$$

Hence $S_g(w_1, \dots, w_m) \notin L_1(\Omega)$ and we have reached a contradiction.

Case (b). Suppose that

$$g(\xi^n, \eta^n) = \alpha_n (|\xi^n|^p + |\eta^n|^p) \rightarrow \infty \quad \text{as } n \rightarrow \infty. \quad (8.14)$$

Then we may assume that

$$g(\xi^n, \eta^n)^{-1} < \lambda^N, \quad n = 1, 2, \dots \quad (8.15)$$

Let

$$\delta_n^N = g(\xi^n, \eta^n)^{-1}. \quad (8.16)$$

Note that $\delta_n < \lambda$.

Let $u_{i,n}$, $i = 1, \dots, m$, be m functions defined in the cube $E_n = \{x: |x_i| < \delta_n/2, i = 1, \dots, N\}$ as in the proof of Theorem 2 (see (5.11)). Set

$$D_n = D = \{x: |x_i| < \lambda, i = 1, \dots, N\}$$

and

$$D_{n,\nu} = \{x: \delta_n/2 \leq |x_\nu| \leq \lambda, |x_\nu| \geq |x_k|, k = 1, \dots, N\}, \quad \nu = 1, \dots, N.$$

Note that $D = (\cup_{\nu=1}^N D_{n,\nu}) \cup E_n$. Now, for every n we define functions $w_{i,n}$, $i = 1, \dots, m$, in D as follows,

$$\begin{cases} w_{i,n}(x) = u_{i,n}(x), & \forall x \in E_n, \\ w_{i,n}(x) = \gamma_{i,n}(\lambda - \delta_n/2)^{-1}(\lambda - |x_\nu|), & \forall x \in D_{n,\nu}, \nu = 1, \dots, N. \end{cases} \quad (8.17)$$

Then, by (5.9)–(5.11),

$$\begin{cases} |w_{i,n}| < |\gamma_{i,n}| + \delta_n|\eta^n| < \text{const}(|\xi^n| + |\eta^n|), & \text{in } D, \\ |\nabla w_{i,n}| < |\eta^n|, & \text{in } E_n, \\ |\nabla w_{i,n}| < |\gamma_{i,n}|(\lambda - \delta_n/2)^{-1} < \text{const}(|\xi^n| + |\eta^n|), & \text{in } D \setminus E_n. \end{cases} \quad (8.18)$$

Here we have used also the fact that $\delta_n < \lambda$.

Let us denote the cubes in \mathcal{Q} by $A_n, n = 1, 2, \dots$, and let x^n be the center of A_n . Then we define functions w_1, \dots, w_m in $\bar{\Omega}$ as follows,

$$\begin{cases} w_i(x) = w_{i,n}(x - x^n), & \forall x \in A_n, n = 1, 2, \dots \\ w_i(x) = 0, & \forall x \in \bar{\Omega} \setminus \cup_1^\infty A_n, i = 1, \dots, m. \end{cases} \quad (8.19)$$

By (8.18).

$$\int_{A_n} |w_i|^p d\mu_N = \int_D |w_{i,n}|^p d\mu_N < \text{const}(|\xi^n| + |\eta^n|)^p \lambda^N$$

and

$$\int_{A_n} |\nabla w_i|^p d\mu_N = \int_D |\nabla w_{i,n}|^p d\mu_N < \text{const}(|\xi^n| + |\eta^n|)^p \lambda^N.$$

Hence, by (8.8), w_i and $|\nabla w_i|$ belong to $L_p(\Omega)$ and so $w_i \in W_{1,p}(\Omega)$. On the other hand, by (8.19), (8.17), (8.16) and (5.12),

$$\begin{aligned} \int_{A_n} S_g(w_1, \dots, w_m) d\mu_N &= \int_D S_g(w_{1,n}, \dots, w_{m,n}) d\mu_N \\ &\geq \int_{E_n} S_g(u_{1,n}, \dots, u_{m,n}) d\mu_N \\ &\geq b_1 \delta_n^N g(\xi^n, \eta^n) = b_1. \end{aligned}$$

Thus $S_g(w_1, \dots, w_m) \notin L_1(\Omega)$ and again we have reached a contradiction.

To complete the proof of the theorem we note that the sequence $\{(\xi^n, \eta^n)\}$ must contain a subsequence for which either (8.9) or (8.14) holds. Therefore (8.6) and (8.7) are incompatible with the assumption that S_g maps $W_{1,p}(\Omega)^m$ into $L_1(\Omega)$. This establishes (8.5)₁ and (8.5)₂ and hence (8.3) and (8.4).

Finally, since Ω is unbounded and satisfies the cone condition it has infinite measure. Therefore the fact that g vanishes for $(\xi, \eta) = (0, 0)$ is obvious; otherwise $S_g(0, \dots, 0)$ would not belong to $L_1(\Omega)$.

Next we turn to

PROOF OF THEOREM 3. Suppose that T_h maps $W_{1,p}(\Omega)^m$ into $W_{1,r}(\Omega)$. If Ω' is a bounded subdomain of Ω whose boundary is Lipschitz, then T_h maps $W_{1,p}(\Omega')^m$ into $W_{1,r}(\Omega')$. Therefore, by Theorem 2, h is locally Lipschitz in R_m . The fact that $h(0) = 0$ is obvious. In order to prove the growth estimate in the case $p < N$ we apply Theorem 4. Using this theorem one can show, by the same arguments that were used in §6, that

$$\left| \frac{\partial h}{\partial \xi_i}(\xi) \right| \leq \text{const} \inf_{0 < \gamma < \infty} \tilde{H}(|\xi|, \gamma) \quad \text{a.e. in } R_m, i = 1, \dots, m,$$

where

$$\tilde{H}(\beta, \gamma) = (\beta^{p/r} + \beta^{q/r} + \gamma^{p/r})/\gamma, \quad \beta, \gamma > 0.$$

Hence (considering the cases $|\xi| < 1$ and $|\xi| > 1$ separately) one obtains (8.1) by a simple calculation.

The sufficiency of the conditions stated in the theorem and inequality (8.2) can be verified by the same arguments that were used in the proof of Theorem 1 (§7).

9. Further remarks and open questions. In this section we shall discuss two additional problems related to the mappings S_g and T_h , for which only partial results are available. For simplicity we shall consider only the case where Ω is a bounded domain in R_N satisfying the cone condition.

The first problem is to obtain necessary and sufficient conditions in order that S_g map $W_{1,p}(\Omega)^m$ into $L_r(\Omega)$. Theorem 2 provides necessary conditions. However, when $m > 1$, these conditions are obviously not sufficient because of the exceptional sets where the inequalities (2.3) or (2.4) may fail. We describe below several results which may help to clarify the problem stated above. But first a definition.

Given a measurable set B in R_m we shall say that it is a *null projection set* if the projection of B on each of the coordinate axes is a μ_1 -null set.

THEOREM 5. *Let $g: R_m \times R_{mN} \rightarrow R$ be a Borel function and let p, r be two numbers such that $1 < r \leq p < \infty$. Suppose that*

$$g(\xi, 0) = 0 \quad \forall \xi \in R_m. \quad (9.1)$$

Suppose that there exists a null projection set B in R_m such that if $p < N$ then (2.3) holds everywhere in $(R_m \setminus B) \times R_{mN}$ and if $p > N$ (or $N = 1$ and $p > 1$) then (2.4) holds everywhere in $(R_m \setminus B) \times R_{mN}$. Then S_g maps $W_{1,p}(\Omega)^m$ into $L_r(\Omega)$.

PROOF. Let $\mathbf{w} = (w_1, \dots, w_m) \in W_{1,p}(\Omega)^m$. Denote by $B^{(i)}$ the projection of B on the ξ_i axis ($i = 1, \dots, m$). Since $B^{(i)}$ is null it follows that $\nabla w_i = 0$ a.e. in $w_i^{-1}(B^{(i)})$ (see [4]). Therefore, by (9.1), $S_g(\mathbf{w}) = 0$ a.e. in $\mathbf{w}^{-1}(B)$. In view of this fact, applying Sobolev's imbedding theorem, (2.3) and (2.4) imply that $S_g(\mathbf{w}) \in L_r(\Omega)$.

Using Theorems 2 and 5 we obtain,

THEOREM 6. *Suppose that $g: R_m \times R_{mN}$ is a Borel function satisfying (9.1) and suppose that there exists a constant K_0 such that*

$$|g(\xi, \eta)| < K_0 \limsup_{r \rightarrow 0} r^{-m} \int_{|\xi'| < r} |g(\xi + \xi', \eta)| d\xi',$$

$$\forall (\xi, \eta) \in (R_m \setminus B) \times R_{mN}, \quad (9.2)$$

where B is a null projection set.

Then the conditions stated in Theorem 2 are necessary and sufficient in order that S_g should map $W_{1,p}(\Omega)^m$ into $L_r(\Omega)$.

PROOF. We have to verify only the sufficiency of the conditions. In view of (9.2) it is clear that (2.3) or (2.4) hold everywhere in $(R_m \setminus B) \times R_{mN}$. Therefore by Theorem 5 these conditions are sufficient.

THEOREM 7. *Suppose that $g: R \times R_N \rightarrow R$ is a Borel function satisfying (9.1), such that $g(\xi, \cdot)$ is continuous in R_N for a.e. ξ in R . Then the conditions stated in Theorem 2 are necessary and sufficient for S_g to map $W_{1,p}(\Omega)$ into $L_r(\Omega)$.*

PROOF. Again we have to verify only the sufficiency of the conditions stated in Theorem 2. The continuity of g with respect to η implies that (2.3) or (2.4) holds everywhere in $(R \setminus B) \times R_N$, where B is a null subset of R . Therefore, by Theorem 5 these conditions are sufficient.

We conjecture that the result of Theorem 7 holds without the assumption that g is continuous with respect to η . At present we have established this conjecture in the case $N = 1$. However the proof is rather lengthy and we shall not present it here.

The second problem that we wish to mention is that of the continuity of the mapping T_h . It is known that if $h \in C^1(R_m)$ and h satisfies the conditions of Theorem 1 then $T_h: W_{1,p}(\Omega)^m \rightarrow W_{1,p}(\Omega)$ is continuous. However, if $m = 1$, the assumption that $h \in C^1(R_m)$ is not necessary. Indeed, in [6] we showed that, if $m = 1$, T_h is a continuous mapping of $W_{1,p}(\Omega)$ into $W_{1,r}(\Omega)$ whenever h satisfies the conditions of Theorem 1. The question of whether this result is valid when $m > 1$ is still unresolved.

10. An additional result concerning the mapping T_h . In this section we consider superposition mappings of the type (1.1) where h depends also on the space variable x . As before we shall assume that Ω is a domain in R_N satisfying the cone condition.

Let h be a real function on $\Omega \times R_m$ and set

$$(T_h \mathbf{u})(x) = h(x, \mathbf{u}(x)), \quad \forall x \in \Omega, \quad (10.1)$$

for every R_m -valued function \mathbf{u} in Ω . We shall present below an extension of

the *sufficiency* part of Theorem 1 for mappings of this type. The sufficient conditions are quite general and they may provide a good indication of the type of conditions that are also necessary in order that T_h should map $W_{1,p}(\Omega)^m$ into $W_{1,r}(\Omega)$. However the problem of obtaining a full extension of Theorem 1 to mappings of the type (10.1) is still unresolved.

The following notations will be used: $D_{x_i}v$ will denote the distribution derivative with respect to x_i of a function v in $L_1^{loc}(\Omega)$; $\partial V/\partial \xi_i$ will denote the (classical) partial derivative with respect to ξ_i of a function V in R_m .

First we bring,

LEMMA 5. *Let Ω be bounded and let p, r be two numbers such that either $1 < r \leq p < \infty$ and $p \neq N$ or $r = p = 1$. Suppose that h is a real function on $\Omega \times R_m$ satisfying the conditions:*

- (i) $h(\cdot, \xi) \in W_{1,r}(\Omega)$ for every ξ in R_m .
- (ii) $h(x, \cdot)$ is locally Lipschitz in R_m for a.e. x in Ω .

In addition, if $p < N$ assume that:

- (iii) *There exists a countable set E , dense in R_m , such that*

$$|D_{x_i}h(x, \xi)| \leq f(x) + b|\xi|^{q/r}, \quad \forall \xi \in E, \text{ a.e. } x \text{ in } \Omega, i = 1, \dots, N, \quad (10.2)$$

where $f \in L_r(\Omega)$, b is a constant and $q = Np/(N - p)$.

- (iv) *For a.e. x in Ω ,*

$$|(\partial h/\partial \xi_i)(x, \xi)| \leq f^*(x) + b^*|\xi|^r, \quad \text{a.e. in } R_m, i = 1, \dots, m, \quad (10.3)$$

where $f^* \in L_s(\Omega)$ with $s = pr/(p - r)$, b^* is a constant and $v = N(p - r)/(r(N - p))$. (The exceptional null set in R_m where (10.3) does not hold may vary with x .)

If $p > N$ or $p \geq 1$ and $N = 1$, assume that the following conditions hold for every $a > 0$. (Here E and s will be as before.)

- (iii)' *There exists $f_a \in L_r(\Omega)$ such that*

$$|D_{x_i}h(x, \xi)| \leq f_a(x), \quad i = 1, \dots, N, \quad (10.4)$$

for every ξ in E such that $|\xi| \leq a$ and a.e. x in Ω .

- (iv)' *For a.e. x in Ω ,*

$$|(\partial h/\partial \xi_i)(x, \xi)| \leq f_a^*(x), \quad \text{a.e. in } \{\xi \in R_m: |\xi| \leq a\}, i = 1, \dots, m, \quad (10.5)$$

where $f_a^* \in L_s(\Omega)$.

Under these assumptions T_h maps $W_{1,p}(\Omega)^m$ into $W_{1,r}(\Omega)$ and the mapping is bounded. Furthermore, the following pointwise estimates hold for $u \in W_{1,p}(\Omega)^m$.

If $p < N$ then

$$\begin{aligned} |T_h u(x)| &\leq \text{const}(|h(x, 0)| + f^*(x)|u(x)| + |u(x)|^{r+1}); \\ |D_{x_i}(T_h u)(x)| &\leq \text{const}(|f(x)| + |u(x)|^{q/r} + (f^*(x) + |u(x)|^r)|D_{x_i}u|), \end{aligned} \quad (10.6)$$

a.e. in Ω , $i = 1, \dots, N$. If $p > N$ or $p \geq 1$ and $N = 1$, and $\text{ess sup}_\Omega |u(x)| < a$ then

$$\begin{aligned} |T_h u(x)| &\leq \text{const}(|h(x, 0)| + f_a^*(x)|u(x)|); \\ |D_{x_i}(T_h u)(x)| &\leq \text{const}(f_a(x) + f_a^*(x)|D_{x_i} u|) \end{aligned} \tag{10.7}$$

a.e. in Ω , $i = 1, \dots, N$. The constants depend only on m , and b^* .

REMARK 1. In the case $r = 1$, $r < p$ the lemma remains valid if we make the following additional assumptions:

$$\begin{aligned} h &\in C(\Omega \times R_m); h(x, \cdot) \text{ is locally Lipschitz for every } x \text{ in } \Omega; \\ (10.3) \text{ and } (10.5) &\text{ hold for every } x \text{ in } \Omega \text{ and } f_a^*, f_a^* \text{ are finite} \\ &\text{everywhere in } \Omega. \end{aligned}$$

REMARK 2. If Ω is an arbitrary domain in R_N , the assumptions of the lemma imply that for every u in $W_{1,p}(\Omega)^m$, the function $T_h u$ is in $W_{1,r}^{loc}(\Omega)$ and inequalities (10.6), (10.7) hold a.e. in Ω . This is obtained by applying the lemma in an appropriate neighborhood of each point x in Ω .

Using this fact, together with Sobolev's imbedding theorem, we conclude that the lemma is valid also for unbounded domains satisfying the cone condition, provided that $h(x, 0) = 0$ a.e. in Ω .

The lemma is essentially a consequence of [5, Theorem 2.1 and Corollary 2.1] (for the case $1 < r \leq p < \infty$) and [5, Theorem 3.1] (for the case $r = p = 1$). However some details in conditions (iii), (iv) (or(iii)', (iv)') do not formally fit in the general framework of [5]; therefore a few comments are in order. In these comments we shall refer to Theorem 2.1 of [5] which deals with the case $1 < r \leq p < \infty$; but they apply equally well to Theorem 3.1 of [5] which deals with the case $r = p = 1$.

Conditions (i) and (ii) of the lemma are stronger than the parallel conditions in [5, Theorem 2.1]. However condition (iii) (resp. (iv)) is weaker in some respect than the parallel condition I (resp. II) of [5, Theorem 2.1].

In (iii) we require that the inequality holds for every ξ in a countable dense set in R_m , while in I it is required that the parallel inequality holds for every ξ in R_m . An examination of the proof of Theorem 2.1 of [5] reveals that condition I was used, first in order to show that $h(\cdot, \xi) \in W_{1,r}(\Omega)$ for every ξ in R_m (a fact which we assume in the lemma), and secondly in order to obtain an estimate for $|h(x', \xi) - h(x'', \xi)|$, namely inequality (2.19) of [5]. However, for the second application, condition I was used only with respect to a dense countable set in R_m , as we assume here in condition (iii).

In (iv) we require that the inequality holds (for a.e. x in Ω) a.e. in R_m , while in II it is required that the parallel inequality holds (for a.e. x in Ω) at every point ξ where $\partial h / \partial \xi_i$ exists. In the proof of Theorem 2.1 of [5] condition II

was used in order to derive an estimate for $|h(x, \xi) - h(x, \xi')|$, namely inequality (2.15) of [5]. In the special case that is considered in the lemma this estimate will be

$$|h(x, \xi) - h(x, \xi')| \leq \text{const}(f^*(x) + |\xi|^r + |\xi'|^r)|\xi - \xi'|, \quad (10.8)_1$$

for all ξ, ξ' in R_m and every x in $\Omega \setminus M$ (M a null set), when $p < N$;

$$|h(x, \xi) - h(x, \xi')| \leq \text{const} f_a^*(x)|\xi - \xi'|, \quad (\forall a > 0), \quad (10.8)_2$$

for all ξ, ξ' bounded in norm by a and every x in $\Omega \setminus M$, when $p > N$,

where the constants depend only on m, b and b^* . However (10.8)₁ (resp. (10.8)₂) can be derived from conditions (ii) and (iv) (resp. (iv)') of the lemma, by the same argument that was used in §7 to prove inequality (7.5).

We note that inequalities (10.6) and (10.7) follow from (10.8) (for $T_h u$) and from [5, Corollary 2.1] (for $D_{x_i}(T_h u)$). Finally the statement made in Remark 1 follows from [5, Theorem 3.3] and our previous comments here. In fact the additional assumptions mentioned in this remark are stronger than those required in [5].

Using Lemma 5 one can derive a more refined set of sufficient conditions for T_h to map $W_{1,p}(\Omega)^m$ into $W_{1,r}(\Omega)$. The next theorem gives such a set of conditions.

THEOREM 8. *Let Ω be a domain in R_N satisfying the cone condition and let p, r be two numbers such that either $1 < r \leq p < \infty$ and $p \neq N$ or $r = p = 1$. Suppose that h is a real continuous function on $\Omega \times R_m$ satisfying conditions (i) and (ii) of Lemma 5. If Ω is unbounded suppose also that $h(x, 0) = 0$ everywhere in Ω . Further, if $p < N$, suppose that there exist*

- a bounded sequence $\{f_n\}$ in $L_r(\Omega)$,*
- a bounded sequence $\{f_n^*\}$ in $L_s(\Omega)$ with $s = rp / (p - r)$,*
- an increasing sequence of open subsets of Ω , say $\{\Omega_n\}$,*

such that

(A) $v \in W_{1,r}(\Omega) \Leftrightarrow v \in W_{1,r}(\Omega')$ where $\Omega' = \cup_1^\infty \Omega_n$,

(B) h satisfies conditions (iii), (iv) of Lemma 5 in every set Ω_n , with f and f^* replaced by f_n and f_n^* and with constants b and b^* independent of n .

Finally, if $p > N$ or $p \geq 1$ and $N = 1$, suppose that for every $a > 0$ there exist sequences $\{f_{n,a}\}$ and $\{f_{n,a}^*\}$ as before such that (with $\{\Omega_n\}$ as before) h satisfies conditions (iii)', (iv)' in every set Ω_n with f_a, f_a^* replaced by $f_{n,a}, f_{n,a}^*$. Suppose also that condition (A) holds.

Then T_h maps $W_{1,p}(\Omega)^m$ into $W_{1,r}(\Omega)$ and the mapping is bounded.

It is known that, if $r > 1$ and the set $\Omega \setminus \Omega'$ is $(1, r)$ polar and $(1, r')$ polar, then condition (A) holds (see e.g. [3, p. 22]).

PROOF. Let $u \in W_{1,p}(\Omega)^m$. Applying Lemma 5 and Remark 2 to each of the sets Ω_n , $n = 1, 2, \dots$, we conclude that $T_h u \in W_{1,r}^{loc}(\Omega')$ and that $T_h u$ satisfies the inequalities (10.6) or (10.7) with $\Omega, f, f^* \dots$ replaced by $\Omega_n, f_n, f_n^* \dots$. Note that the constants appearing in these inequalities will be independent of n . Further, applying Sobolev's imbedding theorem to the space $W_{1,p}(\Omega)$ and using (10.6) or (10.7) we conclude that $T_h u \in W_{1,r}(\Omega')$ and

$$\begin{aligned} \|T_h u\|_{W_{1,r}(\Omega')} &< \text{const}(1 + \|u\|_{W_{1,p}(\Omega)}^{r+1}), & \text{if } p < N, \\ \|T_h u\|_{W_{1,r}(\Omega')} &< \text{const}(1 + \|u\|_{W_{1,p}(\Omega)}^m), & \text{if } p > N \text{ or } p > 1 \text{ and } N = 1. \end{aligned} \tag{10.9}$$

Finally, in view of condition (A), $T_h u \in W_{1,r}(\Omega)$.

REMARK 3. Using Remark 1 it is easily seen that the result of the theorem remains valid in the case $1 = r < p < \infty$ if we assume the additional conditions mentioned in that remark.

Appendix. The purpose of this Appendix is to give a proof of Lemma 3 (§4). First we introduce some notations. Let k, N be two positive integers, $k < N$. If $x = (x_1, \dots, x_N)$ is a point in R_N we shall denote it also by (x', x'') where $x' = (x_1, \dots, x_k)$ and $x'' = (x_{k+1}, \dots, x_N)$.

Lemma 3 is a consequence of the following more general result.

PROPOSITION. Suppose that f is a measurable function in R_N such that $f(\cdot, x'') \in L_1^{loc}(R_k)$ for a.e. x'' in R_{N-k} . Let $P_{x''}$ denote the set of regular Lebesgue points of $f(\cdot, x'')$ and set $P = \cup \{P_{x''}: x'' \in R_{N-k}\}$. Then P is a set of total measure in R_N .

REMARK. This proposition is probably known, but we do not know of a reference for it in this general form. (In the case where f is the characteristic function of a measurable set, a proof can be found in [8].) Therefore we give its proof here.

PROOF. For a.e. x'' in R_{N-k} , $P_{x''}$ is a set of total measure in R_k . Therefore, to prove the proposition we have to show only that P is measurable. Given $\rho < 0$ and x in R_N , set

$$F(x, \rho) = \mu_k(B'(x, \rho))^{-1} \int_{B'(x, \rho)} f(y, x'') d\mu_k(y) \tag{A.1}$$

where $B'(x, \rho) = \{y \in R_k: |y - x'| < \rho\}$. This formula can be rewritten in the form

$$F(x, \rho) = c_k \int_{B'_1} f(x' + \rho y, x'') d\mu_k(y), \tag{A.2}$$

where B'_1 is the unit ball, centered at the origin, in R_k and c_k^{-1} is the volume

of B'_1 . The function $\tilde{f}_\rho: R_N \times R_k \rightarrow R$ given by $\tilde{f}_\rho(x', x'', y) = f(x' + \rho y, x'')$ is obviously measurable, for every ρ . Therefore, applying Fubini's theorem to the positive part and the negative part of \tilde{f}_ρ , we conclude that $F(\cdot, \rho)$ is measurable in R_N , for every $\rho > 0$. Therefore the set

$$Q = \left\{ x \in R_N: f(x) = \lim_{\substack{\rho \rightarrow 0 \\ \rho \text{ rational}}} F(x, \rho) \right\}$$

is measurable. Obviously $Q \supseteq P$; but if x'' is a point in R_{N-k} such that $f(\cdot, x'') \in L_1^{\text{loc}}(R_k)$ then $F(x, \cdot)$ is continuous for ρ in $(0, \infty)$ and so

$$Q_{x''} = \{x' \in R_k | (x', x'') \in Q\} = P_{x''}.$$

Since this equality holds for a.e. x'' in R_{N-k} it follows that $Q \setminus P$ is a null set and hence that P is measurable.

REFERENCES

1. H. Federer, *Geometric measure theory*, Die Grundlehren der Math. Wissenschaften, Band 153, Springer-Verlag, New York, 1969.
2. E. Gagliardo, *Proprietà di alcune classi di funzioni in piu variabili*, Ricerche Mat. 7 (1958), 102-137.
3. J. L. Lions, *Problèmes aux limites dans les equations aux dérivées partielles*, Presses Univ. Montréal, Montréal, 1965.
4. M. Marcus and V. J. Mizel, *Absolute continuity on tracks and mappings of Sobolev spaces*, Arch. Rational Mech. Anal. 45 (1972), 294-320.
5. _____, *Nemitsky operators on Sobolev spaces*, Arch. Rational Mech. Anal. 51 (1973), 347-370.
6. _____, *Every superposition operator which maps one Sobolev space into another is continuous*, J. Functional Anal. (to appear).
7. C. B. Morrey, *Multiple integrals in the calculus of variations*, Die Grundlehren der Math. Wissenschaften, Band 130, Springer-Verlag, Berlin and New York, 1964.
8. S. Saks, *Theory of the integral*, 2nd ed., Dover, New York, 1964.
9. G. Stampacchia, *Equations elliptiques du second ordre à coefficients discontinus*, Univ. of Montreal Press, Montreal, Quebec, 1966.

DEPARTMENT OF MATHEMATICS, ISRAEL INSTITUTE OF TECHNOLOGY, TECHNION, HAIFA, ISRAEL

DEPARTMENT OF MATHEMATICS, CARNEGIE-MELLON UNIVERSITY, PITTSBURGH, PENNSYLVANIA 15213