

DERIVATIONS AND AUTOMORPHISMS OF NONASSOCIATIVE MATRIX ALGEBRAS¹

BY

G. M. BENKART AND J. M. OSBORN

ABSTRACT. This paper studies the derivation algebra and the automorphism group of $M_n(A)$, $n \times n$ matrices over an arbitrary nonassociative algebra A with multiplicative identity 1. The investigation also includes results on derivations and automorphisms of the algebras obtained from $M_n(A)$ using the Lie product $[xy] = xy - yx$, and the Jordan product $x \circ y = \frac{1}{2}(xy + yx)$.

1. Introduction. Results concerning automorphisms and derivations of $n \times n$ matrices over a finite dimensional central associative division algebra have been known for some time. The Noether-Skolem theorem combined with a theorem due to Jacobson gives that automorphisms as well as derivations of these algebras are inner. This present paper generalizes these theorems by considering derivations and automorphisms of $M_n(A)$, $n \times n$ matrices over an arbitrary nonassociative algebra A with 1. The investigation also determines the derivations and automorphisms of the algebras obtained from $M_n(A)$ using the Lie and Jordan products. With one exception the derivation algebras are shown to all follow the same pattern. They consist of inner derivations by matrices with entries in the nucleus N of A , and of derivations gotten by applying derivations of A to each matrix entry. Every derivation is the sum of these two kinds of derivations. More variability arises in the automorphism groups. The common feature is a subgroup composed of conjugations by invertible matrices in $M_n(N)$, and of automorphisms obtained by applying automorphisms of A entry by entry. Each element of this subgroup is a product of these two kinds of maps.

Our motivation to investigate such derivation algebras and automorphism groups was due in part to conversations with physicists concerned with building algebraic models in particle theories. Of particular interest to them were the cases when A was taken to be an octonion algebra or a certain 7-dimensional noncommutative Jordan algebra [1]. The results we present here have already been used in a negative sense to exclude some algebras from their considerations.

The main results concerning the group of antiautomorphisms and automorphisms of $M_n(A)$ are contained in Corollary 3.14. Theorem 4.8 and Corollaries 4.9, 4.10 describe the derivation algebras of $M_n(A)$, $L_n(A)$, $L'_n(A)$, and $K_n(A)$ where

Received by the editors October 20, 1978.

AMS (MOS) subject classifications (1970). Primary 16A72, 16A68, 17B40, 17C30, 17E05.

Key words and phrases. Derivations, automorphisms, antiautomorphisms, nonassociative algebras, nonassociative matrix algebras.

¹Partially supported by N.S.F. grant MCS 77-01724.

$L_n(A)$ is $M_n(A)$ under the product $[\]$, $L'_n(A) = [L_n(A), L_n(A)]$ and $K_n(A)$ is $L'_n(A)$ with its center factored out. We present results for the automorphism groups of $L_n(A)$, $L'_n(A)$, and $K_n(A)$ in Theorems 4.12 and 4.13, and Corollary 4.14. Finally, if $J_n(A)$ denotes $M_n(A)$ viewed under the Jordan product, then its derivation algebra is obtained in Theorem 5.5, and its automorphism group is described in Theorem 5.6.

2. Basic concepts. Let A be an arbitrary nonassociative algebra with identity 1 over the field Φ , and let N denote the nucleus of A . That is

$$N = \{a \in A \mid (a, b, c) = (b, a, c) = (b, c, a) = 0 \text{ for all } b, c \in A\}$$

where $(a, b, c) = (ab)c - a(bc)$. The algebra N is associative, and it is not difficult to show that if $a \in N$, then $\text{ad}_a: A \rightarrow A$ given by $\text{ad}_a(b) = [ab] = ab - ba$ is a derivation of A .

In the same manner we speak of the nucleus of $M_n(A)$. An easy calculation shows that it is $M_n(N)$, and as above for $x \in M_n(N)$, ad_x is a derivation of $M_n(A)$. If $\text{Der } M_n(A)$ denotes the derivation algebra of $M_n(A)$, then $\text{ad}_{M_n(N)} = \{\text{ad}_x \mid x \in M_n(N)\}$ is a subalgebra of $\text{Der } M_n(A)$. Let e_{ij} denote the matrix with 1 in the (i, j) position and 0 elsewhere. Then these matrix units lie in $M_n(N)$, and the derivations $\text{ad}_{e_{ij}}$ will be particularly useful in what follows.

In addition to the inner derivations $\text{ad}_{M_n(N)}$ one can also obtain a derivation of $M_n(A)$ by beginning with a derivation of A and applying it to each matrix entry. The resulting derivations form a subalgebra of $\text{Der } M_n(A)$, which we denote by $(\text{Der } A)^\#$. The notation is suggestive of the fact that $(\text{Der } A)^\#$ is isomorphic to $\text{Der } A$. Similarly one can obtain an automorphism of $M_n(A)$ by applying an automorphism of A to each entry. The resulting set of automorphisms is a subgroup, called $(\text{Aut } A)^\#$, of the full automorphism group $\text{Aut } M_n(A)$. If $u \in M_n(N)$ is invertible then conjugation by u , denoted by χ_u ($\chi_u(x) = u^{-1}xu$), belongs to $\text{Aut } M_n(A)$. We write $\text{GL}(n, N)$ for the subgroup composed of the mappings χ_u .

One final note, let us observe that if $\varphi \in \text{Aut } M_n(A)$ or $\varphi \in \text{Der } M_n(A)$, then φ maps $M_n(N)$ into $M_n(N)$. This is true since $M_n(N)$ is characterized as the nucleus of $M_n(A)$, and derivations and automorphisms preserve the nucleus.

3. Generalized automorphisms and antiautomorphisms of $M_n(A)$. Given any algebra A with nucleus N let $Z(A) = \{a \in N \mid [ab] = 0 \text{ for all } b \in A\}$ denote the center of A . We consider linear transformations φ of A onto A with $\varphi(1) = 1$ and with the following property:

There exist idempotents $f, g \in Z(A)$ such that $f + g = 1$, $fg = 0$, and if $f' = \varphi(f)$, $g' = \varphi(g)$, then $f', g' \in Z(A)$ and $\varphi: fA \rightarrow f'A$ is an algebra isomorphism and $\varphi: gA \rightarrow g'A$ is an anti-isomorphism.

LEMMA 3.1. *The set of all such mappings, $\text{GAut } A$, is a group.*

PROOF. Given $\varphi \in \text{GAut } A$, then the condition holds for φ^{-1} and the idempotents f', g' . Suppose now φ_1 and φ_2 belong to $\text{GAut } A$ and the corresponding idempotents are f_1, g_1 and f_2, g_2 . Then one can verify the above property is satisfied

for $\varphi_2\varphi_1$ using the idempotents $f = f_2f'_1 + g_2g'_1$ and $g = f_2g'_1 + g_2f'_1$ where $f'_1 = \varphi_1(f_1)$ and $g'_1 = \varphi_1(g_1)$. \square

The group $\text{GAut } A$ contains two distinguished subgroups, $\text{Aut } A$ and $\text{AAut } A$, where $\text{AAut } A$ is the group consisting of all automorphisms and antiautomorphisms of A . Corresponding to each $\varphi \in \text{GAut } A$ is the decomposition of A into the direct sum of the ideals fA and gA . So if A has no proper direct summands (for example if A is simple), then $\text{GAut } A = \text{AAut } A$.

A straightforward calculation shows that each $\varphi \in \text{GAut } A$ maps the nucleus of A onto itself.

Following the procedure above we define $\text{GAut } M_n(A)$. The idempotents corresponding to elements in $\text{GAut } M_n(A)$ lie in $Z(M_n(A))$ which is $Z(A)I$, where I is the identity matrix. For each $\varphi \in \text{GAut } M_n(A)$, $\varphi(M_n(N)) = M_n(N)$ since φ preserves the nucleus, and φ restricted to $M_n(N)$ lies in $\text{GAut } M_n(N)$. In this section we investigate the structure of $\text{GAut } M_n(A)$ under the assumption that N is Artinian. As the preceding remarks indicate the place to begin is with $\text{GAut } M_n(N)$, but first some preliminary results brought to our attention by L. Levy [7, Lemma 3.2, p. 282].

LEMMA 3.2. *Let R be an associative ring with 1 and M be a right R -module such that*

$$M = M_1 \oplus \cdots \oplus M_n = M'_1 \oplus \cdots \oplus M'_n$$

are two direct sum decompositions of M into submodules such that M_i is isomorphic to M'_i for $1 \leq i \leq n$. Then there is an invertible σ in the ring of R -endomorphisms of M such that $\sigma^{-1}\pi'_i\sigma = \pi_i$ for each $1 \leq i \leq n$, where π_i (π'_i) denotes the projection of M onto M_i (M'_i).

PROOF. For each i extend the isomorphism between M_i and M'_i to an R -endomorphism of M by defining it to be 0 on the other summands. Take σ to be the sum of the resulting n endomorphisms. \square

LEMMA 3.3. *Let R be an associative ring with 1 and let $\{e_i\}$ and $\{f_i\}$ for $i = 1, \dots, n$ be two sets of orthogonal idempotents summing to 1. Assume for each i , e_iR is isomorphic to f_iR as R -modules. Then there is a unit $u \in R$ with $u^{-1}f_iu = e_i$ for $i = 1, \dots, n$.*

PROOF. Apply the preceding lemma to

$$R = e_1R \oplus \cdots \oplus e_nR = f_1R \oplus \cdots \oplus f_nR.$$

The projection π_i (π'_i) is just left multiplication by e_i (f_i). The R -isomorphism σ is left multiplication by $u = \sigma(1)$, and σ^{-1} is left multiplication by u^{-1} . Then

$$e_i = \pi_i(1) = \sigma^{-1}\pi'_i\sigma(1) = \sigma^{-1}\pi'_i(u) = \sigma^{-1}(f_iu) = u^{-1}f_iu. \quad \square$$

Given $\varphi \in \text{GAut } M_n(A)$ direct verification shows that φ is an automorphism of $J_n(A)$, where $J_n(A)$ is $M_n(A)$ under the Jordan product $x \circ y = \frac{1}{2}(xy + yx)$ provided $\text{char } \Phi \neq 2$. It is convenient for what follows in §5 to work with the larger group $\text{Aut } J_n(A)$. Though we disallow $\text{char } \Phi = 2$ in this process, this restriction is unnecessary in dealing with $\text{GAut } M_n(A)$.

Jacobson [4, Theorem 7.4, p. 26] has shown that as a consequence of results in either [4] or [6] the following theorem holds:

Let A and B be arbitrary algebras with identities, and let B_J be the algebra obtained from B under the Jordan product. Suppose $\psi: J_n(A) \rightarrow B_J$ is a Jordan homomorphism such that the images of the matrix units e_{ij} lie in the nucleus of the algebra C generated by $\psi(J_n(A))$. Then $C = C_1 \oplus C_2$ where the C_i are ideals of C , and if π_i denotes the projection of C onto C_i , then $\pi_i\psi$ is a homomorphism, and $\pi_2\psi$ an antihomomorphism.

We could use this theorem to reduce our considerations to automorphisms and antiautomorphisms. However, we do not follow this line of attack because one virtue of working with $\text{GAut } M_n(A)$ is that it allows us to study automorphisms and antiautomorphisms simultaneously, and because we can obtain in the process results concerning automorphisms of $J_n(A)$ with the property that $\psi(e_{ii}) = e_{ii}$ for all i , without assuming from the outset that all the $\psi(e_{ij})$ lie in the nucleus.

LEMMA 3.4. *Assume $\text{char } \Phi \neq 2$, and let $\varphi \in \text{Aut } J_n(N)$, where N is an associative Artinian ring with 1. Then there is an invertible $u \in M_n(N)$ such that*

$$u^{-1}\varphi(e_{ii})u = e_{ii}.$$

PROOF. The elements $\varphi(e_{ii})$ for $i = 1, \dots, n$ are orthogonal idempotents summing to 1. Thus,

$$\begin{aligned} M_n(N) &= \varphi(e_{11})M_n(N) \oplus \cdots \oplus \varphi(e_{nn})M_n(N) \\ &= e_{11}M_n(N) \oplus \cdots \oplus e_{nn}M_n(N). \end{aligned}$$

Jacobson and Rickart [6, Lemma 3, p. 487] have shown that there are matrix units $\{g_{ij}\}$ and $\{h_{ij}\}$ such that $\varphi(e_{ij}) = g_{ij} + h_{ij}$ and $g_{ij}h_{kl} = 0 = h_{kl}g_{ij}$ for all i, j, k, l . Therefore, $\varphi(e_{ii})M_n(N)$ is isomorphic as an $M_n(N)$ -module to $\varphi(e_{ij})M_n(N)$ via left multiplication by $g_{ji} + h_{ji}$. Thus, $M_n(N)$ is isomorphic to the direct sum of n -copies of $\varphi(e_{ii})M_n(N)$, and we write this $M_n(N) \approx \varphi(e_{ii})M_n(N)^{(n)}$. Similarly $e_{ii}M_n(N)$ is isomorphic as a right $M_n(N)$ -module to $e_{jj}M_n(N)$ using left multiplication by e_{ji} . Hence, $e_{ii}M_n(N)^{(n)} \approx M_n(N) \approx \varphi(e_{ii})M_n(N)^{(n)}$. Breaking $e_{ii}M_n(N)$ into a finite number of indecomposable $M_n(N)$ -modules, and collecting isomorphic indecomposables, one can use the Krull-Schmidt theorem to argue that $e_{ii}M_n(N)$ is isomorphic to $\varphi(e_{ii})M_n(N)$. Thus by the preceding lemma, there is an invertible $u \in M_n(N)$ so that $u^{-1}\varphi(e_{ii})u = e_{ii}$ as desired. \square

COROLLARY 3.5. *Let A be an algebra with 1 such that the nucleus N of A is Artinian. Then for each $\varphi \in \text{GAut } M_n(A)$, there is an invertible $u \in M_n(N)$ so that $\psi = \chi_u\varphi \in \text{GAut } M_n(A)$ has the property that $\psi(e_{ii}) = e_{ii}$ for $i = 1, \dots, n$.*

Automorphisms of $J_n(A)$ with the property that $\psi(e_{ii}) = e_{ii}$ will be the topic of the next lemma, but one additional piece of notation is necessary.

Suppose $\theta \in \text{GAut } A$ with idempotents f, g . Define $\theta^\# : M_n(A) \rightarrow M_n(A)$ by letting $\theta^\#$ on $fI \cdot M_n(A)$ be θ applied to each entry, and letting $\theta^\#$ on $gI \cdot M_n(A)$ be θ applied to each entry followed by taking the transpose of the resulting matrix.

One can show $\theta^\# \in \text{GAut } M_n(A)$. Let $(\text{GAut } A)^\#$ denote the group of these mappings, and observe it is isomorphic to the group $\text{GAut } A$.

THEOREM 3.6. *Assume $\text{char } \Phi \neq 2$, and suppose $\psi \in \text{Aut } J_n(A)$ with $\psi(e_{ii}) = e_{ii}$ for all $i = 1, \dots, n$. Then there is an invertible $v \in M_n(N)$ such that $\chi_v \psi \in (\text{GAut } A)^\#$.*

PROOF. There are only five types of nonzero products in $J_n(A)$, and for i, j, k distinct they are

$$ae_{ii} \circ be_{ii} = (a \circ b)e_{ii}, \quad (3.1)$$

$$ae_{ii} \circ be_{ij} = \frac{1}{2}abe_{ij}, \quad (3.2)$$

$$ae_{ij} \circ be_{jj} = \frac{1}{2}abe_{ij}, \quad (3.3)$$

$$ae_{ij} \circ be_{jk} = \frac{1}{2}abe_{ik}, \quad (3.4)$$

$$ae_{ij} \circ be_{ji} = \frac{1}{2}(abe_{ii} + bae_{jj}). \quad (3.5)$$

We apply ψ to these relations, and make different specializations of a and b .

Relation (3.1) with $b = 1$ says $\psi(ae_{ii}) = \psi(ae_{ii}) \circ e_{ii}$. But the first three equations above demonstrate the fact that the only elements of $J_n(A)$ which lie in the 1-eigenspace relative to right multiplication by e_{ii} are of the form ce_{ii} . Therefore

$$\psi(ae_{ii}) = \psi_i(a)e_{ii}. \quad (3.6)$$

Again by using (3.1) we see ψ_i is an automorphism of A under the product $a \circ b = \frac{1}{2}(ab + ba)$.

From (3.2) and (3.3) it follows that $\psi(ae_{ij})$ must lie in $Ae_{ij} + Ae_{ji}$, which is the intersection of the $\frac{1}{2}$ -eigenspace relative to multiplication by e_{ii} with the $\frac{1}{2}$ -eigenspace relative to multiplication by e_{jj} . As a special case of this fact:

$$\psi(e_{ij}) = \alpha_{ij}e_{ij} + \beta_{ji}e_{ji}. \quad (3.7)$$

It follows directly from $e_{ij} \circ e_{ij} = 0$ that

$$\alpha_{ij}\beta_{ji} = 0 = \beta_{ji}\alpha_{ij}. \quad (3.8)$$

Now using (3.4) and (3.5) with $a = b = 1$, we deduce further results concerning the α 's and β 's, namely

$$\alpha_{ij}\alpha_{jk} = \alpha_{ik}, \quad \beta_{kj}\beta_{ji} = \beta_{ki}, \quad (3.9)$$

$$\alpha_{ij}\alpha_{ji} + \beta_{ij}\beta_{ji} = 1. \quad (3.10)$$

If ψ belongs to $\text{GAut } M_n(A)$, then it is immediate that $\alpha_{ij}, \beta_{ji} \in N$. We do not make this assumption in order to use Theorem 3.6 in §5. Instead the fact that $\alpha_{ij}, \beta_{ji} \in N$ will be a consequence of the next few steps.

LEMMA 3.7. *α_{ij}, β_{ji} lie in the middle nucleus of A for all $i \neq j$.*

PROOF. If ψ is applied to the relation $0 = (ae_{ii} \circ e_{ij}) \circ be_{jj} - ae_{ii} \circ (e_{ij} \circ be_{jj})$, then the (i, j) component of the resulting equation shows that the associator $(\psi_i(a), \alpha_{ij}, \psi_j(b)) = 0$, and the (j, i) component shows that $(\psi_j(b), \beta_{ji}, \psi_i(a)) = 0$. Since ψ_i, ψ_j are onto, the proof is complete. \square

Let us define $f_i = \alpha_{ij}\alpha_{ji}$ and $g_i = \beta_{ij}\beta_{ji}$. Note as a result of Lemma 3.7 and (3.9), if $n \geq 3$ then

$$f_i = \alpha_{ij}\alpha_{ji} = \alpha_{ik}\alpha_{kj}\alpha_{ji} = \alpha_{ik}\alpha_{ki}.$$

This shows f_i does not depend on the j used to define it as long as $j \neq i$. Similarly g_i is also independent of j .

LEMMA 3.8. *For each i , f_i and g_i are orthogonal idempotents summing to 1, and they commute with the elements of A . Moreover $f_i = f_j$, $g_i = g_j$ for all i, j .*

PROOF. Equation (3.9) says $f_i + g_i = 1$. Now

$$f_i^2 = \alpha_{ij}\alpha_{ji}\alpha_{ij}\alpha_{ji} = \alpha_{ij}(1 - \beta_{ji}\beta_{ij})\alpha_{ji} = \alpha_{ij}\alpha_{ji} = f_i,$$

so that f_i and $g_i = 1 - f_i$ are idempotents. Multiplying $f_i + g_i = 1$ on the right by f_i , then on the left, gives $f_i^2 + g_i f_i = f_i$ and $f_i^2 + f_i g_i = f_i$. Thus $g_i f_i = 0 = f_i g_i$, and they are orthogonal.

To obtain the rest of the conclusions we return to (3.2), set $b = 1$, and act on the equation with ψ . The result is a formula for $\psi(ae_{ij})$

$$\psi(ae_{ij}) = \psi_i(a)\alpha_{ij}e_{ij} + \beta_{ji}\psi_i(a)e_{ji}. \quad (3.11)$$

Using this formula let us calculate the (i, i) -coefficient of both sides of

$$2\psi(ae_{ii} + ae_{ij}) \circ \psi(e_{ji})$$

to establish that $\psi_i(a) = \psi_i(a)\alpha_{ij}\alpha_{ji} + \beta_{ij}\beta_{ji}\psi_i(a) = \psi_i(a)f_i + g_i\psi_i(a)$. Since ψ_i is onto, we can replace $\psi_i(a)$ with a to obtain

$$a = af_i + g_i a \quad \text{for all } a \in A. \quad (3.12)$$

Multiplying $1 = f_i + g_i$ on the right by a shows that $a = f_i a + g_i a$. Comparing this result with (3.12) gives $af_i = f_i a$ for all $a \in A$, and hence $g_i a = ag_i$ also.

Finally $f_i = f_i^2 = \alpha_{ij}\alpha_{ji}\alpha_{ij}\alpha_{ji} = \alpha_{ij}f_j\alpha_{ji} = \alpha_{ij}\alpha_{ji}f_j = f_j f_i$. Since this is true for each i and j , $f_j = f_j f_i$. But f_i commutes with every element of A , so $f_j = f_j f_i = f_i f_j = f_i$. Thus $g_i = 1 - f_i = 1 - f_j = g_j$ and the proof is finished. \square

We shall just write f' for f_i and g' for g_i hereafter.

LEMMA 3.9. *The elements α_{ij} and β_{ji} lie in N for every pair i, j with $i \neq j$.*

PROOF. In analogy with equation (3.11), if ψ is applied to (3.3) with $a = 1$, we obtain

$$\psi(be_{ij}) = \alpha_{ij}\psi_j(b)e_{ij} + \psi_j(b)\beta_{ji}e_{ji}. \quad (3.13)$$

We freely use these two equations in the proof of this lemma. First, let us apply ψ to $(ae_{ij} \circ be_{ii}) \circ e_{ij} - ae_{ij} \circ (be_{ii} \circ e_{ij}) = 0$. The (j, j) coefficient is

$$\begin{aligned} & ((\beta_{ji}\psi_i(a))\psi_i(b))\alpha_{ij} + \beta_{ji}(\psi_i(b)(\psi_i(a)\alpha_{ij})) \\ & - (\beta_{ji}\psi_i(a))(\psi_i(b)\alpha_{ij}) - (\beta_{ji}\psi_i(b))(\psi_i(a)\alpha_{ij}) = 0. \end{aligned}$$

Since ψ_i is onto, we can replace $\psi_i(b)$ with any element c in A , and $\psi_i(a)$ with an arbitrary element of A , so let us use $(\beta_{ij}d)g'$. As a consequence of (3.8) and Lemma 3.8, $g'\alpha_{ij} = \beta_{ij}\beta_{ji}\alpha_{ij} = 0$, and thus, $((\beta_{ij}d)g')\alpha_{ij} = 0$. Now

$$\begin{aligned} \beta_{ji}((\beta_{ij}d)g') &= \beta_{ji}(g'(\beta_{ij}d)) = (\beta_{ji}g')(\beta_{ij}d) \\ &= (g'\beta_{ji})(\beta_{ij}d) = (g'\beta_{ji}\beta_{ij})d = g'd, \end{aligned}$$

so that from this judicious choice of $\psi_i(a)$, we deduce

$$(g'd, c, \alpha_{ij}) = 0. \tag{3.14}$$

If in the above equation the substitution of $f'(d\alpha_{ji})$ for $\psi_i(a)$ is made, the result is

$$(\beta_{ji}, c, f'd) = 0. \tag{3.15}$$

To obtain the other half of the proof that α_{ij} is in the right nucleus and β_{ji} in the left, we apply ψ to the relation $(ae_{ji} \circ be_{ii}) \circ e_{ij} - ae_{ji} \circ (be_{ii} \circ e_{ij}) = \frac{1}{4}[ab]e_{ii}$. The (j, j) coefficient of the left side must be 0, and in the resulting equation for the (j, j) coefficient we perform the replacements of $\psi_i(b)$ with c , and $\psi_i(a)$ with $(\alpha_{ij}d)f'$ then with $g'(d\beta_{ji})$. The effect is

$$(f'd, c, \alpha_{ij}) = 0. \tag{3.16}$$

$$(\beta_{ji}, c, g'd) = 0. \tag{3.17}$$

From relations (3.14)–(3.17) we conclude that α_{ij} is in the right nucleus and β_{ji} in the left. The appropriate equations to use to achieve α_{ij} in the left nucleus and β_{ji} in the right are

$$\begin{aligned} (e_{ij} \circ ae_{jj}) \circ be_{ij} - e_{ij} \circ (ae_{jj} \circ be_{ij}) &= 0, \\ (e_{ij} \circ ae_{jj}) \circ be_{ji} - e_{ij} \circ (ae_{jj} \circ be_{ji}) &= \frac{1}{4}[ba]e_{jj}. \end{aligned}$$

The substitutions needed in the first equation are $g'(d\beta_{ij})$ and $(\alpha_{ji}d)f'$ for $\psi_j(b)$, and in the second $f'(d\alpha_{ji})$ and $(\beta_{ji}d)g'$ for $\psi_j(b)$. \square

COROLLARY 3.10. $f', g' \in Z(A)$.

LEMMA 3.11. *Let $f = \psi_i^{-1}(f')$ and $g = \psi_i^{-1}(g')$. Then f, g are orthogonal idempotents in the center of A which sum to 1. They are independent of the defining i . Moreover, ψ_i is an isomorphism from fA onto $f'A$, and an anti-isomorphism from gA onto $g'A$, so that $\psi_i \in \text{GAut } A$.*

PROOF. That f, g are idempotents summing to 1 follows simply from the fact that ψ_i is an automorphism relative to the \circ -product. Orthogonality is shown as it was for f', g' above. To prove the remaining statements we observe that from $\psi(abe_{ij}) = 2\psi(ae_{ii}) \circ \psi(be_{ij})$ we can establish: $\psi_i(ab)\alpha_{ij} = \psi_i(a)\psi_i(b)\alpha_{ij}$ and $\beta_{ji}\psi_i(ab) = \beta_{ji}\psi_i(b)\psi_i(a)$. Multiplication by α_{ji} and β_{ij} respectively demonstrates

$$\psi_i(ab)f' = \psi_i(a)\psi_i(b)f', \tag{3.18}$$

$$g'\psi_i(ab) = g'\psi_i(b)\psi_i(a). \tag{3.19}$$

But then equation (3.19) with $a = f$ shows $g'\psi_i(fb) = g'\psi_i(b)f' = g'f'\psi_i(b) = 0$. Hence, ψ_i maps fA into $f'A$, and it is an isomorphism on fA according to (3.18). Likewise ψ_i is an anti-isomorphism of gA into $g'A$. We use the fact that $A = fA + gA$, and the fact that ψ_i is onto to conclude it is onto the components fA and gA .

What remains to be shown is that $f, g \in Z(A)$ and f, g are independent of i . We note that $\psi_i(af) = \psi_i(af)f'$, and $\psi_i(fa) = \psi_i(fa)f'$ since f' is the identity on $f'A$.

Therefore (3.18) shows that $\psi_i(af) = \psi_i(a)f'$ and also $\psi_i(fa) = f'\psi_i(a) = \psi_i(a)f'$. Hence $\psi_i(af) = \psi_i(fa)$ for all a , and $af = fa$. To prove that f lies in the nucleus of A , it suffices to show f acts as the identity on fA and $fA \cdot gA = 0$. But this can be accomplished by using (3.18) with af instead of a , and f instead of b , and by using (3.19) with af in place of a , and bg in place of b . Thus, $f \in Z(A)$ and $g = 1 - f \in Z(A)$ also.

Now relations (3.11) and (3.13) demonstrate

$$\alpha_{ij}\psi_j(a) = \psi_i(a)\alpha_{ij}, \tag{3.20}$$

$$\psi_j(a)\beta_{ji} = \beta_{ji}\psi_i(a). \tag{3.21}$$

These equations show that $f'\psi_j(a) = \alpha_{ji}\psi_i(a)\alpha_{ij}$ and $\psi_j(a)g' = \beta_{ji}\psi_i(a)\beta_{ij}$. Substituting $a = f$ we see $f'\psi_j(f) = f'$ and $\psi_j(f)g' = 0$. Thus $\psi_j(f) = f'$ and we are done. \square

LEMMA 3.12. *There is an invertible diagonal matrix v in $M_n(N)$ such that $\chi_v\psi = \psi_1^\#$.*

PROOF. It is an easy consequence of (3.8) that $\beta_{ji}f'A = f'AB_{ji} = g'AB_{ji} = \alpha_{ij}g'A = 0$ for all $i \neq j$, and from the remarks immediately following equations (3.20) and (3.21), it follows that $\psi_j(fa) = f'\psi_j(fa) = \alpha_{j1}\psi_1(fa)\alpha_{ij}$ and $\psi_j(ga) = g'\psi_j(ga) = \beta_{j1}\psi_1(ga)\beta_{1j}$. Therefore writing $a = fa + ga$, we conclude from these observations that

$$\psi_j(a) = (\alpha_{j1} + \beta_{j1})\psi_1(a)(\alpha_{1j} + \beta_{1j}) \quad \text{for all } a. \tag{3.22}$$

Let v denote the matrix with 1 in the $(1, 1)$ position, $\alpha_{j1} + \beta_{j1}$ in the (j, j) slot for $j \geq 2$, and 0 elsewhere. Then v^{-1} is also diagonal with diagonal entries 1, $\alpha_{12} + \beta_{12}, \dots, \alpha_{1n} + \beta_{1n}$. Thus, $\chi_v\psi(ae_{jj}) = v^{-1}\psi_j(a)e_{jj}v = \psi_1(a)e_{jj}$. Calculation using (3.8), (3.10) and (3.22) shows

$$\begin{aligned} \chi_v\psi(ae_{jj}) &= (\alpha_{1i} + \beta_{1i})(\alpha_{i1} + \beta_{i1})\psi_1(a)(\alpha_{1i} + \beta_{1i})\alpha_{ij}(\alpha_{j1} + \beta_{j1})e_{ij} \\ &\quad + (\alpha_{1j} + \beta_{1j})\beta_{ji}(\alpha_{i1} + \beta_{i1})\psi_1(a)(\alpha_{1i} + \beta_{1i})(\alpha_{i1} + \beta_{i1})e_{ji} \\ &= \psi_1(a)\alpha_{1j}\alpha_{j1}e_{ij} + \beta_{1i}\beta_{i1}\psi_1(a)e_{ji} \\ &= \psi_1(a)f'e_{ij} + g'\psi_1(a)e_{ji} \\ &= \psi_1(fa)e_{ij} + \psi_1(ga)e_{ji}. \end{aligned}$$

Hence, $\chi_v\psi = \psi_1^\# \in (\text{GAut } A)^\#$, and this concludes the proof of Lemma 3.12, and Theorem 3.6. \square

As a consequence of this theorem and Corollary 3.5 we have

THEOREM 3.13. *Let A be an arbitrary nonassociative algebra with 1 such that the nucleus N of A is Artinian. If $\text{char } \Phi \neq 2$, then*

$$\text{GAut } M_n(A) = \text{GL}(n, N) \cdot (\text{GAut } A)^\#.$$

REMARK. It should be noted that the subgroup $\text{GL}(n, N)$ of conjugations is a normal subgroup of $\text{GAut } M_n(A)$, hence of $\text{Aut } M_n(A)$ and of $\text{AAut } M_n(A)$ as well,

and it equals

$$\text{GAut}_A M_n(A) = \{ \varphi \in \text{GAut } M_n(A) \mid \varphi(aI) = aI \text{ for all } a \in A \}$$

if $Z(A) = N$.

COROLLARY 3.14. *With assumptions as in Theorem 3.13,*

$$\text{Aut } M_n(A) = \text{GL}(n, N) \cdot (\text{Aut } A)^\#$$

and

$$\text{AAut } M_n(A) = \text{GL}(n, N) \cdot (\text{AAut } A)^\#,$$

so that $M_n(A)$ has antiautomorphisms if and only if A does.

PROPOSITION 3.15. *If N is Artinian, $\text{Aut } M_n(A)$ has finite index in $\text{GAut } M_n(A)$.*

PROOF. Since N is Artinian with 1, we can find orthogonal central idempotents e_1, \dots, e_q such that $1 = e_1 + \dots + e_q$ and each e_i cannot be expressed as the sum of two other central idempotents. Then $M_n(N) = \bigoplus \sum_{i=1}^q S_i$ where $S_i = e_i I \cdot M_n(N)$ is an indecomposable ideal of $M_n(N)$. For any ideal T of $M_n(N)$, $T = \bigoplus \sum_{i=1}^q e_i I \cdot T$, so if T is indecomposable, $T = S_j$ for some j . Given $\varphi \in \text{GAut } M_n(A)$ with idempotents $fI, gI, f'I, g'I$, then $fe_i + ge_i = e_i = f'e_i + g'e_i$ implies $fI \cdot M_n(N), gI \cdot M_n(N), f'I \cdot M_n(N), g'I \cdot M_n(N)$ are all just the sums of certain of the S_i . We conclude that each $\varphi(S_i)$ is an indecomposable ideal of $M_n(N)$ and so equals some S_j . Thus, φ permutes the summands and the subgroup of generalized automorphisms fixing all the S_i has index at most $q!$ in $\text{GAut } M_n(A)$. Since every ψ in that subgroup acts as an automorphism or antiautomorphism on each S_i , the index of $\text{Aut } M_n(A)$ in $\text{GAut } M_n(A)$ is at most $q!2^q$. \square

The next example demonstrates that Corollary 3.14 is false for N arbitrary.

Let $\Phi[t]$ denote the ring of polynomials over Φ , and let B be the ideal of $\Phi[t]$ generated by $t^2 + 1$. Define A to be $\{(a, b) \mid a, b \in \Phi[t] \text{ and } a \equiv b \pmod{B}\}$. If w denotes the matrix

$$\begin{pmatrix} (1, -t) & (0, 1 + t^2) \\ (t, 1) & (1, -t) \end{pmatrix},$$

then w lies in $M_2(\Phi[t] \times \Phi[t])$ but not in $M_2(A)$. The matrix w is invertible, and conjugation by w leaves $M_2(A)$ invariant. If $\chi_w = \chi_v \theta^\#$ for $v \in M_2(A)$, and $\theta^\# \in (\text{Aut } A)^\#$, then for each $c \in A$, $cI = \chi_w(cI) = \chi_v \theta^\#(cI) = \theta(c)I$, which implies $\theta = 1$. Now if $\chi_w = \chi_v$, then

$$w = \begin{pmatrix} (r, s) & 0 \\ 0 & (r, s) \end{pmatrix} v$$

where r, s are nonzero elements of $\Phi(t)$. The determinant of v must be a unit in A , so it has the form (α, α) where $\alpha \in \Phi$. From taking determinants we conclude $(1, -1) = (r^2, s^2)(\alpha, \alpha)$, and hence both r and s are in Φ . We write

$$v = \begin{pmatrix} (u_1, v_1) & (u_2, v_2) \\ (u_3, v_3) & (u_4, v_4) \end{pmatrix}$$

and compute

$$w = \begin{pmatrix} (r, s) & 0 \\ 0 & (r, s) \end{pmatrix} v.$$

Then $1 = ru_1$ where $r \in \Phi$, and since $(u_1, v_1) \in A$, it follows $v_1 = r^{-1}$ also. But then $-t = sv_1 = sr^{-1} \in \Phi$, a contradiction. Consequently $\chi_w \in \text{Aut } M_2(A)$, and $\chi_w \notin \text{GL}(2, A) \cdot (\text{Aut } A)^\#$.

REMARK. We are indebted to L. Levy for his assistance in the creation of this example and for making us aware of work by Rosenberg and Zelinsky [10] on this subject.

4. $M_n(A)$ under the Lie product. Let $L_n(A)$ denote the anticommutative algebra obtained by taking $M_n(A)$ under the product $[xy] = xy - yx$, and let $L'_n(A) = [L_n(A), L_n(A)]$. The algebra $L'_n(A)$ is an ideal of $L_n(A)$ which is invariant under all automorphisms and derivations of $L_n(A)$. It is spanned by the elements

$$\begin{aligned} ae_{ij} &= [ae_{ii}, e_{ij}], & i \neq j, \\ abe_{ii} - bae_{jj} &= [ae_{ij}, be_{ji}]. \end{aligned} \quad (4.1)$$

Two noteworthy cases of the second equation occur when $i = j$ and when $b = 1$. The elements obtained are $[ab]e_{ii}$ and $ae_{ii} - ae_{jj}$. These elements in fact generate all of the ones of the second type since $abe_{ii} - bae_{jj} = [ab]e_{ii} + bae_{ii} - bae_{jj}$. Each element in $L'_n(A)$ has trace in $[A, A]$. Conversely if $x = \sum_{i=1}^n a_{ii}e_{ii}$ where $t = \sum_{i=1}^n a_{ii} \in [A, A]$, then $x = te_{nn} + \sum_{i=1}^{n-1} a_{ii}e_{ii} - a_{ii}e_{nn} \in L'_n(A)$. Thus $L'_n(A)$ is precisely the space of all elements with trace in $[A, A]$. Using the above products one can also verify that if $n \geq 3$ when $\text{char } \Phi = 2$, or if $n \geq 2$ when $\text{char } \Phi \neq 2$, then $L'_n(A) = [L'_n(A), L'_n(A)]$.

Let Z denote the center of $L_n(A)$; that is $Z = \{z \in L_n(A) \mid [z, x] = 0 \text{ for all } x \in L_n(A)\}$. If $C(A) = \{c \in A \mid ca = ac \text{ for all } a \in A\}$, then calculation shows that $Z = \{cI \mid c \in C(A)\}$. The center is an ideal of $L_n(A)$ invariant under automorphisms and derivations of $L_n(A)$. Its intersection with $L'_n(A)$ is the center of $L'_n(A)$ and is also invariant. We define $K_n(A) = L'_n(A)/Z \cap L'_n(A)$. Our goal in this section will be to investigate the derivation algebras of $M_n(A)$, $L_n(A)$, $L'_n(A)$, and $K_n(A)$, and the automorphism groups of the last three. Each derivation (automorphism) of $M_n(A)$ induces a derivation (automorphism) of $L_n(A)$. This assertion remains true for the pairs $L_n(A)$, $L'_n(A)$ and $L'_n(A)$, $K_n(A)$. So we have the natural homomorphisms

$$\text{Der } M_n(A) \xrightarrow{\rho} \text{Der } L_n(A) \xrightarrow{\sigma} \text{Der } L'_n(A) \xrightarrow{\tau} \text{Der } K_n(A).$$

LEMMA 4.1. *Suppose $n \geq 3$ if $\text{char } \Phi = 2$, or $n \geq 2$ if $\text{char } \Phi \neq 2$. Then*

(a) $\sigma\rho$, ρ , and τ are one-to-one, and

(b) the kernel of σ consists of all Lie homomorphisms of $L_n(A)$ into Z , denoted by $\text{LHom}(L_n(A), Z)$.

PROOF. If ∂ is in the kernel of τ , then ∂ maps $L'_n(A)$ into Z . But then $L'_n(A) = [L'_n(A), L'_n(A)]$ is sent to 0, which shows $\partial = 0$. Since $L_n(A) = M_n(A)$ as vector spaces, the kernel of ρ must be 0. Now any derivation ∂ in the kernel of $\sigma\rho$

must be 0 on the elements ae_{ij} for $i \neq j$. Since these elements generate $M_n(A)$ under the usual matrix product, it follows that ∂ is identically 0. Turning our attention to part (b), we assume ∂ is in the kernel of σ . Then ∂ is 0 on the elements in (4.1) so we need only examine the effect of ∂ on elements of the form ae_{ij} . For $k \neq l$ we have $[ae_{ij}be_{kl}] = \delta_{jk}abe_{jl} - \delta_{jl}bae_{kj}$, and ∂ applied to this relation shows $[\partial(ae_{ij}), be_{kl}] = 0$ for all $b \in A$. From this we deduce $\partial(ae_{ij}) \in Z$. Any Lie homomorphism of $L_n(A)$ into Z is a derivation of $L_n(A)$ which necessarily vanishes on $L'_n(A)$. Thus the kernel of σ is $\text{LHom}(L_n(A), Z)$ as claimed. \square

In a similar fashion we have the natural homomorphisms

$$\text{Aut } M_n(A) \xrightarrow{\rho} \text{Aut } L_n(A) \xrightarrow{\sigma} \text{Aut } L'_n(A) \xrightarrow{\tau} \text{Aut } K_n(A).$$

LEMMA 4.2. Assume $n \geq 3$ if $\text{char } \Phi = 2$, or $n \geq 2$ if $\text{char } \Phi \neq 2$. Then

- (a) $\sigma\rho, \rho$, and τ are one-to-one, and
- (b) the kernel of σ consists of all mappings φ such that φ is the identity map plus an element in $\text{LHom}(L_n(A), Z)$.

Since the proof of 4.2 involves only a slight modification of the proof of 4.1, we omit it.

The Lie nucleus, which we introduce next, will play the role that $M_n(N)$ played in our investigations of $\text{GAut } M_n(A)$.

For each anticommutative algebra L with product $[]$ the Lie nucleus of L is

$$\nu(L) = \{x \in L \mid [[xy]z] + [[yz]x] + [[zx]y] = 0 \text{ for all } y, z \in L\}.$$

If ad_x denotes the map $\text{ad}_x(y) = [xy]$, then an alternate description of the Lie nucleus is

$$\nu(L) = \{x \in L \mid \text{ad}_x \in \text{Der } L\}.$$

The space $\nu(L)$ is invariant under automorphisms and derivations of L . As a result, if $x, x' \in \nu(L)$ then $\text{ad}_x(x') \in \nu(L)$, and $\nu(L)$ is a subalgebra of L . It follows then that $\nu(L)$ is a Lie algebra.

Let us specialize now to the case $L = L_n(A)$. We define $D(A) = \{a \in A \mid \text{ad}_a \in \text{Der } A\}$, and note that this amounts to saying $D(A) = \{a \in A \mid [a, bc] = [ab]c + b[ac]\}$.

LEMMA 4.3. Assume $n \geq 3$. Then $\nu(L_n(A)) = L_n(N) + D(A)I$.

PROOF. Using the definition of $D(A)I$ above and the fact that $M_n(N)$ is the nucleus of $M_n(A)$, one can show that $L_n(N) + D(A)I \subseteq \nu(L_n(A))$. For the proof of the reverse containment let

$$\lambda(x, y, z) = [[xy]z] + [[yz]x] + [[zx]y]$$

and assume $x = \sum b_{ij}e_{ij} \in \nu(L_n(A))$. Then since $\nu(L_n(A))$ is a subalgebra containing $L_n(N)$, for k, l, m distinct we have $[[[xe_{lk}]e_{mk}]e_{lm}] = b_{kl}e_{lk} \in \nu(L_n(A))$. Dropping the subscript on b_{kl} for simplicity, we have $b_{lk}e_{lk} \in \nu(L_n(A))$ for $l \neq k$. From $\lambda(be_{lk}, ae_{km}, ce_{mk}) = 0$ for all $a, c \in A$ we obtain $(ba)ce_{lk} - b(ac)e_{lk} = 0$, and hence $(ba)c = b(ac)$. Similarly $\lambda(ae_{ml}, be_{lk}, ce_{kl}) = 0$ and $\lambda(ae_{mk}, ce_{kl}, be_{lk}) = 0$

imply $(ab)c = a(bc)$ and $(ac)b = a(cb)$. We conclude b is in N , and $be_{ik} \in L_n(N) \subseteq \nu(L_n(A))$.

It suffices to suppose $x = \sum_{i=1}^n b_i e_{ii} \in \nu(L_n(A))$. For $k \neq l$ we have

$$[x, e_{kl}] = (b_k - b_l)e_{kl} \in \nu(L_n(A))$$

and by the above $b_k - b_l \in N$. Now a computation of $\lambda(x, ae_{qr}, ce_{st})$ shows

$$\begin{aligned} &\delta_{rs}(b_q a - ab_r)ce_{qt} - \delta_{tq}c(b_q a - ab_r)e_{sr} \\ &\quad + \delta_{rs}\{(ac)b_t - b_q(ac)\}e_{qt} - \delta_{tq}\{b_s(ca) - (ca)b_r\}e_{sr} \\ &\quad - \delta_{rs}a(cb_t - b_s c)e_{qt} + \delta_{tq}(cb_t - b_s c)ae_{sr} = 0. \end{aligned}$$

This expression can be made more transparent by writing for each i and j $b_i = b_q + \beta_i$, $b_j = b_r + \gamma_j$ where $\beta_i, \gamma_j \in N$. The result is

$$\begin{aligned} 0 = &\delta_{rs}\{[b_q a]c + [ac, b_q] + a[cb_q] + (a\beta_s)c - a(\beta_r c)\}e_{qt} \\ &- \delta_{tq}\{c[b_r a] + [ca, b_r] - [cb_r]a + (c\gamma_q)a - c(\gamma_t a)\}e_{sr}. \end{aligned}$$

This proves that $b_q \in D(A)$ for each q and hence, $x = \sum_{i=1}^{n-1} (b_i - b_n)e_{ii} + b_n I \in L_n(N) + D(A)I$. Thus, the calculation shows $\nu(L_n(A)) \subseteq L_n(N) + D(A)I$, and it also demonstrates that $D(A)I \subseteq \nu(L_n(A))$ as was asserted above. \square

The same argument can be used to show

LEMMA 4.4. For $n \geq 3$, $\nu(L'_n(A)) = \nu(L_n(A)) \cap L'_n(A)$ and $\nu(K_n(A)) = \nu(L'_n(A)) + Z \cap L'_n(A)$.

Let us adopt the notation $Z' = Z \cap L'_n(A)$ so that $K_n(A) = L'_n(A)/Z'$.

LEMMA 4.5. Assume $\text{char } \Phi \neq 2, 3$, and let T be a transformation on $K_n(A)$ with the property:

$$[e_{ii} - e_{jj} + Z', T(x + Z')] = T([e_{ii} - e_{jj} + Z', x + Z'])$$

for all $x \in L'_n(A)$ and all i, j . Then for each pair $i \neq j$, T induces a transformation T_{ij} on A such that

$$T(ae_{ij} + Z') = T_{ij}(a)e_{ij} + Z'.$$

PROOF. When $\text{char } \Phi \neq 2, 3$ the space $Ae_{ij} + Z'$ can be characterized in the following way:

$$Ae_{ij} + Z' = \{x + Z' \in K_n(A) \mid [e_{ii} - e_{jj} + Z', x + Z'] = 2x + Z'\}.$$

The definition of T makes this space T -invariant. Thus, for each $a \in A$, there is a unique $b \in A$ such that the coset $be_{ij} + Z'$ equals $T(ae_{ij} + Z')$. The T_{ij} desired is given by $T_{ij}(a) = b$. \square

Lemma 4.5 will be used next to study derivations ∂ with the property that $\partial(e_{ii} - e_{jj} + Z') = 0$, and in the future to investigate automorphisms φ such that $\varphi(e_{ii} - e_{jj} + Z') = e_{ii} - e_{jj} + Z'$.

THEOREM 4.6. Assume $n \geq 3$ and $\text{char } \Phi \neq 2, 3$. Let $\partial \in \text{Der } K_n(A)$ be such that $\partial(e_{ii} - e_{jj} + Z') = 0$ for all i, j . Then there is a $\Delta \in (\text{Der } A)^\#$ and a diagonal matrix $y \in M_n(N)$ such that $\partial = \Delta + \text{ad}_y$.

PROOF. The derivation ∂ satisfies the hypotheses of Lemma 4.5, and so

$$\partial(e_{ij} + Z') = \alpha_{ij}e_{ij} + Z' \quad \text{for } i \neq j.$$

The elements α_{ij} belong to N since $e_{ij} + Z$ is in $\nu(K_n(A))$ which is derivation-invariant, and off-diagonal entries of $\nu(K_n(A))$ lie in N .

Applying ∂ to the following relations

$$[e_{ik} + Z', e_{kj} + Z'] = e_{ij} + Z' \quad \text{for } i, j, k \text{ distinct,} \tag{4.2}$$

$$[e_{ij} + Z', e_{ji} + Z'] = e_{ii} - e_{jj} + Z' \tag{4.3}$$

shows that

$$\alpha_{ik} + \alpha_{kj} = \alpha_{ij}, \quad \alpha_{ij} + \alpha_{ji} = 0. \tag{4.4}$$

Let y be the diagonal matrix with $0, \alpha_{21}, \dots, \alpha_{n1}$ down the diagonal. Then $\text{ad}_y(e_{ij} + Z') = (\alpha_{i1} - \alpha_{j1})e_{ij} + Z' = \alpha_{ij}e_{ij} + Z'$. Thus $\Delta = \partial - \text{ad}_y \in \text{Der } K_n(A)$ has the property that $\Delta(e_{ij} + Z') = 0$ as well as $\Delta(e_{ii} - e_{jj} + Z') = 0$ for each pair i, j with $i \neq j$. According to Lemma 4.5, Δ induces transformations Δ_{ij} of A such that $\Delta(ae_{ij} + Z') = \Delta_{ij}(a)e_{ij} + Z'$ for $i \neq j$. Since

$$\Delta(ae_{ij} + Z') = [\Delta(ae_{ik} + Z'), e_{kj} + Z'] = \Delta_{ik}(a)e_{ik} + Z',$$

it follows that $\Delta_{ij} = \Delta_{ik}$ for $k \neq j$. An analogous argument shows the first subscript can also be altered. Thus, all the induced transformations are equal and we call the common map η .

By applying Δ to the relation

$$[ae_{ik} + Z', be_{kj} + Z'] = abe_{ij} + Z' \quad \text{for } i, k, j \text{ distinct} \tag{4.5}$$

we obtain $\eta \in \text{Der } A$. Thus, at least on elements of the form $ae_{ij} + Z', \Delta = \eta^\#$, the derivation obtained by acting by η on each entry. But these elements generate $K_n(A)$ as the next two relations show:

$$[ae_{ij} + Z', e_{ji} + Z'] = ae_{ii} - ae_{jj} + Z', \tag{4.6}$$

$$[ae_{ii} - ae_{jj} + Z', be_{ii} - be_{kk} + Z'] = [ab]e_{ii} + Z'. \tag{4.7}$$

And these equations can be used to verify that $\Delta = \eta^\# \in (\text{Der } A)^\#$. \square

THEOREM 4.7. *Let ∂ be an arbitrary element of $\text{Der } K_n(A)$ where $\text{char } \Phi \neq 2, 3$ and $n \geq 3$. Then there is a $w \in M_n(N)$ such that $\partial' = \partial - \text{ad}_w$ has the property that $\partial'(e_{ii} - e_{jj} + Z') = 0$ for all i, j .*

PROOF. When $i \neq j$,

$$\partial[e_{ii} - e_{jj} + Z'] = [\partial(e_{ij} + Z'), e_{ji} + Z'] + [e_{ij} + Z', \partial(e_{ji} + Z')],$$

and from this relation it is apparent that $\partial(e_{ii} - e_{jj} + Z')$ has a unique coset representative with nonzero entries only in the i th and j th rows and columns. When $j = n$ let us denote this coset in the following fashion:

$$\begin{aligned} \partial(e_{ii} - e_{nn} + Z') &= \sum_{k \neq i} \beta_{ki}^i e_{ki} + \sum_l \beta_{il}^i e_{il} \\ &\quad + \sum_{\substack{k \neq i \\ k \neq n}} \beta_{kn}^i e_{kn} + \sum_{l \neq i} \beta_{nl}^i e_{nl} + Z'. \end{aligned}$$

Since $\partial(e_{ii} - e_{jj} + Z') = \partial(e_{ii} - e_{nn} + Z') - \partial(e_{jj} - e_{nn} + Z')$, we have

$$\beta_{kn}^i = \beta_{kn}^j, \beta_{nk}^i = \beta_{nk}^j, \text{ for } k \text{ distinct from } i, j, \text{ and } n. \tag{4.8}$$

The elements β_{kl}^i lie in N for $k \neq l$ since the Lie nucleus is preserved by derivations. Showing that the diagonal β 's are 0, and deriving relationships between the various β 's for different i 's, such as (4.8) above, will enable us to construct the matrix w .

Let $\partial(e_{ij} + Z') = \sum \alpha_{kl} e_{kl} + Z'$, and compare coefficients of the (i, i) and (i, j) entries in

$$[\partial(e_{ii} - e_{nn} + Z'), e_{ij} + Z'] + [e_{ii} - e_{nn} + Z', \partial(e_{ij} + Z')] = \partial(e_{ij} + Z'). \tag{4.9}$$

The resulting relations are

$$-\beta_{ji}^i = \alpha_{ii}, \quad -\beta_{ii}^i + \alpha_{ij} = \alpha_{ij}. \tag{4.10}$$

Now if $e_{ii} - e_{nn}$ is replaced by $e_{jj} - e_{nn}$ in (4.9) an analogous argument shows

$$-\beta_{ji}^j = -\alpha_{ii}. \tag{4.11}$$

Therefore it follows that

$$\beta_{ii}^i = 0 \quad \text{and} \quad \beta_{ji}^i = -\beta_{ji}^j. \tag{4.12}$$

Finally, if e_{in} is used in place of e_{ij} in (4.9), one can show $\beta_{nn}^i = 0$ also. We define

$$\begin{aligned} \gamma_{ji} &= \beta_{ji}^i \text{ for } j, i \neq n, \\ \gamma_{kn} &= -\beta_{kn}^j \text{ for } j \neq k, n, \\ \gamma_{nk} &= \beta_{nk}^j. \end{aligned}$$

The equations in (4.8) show that γ_{kn} and γ_{nk} are independent of the defining j . Let $w = \sum \gamma_{kl} e_{kl}$ and note that $w \in M_n(N)$, so that ad_w is a derivation. The image of each $e_{ii} - e_{nn} + Z'$ under ad_w is the same as under ∂ except for the (i, n) and (n, i) entries. Thus, if $\partial' = \partial - \text{ad}_w$, then $\partial'(e_{ii} - e_{nn} + Z') = \xi_{in} e_{in} + \xi_{ni} e_{ni} + Z'$. Comparison of the (n, j) and (i, n) coefficients in (4.9) now shows

$$\xi_{ni} - \alpha_{nj} = \alpha_{nj}, \quad 2\alpha_{in} = \alpha_{in}. \tag{4.13}$$

Similar equations resulting from the use of $e_{jj} - e_{nn}$ are

$$-\xi_{jn} + \alpha_{in} = -\alpha_{in}, \quad -2\alpha_{nj} = -\alpha_{nj}. \tag{4.14}$$

Thus, $\xi_{jn} = 0$ and $\xi_{ni} = 0$ for each i and j , and $\partial' = \partial - \text{ad}_w$ is the desired derivation. \square

Combining Theorems 4.6 and 4.7 gives

THEOREM 4.8. *Assume char $\Phi \neq 2, 3$ and $n \geq 3$. Then for any algebra A with 1, $\text{Der } K_n(A) = (\text{Der } A)^\# + \text{ad}_{M_n(N)}$.*

COROLLARY 4.9. *With assumptions as in the previous theorem,*

$$\text{Der } M_n(A) = \text{Der } L'_n(A) = \text{Der } K_n(A) = (\text{Der } A)^\# + \text{ad}_{M_n(N)}.$$

PROOF. Each element of $\text{Der } K_n(A)$ is a derivation of $M_n(A)$ and $L'_n(A)$. However the mappings $\tau\sigma\rho$ and τ in Lemma 4.1 are the canonical injections of $\text{Der } M_n(A)$ and $\text{Der } L'_n(A)$ into $\text{Der } K_n(A)$. So these algebras can be no larger than $\text{Der } K_n(A)$, and the above equalities hold. \square

Using 4.1 and this result we have

COROLLARY 4.10. $\text{Der } L_n(A) = (\text{Der } A)^\# + \text{ad}_{M_n(N)} + \text{LHom}(L_n(A), Z)$.

REMARKS. Martindale [8] has obtained a result in a similar vein as Corollary 4.10. He showed that if R is a primitive associative ring with a nontrivial idempotent, then any Lie derivation of R is the sum of a derivation of R plus a Lie homomorphism of R into Z .

It should be commented that in showing $\text{Der } M_n(A) = (\text{Der } A)^\# + \text{ad}_{M_n(N)}$ no restrictions on char Φ or on n are necessary. In this case $Ae_{ij} = \{x \in M_n(A) \mid e_{ii}x = xe_{jj} = x\}$ so that any derivation ∂ with the property that $\partial(e_{ii}) = 0$ induces a transformation ∂_{ij} on Ae_{ij} regardless of char Φ . The additional hypotheses are needed in the computation of $\text{Der } K_n(A)$ to determine the structure of the Lie nucleus. But in the calculation of $\text{Der } M_n(A)$, $M_n(N)$ is mapped into itself, so no such contortions are necessary. Rather than make a separate case for $M_n(A)$, we chose to treat $M_n(A)$, $L_n(A)$, $L'_n(A)$, and $K_n(A)$ simultaneously at this small expense.

COROLLARY 4.11 (JACOBSON [3, THEOREM 8, p. 215]). *Let S be a finite dimensional central simple associative algebra over Φ . Then every derivation of S is inner.*

PROOF. Let F be a splitting field for S over Φ . Then $S \otimes_\Phi F \approx M_n(F)$. Thus by Corollary 4.9 and the preceding remarks, $\text{Der}_F(S \otimes_\Phi F) \approx \text{Der}_F(M_n(F)) \approx \text{ad}_{M_n(F)}$. Now if $\partial \in \text{Der } S$, then $\partial \otimes 1 \in \text{Der}_F(S \otimes_\Phi F)$. In particular, for each $x \in S$, $\text{ad}_x \in \text{Der } S$, and since S is central it follows that $\dim_\Phi \text{ad } S = n^2 - 1$. Thus,

$$n^2 - 1 = \dim_F(\text{Der}_F M_n(F)) \geq \dim_\Phi \text{Der } S \geq \dim_\Phi \text{ad}_S = n^2 - 1.$$

Hence, $\text{Der } S = \text{ad}_S$ as claimed. \square

THEOREM 4.12. *Assume $n \geq 3$ and char $\Phi \neq 2, 3$. Let $\varphi \in \text{Aut } K_n(A)$ be such that $\varphi(e_{ii} - e_{jj} + Z') = e_{ii} - e_{jj} + Z'$ for all i, j . Then there is a $\psi \in (\text{Aut } A)^\#$ and an invertible diagonal matrix $u \in M_n(N)$ such that $\varphi = \chi_u\psi$ where χ_u is conjugation by u .*

PROOF. Just as in the proof of Theorem 4.6, by Lemma 4.5 there exist elements $\alpha_{ij} \in N$ for $i \neq j$ such that $\varphi(e_{ij} + Z') = \alpha_{ij}e_{ij} + Z'$. Only this time applying φ to relations (22) and (23) yields

$$\begin{aligned} \alpha_{ik}\alpha_{kj} &= \alpha_{ij} && \text{for } i, k, j \text{ distinct,} \\ \alpha_{ij}\alpha_{ji} &= 1 && \text{for } i \neq j. \end{aligned} \tag{4.15}$$

Let u be the diagonal matrix with $1, \alpha_{12}, \dots, \alpha_{1n}$ down the diagonal. Note that u^{-1} is also diagonal with $1, \alpha_{21}, \dots, \alpha_{n1}$ as its diagonal entries. Then $\chi_u(e_{ij} + Z') = u^{-1}e_{ij}u + Z' = \alpha_{i1}\alpha_{1j}e_{ij} + Z' = \alpha_{ij}e_{ij} + Z'$. Thus $\psi = \chi_u^{-1}\varphi \in \text{Aut } K_n(A)$ has the property that it fixes each $e_{ij} + Z'$ as well as each $e_{ii} - e_{jj} + Z'$ for $i \neq j$.

Beyond this stage the proof is exactly parallel to the proof of Theorem 4.6. One shows the transformations induced from ψ are all equal, and then deduces from (4.5) that the common map, call it θ , belongs to $\text{Aut } A$. Finally then, using (4.6) and (4.7) one verifies $\psi = \theta^\# \in (\text{Autv } A)^\#$. \square

These conclusions remain valid for any $\varphi \in \text{Aut } L_n(A)$ ($\text{Aut } L'_n(A)$) with the property that $\varphi(e_{ii}) = e_{ii}$ ($\varphi(e_{ii} - e_{jj}) = e_{ii} - e_{jj}$).

Given any antiautomorphism φ of $M_n(A)$, $-\varphi$ is an automorphism of $L_n(A)$, thus of $L'_n(A)$ and $K_n(A)$ as well. The collection of all such mappings together with $\text{Aut } M_n(A)$ forms a subgroup of the automorphism group of each of these algebras. It is isomorphic to $\text{AAut } M_n(A)$, so we will use that notation to denote it.

THEOREM 4.13. *If A is finite dimensional over Φ where $\text{char } \Phi \neq 2, 3$, and if $n \geq 3$, then $\text{Aut } M_n(A)$ (hence $\text{AAut } M_n(A)$, also) is of finite index in $\text{Aut } L'_n(A)$ and in $\text{Aut } K_n(A)$.*

PROOF. The group $G = \text{Aut } L'_n(A)$ (or $\text{Aut } K_n(A)$) is an algebraic group (see Seligman [11, p. 34]), and $H = \text{Aut } M_n(A)$ is a closed subgroup of G . The Lie algebra of G is $\text{Der } K_n(A)$ which equals the Lie algebra of H , $\text{Der } M_n(A)$. Therefore, by Proposition 4, p. 33 of Seligman, H is of finite index in G . \square

Now apply Theorem 4.13 and Lemma 4.2 to obtain

COROLLARY 4.14. *With hypotheses as in Theorem 4.13, $\text{Aut } M_n(A) + \text{LHom}(L_n(A), Z)$ has finite index in $\text{Aut } L_n(A)$.*

REMARK. In the associative case it follows from Martindale [9] that the Lie automorphisms of R , a prime ring with a nontrivial idempotent, are precisely $\text{AAut } R + \text{LHom}(R, Z)$.

5. $M_n(A)$ under the Jordan product. Throughout this section we assume $\text{char } \Phi \neq 2$. Let $J_n(A)$ denote the commutative algebra obtained by taking $M_n(A)$ under the Jordan product $x \circ y = \frac{1}{2}(xy + yx)$. In this part we investigate the derivation algebra and automorphism group of $J_n(A)$. The study of derivations of $J_n(A)$ is facilitated by the use of results on transformations in triality. The notion of triality comes up in the study of the Lie algebras of types D_4 and F_4 (see Jacobson [5] for example). There the transformations are defined on an octonion algebra, but here we extend the concept to arbitrary algebras.

Let A be an arbitrary algebra with 1, and assume ζ, η, θ are linear transformations on A with the property that

$$\zeta(ab) = \eta(a)b + a\theta(b) \quad \text{for all } a, b \in A.$$

Then we say ζ, η, θ are in local triality (or for short, triality). For example if $\zeta = \eta = \theta$, and ζ is a derivation of A , then ζ, η, θ are in triality.

LEMMA 5.1. *If $\zeta, \eta, \eta', \theta, \theta'$ are transformations on A such that ζ, η, θ and ζ, η', θ' are in triality then*

(1) $\eta' = \eta - R_s$ where $R_s(a) = as$,

(2) $\theta' = \theta + L_s$ where $L_s(a) = sa$,

and $s = \eta(1) - \eta'(1) = \theta'(1) - \theta(1)$ and s is in the middle nucleus of A .

PROOF. Observe that if ζ, η, θ and ζ', η', θ' are in triality then $\alpha\zeta + \beta\zeta', \alpha\eta + \beta\eta', \alpha\theta + \beta\theta'$ are in triality for every pair of scalars α, β . Thus, if $\eta'' = \eta - \eta', \theta'' = \theta - \theta'$, then $0, \eta'', \theta''$ are in triality. This says that $\eta''(a)b + a\theta''(b) = 0$ for all $a, b \in A$. Specializing b to equal 1 shows that

$$\eta''(a) = -a\theta''(1) \quad \text{for all } a \in A. \tag{5.1}$$

Likewise $a = 1$ gives

$$\theta''(b) = -\eta''(1)b \quad \text{for all } b \in B. \tag{5.2}$$

If both $a = 1$ and $b = 1$, then $\eta''(1) = -\theta''(1)$. Let s be this common value. Then (5.1) and (5.2) shows that $\eta - \eta' = \eta'' = R_s$ and $\theta - \theta' = \theta'' = -L_s$. Finally, the relation $\eta''(a)b + a\theta''(b) = 0$ implies $(as)b - a(sb) = 0$ as desired. \square

LEMMA 5.2. *Let $\partial \in \text{Der}(J_n(A))$ have the property that $\partial(e_{ii}) = 0$ for $i = 1, \dots, n$. Then ∂ induces n^2 transformations ∂_{ij} on A such that $\partial(ae_{ij}) = \partial_{ij}(a)e_{ij}$.*

PROOF. The proof begins just as Lemma 3.5 does. The same argument using the five types of nonzero products and the eigenspaces of the elements e_{ii} demonstrates that $\partial(ae_{ii}) = \partial_{ii}(a)e_{ii}$ and $\partial(ae_{ij}) = \partial_{ij}(a)e_{ij} + \partial'_{ij}(a)e_{ji}$. In particular, $\partial(e_{ij}) = \alpha_{ij}e_{ij} + \beta_{ji}e_{ji}$. But since $e_{ij}^2 = 0$, we have $2e_{ij} \circ \partial(e_{ij}) = 0$. Therefore $\beta_{ji} = 0$ for $j \neq i$. Moreover, $e_{ij} \circ ae_{ij} = 0$ implies $\alpha_{ij}e_{ij} \circ ae_{ij} + e_{ij} \circ (\partial_{ij}(a)e_{ij} + \partial'_{ij}(a)e_{ji}) = 0$, and from this we see $\partial'_{ij}(a) = 0$ for all a . Hence, $\partial(ae_{ij}) = \partial_{ij}(a)e_{ij}$ in all cases. \square

LEMMA 5.3. *Let $\text{char } \Phi \neq 2$, and $n \geq 2$. Assume $\partial \in \text{Der } J_n(A)$ has the property that $\partial(e_{ii}) = 0$ for $i = 1, \dots, n$. Then there is a diagonal matrix $y \in M_n(N)$ such that $\partial - \text{ad}_y \in (\text{Der } A)^\#$.*

PROOF. By the preceding lemma ∂ induces the transformations ∂_{ij} , and for i, j distinct we see from relations (3.2), (3.3), (3.5) that the following transformations are in triality on A :

$$\partial_{ij}, \quad \partial_{ii}, \quad \partial_{ij}, \tag{5.3}$$

$$\partial_{ij}, \quad \partial_{ij}, \quad \partial_{jj}, \tag{5.4}$$

$$\partial_{ii}, \quad \partial_{ij}, \quad \partial_{ji}. \tag{5.5}$$

Let $\alpha_{ij} = \partial_{ij}(1)$. Since relation (3.1) implies ∂_{ii} is a derivation on A under the Jordan product, it follows that $\alpha_{ii} = 0$. From (5.5) we conclude that $\alpha_{ij} = -\alpha_{ji}$ for $j \neq i$, so that by Lemma 5.1 we have

$$\partial_{ij} = \partial_{ii} + R_{\alpha_{ij}}, \tag{5.6}$$

$$\partial_{jj} = \partial_{ij} - L_{\alpha_{ij}} \tag{5.7}$$

and also the fact that α_{ij} is in the middle nucleus of A . Subtracting (5.4) from (5.3) and then adding (5.5) shows that $\partial_{ii}, \partial_{ii}, \partial_{ij} - \partial_{jj} + \partial_{ji}$ are in triality. However, using (5.6) and (5.7) we may express each of the last three transformations in terms of ∂_{ii} . Note that

$$\partial_{jj} = \partial_{ii} + R_{\alpha_{ij}} - L_{\alpha_{ij}} \tag{5.8}$$

and interchanging j and i in (5.7) gives: $\partial_{ii} = \partial_{ji} - L_{\alpha_{ji}} = \partial_{ji} + L_{\alpha_{ij}}$. Therefore, $\partial_{ij} - \partial_{ji} + \partial_{ji} = \partial_{ii} + R_{\alpha_{ij}} - \partial_{ii} - R_{\alpha_{ji}} + L_{\alpha_{ij}} + \partial_{ii} - L_{\alpha_{ij}} = \partial_{ii}$. Hence ∂_{ii} , ∂_{ij} , ∂_{ji} are in triality and ∂_{ii} is a derivation on A .

As a consequence of equations (5.3) and (5.6) we have

$$\partial_{ii}(ab) + (ab)\alpha_{ij} = \partial_{ii}(a)b + a\partial_{ii}(b) + a(b\alpha_{ij})$$

which implies that α_{ij} is in the right nucleus of A . Similarly using (5.4) and (5.7) we deduce that α_{ij} is in the left nucleus of A . Thus α_{ij} is in the nucleus of A . Let y be the diagonal matrix with $0, \alpha_{21}, \dots, \alpha_{n1}$ down the diagonal. From equations (5.6), (5.7), (5.8) it follows that $\partial_{1j}(a) = \partial_{11}(a) - a\alpha_{j1}$, $\partial_{i1}(a) = \partial_{11}(a) + \alpha_{i1}a$ and $\partial_{jj}(a) = \partial_{11}(a) + [\alpha_{j1}, a]$. If $n \geq 3$, then for i, j distinct from 1, ∂_{ij} , ∂_{i1} , ∂_{1j} are in triality and $\partial_{ij}(a) = \partial_{11}(a) + \alpha_{i1}a - a\alpha_{j1}$. We conclude from these relations that $\partial = (\partial_{11})^\# + \text{ad}_y$ where $\partial_{11} \in \text{Der } A$. \square

LEMMA 5.4. *Let char $\Phi \neq 2$, and $n \geq 2$. Suppose $\partial \in \text{Der } J_n(A)$. Then there is a $w \in M_n(N)$ such that $\partial' = \partial - \text{ad}_w$ has the property that $\partial'(e_{ii}) = 0$ for all i .*

PROOF. Apply ∂ to the relation $e_{ii} \circ e_{ii} = e_{ii}$ to conclude that $\partial(e_{ii})$ lies in the $\frac{1}{2}$ -eigenspace relative to multiplication by e_{ii} . Therefore we may write

$$\partial(e_{ii}) = \sum_{k \neq i} \beta_{ki}^i e_{ki} + \sum_{l \neq i} \beta_{il}^i e_{il}.$$

Now if $j \neq i$, then $\partial(e_{ii}) \circ e_{jj} + e_{ii} \circ \partial(e_{jj}) = 0$, and from this we calculate that

$$\beta_{ji}^i = -\beta_{ij}^j. \tag{5.9}$$

Thus, if $\gamma_{kl} = \beta_{kl}^l$ for $k \neq l$ and if $\gamma_{kk} = 0$, then $w = \sum \gamma_{kl} e_{kl}$ has the property that

$$\begin{aligned} \text{ad}_w(e_{ii}) &= \sum_k \gamma_{ki} e_{ki} - \sum_l \gamma_{il} e_{il} = \sum_{k \neq i} \beta_{ki}^i e_{ki} - \sum_{l \neq i} \beta_{il}^i e_{il} \\ &= \sum_{k \neq i} \beta_{ki}^i e_{ki} + \sum_{l \neq i} \beta_{il}^i e_{il} = \partial(e_{ii}). \end{aligned}$$

The proof will be complete once it is proven that each $\gamma_{kl} \in N$, and hence $\text{ad}_w \in \text{Der } J_n(A)$. A computation of ∂ applied to $ae_{ii} \circ e_{ii} = ae_{ii}$ shows that

$$\partial(ae_{ii}) = \sum_{k \neq i} \beta_{ki}^i ae_{ki} + \sum_{l \neq i} a\beta_{il}^i e_{il} + \Delta(a, i)e_{ii}.$$

Using this relation, we conclude from ∂ acting on $ae_{ii} \circ be_{jj} = 0$ that

$$0 = \frac{1}{2} \{ (a\beta_{ij}^i)b + a(\beta_{ij}^j)b \} e_{ij} + \frac{1}{2} \{ b(\beta_{ji}^i)a + (b\beta_{ji}^j)a \} e_{ji}.$$

Since $\beta_{ij}^i = -\beta_{ji}^j$, this implies $\gamma_{ij} = \beta_{ij}^j$ is in the middle nucleus of A .

Let us apply ∂ to both sides of $e_{ii} \circ be_{ij} = \frac{1}{2} be_{ij}$. Now

$$\partial(e_{ii}) \circ be_{ij} = \frac{1}{2} \sum_{k \neq i} \beta_{ki}^i be_{kj} + b\beta_{ji}^i e_{ii},$$

and $e_{ii} \circ \partial(be_{ij})$ has nonzero entries just in the i th row and column. Thus,

$$\partial(be_{ij}) = \sum_{k \neq i} \beta_{ki}^i be_{kj} + \sum_{l \neq i} c_{il} e_{il} + \sum_l c_{li} e_{li}.$$

The next step is to equate coefficients after applying ∂ to $e_{ij} \circ be_{ij} = \frac{1}{2}be_{ij}$. The result obtained is

$$\partial(be_{ij}) = \sum_{k \neq i} \beta_{ki}^i be_{kj} + \sum_{l \neq j} b\beta_{jl}^j e_{il} + \Delta(b, i, j)e_{ij} + \Delta'(b, j, i)e_{ji}.$$

Then $\partial(ae_{ii}) \circ be_{ij} + ae_{ii} \circ \partial(be_{ij}) = \frac{1}{2}\partial(abe_{ij})$ implies $(\beta_{ki}^i a)b = \beta_{ki}^i(ab)$ and $a(b\beta_{jl}^j) = (ab)\beta_{jl}^j$. Thus, the γ 's lie in N , $w = \sum \gamma_{kl}e_{kl}$ in $M_n(N)$, and ad_w in $\text{Der } J_n(A)$. As noted above, then for $\partial' = \partial - \text{ad}_w$, we have $\partial'(e_{ii}) = 0$ for all i . \square

The net effect of Lemmas 5.3 and 5.4 is a proof of

THEOREM 5.5. *Let char $\Phi \neq 2$. Then*

$$\text{Der } J_n(A) = (\text{Der } A)^\# + \text{ad}_{M_n(N)} = \text{Der } M_n(A).$$

REMARKS. Recall that the last equality follows from Corollary 4.9 and the comments preceding Corollary 4.11.

Previous results analogous to Theorem 5.5 have been known to hold in the associative case. Jacobson and Rickart showed in [6, Theorems 7 and 22] that $\text{Der } M_n(A) = \text{Der } J_n(A)$ when A is associative, while Herstein [2, p. 55] proved that every Jordan derivation of a prime associative ring R is a derivation of R , except when R is a commutative integral domain of characteristic 2. These results would have been useful had we known ab initio that $\partial \in \text{Der } J_n(A)$ implies $\partial(M_n(N)) \subseteq M_n(N)$. Indeed the major effort of Lemma 5.4 is in proving the image of each e_{ii} lies in $M_n(N)$.

The work on the next theorem concerning $\text{Aut } J_n(A)$ has largely been done already, but we collect the results for the convenience of the reader.

THEOREM 5.6. *Assume char $\Phi \neq 2$. Let $G = \{\psi \in \text{Aut } J_n(A) | \psi(e_{ii}) = e_{ii} \text{ for all } i\}$ and $H = \{\psi \in \text{Aut } J_n(A) | \psi(J_n(N)) \subseteq J_n(N)\}$. Then*

$$(i) \quad \begin{matrix} G \\ \subseteq \\ \text{Aut } M_n(A) \end{matrix} \subseteq \text{GL}(n, N) \cdot (\text{GAut } A)^\# \subseteq \text{GAut } M_n(A) \subseteq H \subseteq \text{Aut } J_n(A).$$

(ii) $G = X \cdot (\text{GAut } A)^\#$ where X is the subgroup consisting of conjugations by diagonal matrices with entries in N .

(iii) If N is Artinian, then $\text{GL}(n, N) \cdot (\text{GAut } A)^\# = \text{GAut } M_n(A) = H$.

(iv) If A is finite dimensional, then $\text{Aut } M_n(A)$, hence $\text{GAut } M_n(A)$ also, is of finite index in $\text{Aut } J_n(A)$.

PROOF. Assertions (i) and (ii) are consequences of results in §3, notably Theorem 3.6 and Lemma 3.12. Part (iii) follows from Lemma 3.4 and Theorems 3.6 and 3.13. Since $\text{Der } J_n(A) = \text{Der } M_n(A)$, statement (iv) can be concluded by the same algebraic group argument used in §4. \square

REMARKS. For associative rings Herstein [2, p. 50] has shown that if φ is a Jordan homomorphism of R onto a prime ring R' , then φ is a homomorphism or an antihomomorphism. This result is false in the nonassociative case as the next example illustrates.

Choose $\lambda, \mu \in \Phi$ such that $\lambda + \mu = 1$, and let B have basis $\{\varepsilon_{ij}\}$ for $i, j = 1, \dots, n$ where the product in B is given by $\varepsilon_{ij} * \varepsilon_{kl} = \delta_{jk} \lambda \varepsilon_{il} + \delta_{il} \mu \varepsilon_{kj}$. If B_J denotes B under the Jordan product then B_J is Jordan isomorphic to $M_n(\Phi)$ under the correspondence $\varepsilon_{ij} \rightarrow e_{ij}$. Since B is nonassociative when $\lambda \neq 1, 0$, this map is neither an isomorphism nor an anti-isomorphism of B onto $M_n(\Phi)$.

Now if A is any algebra then $M_n(A) \approx A \otimes M_n(\Phi)$. If in addition A is assumed to be commutative then $J_n(A) \approx A \otimes J_n(\Phi)$. If A is any commutative algebra isomorphic to $J_n(\Phi)$, such as B_J above, then there is an automorphism of $J_n(A)$ mapping $A \otimes 1$ to $1 \otimes J_n(\Phi)$. If the nucleus of A is $\Phi 1$, as with B_J , then such an automorphism is not in $\text{Aut } M_n(A)$ or $\text{GAut } M_n(A)$ since it fails to preserve the nucleus which is $1 \otimes M_n(\Phi)$. By taking A to be the tensor product of any number of algebras isomorphic to $J_n(\Phi)$, we see that the index of $\text{Aut } M_n(A)$ in $\text{Aut } J_n(A)$ can be arbitrarily large.

We conclude our remarks with one additional example. Suppose B is the algebra above, and form the algebra $J_n(B)$. The product is given by

$$2\varepsilon_{ij}\varepsilon_{qr} \circ \varepsilon_{kl}\varepsilon_{st} = \delta_{rs}\delta_{jk}\lambda\varepsilon_{il}\varepsilon_{qt} + \delta_{rs}\delta_{il}\mu\varepsilon_{kj}\varepsilon_{qt} + \delta_{iq}\delta_{jk}\mu\varepsilon_{il}\varepsilon_{sr} + \delta_{iq}\delta_{il}\lambda\varepsilon_{kj}\varepsilon_{sr}.$$

Using this formula one can show that the map φ given by $\varphi(\varepsilon_{ij}\varepsilon_{qr}) = \varepsilon_{qr}\varepsilon_{ij}$ belongs to $\text{Aut } J_n(B)$. If $\lambda \neq 1, 0$ then φ interchanges the nonassociative algebra B with $M_n(\Phi)$; hence it fails to belong to $\text{Aut } M_n(B)$. This example illustrates that the coefficient algebra B need not be commutative to make interchanges of this sort possible.

REFERENCES

1. G. Domokos and S. Kovesi-Domokos, *The algebra of color* J. Math. Phys. **14** (1973), 1651–1667.
2. I. N. Herstein, *Topics in ring theory*, Univ. of Chicago Press, Chicago, Ill., 1965.
3. N. Jacobson, *Abstract derivations and Lie algebras*, Trans. Amer. Math. Soc. **42** (1937), 206–224.
4. _____, *Structure of alternative and Jordan bimodules*, Osaka J. Math. **6** (1954), 1–71.
5. _____, *Exceptional Lie algebras*, Marcel Dekker, New York, 1971.
6. N. Jacobson and C. E. Rickart, *Jordan homomorphisms of rings*, Trans. Amer. Math. Soc. **69** (1950), 479–502.
7. L. S. Levy, *Artinian, non-Noetherian rings*, J. Algebra **47** (1977), 276–304.
8. W. S. Martindale III, *Lie derivations of primitive rings*, Michigan Math. J. **11** (1964), 183–187.
9. _____, *Lie isomorphisms of prime rings*, Trans. Amer. Math. Soc. **142** (1969), 437–455.
10. A. Rosenberg and D. Zelinsky, *Automorphisms of separable algebras*, Pacific J. Math. **11** (1961), 1109–1117.
11. G. B. Seligman, *Algebraic groups*, Yale University, mimeographed notes, 1964.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN, MADISON, WISCONSIN 53706