

A GENERAL PRINCIPLE FOR LIMIT THEOREMS IN FINITELY ADDITIVE PROBABILITY

BY

RAJEEVA L. KARANDIKAR

ABSTRACT. In this paper we formulate and prove a general principle which enables us to deduce limit theorems for sequences of independent random variables in a finitely additive setting from their analogues in the conventional countably additive setting.

1. Introduction. The first a.s. limit theorems in the finitely additive strategic setting were proved by Purves and Sudderth [10] and included a version of the strong law and the martingale convergence theorem. Chen [4, 5] proved some a.s. limit theorems for a sequence of independent random variables in the finitely additive strategic setting. Ramakrishnan [11] has proved the central limit theorem in the same set-up. In many of these proofs, the main technique was to 'approximate' the given sequence $\{Y_k\}$ of independent random variables by a sequence $\{Z_k\}$ such that each Z_k takes only finitely many values. Behind these approximations, there is a general principle which we isolate and prove. Given the sequence $\{Y_k\}$ as above, we construct an independent sequence $\{X_k\}$ on a countably additive probability space which is 'very close' to $\{Y_k\}$ in distribution. To be more precise, we show that we can approximate $\{Y_k\}$ by $\{Z_k\}$ and $\{X_k\}$ by $\{W_k\}$ such that $\{Z_k\}$ and $\{W_k\}$ have the same distribution. Our main result is that 'a limit theorem' holds for $\{Y_k\}$ if and only if it holds for $\{X_k\}$. The formulation of this result is similar to Aldous' formulation of the 'subsequence principle' [1]. We show how to deduce the usual results like SLIN, WLLN, LIL, CLT and Donsker's and Strassen's invariance principles from our result. It should be pointed out that our method, although quite general, does not apply to prove results about martingales.

We now describe the finitely additive set-up with which we will be working.

Let I be a nonempty set. Let $\{\gamma_n; n \geq 1\}$ be a sequence of finitely additive probability measures on all subsets of I . Let $H = I^\infty$ be equipped with the product of discrete topologies and let \underline{A} be the Borel σ field of H . Using the methods of Dubins and Savage [6, 7] and Purves and Sudderth [10] it is possible to get a finitely additive probability σ on \underline{A} (unique subject to certain regularity conditions) having the following desirable properties:

$$(i) \sigma(A_1 \times A_2 \times \cdots) = \prod_{k=1}^{\infty} \gamma_k(A_k), A_k \subseteq I.$$

Received by the editors May 7, 1981.

1980 *Mathematics Subject Classification.* Primary 60F05, 60G07.

Key words and phrases. Finitely additive probability, strategy, invariance principles.

©1982 American Mathematical Society
0002-9947/82/0000-0038/\$03.50

(ii) If \underline{A}_k is a sequence of σ fields on I such that $\gamma'_k = \gamma_k$ restricted to \underline{A}_k is countably additive, then $\sigma = \gamma'$ on \underline{A}' where $\underline{A}' = \otimes_k \underline{A}_k$ and $\gamma' = \otimes_k \gamma'_k$ (usual product measure).

(iii) Let $A_n \subset I$ and let $B_n = \{h \in H: h_n \in A_n\}$ ($h = (h_1, h_2, \dots) \in H$). Then $\sigma(\bigcap_{k=1}^\infty \bigcup_{n=k}^\infty B_n) = 0$ if and only if $\sum_{n=1}^\infty \sigma(B_n) < \infty$. Further if $\sum_{n=1}^\infty \sigma(B_n) < \infty$, then $\sigma(\bigcup_{n=k}^\infty B_n) \rightarrow 0$ as $k \rightarrow \infty$.

(i) and (iii) follow from (ii). (ii) is a special case of Theorem 1, Purves and Sudderth [10].

2. Preliminaries and statement of the main result. Let μ be a finitely additive probability measure on (R, \underline{B}_R) which is tight (i.e. $\lim_n \mu([-n, n]) = 1$). We will associate with μ a unique countably additive measure ν as follows:

$f \rightarrow \int f d\mu$ defines a positive linear functional on the space of bounded continuous functions from R into itself. It follows easily from tightness of μ and Dini's theorem that this linear functional is σ -smooth (i.e. $f_n \downarrow 0$ implies $\int f_n d\mu \rightarrow 0$). Thus by Daniell's theorem [9, p. 60] there exists a unique countably additive probability measure ν on \underline{B}_R such that

$$\int f d\mu = \int f d\nu$$

for all bounded continuous functions f . The measure ν thus obtained is called the countably additive probability associated with μ . The referee has pointed out that the countably additive measure ν is the "conventional companion" of Dubins and Savage [7, p. 190].

Let f be a positive continuous function. Then for each n ,

$$\int (f \wedge n) d\mu = \int (f \wedge n) d\nu.$$

But

$$\int f d\mu = \lim_n \int (f \wedge n) d\mu \quad (\text{definition})$$

and

$$\int f d\nu = \lim_n \int (f \wedge n) d\nu \quad (\text{monotone convergence theorem}).$$

Thus $\int f d\mu = \int f d\nu$.

From this it follows that, for a continuous function f , $\int f d\mu$ is well defined if and only if $\int f d\nu$ is and then $\int f d\mu = \int f d\nu$.

Let $F(x) = \mu((-\infty, x])$, and $G(x) = \nu((-\infty, x])$; we claim that $F(x) = G(x)$ if x is a continuity point of F or G . To see this, fix x and for each $n \geq 1$ let f_n, g_n be continuous functions such that

$$\chi_{(-\infty, x-1/n]} \leq f_n \leq \chi_{(-\infty, x]} \leq g_n \leq \chi_{(-\infty, x+1/n]}.$$

Then

$$F\left(x - \frac{1}{n}\right) \leq \int f_n d\mu \leq F(x) \leq \int g_n d\mu \leq F\left(x + \frac{1}{n}\right)$$

and

$$G\left(x - \frac{1}{n}\right) \leq \int f_n d\nu \leq G(x) \leq \int g_n d\nu \leq G\left(x + \frac{1}{n}\right).$$

Further, $\int f_n d\mu = \int f_n d\nu$ and $\int g_n d\mu = \int g_n d\nu$ for all n . Thus if x is a continuity point of F (or G), $F(x) = G(x)$.

Before we proceed to state our main result, we define convergence in distribution for ‘random-elements’ on a finitely additive probability space.

Let H be a nonempty set, \underline{A} be a σ field on H and σ be a finitely additive probability measure on \underline{A} . Let S be a metric space. Let $\xi_k: H \rightarrow S$ be a sequence of random elements in S (i.e. $\xi_k^{-1}B \in \underline{A}$ for all Borel sets in S) and let λ be a countably additive probability measure on S .

DEFINITION. Say ξ_k converge in distribution to λ if $\sigma(\xi_k \in B) \rightarrow \lambda(B)$ for all Borel sets B in S such that $\lambda(\partial B) = 0$.

The proof of the following theorem on equivalent conditions for convergence in distribution is the same as in the countably additive case (see [2, p. 11]) and hence is omitted.

THEOREM 1. *Let $\{\xi_k\}$ be S valued random elements on a finitely additive probability space $(H, \underline{A}, \sigma)$ and let λ be a countably additive probability measure on S .*

Then the following are equivalent:

- (i) $\int f(\xi_k) d\sigma \rightarrow \int f d\lambda$ for all bounded continuous functions f .
- (ii) $\int f(\xi_k) d\sigma \rightarrow \int f d\lambda$ for all bounded uniformly continuous functions f .
- (iii) $\limsup_k \sigma(\xi_k \in F) \leq \lambda(F)$ for all closed sets $F \subseteq S$.
- (iv) $\liminf_k \sigma(\xi_k \in G) \geq \lambda(G)$ for all open sets $G \subseteq S$.
- (v) $\xi_k \rightarrow \lambda$ in distribution.

The following proposition follows easily from the equivalence of (ii) and (v) in Theorem 1.

PROPOSITION 2. *Let $\{\xi_k\}, \{\eta_k\}$ be random elements in S (on $(H, \underline{A}, \sigma)$) and let $d(\xi_k, \eta_k) \rightarrow 0$ in σ -probability. Then ξ_k converges in distribution to λ implies that η_k converges in distribution to λ .*

Let $I, \{\gamma_n\}, H, \underline{A}, \sigma$ be as in the introduction. Let $\{f_k: k \geq 1\}$ be a sequence of mappings from I into R . Let $Y_k(h) = f_k(h_k)$ be the coordinate map on H induced by f_k (here $h = (h_1, h_2, \dots) \in H$). Let μ_k be the distribution of Y_k , i.e. $\mu_k(A) = \sigma(Y_k \in A), A \in \underline{B}_R$. Assume that μ_k is tight for each k (i.e. $\sigma(|Y_k| \geq n) \rightarrow 0 \forall k$). Let ν_k be the countably additive measure associated with μ_k . Let $(\Omega, \underline{F}, P)$ be a (countably additive) probability space supporting a sequence $\{X_k\}$ of independent random variables such that $PX_k^{-1} = \nu_k$ for all k .

Let $Y = (Y_1, Y_2, \dots)$ and $X = (X_1, X_2, \dots)$ be the random vectors taking values in R^∞ corresponding to the sequence $\{Y_k\}$ and $\{X_k\}$ respectively.

THEOREM 3. Fix $0 < p < \infty$.

(a) Let $A \in \underline{B}^\infty$ (Borel σ field of R^∞) be such that

$$(1) \quad x \in A \quad \text{and} \quad \sum_{i=1}^{\infty} |x_i - x'_i|^p < \infty \quad \text{implies} \quad x' \in A.$$

Then $Y \in A$ a.s. σ if and only if $X \in A$ a.s. P .

(b) Let (S, d) be a separable metric space and $\{g_k: k \geq 1\}$ be a sequence of measurable mappings from R^∞ into S such that

$$(2) \quad d(g_k(x), g_k(x')) \leq \sum_{i=1}^{\infty} C_{k,i} |x_i - x'_i|^p$$

where $C_{k,i}$ are positive constants bounded by 1 such that for each i , $\lim_k C_{k,i} = 0$.

Let $s \in S$. Then $g_k(Y) \rightarrow s$ in σ -probability if and only if $g_k(X) \rightarrow s$ in P -probability.

(c) Let λ be a countably additive probability measure on S and $\{g_k\}$ be as in (b) satisfying (2). Then $g_k(Y) \rightarrow \lambda$ in distribution if and only if $g_k(X) \rightarrow \lambda$ in distribution.

REMARK 1. If $\{g_k\}$ satisfy (2), then $A = \{x: g_k(x) \rightarrow s\}$ satisfies (1) and hence (a) implies that $g_k(Y) \rightarrow s$ a.s. σ if and only if $g_k(X) \rightarrow s$ a.s. P .

3. Proofs. The idea of the proof is to approximate $Y = (Y_1, Y_2, \dots)$ by $Z = (Z_1, Z_2, \dots)$ and $X = (X_1, X_2, \dots)$ by $W = (W_1, W_2, \dots)$ such that the distribution of Z under σ is the same as the distribution of W under P .

Let $F_k(x) = \mu_k((-\infty, x])$, $G_k(x) = \nu_k((-\infty, x])$ and $D_k = \{x: x \text{ is a continuity point of } F_k\}$. Then as observed earlier $F_k(x) = G_k(x)$ for all $x \in D_k$.

For each $k \geq 1$, choose numbers $\{a_{k,j}: 0 \leq j \leq n_k\}$ belonging to D_k such that

(i) $F_k(a_{k,n_k}) - F_k(a_{k,0}) \geq 1 - 1/2^k$ and

(ii) $0 < a_{k,j} - a_{k,j-1} < 1/2^k, 1 \leq j \leq n_k$.

This can be done as for each k , $\lim_{x \rightarrow \infty} F_k(x) = 1$, $\lim_{x \rightarrow -\infty} F_k(x) = 0$ and D_k complement is at most countable.

Define $Z = (Z_1, Z_2, \dots)$ and $W = (W_1, W_2, \dots)$ by

$$\begin{aligned} Z_k(h) &= a_{k,0} && \text{if } Y_k(h) \leq a_{k,0}, \\ &= a_{k,j} && \text{if } a_{k,j} < Y_k(h) \leq a_{k,j+1}: 0 \leq j < n_k, \\ &= a_{k,n_k} && \text{if } a_{k,n_k} < Y_k(h), \end{aligned}$$

and

$$\begin{aligned} W_k(w) &= a_{k,0} && \text{if } X_k(w) \leq a_{k,0}, \\ &= a_{k,j} && \text{if } a_{k,j} < X_k(w) \leq a_{k,j+1}: 0 \leq j < n_k, \\ &= a_{k,n_k} && \text{if } a_{k,n_k} < X_k(w). \end{aligned}$$

LEMMA 1. $\sigma(Z \in B) = P(W \in B)$ for all Borel sets B in R^∞ .

PROOF. First observe that, for each k ,

$$\begin{aligned} \sigma(Z_k = a_{k,j}) &= F_k(a_{k,j+1}) - F_k(a_{k,j}) = G_k(a_{k,j+1}) - G_k(a_{k,j}) \\ &= P(W_k = a_{k,j}), \quad 1 \leq j < n_k. \end{aligned}$$

Similarly $\sigma(Z_k = a_{k,0}) = P(W_k = a_{k,0})$ and $\sigma(Z_k = a_{k,n_k}) = P(W_k = a_{k,n_k})$.

For each k , let \underline{A}_k be the finite field on I generated by the sets $\{f_k \leq a_{k,0}\}$, $\{f_k > a_{k,n_k}\}$, $\{a_{k,j} < f_k \leq a_{k,j+1}\}$. Let γ'_k be γ_k restricted to \underline{A}_k . Then (trivially) γ'_k is countably additive. Let $\underline{A}' = \otimes \underline{A}_k$ and $\gamma = \otimes \gamma'_k$ (the usual product measure on (H, \underline{A}')). Then by usual arguments, it follows that $\gamma'(Z \in B) = P(W \in B)$ for all Borel sets B in R^∞ . Since σ agrees with γ' on \underline{A}' (property (ii) of σ) the proof is complete.

The next lemma shows that $Z(W)$ approximates $Y(X)$ ‘nicely’.

LEMMA 2. For $0 < p < \infty$

- (a) $\sum_{k=1}^\infty |Y_k - Z_k|^p < \infty$ a.s. σ .
- (b) $\sum_{k=1}^\infty |X_k - W_k|^p < \infty$ a.s. P .
- (c) $d(g_k(Y), g_k(Z)) \rightarrow 0$ in σ -probability.
- (d) $d(g_k(X), g_k(W)) \rightarrow 0$ in P -probability.

PROOF. (a) $\sigma(|Y_k - Z_k| \geq 1/2^k) \leq F_k(a_{k,n_k}) - F_k(a_{k,0}) \leq 1/2^k$; hence by the Borel-Cantelli lemma (property (iii) of σ)

$$\sigma\left(|Y_k - Z_k| \geq \frac{1}{2^k} \text{ i.o.}\right) = 0.$$

Now (a) follows from this.

- (b) follows similarly as $P(|X_k - W_k| \geq 1/2^k) \leq 1/2^k$.
- (c) By the Borel-Cantelli lemma (property (iii) of σ),

$$\sigma(|Y_k - Z_k| \geq 1/2^k \text{ for some } k \geq n) \rightarrow 0 \text{ as } n \rightarrow \infty.$$

We now show that this and (2) imply (c). Given $\epsilon > 0$ and $\delta > 0$ choose N such that

$$\frac{1}{2^{Np}(1 - 2^{-p})} < \frac{\epsilon}{2} \quad \text{and} \quad \sigma\left(|Y_k - Z_k| > \frac{1}{2^k} \text{ for some } k \geq N\right) < \frac{\delta}{2}.$$

For each i , since the distribution of $Y_i - Z_i$ is tight and $\lim_k C_{k,i} = 0$ for each i ,

$$\sigma\left(\sum_{i=1}^N C_{k,i} |Y_i - Z_i|^p \geq \frac{\epsilon}{2}\right) \rightarrow 0 \text{ as } k \rightarrow \infty.$$

Choose k_0 large enough such that

$$\sigma\left(\sum_{i=1}^N C_{k,i} |Y_i - Z_i|^p \geq \frac{\epsilon}{2}\right) \leq \frac{\delta}{2} \text{ if } k \geq k_0.$$

Then for $k \geq k_0$,

$$\begin{aligned} \sigma(d(g_k(Z), g_k(Y)) \geq \epsilon) &\leq \sigma\left(\sum_{i=1}^\infty C_{k,i} |Y_i - Z_i|^p \geq \epsilon\right) \\ &\leq \sigma\left(\sum_{i=1}^N C_{k,i} |Y_i - Z_i|^p \geq \frac{\epsilon}{2}\right) + \frac{\delta}{2} \leq \delta. \end{aligned}$$

Hence $d(g_k(Z), g_k(Y)) \rightarrow 0$ in σ -probability.

(d) follows similarly.

The theorem now follows easily from these two lemmas.

PROOF OF THE THEOREM.

- (a) $Y \in A$ a.s. $\sigma \Leftrightarrow Z \in A$ a.s. σ by (1) and Lemma 2(a)
 $\Leftrightarrow W \in A$ a.s. P by Lemma 1
 $\Leftrightarrow X \in A$ a.s. P by (1) and Lemma 2(b).
- (b) $g_k(Y) \rightarrow s$ in σ probability $\Leftrightarrow g_k(Z) \rightarrow s$ in σ -probability by Lemma 2(c)
 $\Leftrightarrow g_k(W) \rightarrow s$ in P -probability by Lemma 1
 $\Leftrightarrow g_k(X) \rightarrow s$ in P -probability by Lemma 2(d).
- (c) $g_k(Y) \rightarrow \lambda$ in distribution $\Leftrightarrow g_k(Z) \rightarrow \lambda$ in distribution by Lemma 2(c)
 $\Leftrightarrow g_k(W) \rightarrow \lambda$ in distribution by Lemma 1
 $\Leftrightarrow g_k(X) \rightarrow \lambda$ in distribution by Lemma 2(d).

4. Consequences. We have already observed that $\int f(Y_k) d\sigma = \int f(X_k) dP$ (if either of the two integrals is well defined) for all continuous functions f . Thus limit theorems for the sequence $\{X_k\}$ whose hypotheses involve only the moments of $\{X_k\}$ follow for the sequence $\{Y_k\}$ by virtue of our theorem. (With the additional assumption that each ' Y_k ' is tight. In most of the situations, the other conditions will imply tightness of ' Y_k '.) This includes Kolmogorov's strong law, Khinchin's weak law (in the finitely additive setting, strong law does not imply weak law!) and the Hartman-Wintner law of iterated logarithm. We illustrate by an example.

THEOREM (CHEN) SLLN. *Let $\{Y_k\}$ be identically distributed ($\mu_k = \mu_1$ for all k), $\int |Y_1| d\sigma < \infty$ and $\int Y_1 d\sigma = 0$. Then*

$$\frac{Y_1 + \cdots + Y_n}{n} \rightarrow 0 \quad \text{a.s. } \sigma.$$

PROOF. Take $A = \{x \in R^\infty: (x_1 + \cdots + x_n)/n \rightarrow 0\}$.

Check that A satisfies (1) of Theorem 3. Then, $Y \in A$ a.s. σ if and only if $X \in A$ a.s. P . Further $\mu_k = \mu_1 \forall k$ implies X_k are identically distributed,

$$\int |X_1| dP = \int |Y_1| dP < \infty, \quad \int X_1 dP = \int Y_1 d\sigma = 0.$$

Hence (by Kolmogorov's strong law—countably additive version [3, p. 52]) $X \in A$ a.s. P . This implies that $Y \in A$ a.s. σ .

In many other situations, say the Lindeberg-Feller central limit theorem or Kolmogorov's three series theorem, the conditions on X_k can be transformed into equivalent conditions involving $\int f(X_k) dP$ for some continuous functions f and hence again the finitely additive versions can be derived from their countably additive analogues. We illustrate this by an example.

THEOREM (RAMAKRISHNAN) (CLT). *Let $\int Y_k d\sigma = 0$, and let $v_k^2 = \int Y_k^2 d\sigma$, $s_n^2 = v_1^2 + \cdots + v_k^2$. Then*

- (i) $(Y_1 + \cdots + Y_n)/s_n \rightarrow N(0,1)$ in distribution and
(ii) $\max_{k \leq n} (v_k^2/s_n^2) \rightarrow 0$

if and only if the Lindeberg condition (L) holds for Y_n :

$$(L) \quad \frac{1}{s_n^2} \sum_{k=1}^n \int_{\{|Y_k| > \epsilon s_n\}} |Y_k|^2 d\sigma \rightarrow 0 \quad \text{as } n \rightarrow \infty \text{ for all } \epsilon > 0.$$

PROOF. Let $g_k(x) = (x_1 + \dots + x_k)/s_k$ for $x \in R^\infty$. Then

$$|g_k(x) - g_k(x')| \leq \sum_{i=1}^k \frac{1}{s_k} |x_i - x'_i|.$$

Thus if $s_n^2 \rightarrow \infty$, then (2) holds with $C_{k,i} = (1/s_k)$. Thus, if $s_n^2 \rightarrow \infty$, then by Theorem 3, $g_k(Y) \rightarrow N(0,1)$ in distribution if and only if $g_k(X) \rightarrow N(0,1)$ in distribution.

For each $n \geq 1, \epsilon > 0$, fix a continuous function $f_{n,\epsilon}$ such that

$$x^2 1(x)_{\{|y| > \epsilon s_n\}} \leq f_{n,\epsilon}(x) \leq x^2 1(x)_{\{|y| > \epsilon/2 s_n\}}.$$

Then

$$\int_{\{|Y_k| > \epsilon s_n\}} |Y_k|^2 d\sigma \leq \int f_{n,\epsilon}(Y_k) d\sigma \leq \int_{\{|Y_k| > \epsilon/2 s_n\}} |Y_k|^2 d\sigma$$

and hence (L) holds for Y (respectively X) if and only if (L') holds for Y (respectively X), where (L') is

$$(L') \quad \frac{1}{s_n^2} \sum_{k=1}^n \int f_{n,\epsilon}(Y_k) d\sigma \rightarrow 0 \quad \text{for all } \epsilon > 0.$$

But (L') holds for Y if and only if (L') holds for X . Thus (L) holds for Y if and only if (L) holds for X .

Now, the result follows from the Lindeberg-Feller theorem—countably additive case [8, p. 280].

We now show that Donsker's invariance principle and Strassen's invariance principle can be proved in the finitely additive setting.

THEOREM (DONSKER'S INVARIANCE PRINCIPLE). For each n , define $g_n: R^\infty \rightarrow C[0, 1]$ as follows:

$$g_n(x)(t) = \frac{x_1 + \dots + x_k}{\sqrt{n}} \quad \text{if } t = \frac{k}{n}, \text{ linear elsewhere.}$$

Let Y_1, Y_2, \dots have identical distributions, $\int Y_1 d\sigma = 0, \int Y_1^2 d\sigma = 1$. Then $g_n(Y)$ converges in distribution to the Wiener measure.

PROOF. Observe that

$$d(g_n(x), g_n(x')) = \sup_t |g_n(x)(t) - g_n(x')(t)| \leq \sum_{k=1}^n \frac{1}{\sqrt{n}} |x_k - x'_k|.$$

Thus g_k satisfies (2).

Now, by Donsker's theorem (countably additive version [2, p. 68]), $g_k(X)$ converges in distribution to Wiener measure and hence by Theorem 3, so does $g_k(Y)$.

THEOREM (STRASSEN'S INVARIANCE PRINCIPLE). *Let Y_1, Y_2, \dots have identical distributions, $\int Y_1 d\sigma = 0$, $\int Y_1^2 d\sigma = 1$. Then a.s. σ , the sequence of functions $g_n(Y)/\sqrt{\log \log n}$ is relatively compact in $C[0,1]$ and its set of limit points is $K = \{f \in C[0,1]: f \text{ absolutely continuous, } f(0) = 0 \text{ and } \int_0^1 f'(u)^2 du \leq 1\}$ (here g_n are as in the previous theorem).*

PROOF. Let $A = \{x: g_n(x)/\sqrt{\log \log n} \text{ is relatively compact in } C[0, 1] \text{ and set of its limit point is } K\}$.

Then, it can be seen that A satisfies (1) and thus the result follows from Strassen's invariance principle (countably additive version [12]) and Theorem 3.

REMARK. Let F_k be a sequence of nondecreasing functions on R such that $\lim_{x \rightarrow -\infty} F_k(x) = 0$ and $\lim_{x \rightarrow \infty} F_k(x) = 1$. Let \underline{C} be the field of subsets of R generated by the family $\{(a, b]: -\infty \leq a < b \leq \infty\}$. Each F_k defines a unique finitely additive 'tight' probability measure on the field \underline{C} by the formula $\mu_k((-\infty, b]) = F_k(b)$.

Consider the field \underline{D} of subsets of R^∞ defined by

$$\underline{D} = \bigcup_k \bigotimes_k \underline{A}_k$$

where the union is taken over all sequences $\{\underline{A}_k\}$ of finite subfields of \underline{C} . For such a sequence $\{\underline{A}_k\}$, each μ_k restricted to \underline{A}_k is (trivially) countably additive. Let λ_0 be the usual countably additive product measure on $\bigotimes_k \underline{A}_k$ with marginals μ_k (restricted to \underline{A}_k). It can be easily seen that these measures are consistent and hence determine a unique finitely additive measure λ on \underline{D} . Let σ be any extension of λ to \underline{B}^∞ . Then (by construction of λ) it follows that

(ii)' For any sequence $\{\underline{A}_k\}$ of finite subfields of \underline{C} , σ restricted to $\bigotimes_k \underline{A}_k$ agrees with $\bigotimes_k \mu'_k$ where μ'_k is μ_k restricted to \underline{A}_k .

Now let Y_k be the coordinate random variables on $(R^\infty, \underline{B}^\infty, \sigma)$.

Let $G_k(x) = \lim_{y \downarrow x} F_k(y)$ for $x \in R$, $k \geq 1$, and let $(\Omega, \underline{F}, P)$ be a probability space supporting a sequence $\{X_k\}$ of independent random variables with $P(X_k \leq x) = G_k(x)$ for all $x \in R$, $k \geq 1$.

Then it follows easily (by using the property (ii)' of σ instead of (ii)) that Theorem 3 (and hence its consequences) is true in this setting as well.

5. Let (S, d) be a separable metric space and μ be a finitely additive tight probability measure on \underline{B}_S . $f \rightarrow \int f d\mu$ defines a σ -smooth positive linear functional on the space of bounded continuous functions from S into R and hence by Daniell's theorem [9, p. 60] there exists a unique countably additive probability measure ν on \underline{B}_S such that $\int f d\mu = \int f d\nu$ for all $f \in C_b(S)$. Theorem 1 implies that $\mu(A) = \nu(A)$ if $\nu(\partial A) = 0$.

Let $(H, \underline{A}, \sigma)$ be as in the introduction. Let $f_k: I \rightarrow S$, and $Y_k(h) = f_k(h_k)$. Let $\mu_k(B) = \sigma(Y_k \in B)$. Assume μ_k is tight for each k . Let ν_k be the countably additive measure associated with μ_k as above. Let $\{X_k\}$ be an independent sequence of S valued random elements on a countably additive probability space $(\Omega, \underline{F}, P)$ such that $\nu_k(B) = P(X_k \in B)$.

The following theorem is an analogue of Theorem 3.

THEOREM. Fix $0 < p < \infty$.

(a) Let $A \subseteq S^\infty$ be a Borel set such that

$$(1') \quad x \in A \quad \text{and} \quad \sum_{i=1}^{\infty} (d(x_i, x'_i))^p < \infty \quad \text{implies} \quad x' \in A.$$

Then $Y \in A$ a.s. σ if and only if $X \in A$ a.s. P .

(b) Let (S_1, d_1) be a separable metric space and g_k be a sequence of measurable mappings from S^∞ into S_1 such that

$$(2') \quad d_1(g_k(x'), g_k(x)) \leq \sum_{i=1}^{\infty} C_{k,i} (d(x_i, x'_i))^p, \quad x, x' \in S^\infty,$$

where $C_{k,i}$ are positive constants bounded by 1 such that $\lim_k C_{k,i} = 0$ for each i .

Let $s \in S_1$.

Then $g_k(Y) \rightarrow s$ in σ -probability if and only if $g_k(X) \rightarrow s$ in P -probability.

(c) Let $\{g_k\}$ be as in (b) satisfying (2') and let λ be a countably additive measure on S_1 . Then $g_k(Y) \rightarrow \lambda$ in distribution if and only if $g_k(X) \rightarrow \lambda$ in distribution.

The proof of this theorem is exactly the same as in the case $S = R$ except for the following lemma. The sets B_1, B_2, \dots, B_m satisfying conditions (i), (ii) and (iii) below for $\lambda = \nu_k$ and $\epsilon = 1/2^k$ replace the intervals $(a_{k,j-1}, a_{k,j}]$, $1 \leq j \leq n_k$, in the construction of Z_k and W_k .

LEMMA. Let S be a separable metric space and let λ be a countably additive tight probability measure on S . Then for each $\epsilon > 0$, there exist finitely many disjoint Borel sets B_1, \dots, B_m such that:

- (i) $\lambda(\cup_{i=1}^m B_i) \geq 1 - \epsilon$.
- (ii) $\lambda(\partial B_i) = 0$.
- (iii) $x_1, x_2 \in B_i$ implies $d(x_1, x_2) \leq \epsilon$ for all i .

PROOF. Let K be a compact set such that $\lambda(K) \geq 1 - \epsilon$. Let A_1, \dots, A_m be balls of radius $\epsilon/4$ covering K . For any Borel set B in S let $B^\epsilon = \{x \in S: d(x, B) \leq \epsilon\}$. Then $\partial B^\epsilon \subseteq \{x \in S: d(x, B) = \epsilon\}$. Hence, for each i , choose ϵ_i , $0 < \epsilon_i < \epsilon/4$, such that $\lambda(\partial A_i^{\epsilon_i}) = 0$. Let $B_i = A_i^{\epsilon_i} \cap (\cup_{j=1}^{i-1} A_j^{\epsilon_j})^c$. Then B_i satisfy the required conditions.

REMARK. Let μ_k be a sequence of tight finitely additive measures on R . Consider the restriction μ of the independent strategic measure with marginals μ_k to \underline{B}^∞ . Then μ is tight. Let ν, ν_k be the countably additive measures associated with μ, μ_k respectively. Then $\nu = \otimes_k \nu_k$.

ACKNOWLEDGEMENTS. The author thanks Professor A. Maitra, Professor B. V. Rao and Dr. S. Ramakrishnan for their valuable suggestions and the discussions he had with them.

REFERENCES

1. D. J. Aldous, *Limit theorems for sub-sequences of arbitrarily dependent sequences of random variables*, Z. Wahrsch. Verw. Gebeite **40** (1977), 59–82.
2. P. Billingsley, *Convergence of probability measure*, Wiley, New York, 1968.
3. L. Breiman, *Probability*, Addison-Wesley, California, 1968.
4. R. Chen, *A finitely additive version of Kolmogorov's law of iterated logarithm*, Israel J. Math. **23** (1976), 209–220.

5. _____, *On almost sure convergence in a finitely additive setting*, *Z. Wahrsch. Verw. Gebiete* **37** (1977), 341–356.
6. L. E. Dubins, *On Lebesgue like extensions of finitely additive measures*, *Ann. Probab.* **2** (1974), 456–463.
7. L. E. Dubins and L. J. Savage, *How to gamble if you must: Inequalities for stochastic processes*, McGraw-Hill, New York, 1965.
8. M. Loeve, *Probability theory*, 3rd ed., Van Nostrand, Princeton, N. J., 1963.
9. J. Neveu, *Mathematical foundations of the calculus of probability*, Holden-Day, San Francisco, Calif., 1965.
10. R. A. Purves and W. D. Sudderth, *Some finitely additive probability*, *Ann. Probab.* **4** (1976), 259–276.
11. S. Ramakrishnan, *Central limit theorems in a finitely additive setting* (preprint).
12. V. Strassen, *An invariance principle for the law of iterated logarithm*, *Z. Wahrsch. Verw. Gebiete* **3** (1964), 211–226.

STATISTICS-MATHEMATICS DIVISION, INDIAN STATISTICAL INSTITUTE, CALCUTTA 700035, INDIA

Current address: Department of Statistics, The University of North Carolina at Chapel Hill, Chapel Hill, North Carolina 27514