

HARDY SPACES AND JENSEN MEASURES¹

BY

TAKAHIKO NAKAZI

ABSTRACT. Suppose A is a subalgebra of $L^\infty(m)$ on which m is multiplicative. In this paper, we show that if m is a Jensen measure and $A + \bar{A}$ is dense in $L^2(m)$, then $A + \bar{A}$ is weak-* dense in $L^\infty(m)$, that is, A is a weak-* Dirichlet algebra. As a consequence of it, it follows that if $A + \bar{A}$ is dense in $L^4(m)$, then A is a weak-* Dirichlet algebra. (It is known that even if $A + \bar{A}$ is dense in $L^3(m)$, A is not a weak-* Dirichlet algebra.) As another consequence, it follows that if B is a subalgebra of the classical Hardy space H^∞ containing the constants and dense in H^2 , then B is weak-* dense in H^∞ .

1. Introduction. Let (X, \mathcal{Q}, m) be a nontrivial probability measure space and A a subalgebra of $L^\infty = L^\infty(m)$ containing the constants. Suppose m is multiplicative on A , that is, for $f, g \in A$ we have $\int_X fg \, dm = \int_X f \, dm \int_X g \, dm$. The abstract Hardy space $H^p = H^p(m)$, $0 < p \leq \infty$, associated with A is defined as follows. For $0 < p < \infty$, H^p is the $L^p = L^p(m)$ -closure of A , while H^∞ is defined to be the weak-* closure of A . If $A + \bar{A}$ is weak-* dense in L^∞ , A is called a weak-* Dirichlet algebra, which was introduced by T. P. Srinivasan and J. K. Wang [8]. The theory of weak-* Dirichlet algebras has emerged as the correct setting for many of the central results of abstract analytic function theory.

K. Hoffman and H. Rossi [5] gave an example such that even if $A + \bar{A}$ is dense in L^3 , A is not a weak-* Dirichlet algebra. While G. Lumer [6] showed that if $A + \bar{A}$ is dense in L^p for all finite p , $H^p \cap L^\infty$ is a weak-* Dirichlet algebra. Recently, the author [7] proved that if $A + \bar{A}$ is dense in L^4 , then $H^4 \cap L^\infty$ is a weak-* Dirichlet algebra. Hence if $f \in A$, then [8]

$$\int_X \log |f| \, dm \geq \log \left| \int_X f \, dm \right|.$$

We say m a Jensen measure when functions in A satisfy the inequality above.

In this paper, we show that if m is a Jensen measure and $A + \bar{A}$ is dense in $L^2(m)$, then A is a weak-* Dirichlet algebra. If $A + \bar{A}$ is dense in L^4 , then m is a Jensen measure by the remark above and so A is a weak-* Dirichlet algebra. Hence we answer affirmatively the question left open by Hoffman-Rossi [5], Lumer [6] and the author [7]. Moreover the result is applied to give another proof of a theorem of S. D. Fisher for backward shift invariant subalgebras [1].

Received by the editors November 17, 1981.

1980 *Mathematics Subject Classification.* Primary 46J10, 46J15, 47A15.

Key words and phrases. Hardy spaces, Jensen measures, weak-* Dirichlet algebras, backward shift invariant subalgebras.

¹ This research is partially supported by KAKENHI.

©1982 American Mathematical Society
0002-9947/82/0000-0301/\$02.25

2. Jensen measure. Let A be a subalgebra of L^∞ containing the constants and m a multiplicative measure on it. Set $A_0 = \{f \in A: \int f dm = 0\}$.

LEMMA 1. *Suppose $A + \bar{A}$ is dense in L^2 . If $w \in L^\infty$ is a nonnegative function with $w^{-1} \in L^\infty$, then*

$$\inf_{f \in A_0} \int |1 - f|^2 w dm = \left(\inf_{g \in A_0} \int |1 - g|^2 w^{-1} dm \right)^{-1}.$$

PROOF. If $f, g \in A_0$, by Schwarz's inequality,

$$\int |1 - f|^2 w dm \geq \left(\int |1 - g|^2 w^{-1} dm \right)^{-1}.$$

There exists a unique f_0 in the closure of A_0 in $L^2(w dm)$ such that

$$\inf_{f \in A_0} \int |1 - f|^2 w dm = \int |1 - f_0|^2 w dm.$$

Then, by the minimum property of $\int |1 - f_0|^2 w dm$, $1 - f_0$ is orthogonal to A_0 in $L^2(w dm)$. Set $h_0 = (1 - f_0)w$. Since the infimum is positive, then

$$\begin{aligned} \int |1 - f_0|^2 w dm &= \int (1 - \bar{f}_0)(1 - f_0) w dm \\ &= \int (1 - f_0) w dm = \int h_0 dm > 0. \end{aligned}$$

By the hypothesis, \bar{H}^2 is the orthogonal complement of A_0 in L^2 and so $h_0 \in \bar{H}^2$. Hence $g_0 = \bar{h}_0 / \int \bar{h}_0 dm$ belongs to H^2 and

$$\left(\int h_0 dm \right)^2 \int |1 - (1 - g_0)|^2 w^{-1} dm = \int |1 - f_0|^2 w dm = \int h_0 dm.$$

Since $w, w^{-1} \in L^\infty$, $1 - g_0$ belongs to the closure of A_0 in $L^2(w^{-1} dm)$ and

$$\int h_0 dm = \left(\int |1 - (1 - g_0)|^2 w^{-1} dm \right)^{-1}.$$

This implies the lemma.

LEMMA 2 (SZEGŐ'S THEOREM). *Suppose $A + \bar{A}$ is dense in L^2 and m is a Jensen measure. If $w \in L^1$ is a nonnegative function, then*

$$\inf_{f \in A_0} \int |1 - f|^2 w dm = \exp \int \log w dm.$$

PROOF. By the inequality of arithmetic and geometric means and Jensen's inequality, for any $f, g \in A_0$,

$$\int |1 - f|^2 w dm \geq \exp \int \log w dm$$

and

$$\int |1 - g|^2 w^{-1} dm \geq \exp \int \log w^{-1} m$$

if m is a Jensen measure. If $w, w^{-1} \in L^\infty$, by Lemma 1

$$\inf_{f \in A_0} \int |1 - f|^2 w \, dm = \exp \int \log w \, dm.$$

If $w^{-1} \notin L^\infty$ with $w \in L^\infty$, for any $\epsilon > 0$,

$$\begin{aligned} \exp \int \log(w + \epsilon) \, dm &= \inf \int |1 - f|^2 (w + \epsilon) \, dm \\ &\geq \inf \int |1 - f|^2 w \, dm \geq \exp \int \log w \, dm. \end{aligned}$$

and so letting ϵ tend to zero, the lemma follows. For any $w \in L^1$, let $w_n = \min\{w, n\}$, then

$$\begin{aligned} \exp \int \log w \, dm &\geq \left(\inf \int |1 - g|^2 w^{-1} \, dm \right)^{-1} \\ &\geq \left(\inf \int |1 - g|^2 w_n^{-1} \, dm \right)^{-1} = \exp \int \log w_n \, dm \end{aligned}$$

and so letting n tend to infinity, the lemma follows.

THEOREM 1. $A + \bar{A}$ is dense in L^2 and m is a Jensen measure if and only if A is a weak-* Dirichlet algebra.

PROOF. If A is a weak-* Dirichlet algebra, then $A + \bar{A}$ is dense in L^2 clearly and it is known [8] that m is a Jensen measure. If $A + \bar{A}$ is dense in L^2 and m is a Jensen measure, then Szegő's theorem is valid by Lemma 2. Srinivasan and Wang [8] imply that Szegő's theorem is equivalent to that $A + \bar{A}$ is weak-* dense in L^∞ .

THEOREM 2. Let A be a subalgebra of L^∞ containing the constants and m a multiplicative measure on it. If $A + \bar{A}$ is dense in L^2 , then the following (1) ~ (6) are equivalent.

- (1) $A + \bar{A}$ is weak-* dense in L^∞ , that is, A is a weak-* Dirichlet algebra.
- (2) $A + \bar{A}$ is dense in L^4 .
- (3) m is a Jensen measure.
- (4) If $f \in H^1$ is a real function, then f is a constant.
- (5) If $f \in H^{1/2}$ is a nonnegative function, then f is a constant.
- (6) There is a constant γ_p , defined for $0 < p < 1$, such that

$$\|f\|_p \leq \gamma_p \|f + \bar{g}\|_1, \quad f \in A, g \in A_0.$$

PROOF. (1) \Leftrightarrow (3) is equivalent to Theorem 1. (1) \Leftrightarrow (2) is clear by the remark in Introduction and (1) \Leftrightarrow (3). (1) \Rightarrow (6) is known (cf. [2, p. 107]). (6) \Rightarrow (4) is clear. (4) \Rightarrow (3) Suppose $w \in L^\infty$ is a nonnegative function with $w^{-1} \in L^\infty$. Let f_0 (resp. g_0) be the orthogonal projection of 1 into the closure of A_0 in $L^2(w \, dm)$ (resp. $L^2(w^{-1} \, dm)$). Then by Lemma 1 and Schwarz's lemma, $|1 - f_0|^2 w = |1 - g_0|^2 w^{-1}$ and $(1 - f_0)(1 - g_0) = k \geq 0$. If $f = 1 - g_0$, then $kw = |f|^2$ and $f \in H^2$ and $k \in H^1$ because $w, w^{-1} \in L^\infty$. By the hypothesis, k is a constant 1. Thus $w = |f|$ for $f, f^{-1} \in H^2 \cap L^\infty$. This implies that $H^2 \cap L^\infty$ is a logmodular algebra and so m is a Jensen measure [4]. (1) \Rightarrow (5) is known [9]. (5) \Rightarrow (4) is clear.

(5) is Neuwirth-Newman's theorem and (6) is Kolmogoroff's theorem. Even if $A + \bar{A}$ is not dense in L^2 , (3) implies (5) (cf. [3, pp. 135 ~ 136]).

3. Subalgebras of H^∞ on the unit circle. Let T be the unit circle in the complex plane and $d\theta/2\pi$ the normalized Lebesgue measure on T . In this section, we consider A, L^∞ in case $X = T$ and $m = d\theta/2\pi$. If A is the set of all analytic polynomials on T , then A is a weak-* Dirichlet algebra of $L^\infty(T) = L^\infty(d\theta/2\pi)$. Then, for $0 < p \leq \infty$, $H^p(T) = H^p(d\theta/2\pi)$ is the classical Hardy space.

THEOREM 4. *Let B be a subalgebra of $H^\infty(T)$ containing the constants. If B is L^2 -dense in $H^2(T)$, then B is weak-* dense in $H^\infty(T)$.*

PROOF. Since $L^2(T) = H^2(T) + e^{-i\theta}\overline{H^2(T)}$ and $d\theta/2\pi$ is a Jensen measure, and B is L^2 -dense in $H^2(T)$, so it is a weak-* Dirichlet algebra of $L^\infty(T)$ by Theorem 1, then $H^2(T) \cap L^\infty(T) = H^\infty(T)$ is a weak-* closure of B by Theorem 2.4.1 of [8].

COROLLARY (S. D. FISHER). *Let B be a nontrivial subalgebra of $H^\infty(T)$ which (i) contains the constants, (ii) is weak-* closed, and (iii) contains $e^{-i\theta}f$ whenever $f \in B$ and $\hat{f}(0) = \int_0^{2\pi} f d\theta/2\pi = 0$. Then $B = H^\infty(T)$.*

PROOF. Let \mathfrak{N} be a closure of B in $L^2(T)$, then the orthogonal complement \mathfrak{N}^\perp of \mathfrak{N} in $H^2(T)$ is a shift invariant subspace, that is, $e^{i\theta}\mathfrak{N}^\perp \subset \mathfrak{N}^\perp$. By the well-known theorem of Beurling, $\mathfrak{N}^\perp = qH^2(T)$ for some inner function q with $\hat{q}(0) = 0$ if $\mathfrak{N}^\perp \neq \{0\}$. $e^{-i\theta}q$ is orthogonal to $qH^2(T)$ and so $e^{-i\theta}q \in \mathfrak{N} \cap L^\infty(T)$. Since $\mathfrak{N} \cap L^\infty$ is an algebra, $e^{-i2\theta}q^2$ is orthogonal to $qH^2(T)$ and so $\int qe^{-in\theta}d\theta/2\pi = 0$ for $n \geq 1$. Thus q is a zero constant and so this implies $\mathfrak{N}^\perp = \{0\}$ and $\mathfrak{N} = H^2(T)$. Theorem 4 implies $B = H^\infty(T)$.

REFERENCES

1. S. D. Fisher, *Algebras of bounded functions invariant under the restricted backward shift*, J. Funct. Anal. **12** (1973).
2. T. W. Gamelin, *Uniform algebras*, Prentice-Hall, Englewood Cliffs, N. J., 1969.
3. ———, *Uniform algebras and Jensen measures*, London Math. Soc. Lecture Note Series 32, Cambridge Univ. Press, London, 1978.
4. K. Hoffman, *Analytic functions and logmodular Banach algebras*, Acta Math. **108** (1962), 271–317.
5. K. Hoffman and H. Rossi, *Function theory from a multiplicative linear functional*, Trans. Amer. Math. Soc. **116** (1965), 536–543.
6. G. Lumer, *Herglotz transformation and H^p theory*, Bull. Amer. Math. Soc. **71** (1965), 725–730.
7. T. Nakazi, *A note on weak-* Dirichlet algebras*, Bull. Acad. Polon. Sci. Sér. Sci. Math. **27** (1979), 553–555.
8. T. P. Srinivasan and J. K. Wang, *Weak-* Dirichlet algebras*, Proc. Internat. Sympos. Function Algebras (Tuland Univ. 1965), Scott-Foresman, Chicago, Ill., 1966, pp. 216–249.
9. K. Yabuta, *A note on extremum problems in abstract Hardy spaces*, Arch. Math. **23** (1972), 54–57.

DIVISION OF APPLIED MATHEMATICS, RESEARCH INSTITUTE OF APPLIED ELECTRICITY, HOKKAIDO UNIVERSITY, SAPPORO 060, JAPAN