

CONVEXITY AND TIGHTNESS FOR RESTRICTIONS OF  
HAMILTONIAN FUNCTIONS TO FIXED POINT SETS OF  
AN ANTISYMPLECTIC INVOLUTION

BY

J. J. DUISTERMAAT<sup>1</sup>

**ABSTRACT.** The Kostant convexity theorem for real flag manifolds is generalized to a Hamiltonian framework. More precisely, it is proved that if  $f$  is the momentum mapping for a Hamiltonian torus action on a symplectic manifold  $M$  and  $Q$  is the fixed point set of an antisymplectic involution of  $M$  leaving  $f$  invariant, then  $f(Q) = f(M)$  is a convex polytope. Also it is proved that the coordinate functions of  $f$  are tight, using "half-turn" involutions of  $Q$ .

**1. Introduction.** Recently the theorem of Kostant [10], on convexity of certain projections of complex flag manifolds, has been generalized to a Hamiltonian framework by Guillemin and Sternberg [7], and, independently, by Atiyah [2]. However, Kostant proved his theorem for the real flag manifolds as well, and it is the first purpose of this paper to show that also this real version has a generalization in a Hamiltonian setting. More specifically, let  $M$  be a compact connected smooth manifold of dimension  $2n$ , provided with a symplectic form  $\sigma$ . Let  $T$  be a torus acting on  $M$  in a Hamiltonian way, with Hamiltonian functions  $f_X$ ,  $X \in \mathfrak{t}$ , and momentum mapping  $f: M \rightarrow \mathfrak{t}^*$ . Here  $\mathfrak{t}$  is the Lie algebra of  $T$  and  $\mathfrak{t}^*$  its dual, see §2 for more details. Furthermore, let  $\tau$  be a smooth involution of  $M$  such that  $\tau^*\sigma = -\sigma$  and such that  $f_X \circ \tau = f_X$  for all  $X \in \mathfrak{t}$ . (One can always arrange the latter condition by passing to a suitable subtorus  $T_0$  of  $T$ , the new momentum mapping then being equal to  $f$  followed by the natural projection  $\mathfrak{t}^* \rightarrow \mathfrak{t}_0^*$ .) Let  $Q$  be the fixed point set of  $\tau$ , which we assume to be nonvoid. Then  $f(Q) = f(M)$ , the convex hull of finitely many points in  $\mathfrak{t}^*$ . For a more detailed description of the extremal points of  $f(Q) = f(M)$ , see Theorem 2.5 and formula (2.32). The proof follows the pattern of Guillemin and Sternberg [7], which in turn was inspired by Heckman [8].

If  $M$  is a Kähler manifold, then a more refined result is true in terms of gradient flows, this will be discussed in §4.

Secondly, Atiyah [2], following Frankel [6], observed that the Hamiltonian functions  $f_X$ ,  $X \in \mathfrak{t}$ , are tight in the sense that the sum of the Betti numbers of  $M$  is equal to the sum of the Betti numbers of the critical set of  $f_X$ . In the case of isolated critical points this implies that  $f_X$  is a Morse function on  $M$  with the minimal number of critical points. This too generalizes a known result for complex flag manifolds, but

---

Received by the editors December 7, 1981 and, in revised form, January 21, 1982.

1980 *Mathematics Subject Classification.* Primary 53C15; Secondary 53C55, 58F05, 55M20.

<sup>1</sup>Partially supported by NSF Grant MCS 80-23356, while visiting the University of California at Berkeley.

© 1983 American Mathematical Society  
0002-9947/82/0000-0437/\$04.50

which actually is true for the real flag manifolds as well, due to Takeuchi and Kobayashi [13], see also Duistermaat, Kolk and Varadarajan [3, §4]. In §3 it will be shown that also the tightness generalizes to our setting. The theorem is that  $\dim H^*(Q; \mathbf{Z}/2\mathbf{Z}) = \dim H^*(C_X; \mathbf{Z}/2\mathbf{Z})$  if  $C_X$  is the critical set of the function  $f_X|_Q$ ,  $X \in \mathfrak{t}$ .

Finally in §5 we describe how the real flag manifolds fit into the framework described above.

It is a pleasure for me to thank Michael Atiyah for his suggestion, made to me at the Arbeitstagung in Bonn in June 1981 (and in [2]), that the real flag manifolds really should be treated as the fixed point set of an involution in a symplectic manifold. I also thank John Millson, resp. Alan Weinstein for some discussions at UCLA, resp. Berkeley, which stimulated me further.

**2. Convexity.** We recall that  $(M, \sigma)$  is a compact connected symplectic manifold with a Hamiltonian action of a torus  $T$  on it. That is, there is a linear map  $X \mapsto f_X$  from the Lie algebra  $\mathfrak{t}$  of  $T$  to the space of smooth functions on  $M$ , such that

(2.1) For each  $X \in \mathfrak{t}$ , the infinitesimal action  $\tilde{X}$  of  $X$  on  $M$  is equal to the Hamilton vector field of the function  $f_X$ , and

(2.2) The functions  $f_X$ ,  $X \in \mathfrak{t}$ , are in involution.

In formula, (2.1) reads

$$(2.1') \quad \tilde{X} \lrcorner \sigma = -df_X, \quad X \in \mathfrak{t},$$

whereas assuming this, (2.2) is equivalent to

$$(2.2') \quad \tilde{X} \lrcorner df_Y = 0 \quad \text{for all } X, Y \in \mathfrak{t},$$

that is,  $f_Y$  is constant along the  $T$ -orbits in  $M$ .

The mapping  $f: M \rightarrow \mathfrak{t}^*$ , defined by

$$(2.3) \quad \langle X, f(m) \rangle = f_X(m), \quad m \in M, X \in \mathfrak{t},$$

is called the *momentum mapping* of the Hamiltonian  $T$ -action.

The next ingredient which we introduce is a smooth map  $\tau: M \rightarrow M$  which is an *involution* of  $M$ , that is

$$(2.4) \quad \tau \circ \tau = \text{identity on } M,$$

and which is *antisymplectic*, that is

$$(2.5) \quad \tau^* \sigma = -\sigma.$$

We will assume that the Hamiltonian  $T$ -action and the involution  $\tau$  are related to each other by the condition that the functions  $f_X$  are  $\tau$ -invariant, that is

$$(2.6) \quad \tau^* f_X = f_X \quad \text{for all } X \in \mathfrak{t}.$$

Assuming (2.5), this is equivalent to

$$(2.7) \quad \tau^* \tilde{X} = -\tilde{X}, \quad X \in \mathfrak{t},$$

in view of (2.1'). In turn, (2.7) is equivalent to

$$(2.8) \quad \tau \tilde{g} \tau^{-1} = \tilde{g}^{-1}, \quad g \in T,$$

if  $\tilde{g}$  denotes the action of  $g$  on  $M$ . That is,  $\tau$  maps  $T$ -orbits to  $T$ -orbits, but at the same time reverses the time on the orbits of the 1-parameter subgroups of  $T$ .

If  $G$  is a compact group acting by smooth mappings on a manifold  $M$ , then averaging over  $G$  of an arbitrary Riemannian metric on  $M$  leads to a  $G$ -invariant Riemannian metric  $\beta$  on  $M$ . If  $m$  is a fixed point for the  $G$ -action, then the exponential map, centered at  $m$ , with respect to  $\beta$ , intertwines the linear action of  $G$  on  $T_m M$  with the local action of  $G$  around  $m$ . That is, the  $G$ -action is linear orthogonal on suitable local coordinates. In particular the fixed point set for the action of any subset of  $G$  has finitely many components, each of which is a closed smooth submanifold of  $M$ .

We assume that the fixed point set of  $\tau$  in  $M$ ,

$$(2.9) \quad Q = \{m \in M; \tau(m) = m\},$$

is nonvoid. It has finitely many smooth, compact, connected components. From (2.5) we obtain that if  $m \in Q$ , then both  $T_m M = \text{Ker}(D\tau(m) - I)$  and

$$(2.10) \quad P_m = \text{Ker}(D\tau(m) + I)$$

are isotropic subspaces for  $\sigma_m$ . Because also

$$(2.11) \quad T_m M = T_m Q \oplus P_m,$$

the conclusion is that  $T_m Q$  and  $P_m$  are Lagrange subspaces of  $T_m M$ . That is,  $Q$  is a Lagrange submanifold of  $M$ , and the  $P_m, m \in Q$ , form a Lagrange subbundle of  $T_Q M$ , complementary to  $TQ$ . In fact, Meyer [11] showed that there is an open  $\tau$ -invariant neighborhood  $U$  of  $Q$  in  $M$  and a symplectic diffeomorphism  $\Phi$  from  $U$  onto an open neighborhood  $V$  of the zero section in  $T^*Q$ , such that  $\Phi\tau\Phi^{-1}$  maps  $p \in (T_q Q)^* \cap V$  to  $-p$ , for all  $q \in Q$ .

Another rigidity property of compact Lie group actions on a compact manifold  $M$  is that there are only finitely many orbit types (Mostow [12], Yang [15]). In the case of the action of a torus  $T$  this means that there are only finitely many possibilities for the stabilizer groups

$$(2.12) \quad T_m = \{g \in T; g(m) = m\},$$

as  $m$  ranges over  $M$ . This fact will be used both in the proof of the convexity and of the tightness theorem mentioned in the introduction.

2.1. LEMMA. *Let  $m \in Q, X \in \mathfrak{t}$ . Then  $d(f_X|Q)(m) = 0$  implies that  $df_X(m) = 0$ , which in turn is equivalent to the condition that  $m$  is a fixed point for the action of  $\exp tX, t \in \mathbf{R}$ .*

PROOF.  $T_m Q = \text{Ker}(D\tau(m) - I) = \{u + D\tau(m)(u); u \in T_m M\}$ . So using (2.6), we get  $0 = df_X(m)(u + D\tau(m)(u)) = 2df_X(m)(u)$  for all  $u \in T_m M$ .

2.2. PROPOSITION. *Let  $m \in Q, D(f|Q)(m) = 0$ . Then there exist smooth symplectic local coordinates  $q_j, p_j$  for a neighborhood  $U$  of  $m$  (zero at  $m$ ), such that, for suitable  $\omega_j \in \mathfrak{t}^*$ ,*

$$(2.13) \quad f_X = f_X(m) + \sum_{j=1}^n \omega_j(X) \cdot (q_j^2 + p_j^2)/2,$$

and moreover

$$(2.14) \quad \tau: (q, p) \mapsto (q, -p)$$

in  $U$ . In particular  $U \cap Q = \{p = 0\}$ .

PROOF. Although the proof is a combination of the by now standard arguments in Meyer [11] and Guillemin-Sternberg [7, §4], we will give it in some detail for the convenience of the reader. We begin with the result on the tangent level.

In view of Lemma 2.1,  $m$  is not only a fixed point for  $\tau$ , but also for the  $T$ -action. Then the

$$(2.15) \quad X_m = D\tilde{X}(m): T_m M \rightarrow T_m M, \quad X \in \mathfrak{t},$$

form a commuting family of infinitesimally symplectic transformations, antisymmetric with respect to  $\beta_m$ . In particular they are simultaneously diagonalizable over  $\mathbb{C}$  with purely imaginary eigenvalues. That is,

$$(2.16) \quad T_m M \otimes \mathbb{C} = \sum_{\lambda}^{\oplus} E_{\lambda}, \quad X_m|_{E_{\lambda}} = \lambda(X) \cdot \text{identity on } E_{\lambda};$$

here  $X_m$  is extended to a complex linear endomorphism of  $T_m M \otimes \mathbb{C}$  and  $\lambda$  is real linear:  $\mathfrak{t} \rightarrow \mathbb{C}$ , taking only purely imaginary values. The complex conjugation maps  $E_{\lambda}$  to  $E_{\bar{\lambda}} = E_{-\lambda}$ . Because the  $X_m$  are infinitesimally symplectic, the spaces

$$(2.17) \quad E_{\lambda, -\lambda} = (E_{\lambda} + E_{-\lambda}) \cap T_m M$$

are mutually  $\sigma_m$ -orthogonal. Because they span  $T_m M$ , they form a symplectic vector space decomposition of  $T_m M$ .

Now (2.7) implies that

$$(2.18) \quad D\tau(m) \circ X_m = -X_m \circ D\tau(m),$$

showing that  $D\tau(m)$  maps  $E_{\lambda}$  to  $E_{-\lambda}$ , so it leaves  $E_{\lambda, -\lambda}$  invariant. This reduces the problem to the case that all  $X_m$  are real multiples of one infinitesimally symplectic mapping  $J$ , the square of which we can take equal to  $-I$ . Moreover,  $J$  maps  $T_m Q$  to  $P_m$  and  $P_m$  to  $T_m Q$ . Now

$$(2.19) \quad \langle u, v \rangle = \sigma_m(u, Jv), \quad u, v \in T_m Q,$$

defines a nondegenerate symmetric bilinear form on  $T_m Q$ . We can write  $T_m Q = Q_m^+ \oplus Q_m^-$ ,  $\langle Q_m^+, Q_m^- \rangle = 0$ , with (2.19) being positive, resp. negative definite, on  $Q_m^+$ , resp.  $Q_m^-$ . The spaces  $Q_m^+ + J(Q_m^+)$  and  $Q_m^- + J(Q_m^-)$  are  $\sigma_m$ -orthogonal to each other, leading to a  $D\tau(m)$ -invariant symplectic vector space decomposition. This reduces the problem to the case that (2.19) is definite.

In the positive case, let  $e_1, \dots, e_n$  be an orthonormal basis of  $T_m Q$  with respect to (2.19). Then

$$(2.20) \quad (q, p) \mapsto \sum_j (q_j \cdot e_j - p_j \cdot J e_j)$$

is a symplectic mapping, with respect to the symplectic form  $\sum_j dp_j \wedge dq_j$  on  $(q, p)$ -space. It intertwines  $J$  with the mapping  $(q, p) \mapsto (p, -q)$ , which, as a linear vector field, is Hamiltonian with Hamiltonian function

$$(2.21) \quad \sum_j (q_j^2 + p_j^2)/2.$$

In the negative definite case we get (2.21) with a minus sign in front. This proves the proposition on the tangent level.

For the *local* normal form we already know that there are local coordinates linearizing  $\tau$  and the  $T$ -action, hence also the vector fields  $\tilde{X}$ ,  $X \in \mathfrak{t}$ . However, in these coordinates the symplectic form  $\sigma$  will in general not be equal to the constant symplectic form  $\sigma_m = \sum_j dp_j \wedge dq_j$ . The proof is finished by applying the following version (also known to Weinstein) of the equivariant Darboux lemma of Weinstein [14].

2.3. LEMMA. *Let  $G$  be a compact group acting by diffeomorphisms on a symplectic manifold  $(M, \sigma)$ , such that  $\tilde{g}^*\sigma = \epsilon(g) \cdot \sigma$ ,  $\epsilon$  a continuous homomorphism:  $G \rightarrow \{-1, +1\}$ . Let  $m$  be a fixed point for the  $G$ -action,  $\sigma'$  another smooth symplectic form defined on a neighborhood of  $m$ , such that  $\tilde{g}^*\sigma' = \epsilon(g) \cdot \sigma'$  for all  $g \in G$ , and  $\sigma'_m = \sigma_m$ . Then there exists a  $G$ -equivariant local diffeomorphism  $\Psi$  around  $m$ , such that  $\Psi(m) = m$ ,  $D\Psi(m) = I$ , and  $\Psi^*\sigma' = \sigma$ .*

PROOF. Write  $\sigma_t = \sigma + t \cdot (\sigma' - \sigma)$ , so that  $\sigma_0 = \sigma$ ,  $\sigma_1 = \sigma'$ ,  $\tilde{g}^*\sigma_t = \epsilon(g) \cdot \sigma_t$  for all  $g \in G$ ,  $t \in [0, 1]$ . One attempts to find local diffeomorphisms  $\Psi_t$ ,  $G$ -equivariant, depending smoothly on  $t$ , such that  $\Psi_0 = \text{identity}$ , and  $\Psi_t(m) = m$ ,  $D\Psi_t(m) = I$ ,  $\Psi_t^*\sigma_t = \sigma$  for all  $t \in [0, 1]$ . Differentiating with respect to  $t$  one finds that the velocity field  $v_t$  has to satisfy the equations

$$(2.22) \quad d(v_t \lrcorner \sigma_t) = \sigma - \sigma', \quad v_t(m) = 0, \quad Dv_t(m) = 0,$$

for all  $t \in [0, 1]$ . By the Poincaré lemma there is a smooth 1-form  $\alpha$  such that  $d\alpha = \sigma - \sigma'$ ,  $\alpha(m) = 0$ ,  $D\alpha(m) = 0$ . Let  $v_t$  be the unique vector field such that  $v_t \lrcorner \sigma_t = \alpha$ . Then  $v_t$  depends smoothly on  $t$  and satisfies (2.22). Each  $\tilde{g}^*v_t$ ,  $g \in G$ , will again satisfy (2.22), because

$$\begin{aligned} \epsilon(g) \cdot (\sigma - \sigma') &= \tilde{g}^*(\sigma - \sigma') = \tilde{g}^*d(v_t \lrcorner \sigma_t) \\ &= d\tilde{g}^*(v_t \lrcorner \sigma_t) = d(\tilde{g}^*v_t \lrcorner \tilde{g}^*\sigma_t) \\ &= d(\tilde{g}^*v_t \lrcorner \epsilon(g) \cdot \sigma_t) = \epsilon(g) \cdot d(\tilde{g}^*v_t \lrcorner \sigma_t). \end{aligned}$$

So averaging over  $G$  we get a smooth  $G$ -invariant vector field  $v_t$ , depending smoothly on  $t$ , and satisfying (2.22). Integrating it we get a  $G$ -equivariant 1-parameter family of local diffeomorphisms  $\Psi_t$ , such that  $\Psi_t(m) = m$ ,  $D\Psi_t(m) = I$ , and  $t \mapsto \Psi_t^*\sigma_t$  is constant, hence equal to  $\sigma$ . Taking  $\Psi = \Psi_1$ , the lemma is proved.

Ignoring the  $T$ -action in Proposition 2.2, we have, locally, recovered the theorem of Meyer mentioned before. Moreover, the set

$$(2.23) \quad C = \{m \in Q; D(f|Q)(m) = 0\}$$

is equal to  $F \cap Q$ , where  $F$  is the fixed point set of the  $T$ -action in  $M$ . Each connected component  $C_k$  of  $C$ , which in the local coordinates of Proposition 2.2 reads as

$$(2.24) \quad \{(q, p); p = 0 \text{ and } q_j = 0 \text{ whenever } \omega_j \neq 0\},$$

is a Lagrange submanifold of some connected component  $F_{j(k)}$  of  $F$ , the connected components  $F_j$  of  $F$  being symplectic submanifolds of  $M$ .  $F_j$  is  $\tau$ -invariant if it meets

$Q$ , and then the  $C_k$  with  $j(k) = j$  are the connected components of the fixed point set for the involution  $\tau$  in  $F_j$ .

We now want to prove that  $f(Q)$  is equal to the convex hull of  $f(C)$ , which consists of only finitely many points because  $f$  is obviously constant on each  $C_k$ . As in Guillemin-Sternberg [7, §5], the first step is

2.4. LEMMA. *For each  $X \in \mathfrak{t}$ , the function  $f_X|_Q$  has a unique local maximal value.*

PROOF. Fix  $X \in \mathfrak{t}$ . Let  $m \in Q$  be a critical point of  $f_X|_Q$ . From Lemma 2.1 we know that  $m$  is a fixed point for the action of  $\exp tX$ ,  $t \in \mathbf{R}$ . Replacing, in this proof,  $T$  by the closure of  $\{\exp tX, t \in \mathbf{R}\}$  in  $T$ , we get that  $D(f|_Q)(m) = 0$ . A glance at (2.13) and the local characterization of  $Q$  shows that  $m$  is a local maximum for  $f_X|_Q$  if and only if  $m$  is a local maximum for  $f_X$ , both conditions being equivalent to  $\omega_j(X) \leq 0$  for all  $j$ . A Morse theoretic argument gives that the set of points in  $M$  where  $f_X$  has a local maximum is connected, see Atiyah [2] and Guillemin-Sternberg [7]. So  $f_X$  on  $M$  has only one local maximal value, and therefore the same must hold for  $f_X|_Q$ .

Now, let  $\xi \in \mathfrak{t}^*$  be a boundary point of  $f(Q)$ , and let  $m \in Q$  be such that  $f(m) = \xi$ . Then

$$(2.25) \quad \mathfrak{t}_m = \{X \in \mathfrak{t}; d(f_X|_Q)(m) = 0\}$$

is nonzero, otherwise  $D(f|_Q)(m)$  would be surjective, contradicting that  $\xi$  is a boundary point. In view of Lemma 2.1 we may replace  $T$  by the subtorus  $\exp \mathfrak{t}_m$ , fixing  $m$ , and apply Proposition 2.2. Write

$$(2.26) \quad \gamma_m = \sum_{j=1}^n \vartheta_j \cdot \omega_j, \quad \vartheta_j \geq 0,$$

for the convex cone in  $\mathfrak{t}_m^*$  generated by the vectors  $\omega_j$  of (2.13), with  $\mathfrak{t}$  replaced by  $\mathfrak{t}_m$ . Then (2.13) shows that there is a neighborhood  $U$  of  $m$  in  $Q$  and a neighborhood  $V$  of  $\pi_m(\xi)$  in  $\mathfrak{t}_m^*$ , such that

$$(2.27) \quad \pi_m(f(U)) = V \cap (\pi_m(\xi) + \gamma_m).$$

Here  $\pi_m$  is the projection:  $\mathfrak{t}^* \rightarrow \mathfrak{t}_m^*$  obtained by restriction of linear forms to  $\mathfrak{t}_m$ . This implies of course that

$$(2.28) \quad f(U) \subset \xi + \pi_m^{-1}(\gamma_m).$$

Note that  $\text{Ker } \pi_m = \text{Im } D(f|_Q)(m)$ . The fact that there are only finitely many possibilities for the  $T_m$  in (2.12), and that  $\mathfrak{t}_m$  is equal to the Lie algebra of  $T_m$ , leads to  $\text{Im } D(f|_Q)(m') \supset \text{Im } D(f|_Q)(m)$  for all  $m'$  in a neighborhood of  $m$ , so (2.27) can actually be strengthened to

$$(2.29) \quad f(U) = W \cap (\xi + \pi_m^{-1}(\gamma_m))$$

for some neighborhood  $W$  of  $\xi$  in  $\mathfrak{t}^*$ .

Because  $\xi$  is a boundary point of  $f(Q)$ , we see from (2.13) and (2.29) that  $\gamma_m \neq \mathfrak{t}_m^*$ , so the dual cone

$$(2.30) \quad \delta_m = \{X \in \mathfrak{t}_m; \omega_j(X) \leq 0 \text{ for all } j\}$$

must contain nonzero  $X$ . However,  $X \in \mathfrak{t}_m$  means in view of (2.28) that  $\langle X, f(U) \rangle \leq \langle X, \xi \rangle$ , that is,  $\langle X, \xi \rangle$  is a local maximal value for  $f_X|_Q$ . Since Lemma 2.4 implies that local maximal values are global maximal values,  $\langle X, f(Q) \rangle \leq \langle X, \xi \rangle$  for all  $X \in \delta_m$ . That is, (2.28) can be strengthened to

$$(2.31) \quad f(Q) \subset \xi + \pi_m^{-1}(\gamma_m).$$

In particular, since  $\delta_m \neq \{0\}$ , each boundary point of  $f(Q)$  is on the boundary of a half-space containing  $f(Q)$ , so  $f(Q)$  is convex. Because  $f(Q)$  is compact, it is equal to the convex hull of its extremal points. But (2.29) shows that  $\xi$  can only be extremal if  $\mathfrak{t}_m = \mathfrak{t}$ , that is  $m \in C$ , and  $\gamma_m$  is a proper cone in  $\mathfrak{t}^*$ . Also, (2.13) shows that  $\gamma_m$  is constant, say equal to  $\gamma(C_k)$ , if  $m$  runs along a connected component  $C_k$  of  $C$ . Since a nonvoid compact convex set is equal to the intersection of the half-spaces containing it and supported at the extremal points, the local description in (2.29), (2.31) of  $f(Q)$  leads to

$$(2.32) \quad f(Q) = \bigcap_{\gamma(C_k) \text{ a proper cone}} (f(C_k) + \gamma(C_k)).$$

Working on  $M$  and ignoring the involution  $\tau$ , the above arguments also lead to

$$(2.33) \quad f(M) = \bigcap_{\gamma(F_j) \text{ a proper cone}} (f(F_j) + \gamma(F_j)),$$

where the  $F_j$  are the connected components of  $F$ , the fixed point set of the  $T$ -action on  $M$ , and  $\gamma(F_j)$  is the convex cone in  $\mathfrak{t}^*$  generated by the  $\omega_j \in \mathfrak{t}^*$  as in (2.13), in the analogue of Proposition 2.2. disregarding the involution  $\tau$ . Now each  $C_k$  was contained in an  $F_{j(k)}$ , of course  $f(C_k) = f(F_{j(k)})$ . However, (2.13) shows that also  $\gamma(C_k) = \gamma(F_{j(k)})$ . So comparing (2.32) and (2.33) we get  $f(Q) \supset f(M)$ . Because the other inclusion is trivial, we have proved

2.5. THEOREM.  $f(Q) = f(M)$ , and is equal to the convex hull of the finitely many values  $f(m)$  of  $f$  at points  $m \in Q$  such that

$$(2.34) \quad D(f|_Q)(m) = 0 \text{ and the convex cone } \gamma_m \text{ generated by the } \omega_j \in \mathfrak{t}^* \text{ in (2.13) is proper.}$$

Moreover, if  $m' \in M$ ,  $Df(m') = 0$  and the corresponding cone  $\gamma_{m'}$  is proper, then there exists  $m \in Q$  such that (2.34) holds, and  $f(m') = f(m)$ ,  $\gamma_{m'} = \gamma_m$ .

REMARK. The proof shows that the same statements are true if  $Q$  is replaced by any connected component  $Q_0$  of  $Q$ .

3. Tightness. We make the same assumptions as in §2, namely that  $(M, \sigma)$  is a compact symplectic manifold with on it a Hamiltonian action of a torus  $T$ , with Hamilton functions  $f_X$ ,  $X \in \mathfrak{t}$ . Also,  $\tau$  is a smooth involution of  $M$  such that  $\tau^*\sigma = -\sigma$ ,  $\tau^*f_X = f_X$  for all  $X \in \mathfrak{t}$ . In this section we prove

3.1. THEOREM. Let  $C_X$  be the critical set of  $f_X|_Q$ ,  $X \in \mathfrak{t}$ . Then

$$(3.1) \quad \dim H^*(Q; \mathbf{Z}/2\mathbf{Z}) = \dim H^*(C_X; \mathbf{Z}/2\mathbf{Z}).$$

We begin the proof with a reduction to the case of a circle action. By restricting to the closure of  $\{\exp tX; t \in \mathbf{R}\}$  in  $T$ , we may assume that  $\{\exp tX; t \in \mathbf{R}\}$  is dense in

$T$ . In view of Lemma 2.1 we have now  $C_X = F \cap Q$ , where  $F$  is the fixed point set of the action of  $T$  on  $M$ . Because there are only finitely many possibilities for the stabilizer groups  $T_m$  in (2.12), it follows that the collection of  $Y \in \mathfrak{t}$  such that  $C_Y = C_X$  is open and dense in  $\mathfrak{t}$ . Because the set of  $Y$ , such that  $t \mapsto \exp tY$  is periodic, is dense in  $\mathfrak{t}$ , we may assume that  $t \mapsto \exp tX$  is periodic, and multiplying  $X$  by a suitable factor we get  $\exp X = 1$ . All this without changing the critical set.

Now (2.8) implies that  $\tau$  maps  $T$ -orbits to  $T$ -orbits, reversing the time order on them. In particular the  $T$ -orbits through  $m \in Q$  are  $\tau$ -invariant, and

$$(3.2) \quad \tau(\exp t\tilde{X}(m)) = \exp -t\tilde{X}(m).$$

But then  $\tau(\exp \frac{1}{2}\tilde{X}(m)) = \exp -\frac{1}{2}\tilde{X}(m) = \exp \frac{1}{2}\tilde{X}(m)$ , that is

3.2. LEMMA.  $\exp \frac{1}{2}\tilde{X}$  maps  $Q$  to itself,<sup>2</sup> and thereby defines a smooth involution of  $Q$ .

Examples (see the remark below) show that the fixed point set in  $Q$  of  $\exp \frac{1}{2}\tilde{X}$ , which contains  $C_X$ , is not necessarily equal to  $C_X$  (assuming that the minimal period is equal to 1). For this reason we now define, by induction over  $k \in \mathbf{N}$ ,

$$(3.3) \quad Q_{(0)} = Q, \quad Q_{(k)} = \{m \in Q_{(k-1)}; \exp 2^{-k}\tilde{X}(m) = m\}.$$

By induction over  $k$  one proves

3.3. LEMMA.  $\exp 2^{-k}X$  is an involution of  $Q_{(k-1)}$ . The  $Q_{(k)}$  form a decreasing family of subspaces of  $Q$ .  $Q_{(k)}$  has finitely many connected components, each of which is a smooth compact submanifold of  $Q_{(k-1)}$ , resp. of  $Q$ . There exists an  $N \in \mathbf{N}$  such that  $Q_{(N)} = C_X$ .

The last statement follows because such decreasing sequences of submanifolds must stabilize, that is, there is an  $N \in \mathbf{N}$  such that  $Q_{(k)} = Q_{(N)}$  for all  $k \geq N$ . But  $m \in Q_{(k)}$  for all  $k$  implies that  $\tilde{X}(m) = 0$ , hence  $dJ_X(m) = 0$ , or  $m \in C_X$ .

It is known (Floyd [4 or 5, §4]) that if  $B$  is the fixed point set of a periodic transformation of prime period  $p$  in a compact manifold  $A$ , then

$$\dim H^*(A; \mathbf{Z}/p\mathbf{Z}) \geq \dim H^*(B; \mathbf{Z}/p\mathbf{Z}).$$

Lemma 3.3 therefore implies

$$(3.4) \quad \dim H^*(Q; \mathbf{Z}/2\mathbf{Z}) \geq \dim H^*(Q_{(1)}; \mathbf{Z}/2\mathbf{Z}) \geq \dots \geq \dim H^*(C_X; \mathbf{Z}/2\mathbf{Z}).$$

But the opposite inequality  $\dim H^*(Q; R) \leq \dim H^*(C_X; R)$ , valid for general coefficient rings  $R$ , follows from the Bott-Morse inequalities. This proves Theorem 3.1. The idea to conclude tightness by exhibiting the critical set as the fixed point set for a periodic map has been introduced by Frankel [6] in a Kähler framework.

REMARK. If  $m$  is an isolated fixed point for  $\exp \frac{1}{2}\tilde{X}: Q \rightarrow Q$ , then this ‘‘half-turn’’ is the Cartan involution around  $m$  with respect to any invariant Riemannian metric. Suppose now that the isometry group of the connected component  $Q_0$  of  $Q$  acts transitively, as is the case for the real flag manifolds; see §5. Then the existence of isolated fixed points for the half-turn on  $Q_0$  makes  $Q_0$  into a Riemannian symmetric

<sup>2</sup>Note that  $Q$  is not invariant under the  $\tilde{X}$ -flow. In fact  $\tilde{X}(m) \in P_m$  where  $P_m$  is the space complementary to  $T_mQ$ , defined in (2.10).

space. For symmetric real flag manifolds the tightness of  $f_X|_{Q_0}$  was proved in Takeuchi [16, pp. 167–168] by identifying the critical set with the fixed point set of a Cartan involution. Conversely, if  $Q_0$  is a real flag manifold which is not a symmetric space, and if  $f_X|_{Q_0}$  has isolated critical points (as generically is the case), then the fixed point set of  $\exp \frac{1}{2}\tilde{X}$  in  $Q$  cannot be equal to  $C_X$ .

**4. Gradient flows.** If  $\beta$  is a Riemannian metric on  $M$  which is  $T$ - and  $\tau$ -invariant, then the gradient vector fields of the functions  $f_X$ ,  $X \in \mathfrak{t}$ , are also  $T$ - and  $\tau$ -invariant. In particular they are tangent to the fixed point set  $Q$  of  $\tau$ , the gradient flows leaving  $Q$  invariant. The formula  $\beta(u, v) = \sigma(u, Jv)$  defines a tensor field  $J$  on  $M$  which is  $T$ -invariant and  $\tau$ -anti-invariant, and  $\text{grad } f_X$  is equal to  $J$  times the Hamiltonian vector field  $X$  of the function  $f_X$ .

If  $J$  is an integrable almost complex structure, that is if  $M$  is a complex analytic manifold with complex structure equal to  $J$ , then  $\beta + i\sigma$  is a Kähler metric on  $M$ . The  $T$ -action consists of the holomorphic mappings, whereas the involution  $\tau$  is anti-holomorphic, making  $Q$  into a “real subspace” of  $M$  in a strong sense. Because the automorphism group of a compact complex manifold is a complex Lie group, the action of  $T$  extends to a holomorphic action of the complexification  $T_c$  of  $T$ . Its Lie algebra  $\mathfrak{t} \otimes \mathbb{C}$  can be written as  $\mathfrak{t} \oplus \mathfrak{a}$  where  $\mathfrak{a} = i\mathfrak{t}$ . The exponential map is taken to be injective on  $\mathfrak{a}$ , making  $A = \exp \mathfrak{a}$  into a vector subgroup of  $T_c$ , and  $(t, a) \mapsto t \cdot a$  is a diffeomorphism from  $T \times A$  onto  $T_c$ . Now  $\text{grad } f_X = J \cdot \tilde{X} = \widetilde{i \cdot X}$ , so the gradient vector fields together make up the infinitesimal action of  $A$ . In particular the gradient flows commute with each other, in fact this is the only additional assumption which will be used in the sequel.

**4.1. THEOREM.** *Let  $Y$  be an  $A$ -orbit in  $M$  and  $F_j, j = 1, \dots, p$ , the components of the common critical points of the functions  $f_X, X \in \mathfrak{t}$ , which intersect the closure  $\bar{Y}$  of  $Y$  in  $M$ . Then  $f(\bar{Y})$  is equal to the convex polytope  $P$  with extremal points equal to  $c_j = f(F_j), j = 1, \dots, p$ . For each open face  $\varphi$  of  $P$  the inverse image  $f^{-1}(\varphi)$  in  $Y$  consists of a simple  $A$ -orbit, and  $f$  induces a homeomorphism of  $\bar{Y}$  onto  $P$ .*

This is Theorem 2 of Atiyah [2]. That theorem was phrased in terms of  $T_c$ -orbits in  $M$ , but using the invariance of  $f$  under the  $T$ -action, one readily translates it into the above statement. Because  $Q$  is  $A$ -invariant we can apply Theorem 4.1 to the  $A$ -orbits in  $Q$ . Then Theorem 4.1 is a generalization of the corresponding statement for the real flag manifolds, due to Heckman [8, Chapter 2, Theorem 3]. The following result shows that Theorem 4.1 can be regarded as a refinement of Theorem 2.5 in the case of commuting gradient flows.

**4.2. PROPOSITION.** *For all  $m'$  in an open dense subset  $Q'$  of  $Q$ , the set of  $f(m)$  such that  $m \in Q, D(f|_Q)(m) = 0$ , and  $m$  is in the closure of the  $A$ -orbit through  $m'$ , is equal to the set of extremal points of  $f(Q)$ .*

**PROOF.** Without loss of generality we may assume that  $f(Q) = f(M)$  has a nonempty interior, that is

$$(4.1) \quad Q^{\text{reg}} = \{m \in Q; D(f|_Q)(m) \text{ is surjective}\}$$

is nonvoid.

Recall that  $f(Q_0) = f(M)$  for any connected component  $Q_0$  of  $Q$ . Now  $m \notin Q_0^{\text{reg}}$  means that there exists  $X \in \mathfrak{t}$ ,  $X \neq 0$ , such that  $m$  is a critical point for  $f_X$ . For the description of the critical set  $C_X$  of  $f_X|_{Q_0}$  we may, as in the proof of Theorem 3.1, assume that  $t \mapsto \exp tX$  is periodic on  $M$  with minimal positive period equal to 1 when starting on  $Q_0$ . Replacing  $T$  for the moment by the circle  $\{\exp tX; t \in \mathbf{R}\}$ , we read off from (2.13) that

$$(4.2) \quad C_X = \{q; q_j = 0 \text{ if } \omega_j(X) \neq 0\}, \quad \text{locally.}$$

Here  $\omega_j(X) = 2\pi \cdot k_j$ ,  $k_j \in \mathbf{Z}$ , and not all  $k_j$  are even, noting that the half-turn  $\exp \frac{1}{2}X$  of Lemma 3.2 is a nontrivial involution of  $Q_0$ . If  $\text{codim } C_X = 1$ , locally, then only one of the  $\omega_j(X)$  is nonzero, hence odd. The half-turn maps  $q_j$  to  $-q_j$ , so one side of  $C_X$  to the other. There are only finitely many  $C_X$ 's composing  $Q_0 \setminus Q_0^{\text{reg}}$ . Therefore, writing

$$(4.3) \quad T_{1/2} = \{g \in T; \tilde{g}^2(m) = m \text{ for all } m \in Q_0\},$$

we get that  $Q_0^{\text{reg}}/T_{1/2}$  is connected. (This argument is reminiscent of the transitivity of the action of the Weyl group on the set of Weyl chambers.)

In particular, since  $f$  is  $T$ -invariant,  $f(Q_0^{\text{reg}})$  is connected. It is open; and dense in  $f(Q_0) = f(M)$ , because  $Q_0^{\text{reg}}$  is dense in  $Q_0$ ,  $Q_0 \setminus Q_0^{\text{reg}}$  being equal to the union of finitely many closed submanifolds of codimension  $\geq 1$  of  $Q_0$ .

Since there are only finitely many possibilities for the  $f(\bar{Y})$ ,  $Y$  an  $A$ -orbit in  $Q_0$ , one has that for each  $m' \in Q_0$  there is a neighborhood  $U$  of  $m'$  in  $Q_0$  such that  $f(A \cdot m'') \supset f(A \cdot m')$  for all  $m'' \in U$ . Let  $Q'_0$  be the set of  $m' \in Q_0$  such that  $f(A \cdot m'') = f(A \cdot m')$  for all  $m'' \in Q_0$ ,  $m''$  near  $m'$ . Then  $Q'_0$  is open and dense in  $Q_0$ . Each connected component  $R$  of  $Q'_0$  is  $A$ -invariant, and  $f(\bar{R}) = f(A \cdot m')$  for each  $m' \in R$ . We shall show that  $f(\bar{R}) = f(Q_0) = f(M)$ , thus completing the proof of the proposition in view of Theorem 4.1. Indeed, if  $m \in Q_0^{\text{reg}}$  then  $f$  is a diffeomorphism from  $A \cdot m$  to an open subset of  $\mathfrak{t}^*$ , so  $f(m) \notin \partial f(\bar{R})$ . Because  $R$  meets the open dense subset  $Q_0^{\text{reg}}$  of  $Q_0$ , the open set  $f(Q_0^{\text{reg}})$  meets  $f(\bar{R})^{\text{int}}$ . Since  $\partial f(\bar{R}) \in f(Q_0^{\text{reg}}) = \emptyset$ , it follows that  $f(\bar{R})^{\text{int}}$  contains the connected set  $f(Q_0^{\text{reg}})$ . But this implies that  $f(\bar{R}) = f(Q_0)$ .

Choose  $X \in \mathfrak{t}$ . The stable (or unstable) manifolds of the gradient vector field of  $f_X$  define a decomposition of  $M$  and of  $Q$ . These are cell decompositions if  $f_X$  is a Morse function rather than only Bott-Morse. The stable manifolds in  $Q$  are the connected components of the intersections with  $Q$  of the stable manifolds in  $M$ . *In the case of the flag manifolds*, and  $f_X$  a Morse function, the closures of the stable manifolds in  $M$  are complex algebraic varieties, defining cycles (the Schubert cycles) which form a basis for the homology of  $M$ . The closures of the stable manifolds in  $Q$  are real algebraic varieties, defining cycles modulo 2 which form a basis for the homology mod 2 of  $Q$ . Also, two critical points in  $Q$  are connected by a gradient curve in  $M$  only if they are connected by a gradient curve in  $Q$ . See [3, §4]. Finally it is known that the image under  $f$  of both the real and the complex Schubert cycles are convex polytopes, see Heckman [8, Chapter 2, Corollary 2], and Atiyah [2, §4].

It might be interesting to investigate which of these properties generalize to the present setting.

**5. Flag manifolds.** For any connected Lie group  $U$  with Lie algebra  $\mathfrak{u}$ , Kirillov [9]<sup>3</sup> introduced a symplectic form on each orbit of the coadjoint action of  $U$  in  $\mathfrak{u}^*$ , as follows. For  $\xi \in \mathfrak{u}^*$ , the coadjoint orbit  $\mathcal{O}$  of  $\xi$  can be identified with  $U/U_\xi$ , where

$$(5.1) \quad U_\xi = \{g \in U; (\text{Ad } g)^*(\xi) = \xi\},$$

is the stabilizer of  $\xi$  in  $U$ , which has Lie algebra

$$(5.2) \quad \mathfrak{u}_\xi = \{X \in \mathfrak{u}; (\text{ad } X)^*(\xi) = 0\}.$$

The symplectic form on  $T_\xi \mathcal{O} \cong \mathfrak{u}/\mathfrak{u}_\xi$  is defined by

$$(5.3) \quad \sigma_\xi(X, Y) = \xi([X, Y]), \quad X, Y \in \mathfrak{u}/\mathfrak{u}_\xi,$$

and it is then shown that  $\sigma_\xi$  extends to a unique  $U$ -invariant symplectic form  $\sigma$  on  $\mathcal{O}$ . The action of  $U$  on  $\mathcal{O}$  is Hamiltonian (though nonabelian, see Abraham and Marsden [1] for the definitions in this general case), and the momentum mapping is equal to the inclusion mapping:  $\mathcal{O} \rightarrow \mathfrak{u}^*$ . If  $T$  is a torus in  $U$ , then its action on  $\mathcal{O}$  is Hamiltonian with momentum mapping equal to the projection:  $\mathfrak{u}^* \rightarrow \mathfrak{t}^*$ , restricted to  $\mathcal{O}$ .

Up to coverings, these are the only symplectic manifolds with transitive Hamiltonian group actions. If  $U$  is compact, then the theorem of Atiyah [2] and Guillemin and Sternberg [7] gives that the projection of the coadjoint orbit of  $\xi$  in  $\mathfrak{u}^*$  to  $\mathfrak{s}^*$ ,  $\mathfrak{s}$  = the Lie algebra of a maximal torus in  $U$ , is equal to the convex hull of the Weyl group orbit of  $\xi$ . As we shall see below, these coadjoint orbits are the complex flag manifolds and the convexity theorem is Kostant's for the complex case. Note that because the center of  $U$  is contained in  $U_\xi$ , one may assume here that  $U$  has trivial center.

Now we turn to a description of the real flag manifolds, see also [3, §2]. Let  $G$  be a real connected semisimple Lie group with trivial center, and let  $G = KAN$  be its Iwasawa decomposition. We may think of  $G = \text{Ad } G$  as a matrix group, then  $K, A, N$  are the groups of respectively the orthogonal, diagonal with positive eigenvalues, upper triangular unipotent elements of  $G$ . Let  $\mathfrak{g}, \mathfrak{k}, \mathfrak{a}$  be the Lie algebras of  $G, K, A$  respectively. For any  $H \in \mathfrak{a}$ , the  $\text{Ad } K$ -orbit of  $H$  in  $\mathfrak{g}$  (actually contained in a sphere in the orthogonal complement of  $\mathfrak{k}$ ) can be identified with  $K/K_H$ , where

$$(5.4) \quad K_H = \{k \in K; \text{Ad } k(H) = H\}$$

is the centralizer of  $H$  in  $K$ . The functions

$$(5.5) \quad f_{H',H}(k) = \langle H', \text{Ad } k(H) \rangle, \quad k \in K/K_H, \quad H, H' \in \mathfrak{a}$$

(the bilinear form here is the Killing form), can be considered as testing the orthogonal projection of  $\text{Ad } K(H)$  to  $\mathfrak{a}$  by linear forms on  $\mathfrak{a}$ .

This orthogonal projection is actually the infinitesimal version of the Iwasawa projection  $\pi: G \rightarrow \mathfrak{a}$  defined by

$$(5.6) \quad x \in K \cdot \exp \pi(x) \cdot N, \quad x \in G.$$

---

<sup>3</sup> The fact that the form is closed and the relation with general homogeneous symplectic manifolds were observed later by Kostant and Souriau.

This projection will be applied to the  $K$ -orbit

$$(5.7) \quad \{k \cdot \exp H \cdot k^{-1}; k \in K\} \cong K/K_H$$

of  $\exp H$  in  $G$ . The full convexity theorem of Kostant now states that both the Iwasawa projection of (5.7), and its infinitesimal version applied to  $\text{Ad } K(H)$ , have their image equal to the convex hull of the Weyl group orbit of  $H$  in  $\mathfrak{a}$ . We shall only discuss the functions  $f_{H',H}$  in (5.5), noting that Heckman [8] showed that the convexity theorem for the Iwasawa projection can be proved from its infinitesimal version by a homotopy argument.

As we shall show below,  $K/K_H$  is a connected component of  $Q$  and  $f_{H',H} = f_X|_Q$ , where  $Q, f_X$  are as in §§2, 3. The symplectic manifold  $M$  is equal to a complex flag manifold  $U/U_\xi$  as above, for a suitable  $U$ , resp.  $\xi$ , and  $X$  is related to  $H'$  by a linear isomorphism. This then puts the infinitesimal version of Kostant's convexity theorem in the framework of Theorem 2.5. Moreover, in Takeuchi and Kobayashi [13] and [3, §4], it is proved that for generic  $H', f_{H',H}$  is a tight Morse function on  $K/K_H$ . So Theorem 3.1 provides a new proof for this, and extends the tightness to arbitrary  $H' \in \mathfrak{a}$ .

Let  $G_C$  be the complexification of  $G$ , with Iwasawa decomposition

$$(5.8) \quad G_C = UB V$$

(we are clearly running out of letters). Here  $U$  is the maximal compact subgroup of  $G_C$ , which in fact is another real form of  $G_C$ . If  $\tau$  is the complex conjugation of  $G_C$  around  $G$ , then  $G$  is the connected component of 1 of the fixed point set of  $\tau$  in  $G_C$ . Moreover, we can arrange that  $U$  is  $\tau$ -invariant and  $K$  is the connected component of 1 of the fixed point set of  $\tau$  in  $U$ . Similarly  $B$ , resp.  $V$ , are  $\tau$ -invariant and  $A$ , resp.  $N$ , are the fixed point sets of  $\tau$  in  $B$ , resp.  $V$ . Of these groups only  $V$  is complex, in general. In fact, the complexification  $C$  of  $B$  is a Cartan subgroup of  $G_C$ ,  $S = C \cap U$  is a maximal torus in  $u$ , its Lie algebra  $\mathfrak{s}$  is equal to  $i \cdot \mathfrak{b}$  if  $\mathfrak{b}$  denotes the Lie algebra of  $B$ .

For  $H \in \mathfrak{b}$ ,  $U/U_H \xrightarrow{\sim} G_C/U_H B V$ , where  $U_H B V$  turns out to be a complex closed subgroup of  $G_C$ . It contains  $C V$ , which is a maximal solvable subgroup of  $G_C$ , called a Borel subgroup. The Borel subgroups of  $G_C$  are all conjugate to each other. The subgroups  $P_C$  of  $G_C$  containing a Borel subgroup are called the parabolic subgroups. They are also characterized as those for which  $G_C/P_C$  is a complex projective variety. The  $G_C/P_C$  are the complex flag manifolds. Since up to conjugacy each parabolic subgroup of  $G_C$  is of the form  $U_H B V$  for some  $H \in \mathfrak{b}$ , this exhibits the  $U/U_H$  as the general complex flag manifolds.

Now  $U_H = U_\xi$  as in (5.1), if we define  $\xi \in \mathfrak{s}^*$  by

$$(5.9) \quad \xi(Z) = \langle iH, Z \rangle, \quad Z \in \mathfrak{s}.$$

This identifies the coadjoint orbits of compact connected Lie groups with the complex flag manifolds.

If  $H \in \mathfrak{a} \subset \mathfrak{b}$ , then  $U_H B V$  is equal to the complexification of  $P = K_H A N$ , which therefore is called a real parabolic subgroup of  $G$ . As a corollary,  $K/K_H \xrightarrow{\sim} G/P$  has

$U/U_H \xrightarrow{\sim} G_C/P_C$  as its complexification. Conversely,  $K/K_H$  is equal to the connected component of  $1 \cdot K_H$  of the fixed point set of  $\tau$  in  $U/U_H$ . The  $K/K_H$  are called the real flag manifolds. For the classical groups they can be identified with spaces of flags of linear subspaces of a vector space, isotropic with respect to the bilinear form (not necessarily symmetric or nondegenerate) of which  $G$  is taken as the isometry group. In particular all (isotropic) Grassmann manifolds are included in the list of examples.

Since  $\tau$  leaves the elements of  $\mathfrak{a}$  fixed, and maps  $\xi = iH$  to  $\overline{iH} = -iH = -\xi$ , we see from (5.3) that  $\tau$  is antisymplectic. On the other hand, taking for  $T \subset S$  the torus in  $U$  generated by  $\mathfrak{t} = i\mathfrak{a} \subset i\mathfrak{b} = \mathfrak{s}$ , we get that  $T$  acts in a Hamiltonian way on  $U/U_H$ , with Hamilton function  $f_X$  of  $\tilde{X}$ ,  $X \in \mathfrak{t}$ , equal to  $f_{H',H}$ , taking  $H' = -iX$ . In particular  $f_X$  is  $\tau$ -invariant.

#### REFERENCES

1. R. Abraham and J. Marsden, *Foundations of mechanics*, 2nd ed., Benjamin, New York, 1978.
2. M. F. Atiyah, *Convexity and commuting Hamiltonians*, preprint, Oxford Univ. Press, London, 1981.
3. J. J. Duistermaat, J. A. C. Kolk and V. S. Varadarajan, *Functions, flows and oscillatory integrals on flag manifolds and conjugacy classes in real semisimple Lie groups*, preprint, Utrecht, 1981.
4. E. E. Floyd, *On periodic maps and the Euler characteristics of associated spaces*, Trans. Amer. Math. Soc. **72** (1952), 138–147.
5. ———, *Periodic maps via Smith theory*, Seminar on Transformation Groups (A. Borel, ed.), Ann. of Math. Studies, no. 46, Princeton Univ. Press, Princeton, N. J., 1960, pp. 35–47.
6. T. Frankel, *Fixed points on Kähler manifolds*, Ann. of Math. (2) **70** (1959), 1–8.
7. V. Guillemin and S. Sternberg, *Convexity properties of the moment mapping*, preprint, M.I.T., 1981.
8. G. J. Heckman, *Projections of orbits and asymptotic behaviour of multiplicities for compact Lie groups*, Thesis, Leiden, 1980.
9. A. A. Kirillov, *Unitary representations of nilpotent Lie groups*, Uspehi Mat. Nauk **17** (1962), 57–110 = Russian Math. Surveys **17** (1962), 53–104.
10. B. Kostant, *On convexity, the Weyl group and the Iwasawa decomposition*, Ann. Sci. École Norm. Sup. (4) **6** (1973), 413–455.
11. K. R. Meyer, *Hamiltonian systems with discrete symmetry*, J. Differential Equations **41** (1981), 228–238.
12. G. D. Mostow, *On a conjecture of Montgomery*, Ann. of Math. (2) **65** (1957), 513–516.
13. M. Takeuchi and S. Kobayashi, *Minimal embeddings of R-spaces*, J. Differential Geom. **2** (1968), 203–215.
14. A. Weinstein, *Symplectic manifolds and their Lagrangian submanifolds*, Adv. in Math. **6** (1971), 329–346.
15. C.-T. Yang, *On a problem of Montgomery*, Proc. Amer. Math. Soc. **8** (1957), 255–275.
16. M. Takeuchi, *Cell decomposition and Morse inequalities on certain symmetric spaces*, J. Fac. Sci. Univ. Tokyo **12** (1965), 81–192.

MATHEMATISCH INSTITUUT, RIJKSUNIVERSITEIT UTRECHT, BUDAPESTLAAN 6, DE UITHOF, 3508 TA UTRECHT, THE NETHERLANDS