

EXTENSIONS OF A THEOREM OF WINTNER ON SYSTEMS WITH ASYMPTOTICALLY CONSTANT SOLUTIONS

BY

WILLIAM F. TRENCH

ABSTRACT. A theorem of Wintner concerning sufficient conditions for a system $y' = A(t)y$ to have linear asymptotic equilibrium is extended to a system $x' = A(t)x + f(t, x)$. The integrability conditions imposed on f permit conditional convergence of some of the improper integrals that occur. The results improve on Wintner's even if $f = 0$.

An $n \times n$ system

$$(1) \quad y' = A(t)y, \quad t > 0,$$

is said to have linear asymptotic equilibrium if for each constant vector c there is a solution of (1) such that $\lim_{t \rightarrow \infty} y(t) = c$. It is well known that (1) has this property if A is continuous and

$$(2) \quad \int_0^{\infty} \|A(t)\| dt < \infty.$$

Wintner [5] attributed this result to Bôcher and improved on it as follows.

THEOREM 1 (WINTNER). *Let A be continuous on $[a, \infty)$ and suppose the integrals*

$$(3) \quad A_j(t) = \int_t^{\infty} A_{j-1}(s)A(s) ds, \quad 1 \leq j \leq k \quad (A_0 = I),$$

converge, and

$$(4) \quad \int_0^{\infty} \|A_k(t)A(t)\| dt < \infty.$$

Then (1) has linear asymptotic equilibrium.

Notice that (3) is vacuous and (4) reduces to (2) when $k = 0$.

Here we apply Wintner's idea to the system

$$(5) \quad x' = A(t)x + f(t, x).$$

We give sufficient conditions for (5) to have a solution x such that

$$(6) \quad \lim_{t \rightarrow \infty} x(t) = c$$

Received by the editors February 29, 1984.

1980 *Mathematics Subject Classification.* Primary 34C11, 34D10.

©1986 American Mathematical Society
0002-9947/86 \$1.00 + \$.25 per page

for a given constant vector c . The assumptions on f in our main theorem apply only “near” the set $\{(t, c) | t \geq a\}$. Our integral smallness conditions permit conditional convergence of some of the improper integrals that occur. This continues a theme developed previously in [3 and 4]. (See also Hallam [1, 2].) The idea of dealing with the iterated integrals (3) is due to Wintner [5].

Throughout the paper k is a nonnegative integer. It is to be understood that conditions stated for $1 \leq j \leq k$ are vacuous if $k = 0$, and that $\sum_{j=1}^0 = 0$. Our results apply with x , A , and f real- or complex-valued. Improper integrals occurring in hypotheses are tacitly assumed to converge, and the convergence may be conditional except when the integrand is obviously nonnegative. We use “ O ” and “ o ” in the standard way to indicate orders of magnitude as $t \rightarrow \infty$.

To indicate the direction of proof of our main theorem, it is convenient to state part of its hypotheses here.

ASSUMPTION A. Let w be continuous and nonincreasing on $[a, \infty)$, $0 < w(t) \leq 1$, and suppose that either $\lim_{t \rightarrow \infty} w(t) = 0$ or $w = 1$. Suppose A is continuous on $[a, \infty)$ and A_j in (3) exists for $1 \leq j \leq k$. Let c be a given constant vector, and suppose there is a constant $M > 0$ such that f is continuous and

$$(7) \quad \|f(t, x) - f(t, c)\| \leq R(t, \|x - c\|)$$

on

$$(8) \quad S = \{(t, x) | t \geq a, \|x - c\| \leq Mw(t)\},$$

where $R(t, \lambda)$ is continuous on $\{(t, \lambda) | t \geq a, 0 \leq \lambda \leq Mw(t)\}$ and nondecreasing in λ for each t .

We define $\Gamma_k = \sum_{j=0}^k A_j$, and observe from (3) that

$$(9) \quad \Gamma'_k = -\Gamma_{k-1}A, \quad k \geq 0 \ (\Gamma_{-1} = 0).$$

Moreover, since $\lim_{t \rightarrow \infty} \Gamma_k(t) = I$, Γ_k is invertible for large t , and $\lim_{t \rightarrow \infty} \Gamma_k^{-1}(t) = I$.

Now let $t_0 \geq a$ be such that Γ_k^{-1} exists on $[t_0, \infty)$. For convenience below, we define

$$(10) \quad \mu_k(t) = \sup_{s \geq t} \{\|\Gamma_k^{-1}(s)\|\} = 1 + o(1),$$

and

$$(11) \quad \nu_k(t) = \mu_k(t) \sup_{s \geq t} \{\|\Gamma_k(s)\|\} = 1 + o(1).$$

Let $C[t_0, \infty)$ be the space of continuous n -vector functions (with real or complex components) on $[t_0, \infty)$, with the topology of uniform convergence on finite intervals. Let $V[t_0, \infty)$ be the closed convex subset of $C[t_0, \infty)$ defined by

$$(12) \quad V[t_0, \infty) = \{x \in C[t_0, \infty) | \|x(t) - c\| \leq Mw(t), t \geq t_0\}.$$

We obtain our results by applying the Schauder-Tychonov theorem to an appropriate transformation T of $V[t_0, \infty)$ (for sufficiently large t_0) into itself. To motivate the choice of T , we observe that if

$$x(t) = c - \int_t^\infty [A(s)x(s) + f(s, x(s))] ds,$$

where the integral is assumed to converge, then x satisfies (5) and (6). Repeated integration by parts, assuming at each step that x satisfies (5), yields the equation

$$(13) \quad \Gamma_k(t)x(t) = c - \int_t^\infty A_k(s)A(s)x(s) ds - \int_t^\infty \Gamma_k(s)f(s, x(s)) ds.$$

Although these manipulations are completely formal, (13) suggests the transformation T defined by

$$(14) \quad (Tx)(t) = \Gamma_k^{-1}(t) \left[c - \int_t^\infty [A_k(s)A(s)x(s) + \Gamma_k(s)f(s, x(s))] ds \right].$$

Assumption A implies that the function $F(t) = f(t, x(t))$ is continuous on $[t_0, \infty)$ if $x \in V[t_0, \infty)$. Hence, if the integrals in (14) converge, differentiation yields

$$(15) \quad (Tx)'(t) = \Gamma_k^{-1}(t) [\Gamma_{k-1}(t)A(t)(Tx)(t) + A_k(t)A(t)x(t)] + f(t, x(t)),$$

where we have used (9) and the fact that $(\Gamma_k^{-1})' = -\Gamma_k^{-1}\Gamma_k'\Gamma_k^{-1}$. Therefore, if T has a fixed point (function) x_0 in $V[t_0, \infty)$, we see on setting $Tx = x = x_0$ in (15) that

$$\begin{aligned} x_0'(t) &= \Gamma_k^{-1}(t) [\Gamma_{k-1}(t) + A_k(t)] A(t)x_0(t) + f(t, x(t)) \\ &= A(t)x_0(t) + f(t, x(t)) \end{aligned}$$

(since $\Gamma_{k-1} + A_k = \Gamma_k$); i.e., x_0 satisfies (5). Moreover, setting $Tx = x = x_0$ in (14) shows that x_0 also satisfies (6).

The following theorem allows the integrals occurring in Tc (the function obtained by setting $x = c$ in (14)) to converge conditionally, and exploits the rapidity with which $Tc - c$ approaches zero for large t to restrict the set $V[t_0, \infty)$ on which T must satisfy the hypotheses of the Schauder-Tychonov theorem. (See Remark 1, below.)

THEOREM 2. *Suppose Assumption A holds. Let*

$$(16) \quad h(t) = (\Gamma_k^{-1}(t) - I)c - \Gamma_k^{-1}(t) \int_t^\infty [A_k(s)A(s)c + \Gamma_k(s)f(s, c)] ds,$$

and suppose that

$$(17) \quad \overline{\lim}_{t \rightarrow \infty} \frac{\|h(t)\|}{w(t)} = \alpha.$$

Suppose also that

$$(18) \quad \overline{\lim}_{t \rightarrow \infty} \int_t^\infty \left[\frac{R(s, Mw(s))}{M} + \|A_k(s)A(s)\|w(s) \right] ds = \theta < 1$$

and

$$(19) \quad \alpha < M(1 - \theta).$$

Then, if t_0 is sufficiently large, there is a solution x_0 of (5) on $[t_0, \infty)$ such that

$$(20) \quad \|x_0(t) - c\| \leq Mw(t), \quad t \geq t_0,$$

and

$$(21) \quad \overline{\lim}_{t \rightarrow \infty} (w(t))^{-1} \|x_0(t) - c\| \leq \alpha + M\theta;$$

or, more precisely,

$$(22) \quad \overline{\lim}_{t \rightarrow \infty} (w(t))^{-1} \|x_0(t) - c - h(t)\| \leq M\theta.$$

PROOF. We can rewrite (14) as

$$(23) \quad (Tx)(t) = c + h(t) - \Gamma_k^{-1}(t) \int_t^\infty [A_k(s)A(s)(x(s) - c) + \Gamma_k(s)(f(s, x(s)) - f(s, c))] ds,$$

where the integral converges if $x \in V[t_0, \infty)$, because of (7), (12), and (18); moreover

$$(24) \quad \|(Tx)(t) - c\| \leq \|h(t)\| + M\mu_k(t) \int_t^\infty \|A_k(s)A(s)\|w(s) ds + \nu_k(t) \int_t^\infty R(s, Mw(s)) ds.$$

From (10), (11), (18), and (19), we can assume henceforth that t_0 is so large that the right side of (24) is $\leq Mw(t)$ if $t \geq t_0$. Then

$$(25) \quad T(V[t_0, \infty)) \subset V[t_0, \infty).$$

Now we show that T is continuous on $V[t_0, \infty)$. Let $\{x_j\}$ be a sequence in $V[t_0, \infty)$ which converges to a limit x in $V[t_0, \infty)$. From (10), (11), and (23).

$$\begin{aligned} \|(Tx_j)(t) - (Tx)(t)\| &\leq \mu_k(t_0) \int_{t_0}^\infty \|A_k(s)A(s)\| \|x_j(s) - x(s)\| ds \\ &\quad + \nu_k(t_0) \int_{t_0}^\infty \|f(s, x_j(s)) - f(s, x(s))\| ds, \quad t \geq t_0. \end{aligned}$$

The integrands here converge to zero on $[t_0, \infty)$, and they are dominated by $2M\|A_k(s)A(s)\|w(s)$ and $2R(s, Mw(s))$, respectively. (See (7) and (12).) Therefore, (18) and Lebesgue's dominated convergence theorem imply that the integrals approach zero as $j \rightarrow \infty$. Hence, $\{Tx_j\}$ converges to Tx uniformly on $[t_0, \infty)$, and therefore T is continuous on $V[t_0, \infty)$.

From (12) and (25), $T(V[t_0, \infty))$ is equibounded on finite intervals. This, (15), and Assumption A imply that $T(V[t_0, \infty))$ is also equicontinuous on finite intervals. Now we have verified the hypotheses of the Schauder-Tychonov theorem, which implies that $Tx_0 = x_0$ for some x_0 in $V[t_0, \infty)$. Setting $Tx = x_0$ in (24) and invoking (10), (11), (17), and (18) yields (20). Similarly, setting $x = Tx = x_0$ in (23) yields (22). This completes the proof.

COROLLARY 1. *In addition to the assumptions of Theorem 2, suppose that*

$$(26) \quad R(t, \lambda_1)/R(t, \lambda_2) \leq \lambda_1/\lambda_2, \quad 0 \leq \lambda_1 < \lambda_2.$$

Then (21) can be replaced by

$$(27) \quad \overline{\lim}_{t \rightarrow \infty} (w(t))^{-1} \|x_0(t) - c\| \leq \alpha/(1 - \theta).$$

PROOF. Let

$$\phi(t) = \sup_{s \geq t} \{ (w(s))^{-1} \|x_0(s) - c\| \}$$

and

$$(28) \quad \delta = \lim_{t \rightarrow \infty} \phi(t) = \overline{\lim}_{t \rightarrow \infty} (w(t))^{-1} \|x_0(t) - c\|.$$

Setting $x = Tx = x_0$ in (23) and using routine estimates yields

$$(29) \quad \|x_0(t) - c\| \leq \|h(t)\| + \nu_k(t) \int_t^\infty \|A_k(s)A(s)\| \|x_0(s) - c\| ds \\ + \mu_k(t) \int_t^\infty R(s, \|x_0(s) - c\|) ds.$$

Applying (26) in the second integral and then dividing (29) by $w(t)$ yields the inequality

$$(30) \quad (w(t))^{-1} \|x_0(t) - c\| \leq (w(t))^{-1} \|h(t)\| + P(t)\phi(t),$$

where

$$P(t) = (w(t))^{-1} \left[\nu_k(t) \int_t^\infty \|A_k(s)A(s)\| w(s) ds + \frac{\mu_k(t)}{M} \int_t^\infty R(s, Mw(s)) ds \right],$$

so that

$$(31) \quad \lim_{t \rightarrow \infty} P(t) = \theta,$$

from (10), (11), and (18). Letting $t \rightarrow \infty$ in (30) and invoking (17), (28), and (31) shows that $\delta \leq \alpha + \theta\delta$, which proves (27).

It is worthwhile to state Theorem 2 separately for the case where $w = 1$, so that α and θ are necessarily zero and (19) is automatic.

THEOREM 3. *Suppose Assumption A holds with $w = 1$. Suppose also that the integral in (16) converges and that*

$$(32) \quad \int^\infty R(t, M) dt < \infty, \quad \int^\infty \|A_k(t)A(t)\| dt < \infty.$$

Then, if t_0 is sufficiently large, there is a solution x_0 of (5) on $[t_0, \infty)$ such that $\|x_0(t) - c\| \leq M$ for $t \geq t_0$ and $\lim_{t \rightarrow \infty} x_0(t) = c$.

REMARK 1. The continuity assumption on f is the most stringent when $w = 1$, since the set S in (8) is maximized in that case. More importantly, if R satisfies (26), then (32) implies (18) with $\theta = 0$ for every admissible $w \neq 1$, while the converse is obviously false. Nevertheless, the conclusions of Theorem 2 are *weakest* when

$w = 1$. (See Theorem 3.) The reason for this is that in Theorem 3 it is assumed only that the integral in (16) converges, while Theorem 2 exploits the rapidity of its convergence.

The hypotheses of Theorem 2 may hold for some constant vectors c and fail to hold for others. In the following theorem c may be chosen arbitrarily.

THEOREM 4. *Suppose f is continuous for $t \geq a$ and all x , and*

$$\|f(t, x_1) - f(t, x_2)\| \leq R(t, \|x_1 - x_2\|),$$

where $R(t, \lambda)$ is continuous on $[a, \infty) \times [0, \infty)$ and nondecreasing in λ . For some integer $k \geq 0$, suppose the integrals A_1, \dots, A_{k+1} converge, and

$$(33) \quad \|A_j(t)\| = O(w(t)), \quad 1 \leq j \leq k + 1.$$

Suppose also that

$$(34) \quad \left\| \int_t^\infty \Gamma_k(s) f(s, c) ds \right\| = O(w(t))$$

for every constant vector c , and that (18) holds for all $M > 0$. Then, if c is a given constant vector, there is a solution x_0 of (5) which is defined for t sufficiently large and satisfies

$$(35) \quad \|x_0(t) - c\| = O(w(t)).$$

Moreover, if $\theta = 0$ in (18) and (33) and (34) hold with “ O ” replaced by “ o ” (which is necessarily true if $w = 1$), then (35) holds with “ O ” replaced by “ o .”

PROOF. The hypotheses imply (17) for some α (which may depend upon c , but is zero if (33) and (34) hold with “ o ”). Simply choose M to satisfy (19) and invoke Theorem 2.

Theorem 4 has the following corollary for the linear system

$$(36) \quad x' = [A(t) + B(t)]x + g(t).$$

COROLLARY 2. *Let A, B , and g be continuous on $[a, \infty)$. Suppose (33) holds.*

$$(37) \quad \left\| \int_t^\infty \Gamma_k(s) g(s) ds \right\| = O(w(t)), \quad \left\| \int_t^\infty \Gamma_k(s) B(s) ds \right\| = O(w(t)).$$

and

$$(38) \quad \overline{\lim}_{t \rightarrow \infty} (w(t))^{-1} \int_t^\infty [\|A_k(s)A(s)\| + \|B(s)\|] w(s) ds = \theta < 1.$$

Then, for any constant c , (36) has a solution which satisfies (35); moreover, if $\theta = 0$ and (33) and (37) hold with “ O ” replaced by “ o ,” then so does (35).

The following special case of Corollary 2 extends Theorem 1.

COROLLARY 3. *Suppose (33) holds and*

$$\overline{\lim}_{t \rightarrow \infty} (w(t))^{-1} \int_t^\infty \|A_k(s)A(s)\| w(s) ds < 1$$

for some w as in Assumption A. Then (1) has linear asymptotic equilibrium.

BIBLIOGRAPHY

1. T. G. Hallam, *Asymptotic integration of second order differential equations with integrable coefficients*, SIAM J. Appl. Math. **19** (1970), 430-439.
2. _____, *Asymptotic integration of a nonhomogeneous differential equation with integrable coefficients*, Czechoslovak Math. J. **21** (1971), 661-671.
3. W. F. Trench, *Systems of differential equations subject to mild integral conditions*, Proc. Amer. Math. Soc. **87** (1983), 263-270.
4. _____, *Asymptotics of differential systems with deviating arguments*, Proc. Amer. Math. Soc. **92** (1984), 219-224.
5. A. Wintner, *On a theorem of Bôcher in the theory of ordinary linear differential equations*, Amer. J. Math. **76** (1954), 183-190.

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, DREXEL UNIVERSITY, PHILADELPHIA, PENNSYLVANIA 19104