

SIMPLE HOMOTOPY TYPE OF FINITE 2-COMPLEXES WITH FINITE ABELIAN FUNDAMENTAL GROUP

BY

M. PAUL LATIOLAIS

ABSTRACT.

THEOREM 1. *Let K be a finite 2-dimensional CW-complex with $\pi_1(K)$ finite and abelian. Then every element of the Whitehead group of K is realizable as the torsion of a self-homotopy equivalence on K .*

THEOREM 2. *Homotopy equivalence and simple homotopy equivalence are the same for finite 2-dimensional CW-complexes with finite abelian fundamental groups.*

0. Introduction. It is known that there exist finite n -complexes for all $n > 2$ which are homotopy equivalent but not simply homotopy equivalent. In this paper, we show that in dimension 2 homotopy type and simple homotopy type are the same when the fundamental group of the finite 2-dimensional complexes is finite and abelian.

The technique used is to show that all the elements of the Whitehead group of a complex K are realizable as torsions of self-equivalences on K .

1. An important example. Dyer and Sieradski [DS] showed in 1973 that two 2-dimensional CW-complexes whose fundamental group was Z_n were homotopy equivalent if and only if they were simple homotopy equivalent. The next case to consider would be $Z_n \times Z_m$. We want to show that any 2-dimensional complex with fundamental group $Z_n \times Z_m$ realizes all of its Whitehead group as torsions of self-equivalences. ($\xi(K)$ denotes the group of self-homotopy equivalences of K .)

THEOREM 1.1. *Let K be the standard 2-complex of the presentation $P = \{a, b | a^n, b^m, [a, b]\}$. Then every element of $\text{Wh}(K)$ is realizable as $\tau(f)$ for some $f \in \xi(K)$.*

To prove the above theorem, we need the following three lemmas.

LEMMA 1.2. *Suppose G is a group with generating set $\{a_i | i \in I\}$ for some index set I . Let $A: Z(G) \rightarrow Z$ be the augmentation map (i.e., the ring homomorphism taking g to 1 for each $g \in G$). Then each element $\theta \in \ker A$ is of the form*

$$\theta = \sum_i \phi_i(a_i - 1) \quad \text{for some elements } \phi_i \in Z(G).$$

PROOF. See [F, p. 549].

Received by the editors June 1, 1984 and, in revised form, February 18, 1985. Presented to the Society in Louisville, Kentucky, on January 28, 1984.

1980 *Mathematics Subject Classification.* Primary 57M20, 57Q10.

©1986 American Mathematical Society
0002-9947/86 \$1.00 + \$.25 per page

LEMMA 1.3. Let K be the standard complex of the presentation $\{a, b \mid a^n, b^m, aba^{-1}b^{-1}\}$ and let \tilde{K} be the universal cover of K with a chosen lift \tilde{e}_0 of the single 0-cell in K . Consider $C_i(\tilde{K})$ as a $Z\pi_1$ -module generated by the preferred lifts of reach i -cell. Then $H_2(\tilde{K})$ is generated by the set

$$(a-1)\tilde{R}_a, \quad (b-1)\tilde{R}_b, \\ (b-1)\tilde{R}_a + \left(\sum_{i=0}^{n-1} a^i\right)\tilde{R}_{[a,b]}, \quad (a-1)\tilde{R}_b - \left(\sum_{i=0}^{m-1} b^i\right)\tilde{R}_{[a,b]}.$$

PROOF. Simple calculations will show that the given chains are cycles and thus represent elements of $H_2(\tilde{K}) = Z_2(\tilde{K})$. To show that they generate all of $H_2(\tilde{K})$ (a fact not actually needed in this paper), use a technique similar to Metzler's in [M2, p. 330]. \square

LEMMA 1.4. Given a finite 2-dimensional complex K and $Z\pi_1$ -module map $\phi: C_2(\tilde{K}) \rightarrow C_2(\tilde{K})$ which commutes with the boundary operator in the sense that $\partial_2\phi = \partial_2$, then there exists a homotopy equivalence $f: K \rightarrow K$ such that $\tilde{f}_2 = \phi$ and $\tilde{f}_1 = \text{identity}$ only if the $Z\pi_1$ -module representation of ϕ is invertible.

PROOF. This is done by modifying the identity map on K using the Puppe action. This technique is patterned after [DS, p. 41]. For the original reference, see [P].

Suppose $f: K \rightarrow K$ is a homotopy equivalence which induces the identity map on $\pi_1(K)$. Let \tilde{K} be the universal cover of K . Since $f_*: \pi_1 \rightarrow \pi_1$ is the identity, then f is homotopic to a map which induces the identity on $C_1(\tilde{K})$. Since we are only interested in homotopy classes of maps, we may assume f is that map. Now f induces the following map on the chain complex of the universe cover \tilde{K} :

$$(1) \quad \begin{array}{ccc} 0 & & 0 \\ \downarrow & & \downarrow \\ H_2(\tilde{K}) & \xrightarrow{f_2^*} & H_2(\tilde{K}) \\ \partial \downarrow & & \partial \downarrow \\ C_2(\tilde{K}) & \xrightarrow{\tilde{f}_2} & C_2(\tilde{K}) \\ \partial \downarrow & & \partial \downarrow \\ C_1(\tilde{K}) & \xrightarrow{\tilde{f}_1 = \text{id}} & C_1(\tilde{K}) \\ \partial \downarrow & & \partial \downarrow \\ C_0(\tilde{K}) & \xrightarrow{\tilde{f}_0 = \text{id}} & C_0(\tilde{K}) \\ \downarrow & & \downarrow \\ 0 & & 0 \end{array}$$

Consider the map $\text{id}: K \rightarrow K$. The induced maps on the chain complex of \tilde{K} will all be the identity. Replace \tilde{f}_2 by ϕ in diagram (1). Now we have two chain maps that agree except on $C_2(\tilde{K})$. That means that $\phi - \text{id}_2$ will commute in the following

diagram:

$$(2) \quad \begin{array}{ccc} C_2(\tilde{K}) & \xrightarrow{\phi \cdot \text{id}} & C_2(\tilde{K}) \\ \partial \downarrow & & \partial \downarrow \\ C_1(\tilde{K}) & \xrightarrow{0} & C_1(\tilde{K}) \end{array}$$

Consequently, the image of $\phi \cdot \text{id}$ will be in $H_2(\tilde{K}) \equiv \pi_2(K)$. Let R be a 2-cell in K (from the CW-decomposition). Then $[\phi \cdot \text{id}](\tilde{R}) \in H_2(\tilde{K}) \equiv \pi_2(K)$. Let p be the representative of the image of \tilde{R} in $\pi_2(K)$. Also let p represent the actual map $p: S^2 \rightarrow K$. Now define g by the composition of the following maps:

$$K \xrightarrow{h} K \vee S^2 \xrightarrow{\text{id} \vee p} K,$$

where h is the identity on $K \setminus R$ and maps R onto $R \vee S^2$ by mapping some $(S^1, e) \subset (R, e_0)$ to e_0 and the interior disk to S^2 . The rest of R gets "stretched" to cover R . Now $\tilde{g}_2(\tilde{R}) = \phi(\tilde{R})$.

We can modify g using the above technique on the other 2-cells in the decomposition of K to obtain our hypothesized f . Using the Five Lemma on diagram (1), f will be a homotopy equivalence if and only if ϕ is an isomorphism.

That $\tau(f) = [M]$ comes from direct computation, using Cohen's §15 and (22.8) in [C]. \square

PROOF OF THEOREM 1.1. Given an element $\phi \in \text{Wh}(K)$, we want to construct a homotopy equivalence $f: K \rightarrow K$, so that $\tau(f) = \phi$. By Lemma 1.4 we merely need to produce an invertible $Z\pi_1$ -matrix M such that M commutes with the boundary operator and $[M] = \phi \in \text{Wh}(K)$. Since in this case K has three 2-cells, we need $M \in \text{GL}_3(Z\pi_1)$.

Let $\bar{a} = a - 1$ and $\bar{b} = b - 1$. Then by Lemma 1.3, any matrix of the following form (where the *rows* represent the images of $\tilde{R}_a, \tilde{R}_b, \tilde{R}_{[a,b]}$) will commute with the boundary operator:

$$\begin{bmatrix} 1 + \phi_{11}\bar{a} + \psi_{11}\bar{b}, & \phi_{12}\bar{a} + \psi_{12}\bar{b}, & \psi_{11}\sum a^i - \phi_{12}\sum b^i \\ \phi_{21}\bar{a} + \psi_{21}\bar{b}, & 1 + \phi_{22}\bar{a} + \psi_{22}\bar{b}, & \psi_{21}\sum a^i - \phi_{12}\sum b^i \\ \phi_{31}\bar{a} + \psi_{31}\bar{b}, & \phi_{32}\bar{a} + \psi_{32}\bar{b}, & 1 + \psi_{31}\sum a^i - \phi_{32}\sum b^i \end{bmatrix}$$

where ϕ_{ij}, ψ_{ij} are appropriate elements of $Z[\pi_1(K)] = Z[Z_n \times Z_m]$.

Now any element of $\text{Wh}(Z(Z_n \times Z_m))$ may be represented by a 2×2 -matrix, see Bass [B, p. 183] and Lam [L, p. 143]. Let $\begin{pmatrix} r & s \\ t & u \end{pmatrix}$ be that matrix, where $r, s, t, u \in Z(Z_n \times Z_m)$. Consider the augmentation map $Z(Z_n \times Z_m) \rightarrow Z$ which maps the elements of $Z_n \times Z_m$ to 1. Call the images of r, s, t and u respectively r', s', t' and $u' \in Z$. Then we may transform the matrix $\begin{pmatrix} r' & s' \\ t' & u' \end{pmatrix}$ to $\begin{pmatrix} \pm 1 & 0 \\ 0 & 1 \end{pmatrix}$ via row and column operators.

We can choose our representative $\begin{pmatrix} r' & s' \\ t' & u' \end{pmatrix}$ without changing the element in $\text{Wh}(Z(Z_n \times Z_m))$ in order that $\begin{pmatrix} r' & s' \\ t' & u' \end{pmatrix}$ transforms to $\begin{pmatrix} +1 & 0 \\ 0 & 1 \end{pmatrix}$.

Now use the same row and column operations on $\begin{pmatrix} t' & s' \\ u' & u' \end{pmatrix}$ that we did on $\begin{pmatrix} t' & s' \\ u' & u' \end{pmatrix}$. We get a matrix which maps to $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ when $a, b \mapsto 1$. Using Lemma 1.2, this new matrix must be of the form

$$\begin{bmatrix} 1 + \alpha_1 \bar{a} + \beta_1 \bar{b}, & \phi_1 \bar{a} + \psi_1 \bar{b} \\ \psi_2 \bar{a} + \beta_2 \bar{b}, & 1 + \phi_2 \bar{a} + \psi_2 \bar{b} \end{bmatrix} \quad \text{where } \alpha_i, \beta_i, \phi_i, \psi_i \in Z(Z_n \times Z_m)$$

Therefore, we may represent any element of $\text{Wh}(Z(Z_n \times Z_m))$ in the form (2). If we choose the constants correctly for our matrix (1), and manipulate that matrix to get the form (2), then we will be done. First, letting $\phi_{31} = \psi_{31} = \phi_{32} = \psi_{32} = 0$, we get

$$\begin{bmatrix} 1 + \phi_{11} \bar{a} + \psi_{11} \bar{b}, & \phi_{12} \bar{b} + \psi_{12}(-\bar{a}), & \psi_{11} \sum a + \phi_{12} \sum b^i \\ \phi_{21} \bar{a} + \psi_{21} \bar{b}, & 1 + \phi_{22} \bar{b} + \psi_{22}(-\bar{a}), & \psi_{21} \sum a + \phi_{22} \sum b^i \\ 0 & 0 & 1 \end{bmatrix}.$$

We then use row 3 to clear column 3 to the equivalent matrix

$$\begin{bmatrix} 1 + \phi_{11} \bar{a} + \psi_{11} \bar{b}, & \phi_{12} \bar{b} + \psi_{12}(-\bar{a}) \\ \phi_{21} \bar{a} + \psi_{21} \bar{b}, & 1 + \phi_{22} \bar{b} + \psi_{22}(-\bar{a}) \end{bmatrix}$$

which, with appropriate choice of constants, is equal to the matrix (2). Consequently, we may produce a matrix M which represents any given element of $\text{Wh}(K)$, by Lemma 1.4, such that M represents \tilde{f}_2 for some map $f: K \rightarrow K$, with $\tilde{f}_1 = \text{identity}$. M is invertible since it represents an element of $\text{Wh}(K)$. Therefore, f is a homotopy equivalence. Since \tilde{f}_1 is the identity map, then $\tau(f) = M$, and therefore all the elements of $\text{Wh}(K)$ are representable. \square

COROLLARY 1.5. *Let L be a finite 2-dimensional complex with $\pi_1(L) = Z_n \times Z_m$ and $\chi(L) = k$. Then L is simple homotopy equivalent to $K \vee (k - 2)(S^2)$, where K is the complex of Theorem 1.1.*

PROOF. Since the complex K of Theorem 1.1 realizes all of its Whitehead group as the torsions of self-equivalence on K , then by Cohen [C, Theorem 24.4] any complex homotopy equivalent to K is simple homotopy equivalent to K . By Dyer [D3], homotopy type of any finite 2-complex with fundamental group $Z_n \times Z_m$ is determined by the Euler characteristic. Since our example K has minimal Euler characteristic (see Swan [Sw, Proposition 2.1]), the other Euler characteristics may be obtained by wedging K with the appropriate number of copies of S^2 . If K realizes all of its torsion by self-equivalences then so does $K \vee S^2$. For if $f: K \rightarrow K$ is a homotopy equivalence, let $f \vee \text{id}: K \vee S^2 \rightarrow K \vee S^2$. Then $\tau(f) = \tau(f \vee \text{id})$.

Consequently, given a 2-complex L with $\pi_1(L) = Z_n \times Z_m$ and $\chi(L) = k$, then by Dyer [D3] L is homotopy equivalent to $K \vee (k - 2)S^2$. But since $K \vee (k - 2)S^2$ realizes all of its torsion by self-equivalence we know by Cohen [C] that L is simple homotopy equivalent to $K \vee (k - 2)S^2$. \square

COROLLARY 1.6. *Let K, L be finite 2-dimensional CW-complexes with fundamental groups $Z_n \times Z_m$. Then the following are equivalent.*

- (a) $\chi(K) = \chi(L)$,
- (b) K is homotopy equivalent to L .
- (c) K is simple homotopy equivalent to L .

2. The general abelian case. We want to prove that simple homotopy type and homotopy type agree for finite 2-complexes with finite abelian fundamental group. This general situation is unlike the $Z_n \times Z_m$ case in that homotopy type does not depend only on Euler characteristic. When the Euler characteristic is minimal, homotopy type depends also on bias (see Metzler [M1], Sieradski [S] and Browning [Br]).

According to Browning [Br, Theorem 1.7] and Sieradski [S, Theorem 2], any finite 2-complex of minimal Euler characteristic with finite abelian fundamental group is homotopy equivalent to the standard complex of some twisted (or “untwisted”) presentation of the form

$$\{a_i | a_i^{n_i}, [a_1^r, a_2], [a_i, a_j], i < j, j \neq 2\},$$

where $r < n_1$ and $(r, n_1) = 1$.

So if we are to use our previous techniques for showing that homotopy type and simple type agree, we not only have to show that the standard complex of the presentation

$$\{a_i | a_i^{n_i}; [a_i, a_j]; i < j; j = 1, \dots, N\}$$

realizes all of its Whitehead torsion by self-equivalences, we also have to show that the standard complex of any twisted presentation realizes all of its torsion by self-equivalences.

LEMMA 2.1. *Let G be a finite abelian group, and let $P = \{a_i | a_i^{n_i}, [a_1^r, a_2], [a_i, a_j]; i < j; j \neq 2; i, j = 1, \dots, N\}$, where $r < n_1$ and $(r, n_1) = 1$, be a twisted presentation of G . Let $\tilde{K}(P) = K$ be the standard complex of P . Then all of the torsion of $\text{Wh}(K)$ is realizable as $\tau(f)$ for some $f \in \xi(K)$.*

PROOF. The proof proceeds as that of Theorem 1.1. Let \tilde{K} be the universal cover of K . Then the following are elements of $H_2(\tilde{K})$:

$$\begin{aligned} (a_i - 1)\tilde{R}_i, & \quad (a_1^r - 1)\tilde{R}_2 - \left(\sum a_2^k\right)\tilde{R}_{12}, \\ (a_j - 1)\tilde{R}_i + \left(\sum a_i^k\right)\tilde{R}_{ij}, & \quad ij \neq 12, i < j, \\ (a_i - 1)\tilde{R}_j - \left(\sum a_j^k\right)\tilde{R}_{ij}, & \quad ij \neq 12, i < j. \end{aligned}$$

The above restrictions on ij tell us that if we try to use these elements of $H_2(\tilde{K})$ to modify the first two rows of the identity matrix, the first column will have no $(a_1 - 1) = \bar{a}_1$ component. So what we will do is ignore the first row and attempt to get our result using the second and third rows.

Consider the map $\phi: C_2(\tilde{K}) \rightarrow C_2(\tilde{K})$ which is the identity except:

$$\begin{aligned}
 \tilde{R}_2 &\mapsto 1 \cdot \tilde{R}_2 + C_{11}[(a_1^r - 1)\tilde{R}_2 - (\sum a_2^k)\tilde{R}_{12}] + C_{12}(a_2 - 1)\tilde{R}_2 \\
 &\quad + \sum_{h=3}^N C_{1h}[(a_h - 1)\tilde{R}_2 - (\sum a_2^k)\tilde{R}_{2h}] \\
 &\quad + C_{21}[(a_1 - 1)\tilde{R}_3 - (\sum a_3^k)\tilde{R}_{13}] \\
 &\quad + C_{22}[(a_2 - 1)\tilde{R}_3 - (\sum a_3^k)\tilde{R}_{23}] + C_{23}(a_3 - 1)\tilde{R}_3 \\
 &\quad + \sum_{h=4}^N C_{2h}[(a_h - 1)\tilde{R}_3 + (\sum a_3^k)\tilde{R}_{3h}], \\
 \tilde{R}_3 &\mapsto 1 \cdot \tilde{R}_3 + C_{31}[(a_1^r - 1)\tilde{R}_2 - (\sum a_2^k)\tilde{R}_{12}] + C_{32}(a_2 - 1)\tilde{R}_2 \\
 &\quad + \sum_{h=3}^N C_{3h}[(a_h - 1)\tilde{R}_2 - (\sum a_2^k)\tilde{R}_{2h}] \\
 &\quad + C_{41}[(a_1 - 1)\tilde{R}_3 - (\sum a_3^k)\tilde{R}_{13}] \\
 &\quad + C_{42}[(a_2 - 1)\tilde{R}_3 - (\sum a_3^k)\tilde{R}_{23}] + C_{43}(a_3 - 1)\tilde{R}_3 \\
 &\quad + \sum_{h=4}^N C_{4h}[(a_h - 1)\tilde{R}_3 + (\sum a_3^k)\tilde{R}_{3h}],
 \end{aligned}$$

where the C 's are arbitrary elements of $Z[\pi_1 K]$.

If the matrix M representing ϕ is invertible, ϕ will (by Lemma 1.4) represent the induced map on $C_2(\tilde{K})$ of a homotopy equivalence which induces the identity on $C_1(\tilde{K})$, since $\partial_2 \circ M = \partial_2$.

M will have the following form:

$$\begin{array}{c}
 \tilde{R}_1 \\
 \tilde{R}_2 \\
 \tilde{R}_3 \\
 \vdots \\
 \tilde{R}_{ij}
 \end{array}
 \left[
 \begin{array}{cccccc}
 \tilde{R}_1 & \tilde{R}_2 & & \tilde{R}_3 & \cdots, & R_{ij} \\
 1, & 0, & & 0, & \cdots, & 0 \\
 0, & 1 + C_{11}\bar{a}_1^r + \sum_{h=2}^N C_{1h}\bar{a}_h, & & \sum_{h=1}^N C_{2h}\bar{a}_h, & 0, \dots, 0, & \phi_{ij_2} \\
 0, & C_{31}\bar{a}_1^r + \sum_{h=2}^N C_{2h}\bar{a}_h, & & 1 + \sum_{h=2}^N C_{2h}\bar{a}_h, & 0, \dots, 0, & \phi_{ij_3} \\
 & & \bigcirc & & & \bigcirc \\
 \hline
 & & \bigcirc & & & I_{\binom{2}{2}}
 \end{array}
 \right]$$

where ϕ_{ij_g} ($g = 1, 2$) is the appropriate row vector, i.e.,

$$\begin{aligned}\phi_{12_1} &= C_{11}(\sum a_2^k), \\ \phi_{2h_1} &= -C_{1h}(\sum a_3^k), \quad 4 \leq h \leq N, \\ \phi_{13_1} &= -C_{21}[\sum a_3^k], \\ \phi_{23_1} &= -C_{22}(\sum a_3^k) - C_{13}(\sum a_2^k), \\ \phi_{3h_1} &= C_{2h}(\sum a_3^k), \quad 4 \leq h \leq N, \\ \phi_{12_2} &= -C_{31}(\sum a_2^k), \\ \phi_{2h_2} &= -C_{3h}(\sum a_2^k), \quad 4 \leq h \leq N, \\ \phi_{13_2} &= -C_{41}(\sum a_3^k), \\ \phi_{3h} &= C_{4h}(\sum a_3^k) \quad \text{and} \\ \phi_{23_2} &= -C_{42}(\sum a_2^k) - C_{33}(\sum a_2^k), \\ \phi_{ij_g} &= 0 \quad \text{otherwise.}\end{aligned}$$

As before, this matrix is Whitehead equivalent to

$$\begin{bmatrix} 1 + C_{11}(a_1^r - 1) + \sum_{h=2}^N C_{1h}(a_h - 1), & \sum_{h=1}^N C_{2h}(a - 1) \\ C_{31}(a_1^r - 1) + \sum_{h=2}^N C_{3h}(a_h - 1), & 1 + \sum_{h=1}^N C_{4h}(a_h - 1) \end{bmatrix}.$$

To show that all elements of $\text{Wh}(Z(G))$ can be represented in the above fashion, we invoke Lemma 1.2 using $\{a_1^r, a_2, \dots, a_N\}$ as the generators of G . \square

Since we only need to consider the above twisted presentations, from Sieradski's Theorem 2 in [2, Lemma 2.1], they give us the following two theorems.

THEOREM 2.2. *Let K be a finite 2-dimensional CW-complex with finite abelian fundamental group. Then all of the elements of $\text{Wh}(K)$ are realizable as the torsions of self-equivalences on K .*

THEOREM 2.3. *Homotopy equivalence and simple homotopy equivalence are the same for finite 2-dimensional CW-complexes with finite abelian fundamental group.*

REFERENCES

- [B] H. Bass, *Algebraic K-theory*, Benjamin, New York, 1968.
- [Br] W. Browning, *Finite CW-complexes of cohomological dimension 2 with finite abelian π_1* (unpublished), Forschungsinstitut für Mathematik, E.T. H., CH-8092, Zurich, Switzerland.
- [C] M. Cohen, *A course in simple-homotopy theory*, Springer-Verlag, 1970.
- [D2] M. Dyer, *Simple homotopy types for $(G - m)$ -complexes*, Proc. Amer. Math. Soc. **81** (1981), 111–115.
- [D3] ———, *An application of homological algebra to the homotopy classification of two-dimensional CW-complexes*, Trans. Amer. Math. Soc. **259** (1980), 505–514.
- [DS] M. Dyer and A. Sieradski, *Trees of homotopy types of two-dimensional CW-complexes*, Comment. Math. Helv. **48** (1973), 31–44.

[F] R. Fox, *Free differential calculus. I*, Ann. of Math. (2) **57** (1953), 547–559.

[L] T.-Y. Lam, *Induction theorems for Grothendieck groups and Whitehead groups of finite groups*, Ann. Sci. Ecole. Norm. Sup. **1** (1968), 91–148.

[M1] W. Metzler, *Über den Homotopietyp zweidimensionaler CW-Komplexe und Elementarttransformationen bei Darstellungen von Gruppen durch Erzeugende und definierende Relationen*, J. Reine Angew. Math. **285** (1976), 7–23.

[M2] ———, *Two-dimensional complexes with torsion values not realizable by self-equivalences*, Homological Group Theory, London Math. Soc. Lecture Notes Series 36, Cambridge Univ. Press, 1977, pp. 327–337.

[P] D. Puppe, *Homotopiemengen und Ihre Induzierten Abbildungen. I*, Math. Z. **69** (1958), 299–344.

[S] A. Sieradski, *A semigroup of simple homotopy types*, Math. Z. **153** (1977), 134–148.

[Sw] R. Swan, *Minimal resolutions for finite groups*, Topology **4** (1965), 193–208.

DEPARTMENT OF MATHEMATICS, TULANE UNIVERSITY, NEW ORLEANS, LOUISIANA 70118

Current address: Department of Mathematics and Computer Science, Dartmouth College, Hanover, New Hampshire 03755