

RANDOM RECURSIVE CONSTRUCTIONS: ASYMPTOTIC GEOMETRIC AND TOPOLOGICAL PROPERTIES

R. DANIEL MAULDIN¹ AND S. C. WILLIAMS

ABSTRACT. We study some notions of "random recursive constructions" in Euclidean m -space which lead almost surely to a particular type of topological object; e.g., Cantor set, Sierpiński curve or Menger curve. We demonstrate that associated with each such construction is a "universal" number α such that almost surely the random object has Hausdorff dimension α . This number is the expected value of the sum of some ratios which in the deterministic case yields Moran's formula.

We introduce the notion of a "random recursive construction" and prove several basic facts about such constructions. We give specific examples which lead to random Cantor sets, Sierpiński curves or Menger universal curves. It is perhaps best to begin with a specific example of such a construction. To this end, let us make some notation. Let \mathbf{N} be the set of positive integers and \mathbf{R} the real numbers. If S is a set, let S^* be the set of all finite sequences of elements of S including \emptyset , the empty sequence. If $\alpha = (a_1, \dots, a_n)$ and $\beta = (b_1, \dots, b_m)$ are elements of S , then $|\alpha| = n$, the length of α , and $\alpha * \beta = (a_1, \dots, a_n, b_1, \dots, b_m)$. Now, consider the following construction of a Cantor subset of $[0, 1]$, the unit interval. (Of course, by a Cantor set we mean a compact, perfect, 0-dimensional metric space.) Set $J_\emptyset = [0, 1]$ and, by recursion, if $J_\sigma = [a, b]$, for $\sigma \in \{0, 1\}^*$, then set $J_{\sigma*0} = [a, a + x(b-a)]$ and $J_{\sigma*1} = [a + y(b-a), b]$, where the point (x, y) is chosen from the triangular region $\Delta = \{(s, t) \mid 0 \leq s \leq t \leq 1\}$ according to the uniform distribution. It follows from the results given in this paper that with probability one, the set

$$K = \bigcap_n \left[\bigcup_{\sigma \in \{0,1\}^n} J_\sigma \right]$$

is a Cantor set and the Hausdorff dimension of K , $\dim_H(K)$, is $(\sqrt{17} - 3)/2$.

The paper is organized into four sections. In §1, we define the notion of a random construction and prove a few basic facts concerning such a construction. We demonstrate that with each construction there is a number α such that with probability one the object constructed has Hausdorff dimension $\leq \alpha$. In this section, we relate our results to some deterministic results of P. A. P. Moran [15].

In §2 (Theorem 2.1), we show that certain commonly occurring constructions have finite moments of all orders. This result is necessary for our proof that with probability one the Hausdorff dimension is α .

Received by the editors December 27, 1984 and, in revised form, July 9, 1985.

1980 *Mathematics Subject Classification.* Primary 54H20, 60B05; Secondary 28C10.

Key words and phrases. Martingale, Hausdorff dimension, Cantor set, Sierpiński curve, Menger curve.

¹Research supported in part by a grant from the National Science Foundation.

In §3, we construct a random measure associated with the construction. For each object K_ω generated by the construction there corresponds a Borel measure ν_ω supported on K_ω such that if $\beta < \alpha$ and E is a set with finite $\beta - m$ measure, then $\nu_\omega(E) = 0$. More accurately, in Theorem 3.6, we show that there is a sequence of measures $\nu_{\omega,n}$ supported by K_ω such that if $\beta < \alpha$ and $\beta - m(E) < \infty$, then there is some n_0 with $\nu_{\omega,n_0}(K_\omega) > 0$ and $\nu_{\omega,n_0}(E) = 0$. This, of course, implies that $\dim_H(K_\omega) = \alpha$. Our methods involve a probabilistic mixture of some deterministic results of Moran and of Rogers and Taylor [15, 16]. We remark that there is at least one major unsolved problem in this context. The problem is that we do not have conditions under which one can be guaranteed that with probability one, $\alpha - m(K_\omega) > 0$, although this seems to be the case with many constructions. We comment on this at the end of §3. By $\alpha - m(E)$, we mean the measure of E with respect to the Hausdorff measure defined by the function $h(x) = x^\alpha$.

In the fourth and last section, we give a number of specific examples of constructions which we hope illustrate some of the possibilities. We give several examples of random Cantor subsets of $[0, 1]$. In Example 4.6, we construct random Cantor subsets of $[0, 1] \times [0, 1]$ with Hausdorff dimension 1. In Example 4.7 we generate random Sierpiński curves. In Example 4.8 we generate random locally connected nonplanar continua. We are unable at this time to show that they are Menger curves. In Example 4.9 we modify the construction to generate random Menger curves.

Our construction and the results we prove have features in common with several other processes of current interest, and we believe our methods may be useful in their development. For example, our construction incorporates two features common to the theses of Mandelbrot [11, 12]. It certainly maintains a degree of randomness and yet at the same time preserves some properties of self-similarity. Perhaps even closer to the heart of the matter is the definition of fractal geometry and a fractal given by Cannon [4]. A random construction codifies certain geometric-algorithmic processes which by their nature exhibit some random behavior. The examples given in the last section indicate what one can say about the asymptotic geometric or topological shape of particular constructions. Falconer gives a more complete listing of recent references and a development of some of the central general issues [7]. Related topics are treated by Zähle [19].

1. Random constructions and the Φ -function. Our general model is as follows: We fix a Euclidean space \mathbf{R}^m and a nonempty compact subset J of \mathbf{R}^m . We further require that J is the closure of its interior in \mathbf{R}^m . We assume we have a probability space (Ω, Σ, P) and are given a family of random subsets of \mathbf{R}^m ,

$$\mathbf{J} = \left\{ J_\sigma \mid \sigma \in \mathbf{N}^* = \bigcup_{n=0}^{\infty} \mathbf{N}^n \right\},$$

satisfying three properties.

(1) $J_\emptyset(\omega) = J$ for almost all $\omega \in \Omega$. For every $\sigma \in \mathbf{N}^*$ and for almost all ω , if $J_\sigma(\omega)$ is nonempty, then $J_\sigma(\omega)$ is geometrically similar to J .

(2) For almost every ω and for every $\sigma \in \mathbf{N}^*$, $J_{\sigma*1}(\omega), J_{\sigma*2}(\omega), J_{\sigma*3}(\omega), \dots$ is a sequence of nonoverlapping subsets of $J_\sigma(\omega)$. (A and B nonoverlapping means $\text{int } A \cap \text{int } B = \emptyset$.)

(3) The random vectors $\tau_\sigma = \langle T_{\sigma^*1}, T_{\sigma^*2}, \dots \rangle$, $\sigma \in \mathbf{N}^*$, are i.i.d., where $T_{\sigma^*n}(\omega)$ equals the ratio of the diameter of $J_{\sigma^*n}(\omega)$ to the diameter of $J_\sigma(\omega)$ if $J_\sigma(\omega)$ is nonempty. (For convenience, let $T_\emptyset(\omega) = \text{diameter of } J$.)

We shall call such a system \mathbf{J} a construction. Our constructions require only a "stochastic ratio self-similarity". We now define the random set K by

$$(1.1) \quad K(\omega) = \bigcap_{n=1}^{\infty} \bigcup_{\sigma \in \mathbf{N}^n} J_\sigma(\omega).$$

Our interest centers on the asymptotic properties of this random set K .

For convenience, let $0^0 = 0$. Then $\sum_{p=1}^{\infty} T_{\sigma^*p}^0(\omega)$ counts the number of nonempty $J_{\sigma^*p}(\omega)$, if $J_\sigma(\omega)$ is itself nonempty.

THEOREM 1.1. *Suppose $E(\sum_{n=1}^{\infty} T_n^0) > 1$. Then, with positive probability, K is nonempty. Moreover, given that K is nonempty, then almost surely K has Hausdorff dimension α , where α is the least $\beta > 0$ such that $E(\sum_{n=1}^{\infty} T_n^\beta) \leq 1$.*

We shall establish this theorem by proving several simpler theorems. But first we shall show that α is well defined, relate our results to some results of P. A. P. Moran [15], and motivate the hypothesis $E(\sum_{n=1}^{\infty} T_n^0) > 1$.

Let λ denote m -dimensional Lebesgue measure and define $\Phi: [0, \infty) \rightarrow [0, \infty]$ by

$$(1.2) \quad \Phi(\beta) = E \left(\sum_{n=1}^{\infty} T_n^\beta \right) = E \left(\sum_{n=1}^{\infty} T_{\sigma^*n}^\beta \right),$$

where σ can be any member of \mathbf{N}^* . Now, using the monotone convergence theorem and the fact $T_i \in [0, 1]$ for each i , we see that Φ is nonincreasing and right continuous. We have

$$(1.3) \quad \sum_{n=1}^{\infty} \lambda(\text{int}(J_n)) \leq \lambda(\text{int}(J))$$

and by the scaling property of λ ,

$$(1.4) \quad \lambda(\text{int}(J_n)) = T_n^m \lambda(\text{int}(J)).$$

Thus,

$$(1.5) \quad \sum_{n=1}^{\infty} T_n^m \leq 1 \quad \text{a.s.}$$

Therefore, $\Phi(m) \leq 1$. From these facts it follows that α is well defined. (An example with $\alpha = m$ is given in Example 4.5.)

Concerning the calculation of α , note if $\Phi(y) < \infty$, then Φ is continuous on $[y, +\infty)$ and if, in addition, $\Phi(0) > 1$, then Φ is strictly decreasing on $[y, \infty)$. Thus, if $\Phi(0) > 1$ and $\Phi(\beta) = 1$, then $\beta = \alpha$. However, as Example 4.5 shows, in the general case, $\Phi(\alpha)$ may be less than 1.

An interesting special case of our construction occurs on a n -ary tree; i.e., for some fixed $n \in \mathbf{N}$, our construction is

$$\mathbf{J} = \left\{ J_\sigma \mid \sigma \in \{1, 2, \dots, n\}^* = \bigcup_{j=1}^{\infty} \{1, 2, \dots, n\}^j \right\},$$

or what is essentially the same in our construction using \mathbf{N}^* , we have $\sum_{k=1}^{\infty} T_k^0 \leq n$ a.s. In this case, Φ is a finite valued continuous function, and if further $\Phi(0) > 1$, then $\Phi(\beta) = 1$ has a solution, and so trivially $\Phi(\alpha) = 1$. This will be used later.

As a further special case, if τ_{\emptyset} is distributed as point mass at $(t_1, t_2, \dots, t_n, 0, 0, \dots)$, where $t_1, t_2, \dots, t_n > 0$, Theorem 1.1 implies K has Hausdorff dimension α where α satisfies the equation $t_1^\alpha + t_2^\alpha + \dots + t_n^\alpha = 1$. This result was proved by P. A. P. Moran in [15]. In this sense, $\alpha = \min\{\beta > 0 \mid E(T_1^\beta + T_2^\beta + \dots) \leq 1\}$ and $E(T_1^\alpha + T_2^\alpha + \dots) = 1$, in the special case when Φ is continuous, are generalizations of Moran's equation.

For each $\sigma \in \mathbf{N}^*$, we define the random variable l_σ by

$$(1.6) \quad l_\sigma = \text{diameter}(J_\sigma) = \prod_{n=0}^{|\sigma|} T_{\sigma|n},$$

where $|\sigma|$ is the length of the finite sequence σ and $\sigma|n$ is the sequence obtained by restricting σ to its first n terms. ($\sigma|\emptyset = \emptyset =$ empty sequence.) For each $n \in \mathbf{N}$, we let \mathcal{F}_n denote the σ -algebra $\sigma - (\{T_\gamma: |\gamma| \leq n\}) = \sigma - (\{\tau_\gamma: |\gamma| < n\})$. For each $n \in \mathbf{N}$ and $\beta \geq 0$, $S_{\beta,n}$ is the random variable

$$(1.7) \quad S_{\beta,n} = \sum_{\sigma \in \mathbf{N}^n} l_\sigma^\beta.$$

We note the useful fact concerning conditional expectations:

$$(1.8) \quad E(S_{\beta,n+1} | \mathcal{F}_n) = \Phi(\beta) S_{\beta,n},$$

where $\infty \cdot 0 = 0$. In order to derive (1.8), we use (1.7) and (1.6) to obtain the formula

$$(1.9) \quad S_{\beta,n+1} = \sum_{\sigma \in \mathbf{N}^{n+1}} l_\sigma^\beta \left[\sum_{p=1}^{\infty} T_{\sigma^*p}^\beta \right].$$

We obtain (1.8) by taking conditional expectations on both sides of (1.9), noting that l_σ^β is \mathcal{F}_n -measurable for $|\sigma| \leq n$ and noting

$$E \left(\sum_{p=1}^{\infty} T_{\sigma^*p}^\beta | \mathcal{F}_n \right) = E \left(\sum_{p=1}^{\infty} T_{\sigma^*p}^\beta \right) = \Phi(\beta)$$

since τ_σ is independent of \mathcal{F}_n for $n = |\sigma|$.

Now, we shall motivate the hypothesis $\Phi(0) > 1$.

Clearly, $S_{0,n}$ counts the number of nonempty J_σ for $\sigma \in \mathbf{N}^n$. In fact, $\{S_{0,n}\}_{n=1}^{\infty}$ is a classical Galton-Watson branching process, in case $P(\sum T_n^0 < \infty) = 1$ (see [1, pp. 7-8]). Further, $\Phi(0)$ is the mean number of offspring of a single parent in the branching process. It is well known that if $\Phi(0) \leq 1$, then either $\{S_{0,n}\}_{n=1}^{\infty}$ is a process bound for extinction ($S_{0,n} \rightarrow 0$ as $n \rightarrow \infty$ a.s.) or it is trivial ($S_{0,n} = 1$ a.s., for each n). Thus, if $\Phi(0) \leq 1$, either $K = \emptyset$ a.s. or K is a point a.s. or $K = J$ a.s. On the other hand, if $\Phi(0) > 1$, a slight generalization of the well-known result is $S = \lim S_{0,n}$ exists a.s. and $S \in \{0, \infty\}$ a.s.

If $\Phi(0) > 1$, then $P(S = \infty) > 0$. It is clear from what we have said that the only interesting case is $\Phi(0) > 1$. This will be assumed throughout the remainder of the paper but often stated for emphasis.

THEOREM 1.2. *Suppose $\Phi(0) > 1$. Then for almost all ω ,*

$$(1.10) \quad "S(\omega) = \infty \text{ if and only if } K(\omega) \neq \emptyset".$$

PROOF. Clearly, if $S(\omega) \neq \infty$, then almost surely $S(\omega) = 0$. Thus, there is some $n \in \mathbf{N}$ such that $S_{0,n}(\omega) = 0$. This means $K(\omega) \subseteq \bigcup_{\sigma \in \mathbf{N}^n} J_\sigma(\omega) = \emptyset$.

For the converse we consider two cases: Case A: $P(\sum_{n=1}^\infty T_n^0 < \infty) = 1$ and Case B: $P(\sum_{n=1}^\infty T_n^0 = \infty) > 0$.

Case A. Since $P(\sum_{n=1}^\infty T_n^0 < \infty) = 1$, then a.s., for each n , $\bigcup_{\sigma \in \mathbf{N}^n} J_\sigma(\omega)$ is a finite union of compact sets and thus is itself compact. Therefore, if $S(\omega) = \infty$, then $\bigcap_{n=1}^\infty \bigcup_{\sigma \in \mathbf{N}^n} J_\sigma(\omega) \neq \emptyset$, by the intersection property of a nested sequence of nonempty compact sets.

Case B. Suppose $P(\sum_{n=1}^\infty T_n^0 = \infty) = \delta > 0$. We claim that for any σ , the following statement is true for almost all ω :

$$(1.11) \quad \text{"If } \sum_{n=1}^\infty T_{\sigma^{**}n}^0(\omega) = \infty, \text{ then there exists } k \in \mathbf{N} \text{ such that } T_{\sigma^{**}k}^0(\omega) = 1 \text{ and } \sum_{n=1}^\infty T_{\sigma^{**}k^{**}n}^0(\omega) = \infty."$$

To see this, let A be the event $\sum_{n=1}^\infty T_{\sigma^{**}n}^0 = \infty$, B_k the event $\forall n \leq k$, if $T_{\sigma^{**}n}^0 = 1$, then $\sum_{j=1}^\infty T_{\sigma^{**}n^{**}j}^0 < \infty$ and B_∞ be the event $\forall n \in \mathbf{N}$, if $T_{\sigma^{**}n}^0 = 1$, then $\sum_{j=1}^\infty T_{\sigma^{**}n^{**}j}^0 < \infty$. Let H consist of all finite subsets of \mathbf{N} and G_k be the random set $\{n \leq k | T_{\sigma^{**}n}^0 = 1\}$. Thus,

$$\begin{aligned} P(A \cap B_\infty) &\leq P(A \cap B_k) = \sum_{D \in H} P(A \cap B_k \text{ and } D = G_k) \\ &= \sum_{D \in H} P \left(A, D = G_k, \forall n \in D \left[\sum_{j=1}^\infty T_{\sigma^{**}n^{**}j}^0 < \infty \right] \right), \end{aligned}$$

and, by independence,

$$= \sum_{D \in H} P(A \text{ and } D = G_k)(1 - \delta)^{\#D},$$

where $\#E$ denotes the cardinality of E . Thus,

$$P(A \cap B_\infty) \leq \sum_{i=1}^\infty P(A \text{ and } \#G_k = i)(1 - \delta)^i.$$

Fix $p \in \mathbf{N}$. Then

$$P(A \cap B_\infty) \leq \sum_{i=1}^p P(A \text{ and } \#G_k = i) + (1 - \delta)^{p+1} P(A \text{ and } \#G_k \geq p + 1)$$

or,

$$P(A \cap B_\infty) \leq P(A \text{ and } \#G_k \leq p) + (1 - \delta)^{p+1}.$$

Letting $k \rightarrow \infty$, we obtain

$$P(A \cap B_\infty) \leq (1 - \delta)^{p+1}.$$

Now, letting $p \rightarrow \infty$, we find $P(A \cap B_\infty) = 0$ or $P(B_\infty | A) = 0$. This yields (1.11).

From (1.11), it is easy to see that the following statement is true almost surely:

$$(1.12) \quad \text{“If } J_\sigma(\omega) \neq \emptyset \text{ and } \sum_{n=1}^\infty T_{\sigma^{*n}}^0(\omega) = \infty, \text{ then there exists a sequence of natural numbers } k_1, k_2, k_3, \dots \text{ which depend on } \omega, \text{ so that } J_{\sigma^{*k_1 * k_2 * \dots * k_t}}(\omega) \neq \emptyset \text{ for each } t. \text{ Thus, } \emptyset \neq \bigcap_{t=1}^\infty J_{\sigma^{*k_1 * \dots * k_t}}(\omega) \subseteq K(\omega)\text{”}.$$

Thus, we only need to establish that

$$(1.13) \quad P \left(\exists \sigma \in \mathbf{N}^* \left[J_\sigma \neq \emptyset \text{ and } \sum_{n=1}^\infty T_{\sigma^{*n}}^0 = \infty \right] \mid S = \infty \right) = 1,$$

in order to complete the proof in Case B.

Let A be the event $(\forall \sigma \in \mathbf{N}^* [J_\sigma \neq \emptyset \rightarrow \sum_{n=1}^\infty T_{\sigma^{*n}}^0 < \infty])$. For each k , let $A(k)$ denote the event $(\forall \sigma \in \mathbf{N}^k [J_\sigma \neq \emptyset \rightarrow \sum_{n=1}^\infty T_{\sigma^{*n}}^0 < \infty])$. Then, concerning indicator functions, we have $1_A \leq 1_{A(n)}$ and $E(1_{A(n)} | \mathcal{F}_n) = (1 - \delta)^{S_{0,n}}$. Now, $E(1_A | \mathcal{F}_n) \rightarrow 1_A$ as $n \rightarrow \infty$. But, also, $E(1_A | \mathcal{F}_n) \leq (1 - \delta)^{S_{0,n}} \rightarrow (1 - \delta)^S$ as $n \rightarrow \infty$ almost surely. Therefore, for almost all ω , if $S(\omega) = \infty$, then $\omega \notin A$. Q.E.D.

To begin the demonstration of Theorem 1.1, we know, according to (1.5), $\Phi(m) \leq 1$. Thus, if $\Phi(0) > 1$, $E(S_{m+1,n}) = (\text{diam}(J))^{m+1} (\Phi(m+1))^n$ and $E(\sum_{n=1}^\infty S_{m+1,n}) = (\text{diam}(J))^{m+1} / (1 - \Phi(m+1)) < +\infty$. This implies $\sum_{\sigma \in \mathbf{N}^n} l_\sigma^{m+1} \rightarrow 0$ a.s. So, $\sup_{\sigma \in \mathbf{N}^n} l_\sigma^{m+1} \rightarrow 0$ a.s. Therefore,

$$(1.14) \quad \text{if } \Phi(0) > 1, \quad \sup_{\sigma \in \mathbf{N}^n} l_\sigma \rightarrow 0 \quad \text{a.s.}$$

THEOREM 1.3. *Suppose $\Phi(0) > 1$. Then almost surely the Hausdorff dimension of K is $\leq \alpha$.*

PROOF. Since $\Phi(\alpha) \leq 1$, by (1.8), $\{S_{\alpha,n}\}_{n=1}^\infty$ is a positive supermartingale and thus converges to some real random variable X . For each $\sigma \in \mathbf{N}^*$, there exists a random m -sphere M_σ so that $J_\sigma \subseteq M_\sigma$ and the radius of M_σ is l_σ . According to (1.14), $\sup_{\sigma \in \mathbf{N}^n} l_\sigma \rightarrow 0$ as $n \rightarrow \infty$. So,

$$(1.15) \quad \alpha - m(K) \leq \lim_{n \rightarrow \infty} \sum_{\sigma \in \mathbf{N}^n} (\text{diam}(M_\sigma))^\alpha \leq 2^\alpha X < \infty,$$

where $\alpha - m$ is the α -dimension Hausdorff measure. Q.E.D.

REMARK. If $\Phi(\alpha) = 1$, then $\{S_{\alpha,n}\}_{n=1}^\infty$ is a martingale.

We will now begin consideration towards showing the Hausdorff dimension of K is almost surely $\geq \alpha$. First, some involved calculations concerning moments seem necessary.

2. The moments of X .

THEOREM 2.1. *Suppose $\Phi(0) > 1$, $k_0 \in \mathbf{N}$ and $E((\sum_{n=1}^\infty T_n^\alpha)^{k_0}) < \infty$. Then $\{(S_{\alpha,n}, \mathcal{F}_n)\}_{n=1}^\infty$ is L^p -bounded for all $p \in [1, k_0]$. Consequently, $X = \lim_n S_{\alpha,n}$ has a finite moment of order k_0 .*

It is sufficient to establish the following lemma.

LEMMA 2.2. For every $k \in [1, k_0]$, $\{E(S_{\alpha,n}^k)\}_{n=1}^\infty \in l^\infty$ and for any $\gamma > \alpha$ and any $z \in (0, 1]$, $\{|E(S_{\gamma,n}^k)|^z\}_{n=1}^\infty \in l^1$.

PROOF. For each $p, n \in \mathbf{N}$, let $S_{p;\gamma,n}$ denote $\sum_{\sigma \in \{1, \dots, p\}^n} l_\sigma^\gamma$. For any $k \in (0, \infty)$, $E(S_{p;\gamma,n}^k) \leq (\text{diam}(J))^{\gamma k p^{nk}}$. Consider

$$(2.1) \quad E(S_{p;\gamma,n+1}^k | \mathcal{F}_n) = E \left(\left(\sum_{\sigma \in \{1, \dots, p\}^n} l_\sigma^\gamma \left(\sum_{j=1}^p T_{\sigma^* j}^\gamma \right) \right)^k \mid \mathcal{F}_n \right).$$

Thus, if k is also an integer,

$$(2.2) \quad E(S_{p;\gamma,n+1}^k | \mathcal{F}_n) = \sum_{f: \{1, \dots, k\} \rightarrow \Delta} \prod_{i=1}^k x_{f(i)} \prod_{j=1}^k m_{p;\gamma,j}^{r_j(f)},$$

where $\Delta = \{1, \dots, p\}^n$, $x_\sigma = l_\sigma^\gamma$ for each $\sigma \in \Delta$, $m_{p;\gamma,j} = E[(\sum_{i=1}^p T_i^\gamma)^j]$ and $r_j(f) = \#\{A \in \ker(f) \mid \#A = j\}$ with $\ker(f) = \{f^{-1}(i) \mid i \in \Delta\}$. Notice that $\prod_{j=1}^k m_{p;\gamma,j}^{r_j(f)}$ depends only on $\ker(f)$, the partition of $\{1, \dots, k\}$ induced by f . So, for each partition π , let $m_{p;\gamma,\pi}$ denote this number. Thus,

$$(2.3) \quad E(S_{p;\gamma,n+1}^k | \mathcal{F}_n) = \sum_{\pi \in P} m_{p;\gamma,\pi} k_\pi,$$

where P is the set of all partitions and $k_\pi = \sum_{\ker(f)=\pi} \prod_{i=1}^k x_{f(i)}$. Our notation is consistent with that given by Doubilet [6]. So, according to [6, Theorem 2, equation 13], we have

$$(2.4) \quad E(S_{p;\gamma,n+1}^k | \mathcal{F}_n) = \sum_{\pi \in P} m_{p;\gamma,\pi} \sum_{\sigma \geq \pi} \mu(\pi, \sigma) s_\sigma;$$

interchanging order of summation,

$$(2.5) \quad = \sum_{\sigma \in P} s_\sigma \sum_{\pi \leq \sigma} \mu(\pi, \sigma) m_{p;\gamma,\pi},$$

where the Möbius function μ , the order \leq and the s_σ 's are as given in [6].

In our particular case,

$$(2.6) \quad s_\sigma = \prod_{j=1}^k S_{p;j\gamma,n}^{\hat{\sigma}(j)},$$

where $\hat{\sigma}(j) = \#\{A \in \sigma \mid \#(A) = j\}$. Now, set $B = \{\eta: \{1, \dots, k\} \rightarrow \mathbf{Z}_+ \mid \sum_{j=1}^k j\eta(j) = k\}$, and

$$(2.7) \quad c_{p;\gamma,\eta} = \sum_{\substack{\sigma \in P \\ \hat{\sigma} = \eta}} \sum_{\pi \leq \sigma} \mu(\pi, \sigma) m_{p;\gamma,\pi}.$$

We have

$$(2.8) \quad E(S_{p;\gamma,n+1}^k | \mathcal{F}_n) = \sum_{\eta \in B} c_{p;\gamma,\eta} \prod_{j=1}^k S_{p;j\gamma,n}^{\eta(j)}.$$

Taking expectations, and using the triangle inequality we find

$$(2.9) \quad E(S_{p;\gamma,n+1}^k) \leq \sum_{\eta \in B} |c_{p;\gamma,\eta}| E \left(\prod_{j=1}^k S_{p;j\gamma,n}^{\eta(j)} \right).$$

Let $m_{\gamma,j} = E((\sum_{i=1}^{\infty} T_i^\gamma)^j)$. If $\gamma \geq \alpha$ and $j \in [0, k_0]$, then $m_{\gamma,j} < \infty$ and $m_{p;\gamma,j} \rightarrow m_{\gamma,j}$ as $p \rightarrow \infty$ for $j = 1, \dots, k$. Thus, $c_{p;\gamma,\eta} \rightarrow c_{\gamma,\eta}$ (where $c_{\gamma,\eta}$ is defined similarly to $c_{p;\gamma,\eta}$). But, $S_{p;\gamma,q}^j$ converges upward to $S_{\gamma,q}^j$ as $p \rightarrow \infty$. Taking limits in (2.9) as $p \rightarrow \infty$, we find

$$(2.10) \quad E(S_{\gamma,n+1}^k) \leq \sum_{\eta \in B} |c_{\gamma,\eta}| E \left[\prod_{j=1}^k S_{j\gamma,n}^{\eta(j)} \right].$$

Let ξ denote the unique element of B such that $\xi(1) = k$. We calculate $c_{\gamma,\xi}$. Note there is only one $\sigma \in P$ such that $\hat{\sigma} = \xi$, namely the discrete partition. Also, if $\pi \leq \sigma$, then $\pi = \sigma$. So, according to (2.7) $c_{\gamma,\xi} = \mu(\sigma, \sigma) m_{\gamma,\sigma} = (\Phi(\gamma))^k$. We can rewrite (2.10) as

$$(2.11) \quad E(S_{\gamma,n+1}^k) \leq \Phi(\gamma)^k E(S_{\gamma,n}^k) + \sum_{\substack{\eta \in B \\ \eta \neq \xi}} |c_{\gamma,\eta}| E \left[\prod_{j=1}^k S_{j\gamma,n}^{\eta(j)} \right].$$

From this we derive by backwards recursion,

$$(2.12) \quad E(S_{\gamma,n+1}^k) \leq \Phi(\gamma)^{k(n+1)} l_0^k + \sum_{\substack{\eta \in B \\ \eta \neq \xi}} |c_{\gamma,\eta}| \sum_{t=0}^n \Phi(\gamma)^{k(n-t)} E \left[\prod_{j=1}^k S_{j\gamma,t}^{\eta(j)} \right].$$

Easily, $E(S_{\gamma,n}^1) = \Phi(\gamma)^n l_0^1$ implies Lemma 2.2 for $k = 1$ since $\Phi(\alpha) \leq 1$ and $\Phi(\gamma) \in (0, 1)$ if $\gamma > \alpha$. As another special case, let $k = 2$. Then B has only two elements $(2, 0)$ and $(0, 1)$. Thus,

$$(2.13) \quad E(S_{\gamma,n+1}^2) \leq l_0^{2\gamma} \left[\Phi(\gamma)^{2(n+1)} + |c_{\gamma,(0,1)}| \sum_{t=0}^n \Phi(\gamma)^{2(n-t)} \Phi(2\gamma)^t \right].$$

In particular,

$$(2.14) \quad \begin{aligned} E(S_{\alpha,n+1}^2) &\leq l_0^{2\alpha} \left[1 + |c_{\alpha,(0,1)}| \sum_{t=0}^n \Phi(2\alpha)^t \right] \\ &\leq l_0^{2\alpha} \left[1 + \frac{|c_{\alpha,(0,1)}|}{1 - \Phi(2\alpha)} \right] < \infty, \end{aligned}$$

since $\Phi(2\alpha) < 1$ (because $\Phi(0) > 1$). Actually, from (2.7) we have

$$c_{\alpha,(0,1)} = \text{Var} \left(\sum_{n=1}^{\infty} T_n^\alpha \right).$$

If $\gamma > \alpha$, then $\Phi(\gamma) \in (0, 1)$ and $\Phi(2\gamma) \in (0, 1)$. Using the fact if $a, b > 0$, then $(a + b)^z \leq a^z + b^z$ for any $z \in [0, 1]$, we find

$$(2.15) \quad \sum_{n=0}^{\infty} [E(S_{\gamma,n}^2)]^z \leq \sum_{n=0}^{\infty} \Phi(\gamma)^{2zn} l_0^{2z\gamma} + |c_{\gamma,(0,1)}|^z \sum_{n=0}^{\infty} \sum_{t=0}^n \Phi(\gamma)^{2z(n-t)} \Phi(2\gamma)^{zt} l_0^{2z\gamma} \\ \leq l_0^{2z\gamma} \left[\frac{1}{1 - \Phi(\gamma)^{2z}} + |c_{\gamma,(0,1)}|^z \frac{1}{1 - \Phi(\gamma)^{2z}} \cdot \frac{1}{1 - \Phi(2\gamma)^z} \right] < \infty.$$

If $1 < k < 2$, and $z \in (0, 1]$, then $[E(S_{\gamma,n}^k)]^z \leq [E(S_{\gamma,n})]^z + [E(S_{\gamma,n}^2)]^z$. Thus Lemma 2.2 is true for $k \in [1, 2]$.

Suppose Lemma 2.2 is true for $\tilde{k} \in [1, k - 1]$, where $k \leq k_0$ and k is an integer. If $\eta \in B$ with $\eta \neq \xi$, then $\eta(1) \leq k - 2$ and $\sum_{j=1}^k \eta(j) \leq k - 1$. Thus there are nonnegative reals r_1, \dots, r_k , depending on η , so that $\eta(j)/k - 1 \leq r_j \leq 1$ and $\sum_{j=1}^k r_j = 1$. Let $p(\eta, j) = 1/r_j$. Then for each $\eta \in B$ with $\eta \neq \xi$, $\sum_{j=1}^k 1/p(\eta, j) = 1$, $p(\eta, j)\eta(j) \leq k - 1$, and if $\eta(j) \neq 0$, then $\eta(j)p(\eta, j) \geq 1$, i.e., $p(\eta, j)\eta(j) \in [1, k - 1]$ if $\eta(j) \neq 0$. By Hölder's inequality,

$$(2.16) \quad E \left(\prod_{j=1}^k S_{j,\gamma,n}^{\eta(j)} \right) \leq \prod_{j=1}^k \|S_{j,\gamma,n}^{\eta(j)}\|_{p(\eta,j)}.$$

Substituting into (2.12), we find for $z \in (0, 1]$

$$(2.17) \quad [E(S_{\gamma,n}^k)]^z \leq \Phi(\gamma)^{zkn} l_0^{zk} + \sum_{\substack{\eta \in B \\ \eta \neq \xi}} |c_{\gamma,\eta}|^z \sum_{t=0}^{n-1} \Phi(\gamma)^{zk(n-1-t)} \prod_{j=1}^k \|S_{j,\gamma,t}^{\eta(j)}\|_{p(\eta,j)}^z.$$

Let $W_{\eta,j} = \{\|S_{j,\gamma,t}^{\eta(j)}\|_{p(\eta,j)}^z\}_{t=0}^{\infty}$. If $\eta(j) = 0$, then obviously $W_{\eta,j} \in l^\infty$. If $\eta(j) > 0$, then $W_{\eta,j} \in l^\infty$ by the induction hypothesis and $\eta(j)p(\eta, j) \in [1, k - 1]$. But, since $\eta \neq \xi$, there is some $j \geq 2$ so that $\eta(j) \geq 1$ which implies $W_{\eta,j} \in l^1$ since $j\gamma > \alpha$ if $\gamma \geq \alpha$ and $j \geq 2$. Thus if $\gamma \geq \alpha$, then

$$(2.18) \quad \left\{ \prod_{j=1}^k \|S_{j,\gamma,t}^{\eta(j)}\|_{p(\eta,j)}^z \right\}_{t=0}^{\infty} \in l^1.$$

Obviously, using $\Phi(\alpha) \leq 1$,

$$(2.19) \quad \sup_n ([E(S_{\alpha,n}^k)]^z) \leq l_0^{\alpha zk} + \sum_{\substack{\eta \in B \\ \eta \neq \xi}} |c_{\alpha,\eta}|^z \left(\sum_{t=0}^z \prod_{j=1}^k \|S_{j,\alpha,t}^{\eta(j)}\|_{p(\eta,j)}^z \right) < \infty.$$

If $\gamma > \alpha$, then $\Phi(\gamma) \in (0, 1)$. Using this fact, summing both sides of (2.17), making a change of variables and an obvious approximation, we obtain

$$(2.20) \quad \sum_{n=0}^{\infty} [E(S_{\gamma,n}^k)]^z \leq \frac{l_0^{\gamma zk}}{1 - \Phi(\gamma)^{zk}} + \sum_{\substack{\eta \in B \\ \eta \neq \xi}} |c_{\gamma,\eta}|^z \frac{\sum_{t=0}^{\infty} \prod_{j=1}^k \|S_{j,\gamma,t}^{\eta(j)}\|_{p(\eta,j)}^z}{1 - \Phi(\gamma)^{zk}} < \infty.$$

Since for any $r \in [k - 1, k]$ and $z \in (0, 1]$,

$$(2.21) \quad [E(S_{\gamma,n}^r)]^z \leq [E(S_{\gamma,n}^{k-1})]^z + [E(S_{\gamma,n}^k)]^z,$$

we have extended Lemma 2.2 to $[1, k]$. This completes the induction. Q.E.D.

Thus,

$$(3.6) \quad \begin{aligned} \overline{\lim}_{p,q \rightarrow \infty} \varepsilon_{p,q}(\omega) &\leq 2 \sum_{\sigma \in \mathbf{N}^k} l_\sigma^\alpha(\omega) \text{diam}(f(J_\sigma(\omega))) X_\sigma(\omega) \\ &\leq 2 \sup_{\sigma \in \mathbf{N}^k} \text{diam}(f(J_\sigma(\omega))) X(\omega) \end{aligned}$$

if $\omega \in \Omega'$. But, if $\omega \in \Omega'$, $\sup_{\sigma \in \mathbf{N}^k} \text{diam}(f(J_\sigma(\omega))) \rightarrow 0$ as $k \rightarrow \infty$. Thus,

$$(3.7) \quad \overline{\lim}_{p,q \rightarrow \infty} \varepsilon_{p,q}(\omega) = 0 \quad \text{if } \omega \in \Omega'.$$

Obviously, F_ω is linear positive and for any $f \in C_c(\mathbf{R}^m)$ so that $J \subseteq f^{-1}(1)$, we have $F_\omega(f) = \lim_{n \rightarrow \infty} \sum_{\sigma \in \mathbf{N}^n} l_\sigma^\alpha(\omega) = K(\omega)$. Thus, $\|F_\omega\| = X(\omega)$. We let ν_ω denote the Borel measure on \mathbf{R}^m so that $F_\omega(f) = \int_{\mathbf{R}^m} f(x) d\nu_\omega(x)$.

THEOREM 3.2. *Suppose $\Phi(0) > 1$. If A is a compact subset of \mathbf{R}^m , then*

$$(3.8) \quad \nu(A) = \lim_{n \rightarrow \infty} \sum_{\substack{\sigma \in \mathbf{N}^n \\ J_\sigma \cap A \neq \emptyset}} l_\sigma^\alpha X_\sigma \quad \text{a.s.}$$

In fact,

$$(3.9) \quad \sum_{\substack{\sigma \in \mathbf{N}^n \\ J_\sigma \cap A \neq \emptyset}} l_\sigma^\alpha X_\sigma \downarrow \nu(A) \quad \text{as } n \rightarrow \infty \text{ a.s.}$$

PROOF. Fix $\omega \in \Omega'$ and $k \in \mathbf{N}$. Let $\varepsilon > 0$. Since $\sum_{\sigma \in \mathbf{N}^k} l_\sigma^\alpha(\omega) X_\sigma(\omega)$ is finite, there is a finite set $M \subseteq \mathbf{N}^k$ such that $\sum_{\sigma \in \mathbf{N}^k \setminus M} l_\sigma^\alpha(\omega) X_\sigma(\omega) < \varepsilon$. Let f be a continuous map of \mathbf{R}^m into $[0, 1]$ such that $f^{-1}(1) = A$ and $J_\sigma \subseteq f^{-1}(0)$ if $\sigma \in M$ and $J_\sigma \cap A = \emptyset$. Now

$$(3.10) \quad \begin{aligned} \nu_\omega(A) &\leq \int_{\mathbf{R}^m} f d\nu_\omega = \lim_{n \rightarrow \infty} \sum_{\sigma \in \mathbf{N}^n} f(s_\sigma(\omega)) l_\sigma^\alpha(\omega) \\ &= \lim_{n \rightarrow \infty} \sum_{\sigma \in \mathbf{N}^k} \sum_{\eta \in \mathbf{N}^{n-k}} f(s_{\sigma*\eta}(\omega)) l_{\sigma*\eta}^\alpha(\omega) \\ &\leq \lim_{n \rightarrow \infty} \sum_{\substack{\sigma \in \mathbf{N}^k \\ J_\sigma(\omega) \cap A \neq \emptyset}} \sum_{\eta \in \mathbf{N}^{n-k}} l_{\sigma*\eta}^\alpha(\omega) \\ &\quad + \lim_{n \rightarrow \infty} \sum_{\substack{\sigma \in \mathbf{N}^k \setminus M \\ J_\sigma(\omega) \cap A = \emptyset}} \sum_{\eta \in \mathbf{N}^{n-k}} l_{\sigma*\eta}^\alpha(\omega). \end{aligned}$$

Note

$$\lim_{n \rightarrow \infty} \sum_{\sigma \in D \subseteq \mathbf{N}^k} \sum_{\eta \in \mathbf{N}^{n-k}} l_{\sigma*\eta}^\alpha(\omega) = \sum_{\sigma \in D} l_\sigma(\omega) X_\sigma(\omega).$$

So,

$$\nu_\omega(A) \leq \sum_{\substack{\sigma \in \mathbf{N}^k \\ J_\sigma(\omega) \cap A \neq \emptyset}} l_\sigma(\omega) X_\sigma(\omega) + \varepsilon.$$

Or,

$$(3.11) \quad \nu_\omega(A) \leq \sum_{\substack{\sigma \in \mathbf{N}^k \\ J_\sigma(\omega) \cap A \neq \emptyset}} l_\sigma(\omega) X_\sigma(\omega).$$

Since, for any $\sigma \in \mathbf{N}^*$, $l_\sigma^\alpha X_\sigma = \sum_{n=1}^\infty l_{\sigma^{**n}}^\alpha X_{\sigma^{**n}}$, we have

$$(3.12) \quad \sum_{\substack{\sigma \in \mathbf{N}^k \\ J_\sigma(\omega) \cap A \neq \emptyset}} l_\sigma^\alpha(\omega) X_\sigma(\omega) \geq \sum_{\substack{\sigma \in \mathbf{N}^{k+1} \\ J_\sigma(\omega) \cap A \neq \emptyset}} l_\sigma^\alpha(\omega) X_\sigma(\omega).$$

Let $\mu_\omega(A) = \lim_n \sum_{\sigma \in \mathbf{N}^n; A \cap J_\sigma(\omega) \neq \emptyset} l_\sigma^\alpha(\omega) X_\sigma(\omega)$. Suppose B is a compact subset of \mathbf{R}^m disjoint from A ; then there is a $k \in \mathbf{N}$ so that if $\sigma \in \mathbf{N}^k$, then J_σ meets at most one of A and B . Thus, $\mu_\omega(A) + \mu_\omega(B) \leq X(\omega)$. Further,

$$(3.13) \quad \nu_\omega(A) + \nu_\omega(B) \leq \mu_\omega(A) + \nu_\omega(B) \leq \mu_\omega(A) + \mu_\omega(B) \leq X(\omega).$$

Find compact sets $B_1 \subseteq B_2 \subseteq B_3 \subseteq \dots \subseteq A^c$ so that $\nu(B_n) \uparrow \nu(A^c)$. We see

$$(3.14) \quad X(\omega) = \nu_\omega(A) + \nu_\omega(A^c) \leq \mu_\omega(A) + \nu_\omega(A^c) \leq X(\omega),$$

i.e., $\mu_\omega(A) = \nu_\omega(A)$. Q.E.D.

THEOREM 3.3. *Suppose $\Phi(0) > 1$. Almost surely $\nu(K) = X$.*

PROOF. Since $K = \bigcap_{n=1}^\infty \bigcup_{\sigma \in \mathbf{N}^n} J_\sigma$ is a nested intersection,

$$\nu(K) = \lim_{n \rightarrow \infty} \nu \left(\bigcup_{\sigma \in \mathbf{N}^n} J_\sigma \right).$$

Temporarily fix n and let $\Delta_1 \subseteq \Delta_2 \subseteq \Delta_3 \subseteq \dots$ be an increasing sequence of finite subsets of \mathbf{N}^n such that $\bigcup \Delta_j = \mathbf{N}^n$. According to Theorem 3.2, we have, for each i ,

$$\begin{aligned} \nu \left(\bigcup_{\sigma \in \Delta_i} J_\sigma \right) &= \lim_{k \rightarrow \infty} \sum_{\substack{\eta \in \mathbf{N}^k \\ J_\eta \cap J_\sigma \neq \emptyset \\ \text{for some } \sigma \in \Delta_i}} l_\eta^\alpha X_\eta \\ &\geq \overline{\lim}_{k \rightarrow \infty} \sum_{\sigma \in \Delta_i} \sum_{\eta \in \mathbf{N}^k} l_{\sigma^{**\eta}}^\alpha X_{\sigma^{**\eta}} = \sum_{\sigma \in \Delta_i} l_\sigma^\alpha X_\sigma. \end{aligned}$$

Thus,

$$X \geq \nu \left(\bigcup_{\sigma \in \mathbf{N}^n} J_\sigma \right) \geq \sum_{\sigma \in \Delta_j} l_\sigma^\alpha X_\sigma \xrightarrow{j \rightarrow \infty} \sum_{\sigma \in \mathbf{N}^n} l_\sigma^\alpha X_\sigma = X. \quad \text{Q.E.D.}$$

THEOREM 3.4. *If $E(X) > 0$, then $P(\nu(K) > 0 \mid K \neq \emptyset) = 1$.*

PROOF. By hypothesis, $E(X) > 0$ and so $P(X > 0) = \delta > 0$. Recall that for each $\sigma \in \mathbf{N}^n$, X_σ exists a.s. and has the same distribution as $X/(\text{diam } J)^\alpha$. Further, for each $n \in \mathbf{N}$, $\{X_\sigma \mid \sigma \in \mathbf{N}^n\}$ is an independent family and, as a family, also independent of \mathcal{F}_n . We note again

$$(3.15) \quad X = \sum_{\sigma \in \mathbf{N}^n} l_\sigma^\alpha X_\sigma.$$

As in the proof of Theorem 1.2 we consider two cases:

$$\text{A: } P\left(\sum_{n=1}^{\infty} T_n^0 < \infty\right) = 1 \quad \text{and} \quad \text{B: } P\left(\sum_{n=1}^{\infty} T_n^0 = \infty\right) > 0.$$

Case A. We will establish that for each n and q ,

$$(3.16) \quad P(X > 0 \text{ and } S_{0,n} \geq q) \leq (1 - \delta)^q P(S_{0,n} \geq q).$$

Since $P(\sum_{n=1}^{\infty} T_n^0 < \infty) = 1$, the random index set $D_n = \{\sigma \in \mathbf{N}^n \mid l_\sigma > 0\}$ is finite. Let F be a finite subset of \mathbf{N}^n . Using (3.15) we see

$$\begin{aligned} P(X = 0 \text{ and } F = D_n) &= P(X_\sigma = 0 \text{ for } \sigma \in F \text{ and } F = D_n) \\ &= (1 - \delta)^{\#F} P(D_n = F), \end{aligned}$$

due to independence. Now, inequality (3.16) follows by summing over finite F such that $\#F \geq q$ and using $(1 - \delta)^{\#F} \leq (1 - \delta)^q$.

From (3.16), we calculate

$$(3.17) \quad P(X > 0) \geq P(X > 0 \text{ and } S_{0,n} \geq q) \geq [1 - (1 - \delta)^q] P(S_{0,n} \geq q).$$

Noting $P(S_{0,n} \geq q) \rightarrow P(S = \infty)$ as $n \rightarrow \infty$, and letting $n \rightarrow \infty$ in (3.17), and then letting $q \rightarrow \infty$, we find

$$(3.18) \quad P(X > 0) \geq P(S = \infty).$$

But a.s. "if $S = 0$, then there is some $n \in \mathbf{N}$ so that $S_{0,n} = 0$; i.e., $X = \sum_{\sigma \in \mathbf{N}^n} l_\sigma X_\sigma = 0$ ". Thus,

$$(3.19) \quad \text{"if } X > 0, \text{ then } S = \infty\text{" a.s.}$$

Combining (3.18), (3.19), Theorems 1.2 and 1.3, we find

$$\text{"}K \neq \emptyset \text{ if and only if } X > 0 \text{ if and only if } \nu(K) > 0\text{" a.s.}$$

Case B. Suppose $P(\sum_{n=1}^{\infty} T_n^0 = \infty) > 0$. As in the proof of Theorem 1.2, with probability one, "if $S = \infty$, then there is some $\sigma \in \mathbf{N}^k$ such that $J_\sigma \neq \emptyset$ and $\sum_{n=1}^{\infty} T_{\sigma*n}^0 = \infty$ ". But, since $\{X_{\sigma*n}\}_{n=1}^{\infty}$ is an independent family with common distribution, the distribution of $X/(\text{diam } J)^\alpha$, and since this family is independent of $\mathcal{F}_{|\sigma|+1}$, we have with probability one "if $J_\eta \neq \emptyset$ and $\sum_{n=1}^{\infty} T_{\eta*n}^0 = \infty$, then there is some $k \in \mathbf{N}$ such that $T_{\eta*k}^0 = 1$ and $X_{\eta*k} > 0$, i.e., $X = \sum_{\sigma \in \mathbf{N}^{|\eta|}} l_\sigma^\alpha X_\sigma \geq l_{\sigma*k}^\alpha X_{\eta*k} = l_\eta^\alpha T_{\eta*k}^\alpha X_{\eta*k} > 0$ ". Thus, we may conclude with the following statement: Almost surely "if $S = \infty$, then $X > 0$ ". Q.E.D.

The next theorem is a probabilistic mixing of the deterministic methods of Rogers and Taylor [16] and Moran [15].

THEOREM 3.5. *Suppose the construction \mathbf{J} is such that there exists $\delta > 0$ so that with probability one, if $T_i(\omega) > 0$, then $T_i(\omega) > \delta$. If $\Phi(0) > 1$, then, for almost all ω , for all $\beta < \alpha$ and for all Borel $E \subseteq \mathbf{R}^m$, if $\beta - m(E) < \infty$, then $\nu_\omega(E) = 0$.*

PROOF. Since for any $\gamma \geq 0$, $\delta^m \cdot \sum_{i=1}^{\infty} T_i^\gamma \leq \sum_{i=1}^{\infty} T_i^m \leq 1$ a.s., we have $\sum_{i=1}^{\infty} T_i^\gamma \leq 1/\delta^m$ a.s. It follows that

$$(3.20) \quad E \left(\left(\sum_{i=1}^{\infty} T_i^\gamma \right)^t \right) \leq (1/\delta^m)^t < \infty.$$

Consequently, by Theorem 2.1, $E(X^t) < \infty$ for all $t \in (0, \infty)$. Fix $k > 0$, and $\beta < \alpha$; then for any $t > 0$,

$$P(l_\sigma^\alpha X_\sigma > kl_\sigma^\beta) \leq \frac{E([l_\sigma^{\alpha-\beta}]^t X_\sigma^t)}{k^t}$$

and, by independence,

$$= \frac{E(l_\sigma^{(\alpha-\beta)t})E(X^t)}{\text{diam}(J)^{\alpha t} k^t}$$

which implies

$$(3.21) \quad \sum_{\sigma \in \mathbf{N}^n} P(l_\sigma^\alpha X_\sigma > kl_\sigma^\beta) \leq \frac{E(S_{(\alpha-\beta)t,n})E(X^t)}{\text{diam}(J)^{\alpha t} k^t} = \frac{\Phi((\alpha-\beta)t)^n E(X^t)}{k^t \text{diam}(J)^{\alpha t}}.$$

Choose t_0 such that $\Phi((\alpha-\beta)t_0) < 1$. Thus,

$$(3.22) \quad \sum_{n=0}^\infty P(\exists \sigma \in \mathbf{N}^n \ni l_\sigma^\alpha X_\sigma > kl_\sigma^\beta) \leq \sum_{n=0}^\infty \frac{\Phi((\alpha-\beta)t_0)^n E(X^{t_0})}{k^{t_0} \text{diam}(J)^{\alpha t_0}} < \infty.$$

Thus by the Borel-Cantelli lemma, $P(\exists N \ni \forall n \geq N, \text{ if } \sigma \in \mathbf{N}^n, \text{ then } l_\sigma^\alpha X_\sigma \leq kl_\sigma^\beta) = 1$. Fix ω so that ν_ω is defined and for which there is $N(\omega) \in \mathbf{N}$ so that if $\sigma \in \mathbf{N}^n, n \geq N(\omega)$, then $l_\sigma^\alpha(\omega) X_\sigma(\omega) \leq kl_\sigma^\beta(\omega)$. For each $x \in K(\omega)$ and $p \in \mathbf{N}$, let $\Delta_{x,p} \equiv \{\sigma \in \mathbf{N}^* | x \in J_\sigma(\omega), l_\sigma(\omega) = \text{diam}(J_\alpha(\omega)) < 2^{-p} \text{ and } \text{diam}(J_{\sigma|\sigma|-1}(\omega)) \geq 2^{-p}\}$.

Suppose E is a compact subset of \mathbf{R}^m with $\beta - m(E) < \infty$. Suppose L_1, L_2, \dots is a cover of E by closed m -spheres so that $\text{diam}(L_i) < \varepsilon/2$ for each i , where $\varepsilon = 1 \wedge \min\{l_\sigma(\omega) > 0 | \sigma \in \mathbf{N}^{N(\omega)}\}$. For each i , there exists a $p_i \in \mathbf{N}$ so that $2^{-1-p_i} \leq \text{diam}(L_i) < 2^{-p_i}$. For each i , consider $D_i = \bigcup \{\Delta_{x,p_i} | x \in L_i \cap E \cap K(\omega)\}$. Now, D_i is an antichain in the lattice \mathbf{N}^* with the natural partial order: $\sigma \leq \eta$ iff $\exists \xi \ni \sigma * \xi = \eta$. Thus, the set $\{J_\sigma(\omega) : \sigma \in D_i\}$ is a nonoverlapping collection. For any $y \in L_i, \bigcup_{\sigma \in D_i} J_\sigma \subseteq B(y, 2^{1-p_i})$. By the definition of Δ_{x,p_i} , if $\sigma \in \Delta_{x,p_i}$, then $\text{diam}(J_\sigma(\omega)) \geq \delta 2^{-p_i}$, implying

$$\lambda(\text{Int } J_\sigma(\omega)) \geq \left[\frac{\delta \cdot 2^{-p_i}}{\text{diam}(J)} \right]^m \lambda(\text{Int } J).$$

Let N_i denote the cardinality of D_i ; then

$$N_i \left[\frac{\delta \cdot 2^{-p_i}}{\text{diam}(J)} \right]^m \lambda(\text{int}(J)) \leq \sum_{\sigma \in D_i} \lambda(\text{Int } J_\sigma(\omega)) \leq \lambda(B(y, 2^{1-p_i})) = 2^{-p_i m} \lambda(B(0, 2)),$$

i.e.,

$$(3.23) \quad N_i \leq \left[\frac{\text{diam}(J)}{\delta} \right]^m \frac{\lambda(B(0, 2))}{\lambda(\text{Int}(J))} \equiv M < \infty.$$

For any $\sigma \in D_i$, $l_\sigma^\alpha(\omega)X_\sigma(\omega) \leq kl_\sigma^\beta \leq k[2^{-p_i}]^\beta \leq 2^\beta k \text{diam}(L_i)^\beta$, implying,

$$(3.24) \quad \sum_{\sigma \in D_i} l_\sigma^\alpha(\omega)X_\sigma(\omega) \leq 2^\beta kM \text{diam}(L_i)^\beta.$$

For each i , let $n(i)$ denote $\max\{|\sigma|: \sigma \in D_i\}$. Then

$$(3.25) \quad \nu_\omega(E \cap L_i) \leq \sum_{\substack{\sigma \in \mathbf{N}^{n(i)} \\ E \cap L_i \cap J_\sigma \neq \emptyset}} l_\sigma^\alpha(\omega)X_\sigma(\omega) \leq \sum_{\sigma \in D_i} l_\sigma^\alpha(\omega)X_\sigma(\omega),$$

by Theorem 3.2 and the fact $l_\sigma^\alpha(\omega)X_\sigma(\omega) = \sum_{n=1}^\infty l_{\sigma^{**n}}^\alpha(\omega)X_{\sigma^{**n}}(\omega)$. Combining (3.24) and (3.25) we find $\nu_\omega(E) \leq 2^\beta kM \sum_i \text{diam}(L_i)^\beta$. Therefore, $\nu_\omega(E) \leq 2^\beta kM\beta - m(E) \rightarrow 0$ as $k \rightarrow 0$. Q.E.D.

THEOREM 3.6. *Suppose $\Phi(0) > 1$. Then $P(K \text{ has Hausdorff dim } \geq \alpha|K \neq \emptyset) = 1$.*

PROOF. For each $n \in \mathbf{N}$, we define an auxiliary construction $\mathbf{J}_n = \{J_{n;\sigma} | \sigma \in \mathbf{N}^*\}$ by

$$J_{n;\sigma} = \begin{cases} J_\sigma, & \text{if } l_{|\sigma|k} \geq (1/n)l_{|\sigma|k-1} \text{ for } k = 1, \dots, |\sigma|, \\ \emptyset, & \text{otherwise,} \end{cases}$$

and $T_{n;\sigma} = 1_{[1/n, \infty)}(T_\sigma) \cdot T_\sigma$.

Now,

$$(3.26) \quad \Phi_n(\beta) = E \left(\sum_{p=1}^\infty 1_{[1/n, \infty)}(T_\sigma) T_\sigma^\beta \right) \uparrow \Phi(\beta),$$

by the monotone convergence theorem. Thus, there is some $N_0 \in \mathbf{N}$ so that for all $n \geq N_0$, $\Phi_n(0) > 1$.

Let $K_n(\omega) = \bigcap_{p=1}^\infty \bigcup_{\sigma \in \mathbf{N}^p} J_{n;\sigma}(\omega)$. Obviously,

$$(3.27) \quad K_n(\omega) \subseteq K_{n+1}(\omega) \subseteq K(\omega)$$

for all n . (For later use, note $S_{n;\gamma,k} = \sum_{\sigma \in \mathbf{N}^k} l_{n;\sigma}^\gamma$.)

For each n , let $\gamma_n = P(K_n \text{ is empty})$ and let $\gamma_0 = P(K \text{ is empty})$. For each $n, p \in \mathbf{N}$, let $C_{n;p}$ denote $P(\sum_{q=1}^\infty T_{n;q}^0 = p)$, and $C_{0;p}$ denote $P(\sum_{q=1}^\infty T_q^0 = p)$. By a well-known formula (see [1]):

$$(3.28) \quad \gamma_p = \sum_{q=0}^\infty C_{p;q} \gamma_p^q.$$

(This formula is correct even if $P(\sum T_n^0 = \infty) > 0$.)

Set $\psi_p(x) = -x + \sum_{q=0}^\infty C_{p;q} x^q$. Clearly, ψ_p is defined on $[0, 1]$ for each p . Also, for $p \geq N_0$ or $p = 0$, γ_p is the unique root of ψ_p in $[0, 1]$ (see [1, p. 4]). Moreover, ψ_p is strictly convex for $p \geq N_0$ and ψ_0 is either strictly convex or linear with negative slope. By (3.27), $\gamma_p \geq \gamma_{p+1} \geq \gamma_0$ for $p \in \mathbf{N}$. Thus, $\gamma_\infty = \lim_{p \rightarrow \infty} \gamma_p$ exists and $\gamma_\infty \geq \gamma_0$. Now $C_{p;q} \rightarrow C_{0;q}$ as $p \rightarrow \infty$ if $q < \infty$. Easily, $\{\psi_n\}_{n=1}^\infty$ converges uniformly to ψ_0 on $[0, \gamma_{N_0}]$. Thus, $0 = \psi_n(\gamma_n) \rightarrow \psi_0(\gamma_\infty)$, i.e., $\psi_0(\gamma_\infty) = 0$. This implies $\gamma_\infty = \gamma_0$ or

$$(3.29) \quad \lim_{n \rightarrow \infty} P(K_n(\omega) = \emptyset) = P(K(\omega) = \emptyset).$$

Now (3.29) together with (3.27) imply:

$$(3.30) \quad \text{For almost all } \omega, "K(\omega) \neq \emptyset \text{ iff for some } n \geq N, K_n(\omega) \neq \emptyset".$$

For each n , let α_n be the “ α ” for the construction \mathbf{J}_n . Note that since $T_{n;i} \leq 1$, for all i and $\beta > 0$,

$$(3.31) \quad \sum_{k=1}^{\infty} T_{n;k}^{\beta} \leq \sum_{k=1}^{\infty} T_{n;k}^0.$$

Since $T_{n;k} \notin (0, 1/n)$, we have according to (1.5)

$$(3.32) \quad (1/n)^m \sum_{k=1}^{\infty} T_{n;k}^0 \leq \sum_{k=1}^{\infty} T_{n;k}^m \leq 1.$$

Thus, for each n , $\Phi_n(0) \leq n^m$. So, for all $n \geq N_0$, Φ_n is finite valued, continuous, strictly decreasing, and $\Phi_n(\alpha_n) = 1$. Also, $\Phi_n \leq \Phi_{n+1} \leq \Phi$ for each n . Thus, $\alpha_n \leq \alpha_{n+1} \leq \alpha$ for all $n \geq N_0$. This implies $\alpha_{\infty} = \lim_{n \rightarrow \infty} \alpha_n \leq \alpha$. But,

$$1 = \Phi_n(\alpha_n) \geq \Phi_n(\alpha_{\infty}) = E \left(\sum_{p=1}^{\infty} T_p^{\alpha_{\infty}} 1_{[1/n, \infty)}(T_p) \right) \uparrow \Phi(\alpha_{\infty})$$

as $n \rightarrow \infty$. Therefore, $1 \geq \Phi(\alpha_{\infty})$ which implies $\alpha_{\infty} \geq \alpha$, i.e., $\alpha = \alpha_{\infty}$.

We derive from (3.31) and (3.32)

$$(3.33) \quad E \left(\left(\sum_{k=1}^{\infty} T_{n;k}^{\alpha_n} \right)^2 \right) \leq n^{2m}.$$

Thus, by Theorem 2.1, $\{S_{n;\alpha_n,k}\}_{k=1}^{\infty}$ is an L_2 -bounded martingale. This implies $E(X_n) = E(S_{n;\alpha_n,0}) = (\lambda(J))^{\alpha_n} > 0$.

For each $n \geq N_0$, \mathbf{J}_n satisfies the hypothesis of Theorem 3.5. Suppose $\beta < \alpha$ and $K(\omega) \neq \emptyset$. Then almost surely there exists an $n \geq N_0$ so that $\alpha_n > \beta$ and $K_n(\omega) \neq \emptyset$. By definition of $\nu_{n;\omega}$ and Theorem 3.4, $\nu_{n;\omega}(K(\omega)) = \nu_{n;\omega}(K_n(\omega)) = X_n(\omega) > 0$. Thus, by Theorem 3.5, almost surely $\beta - m(K(\omega)) = \infty$ if $K(\omega) \neq \emptyset$. Q.E.D.

In what has preceded, we have required of our construction only a “stochastic ratio self-similarity”. In order to ask a question of interest, we will now introduce a version of a construction being “stochastically geometrically self-similar”.

We let G denote the set of geometric similarity maps with domain J . In other words,

$$G = \{f: J \rightarrow \mathbf{R}^n | \exists \lambda \in (0, \infty) \text{ so that for all } x, y \in J, \|f(x) - f(y)\| = \lambda \|x - y\|\},$$

where $\|\cdot\|$ is the Euclidean norm in \mathbf{R}^m . Also, let \mathcal{J}_p be the σ -algebra of subsets of Ω generated by $\{J_{\sigma} | |\sigma| \leq p\}$.

Note 3.7. Suppose $\sigma \in \mathbf{N}^p$, $P(J_{\sigma} \neq \emptyset) > 0$ and $F_{\sigma}: \Omega \times J \rightarrow \mathbf{R}^m$ satisfies:

- (1) F_{σ} is $\mathcal{J}_p \times \mathcal{B}(J)$ -measurable.
- (2) $F_{\sigma}(\omega, J) = J_{\sigma}(\omega)$, if $J_{\sigma}(\omega) \neq \emptyset$ for almost all ω .
- (3) $F_{\sigma} \in G$ a.s.

Define $\tilde{\mathbf{J}}_{\sigma} = \{\tilde{J}_{\sigma;\eta} | \eta \in \mathbf{N}^*\}$ by $\tilde{J}_{\sigma;\eta}(\omega) = [F_{\sigma}(\omega, \cdot)]^{-1}(J_{\sigma;\eta}(\omega))$. Then given $J_{\sigma} \neq \emptyset$, we have $\tilde{\mathbf{J}}_{\sigma}$ is a construction (in fact $\tilde{\tau}_{\sigma;\emptyset}$ is equal in distribution to τ_{\emptyset}). We

say a construction \mathbf{J} is *stochastically geometrically self-similar* if for each $\sigma \in \mathbf{N}^*$ with $P(J_\sigma \neq \emptyset) > 0$, there is a F_σ satisfying (1)–(3) of Note 3.7 so that $\tilde{\mathbf{J}}_\sigma$ given $\tau_\sigma \neq \emptyset$ is distributed the same as J , i.e., if B is a Borel subset of $[2^J]^{\mathbf{N}^*}$, then $P(\tilde{\mathbf{J}}_\sigma \in B | J_\sigma \neq \emptyset \text{ and } \mathcal{F}_n) = P(\mathbf{J} \in B)$. All the examples given in §4 have this property.

Question 3.8. Suppose \mathbf{J} is a stochastically geometrically self-similar construction indexed by an n -ary $\{1, \dots, n\}^*$, $n < \infty$, and $P(T_i > 0; i = 1, \dots, n) = 1$ and $P(J_i \cap J_k \neq \emptyset \text{ if } (i \neq j)) = 0$. Then is it true for almost all ω that

$$0 < \alpha - m(K(\omega))?$$

For the deterministic case, this was proved by Moran [15]. At least we know the following facts in this case. Let $a = E(\alpha - m(K))$. Since our construction is geometrically self-similar, for $|\sigma| = k$, we have

$$a = E(\alpha - m(\tilde{K}_\sigma) | J_k \text{ and } J_\sigma \neq \emptyset).$$

Also, since for these constructions $J_\sigma \neq \emptyset$ a.s., we have

$$a = E(\alpha - m(\tilde{K}_\sigma) | J_k).$$

We now claim

$$\alpha - m(K) = \frac{aX}{(\text{diam}(J))^\alpha}.$$

To see this, note

$$\alpha - m(K) = \sum_{\sigma \in \{1, \dots, n\}^k} \alpha - m(K \cap J_\sigma) = \sum_{\sigma \in \{1, \dots, n\}^k} \alpha - m(K_\sigma),$$

where K_σ is the set obtained by pruning the tree to start at σ with J_σ . For $\sigma \in \{1, \dots, n\}^k$, let F_σ be a random geometric self-similarity map,

$$\alpha - m(K(\omega)) = \frac{1}{[\text{diam}(J)]^\alpha} \sum_{\sigma \in \{1, \dots, n\}^k} l_\sigma^\alpha \alpha - m(F_\sigma(\omega, \cdot))^{-1}(K_\sigma(\omega)).$$

This implies

$$\begin{aligned} E(\alpha - m(K) | J_k) &= \frac{1}{[\text{diam}(J)]^\alpha} \sum_{\sigma \in \{1, \dots, n\}^k} l_\sigma^\alpha E[\alpha - m(\tilde{K}_\sigma) | J_k] \\ &= \frac{a}{[\text{diam}(J)]^\alpha} S_{\alpha, k}. \end{aligned}$$

Now, letting $k \rightarrow \infty$, we find

$$\alpha - m(K) = \frac{aX}{[\text{diam}(J)]^\alpha}.$$

Using Theorem 2.1, it is easy to see $\{S_{\alpha, n}\}_{n=1}^\infty$ is a L^2 -bounded martingale and $E(X) > 0$. Thus, by Theorem 3.4, we see $X > 0$ a.s., so either $\alpha - m(K) > 0$ a.s. or $\alpha - m(K) = 0$ a.s.

4. Examples.

EXAMPLE 4.1. Let us note that the following simple deterministic process shows that $X(\omega)$ may not yield $\alpha - m(K(\omega))$. Set $J_\emptyset = [0, 1]$, $J_0 = [\frac{1}{8}, \frac{3}{8}]$ and $J_1 = [\frac{5}{8}, \frac{7}{8}]$. Given $J_\sigma = [a, b]$, set

$$J_{\sigma*0} = [a + (b - a)/8, a + 3(b - a)/8]$$

and

$$J_{\sigma*1} = [a + 5(b - a)/8, a + 7(b - a)/8].$$

The Hausdorff dimension of the final Cantor set is $\alpha = \frac{1}{2}$. In this construction $X(\omega) = 1$ a.s., whereas it can be shown that $\frac{1}{2} - m(K(\omega)) = \sqrt{\frac{2}{3}}$.

EXAMPLE 4.2. We return to the example described in the introduction of a random Cantor subset of $[0, 1]$. It follows from Theorem 1.1 that the Hausdorff dimension of such a set in this construction is the number α such that

$$1 = E(T_1^\alpha + T_2^\alpha) = 2 \int_0^1 \int_x^1 x^\alpha + (1 + y)^\alpha dy dx.$$

In this case $\alpha = (\sqrt{17} - 3)/2$.

EXAMPLE 4.3. Another method of constructing a Cantor subset of $[0, 1]$ may be described as follows.

Choose a number u from $[0, 1]$ with respect to the uniform distribution, then choose x from $[0, u]$ with respect to the uniform distribution on $[0, u]$ and independently choose a number y from $[u, 1]$ with respect to the uniform distribution on $[u, 1]$. Set $J_0 = [0, x]$ and $J_1 = [y, 1]$. Continue this process by rescaling to each interval already determined. Again, according to Theorem 1.1, with probability one, we obtain a Cantor set with Hausdorff dimension $\alpha = \sqrt{2} - 1$, where

$$1 = \Phi(\alpha) = \int_0^1 \left[\frac{1}{u} \int_0^u x^\alpha dx + \frac{1}{1 - u} \int_u^1 (1 - y)^\alpha dy \right] du.$$

This example arose naturally in the course of our study of random homeomorphisms of $[0, 1]$ [6]. In that paper, a measure P is constructed on H , the space of orientation preserving homeomorphisms of $[0, 1]$, which has the property (Theorem 4.16) of [8] that if B is a Borel subset of $[0, 1]$, then

$$(*) \quad \lambda(B) = \int_H \lambda(h(B)) dP(h).$$

In particular, if B is the Cantor subset of $[0, 1]$ constructed by repeatedly removing the middle half interval, then $\lambda(B)$, the Lebesgue measure of B , is zero. Therefore, according to (*) for P -a.e. h , $\lambda(h(B)) = 0$. Example 4.3 shows that for P -a.e. h , the Hausdorff dimension of the Cantor set $h(B)$ is $\sqrt{2} - 1$. The measure P was first studied by Dubins and Freedman [5].

EXAMPLE 4.4. Choose x from $[0, 1]$ according to the uniform distribution and then choose y from $[x, 1]$ according to the uniform distribution on $[x, 1]$. Set $J_0 = [0, x]$ and $J_1 = [y, 1]$. Continue this procedure by rescaling to each of the intervals already obtained. With probability one, we obtain a Cantor set with Hausdorff dimension α , where

$$1 = E(T_1^\alpha + T_2^\alpha) = \int_0^1 \left[x^\alpha + \frac{1}{1 - x} \int_x^1 (1 - y)^\alpha dy \right] dx.$$

In this case $\alpha = (\sqrt{5} - 1)/2$.

EXAMPLE 4.5. We give a construction to show that the final Cantor set may have the same dimension as the dimension of J_\emptyset : Choose the positive integer n with probability 2^{-n} . Then partition the interval $J_\emptyset = [0, 1]$ into 2^{n^2+n} intervals of equal length, set $J_i = [2i/2^{n^2+n}, (2i + 1)/2^{n^2+n}]$, $i = 0, \dots, 2^{n^2+n} - 1$. Continue this construction. We have

$$\Phi(\beta) = \sum_{n=1}^{\infty} 2^{-n} \left[\frac{1}{2^{n^2+n}} \right]^\beta 2^{n^2+n-1}.$$

Thus, $\Phi(\beta) = +\infty$ if $\beta < 1$ and $\Phi(1) = \frac{1}{2}$.

EXAMPLE 4.6. We construct at random a Cantor subset of $[0, 1] \times [0, 1]$ with Hausdorff dimension 1. Choose four numbers $\mu_1, \mu_2, \mu_3, \mu_4$ independently from $[0, 1]$. Let $x_1 = \min(\mu_1, \mu_2)$, $x_2 = \max(\mu_1, \mu_2)$, $x_3 = \min(\mu_3, \mu_4)$ and $x_4 = \max(\mu_3, \mu_4)$. Let $J_1 = [0, s_1] \times [0, s_1]$, where $s_1 = \min(x_1, x_3)$, let $J_2 = [1 - s_2, 1] \times [0, s_2]$, where $s_2 = \min(x_3, 1 - x_2)$. Let J_3 be the largest square with one vertex at $(1, 1)$ which lies in the rectangle $[x_2, 1] \times [x_4, 1]$. Similarly, let J_4 be the largest square with one vertex at $(0, 1)$ which lies in the rectangle $[0, x_1] \times [x_4, 1]$. Finally, consider the rectangle $[x_1, x_2] \times [x_3, x_4]$ and its center $((x_1 + x_2)/2, (x_3 + x_4)/2)$. Let J_5 be the largest square lying in this rectangle and having the same center. Iterate this process. According to our results, with probability one the final object is a Cantor subset of $[0, 1] \times [0, 1]$ with Hausdorff dimension α where

$$1 = \Phi(\alpha) = 5 \iiint\limits_{[0,1]^4} (\mu_1 \wedge \mu_2 \wedge \mu_3 \wedge \mu_4)^\alpha d\mu_1 d\mu_2 d\mu_3 d\mu_4.$$

One can check


$$1 = \Phi(\alpha) = 5 \left[4! \int_0^1 \int_0^{\mu_4} \int_0^{\mu_3} \int_0^{\mu_2} \mu_1^\alpha d\mu_1 d\mu_2 d\mu_3 d\mu_4 \right],$$

or

$$1 = \Phi(\alpha) = 5! / (\alpha + 1)(\alpha + 2)(\alpha + 3)(\alpha + 4).$$

Obviously, $\alpha = 1$.

EXAMPLE 4.7 (SIERPIŃSKI UNIVERSAL CURVE). Set $J_\emptyset = [0, 1] \times [0, 1]$. Choose x_1, x_2, x_3 , and x_4 from $(\frac{1}{3}, \frac{1}{2})$ independently and each according to the uniform distribution. Set $J_1 = [0, x_1] \times [0, x_1]$, $J_2 = [1 - x_2, 1] \times [0, x_2]$, $J_3 = [0, x_3] \times [1 - x_3, 1]$ and $J_4 = [1 - x_4, 1] \times [1 - x_4, 1]$. Since $\frac{1}{3} < x_1, x_2 < \frac{1}{2}$, we have $1 - (x_1 + x_2) < \min(x_1, x_2)$. Set $J_5 = [x_1, 1 - x_2] \times [0, 1 - (x_1 + x_2)]$. Thus, J_5 is a square having one side a subset of a side of J_1 and one side a subset of J_2 . Similarly, let J_6 be the square having one side in common with J_1 , one side in common with J_3 and one side on the Y -axis. J_7 is a square similarly placed in relation to J_2 and J_4 , and J_8 is similarly placed in relation to J_3 and J_4 . Thus, the first stage in this construction is analogous to the first stage in the construction of Sierpiński's carpet [17]. Of course, in Sierpiński's construction there is no random element, one simply partitions the unit square into nine congruent subsquares and deletes the middle square. Our construction is more like a Swiss flag:



Now, one iterates this process. In order to see that the final set is a topological Sierpiński curve, we use Whyburn’s topological characterization [18]. It is easily seen that our final set is locally connected (since for each $\varepsilon > 0$, it can be expressed as the union of finitely many subcontinua of diameter $< \varepsilon$) and its complement with respect to S^2 , the two-sphere, is the union of countably many open disks D_1, D_2, D_3, \dots such that for each i and j , the boundary of D_i is a simple closed curve and if $i \neq j$, $\overline{D_i} \cap \overline{D_j} = \emptyset$. These properties characterize Sierpiński’s curve.

According to Theorem 1.1, the Hausdorff dimension of the final Sierpiński curve is α , where

$$\begin{aligned} 1 = \Phi(\alpha) &= 24 \int_{1/3}^{1/2} x^\alpha dx + 144 \int_{1/3}^{1/2} \int_{1/3}^{1/2} (1 - (x + y))^\alpha dy dx \\ &= \frac{24}{\alpha + 1} \left[\frac{1}{2^{\alpha+1}} - \frac{1}{3^{\alpha+1}} + \left(\frac{2}{\alpha + 2} \right) \left[\frac{1}{3^{\alpha+1}} - \frac{1}{6^{\alpha+1}} \right] \right]. \end{aligned}$$

We know that $\Phi(1) \geq 1$ [7, p. 107] (actually $\Phi(1) = \frac{7}{3}$ and $\Phi(2) < 1$). Numerical studies show that $\alpha \cong 1.8947$.

EXAMPLE 4.8. Set $J_\emptyset = [0, 1] \times [0, 1] \times [0, 1]$. Choose $x_i, i = 1, 2, \dots, 8$, independently from the open interval $(\frac{1}{3}, \frac{1}{2})$ and each according to the uniform distribution. Let J_i be a cube lying in J with edge length x_i , with faces parallel to the coordinate planes and with one vertex at (i_1, i_2, i_3) , where $i = 1 + i_1 + i_2 2 + i_3 2^2$. Let J_9 be a cube lying in J with edge length $1 - (x_1 + x_2)$ having one face a subset of a face of J_1 , one face a subset of a face of J_2 , and one edge on the x -axis. Continue placing cubes in this manner, until there are a total of twenty cubes. Then iterate this construction inside each of these twenty cubes. The first stage of this construction is somewhat like the first stage in the construction of Menger’s universal curve [14, p. 345]. (There is a sketch of this curve in the second edition between pp. 346 and 347.) Of course, in Menger’s construction, one partitions the unit cube into 27 congruent subcubes and then one deletes the center cube and the six other cubes having a face in common with it. It is easy to see that the final set K is a locally connected continuum and that every nonempty open subset of K contains K_5 , the complete graph on five vertices and also contains $K_{3,3}$. Thus, our final object is certainly nonplanar. If we knew K were one-dimensional, then we could use R. D. Anderson’s result [2, Theorem XII] to conclude that K is a Menger curve, but we have been unable to show this.

According to Theorem 1.1, the Hausdorff dimension of the final universal curve is α , where

$$\begin{aligned} 1 = \Phi(\alpha) &= 48 \int_{1/3}^{1/2} x^\alpha dx + 432 \int_{1/3}^{1/2} \int_{1/3}^{1/2} (1 - (x + y))^\alpha dy dx \\ &= \frac{48}{\alpha + 1} \left[\frac{1}{2^{\alpha+1}} - \frac{1}{3^{\alpha+1}} + \left(\frac{3}{\alpha + 2} \right) \left[\frac{1}{3^{\alpha+1}} - \frac{1}{6^{\alpha+1}} \right] \right]. \end{aligned}$$

We have $2 > \Phi(2) > 1$ and $\Phi(3) < 1$. Numerical studies show that $\alpha \cong 2.5968$.

REMARK. Our Example 4.7, concerning Sierpiński curves, is in accordance with the category version. Mazurkiewicz showed that for almost all (in the sense of category) continuous maps f of $[0, 1]$ into $[0, 1] \times [0, 1]$, the image $f([0, 1])$ is a Sierpiński curve [13]. On the other hand, our Example 4.9, concerning Menger

curves, stands in contrast with the category version for almost all continuous maps f of $[0, 1]$ into $[0, 1]^3$, the image set is an arc [10].

EXAMPLE 4.9 (MENGER UNIVERSAL CURVE). Let $J_\emptyset = [0, 1] \times [0, 1] \times [0, 1]$. Choose x_i , $i = 1, 2, \dots, 8$, independently from the open interval $(0, \frac{1}{2})$ with common uniform distribution. Construct subcubes J_1, \dots, J_8 as in Example 4.8. Let $k(x_1, x_2)$ be the least positive integer greater than $3(1 - x_1 - x_2)/(x_1 \wedge x_2)$. Now, place $k(x_1, x_2)$ nonoverlapping subcubes of J with equal edge length, each with an edge on the x -axis and forming a chain from J_1 to J_2 . Similarly, join each pair of the original eight cubes which are adjacent. This completes the first stage in the construction. Iterate this process in each of the subcubes obtained so far. As in Example 4.8, it is easy to see that K is a locally connected continuum and every nonempty open subset of K is nonplanar. To conclude that K is a Menger curve, we first estimate the Hausdorff dimension of K . We have

$$\Phi(\beta) = 16 \int_0^{1/2} x^\beta dx + 48 \int_0^{1/2} \int_0^{1/2} k(x_1, x_2) \left(\frac{1 - x_1 - x_2}{k(x_1, x_2)} \right)^\beta dx_1 dx_2.$$

In particular,

$$\begin{aligned} \Phi(2) &\leq \frac{2}{3} + 48 \int_0^{1/2} \int_0^{1/2} \frac{(1 - x_1 - x_2)^2}{k(x_1, x_2)} dx_1 dx_2 \\ &\leq \frac{2}{3} + 48 \int_0^{1/2} \int_0^{1/2} \frac{(x_1 \wedge x_2)}{3} (1 - x_1 - x_2) dx_1 dx_2 \\ &\leq \frac{2}{3} + 16 \left[\int_0^{1/2} \int_0^{x_2} x_1 (1 - x_1 - x_2) dx_1 dx_2 \right. \\ &\quad \left. + \int_0^{1/2} \int_{x_2}^{1/2} x_2 (1 - x_1 - x_2) dx_1 dx_2 \right] \\ &\leq \frac{2}{3} + 32 \int_0^{1/2} \int_0^{x_2} x_1 - x_1^2 - x_2 x_1 dx_1 dx_2. \end{aligned}$$

So, $\Phi(2) \leq \frac{11}{12}$.

Since Φ is finite valued, we have that $\dim_H(K) = \alpha < 2$. This means that the topological dimension of K is ≤ 1 [9, p. 8]. Thus, with probability one, K has dimension one and according to Anderson's theorem, K is a Menger curve.

REFERENCES

1. K. B. Athreya and P. E. Ney, *Branching processes*, Springer-Verlag, Berlin, 1972.
2. R. D. Anderson, *A characterization of the universal curve and a proof of its homogeneity*, Ann. of Math. (2) **67** (1958), 313-324.
3. —, *One-dimensional continuous curves and a homogeneity theorem*, Ann. of Math. (2) **68** (1958), 1-16.
4. J. W. Cannon, *Book review*, Amer. Math. Monthly **91** (1984), 594-598.
5. L. E. Dubins and D. A. Freedman, *Random distribution functions*, Proc. Fifth Berkeley Sympos. on Math. Statistics and Probability (L. M. LeCam and J. Neyman, eds.), Univ. of California Press, Berkeley, Calif., 1967, pp. 183-214.
6. P. Doubilet, *On the foundations of combinatorial theory. VIII: Symmetric functions through the theory of distribution and occupancy*, Stud. Appl. Math. **51** (1972), 377-396.
7. K. J. Falconer, *The geometry of fractal sets*, Cambridge Tracts in Math., vol. 85, Cambridge Univ. Press, Cambridge, 1985.

8. S. Graf, R. D. Mauldin and S. C. Williams, *Random homeomorphisms*, Adv. in Math. (to appear).
9. W. Hurewicz and H. Wallman, *Dimension theory*, Princeton Math. Series, vol. 4, Princeton Univ. Press, Princeton, N.J., 1941.
10. V. Jarník, *Über die stetigen Abbildungen der Strecke*, Monatsh. Math. Phys. **41** (1934), 408–428.
11. B. B. Mandelbrot, *The fractal geometry of nature*, Freeman, San Francisco, Calif., 1982.
12. ———, *Fractals: form, chance and dimension*, Freeman, San Francisco, Calif., 1977.
13. S. Mazurkiewicz, *Über die stetigen Abbildungen der Strecke*, Fund. Math. **25** (1935), 253–260.
14. K. Menger, *Kurventheorie*, Chelsea, New York, 1967.
15. P. A. P. Moran, *Additive functions of intervals and Hausdorff measure*, Proc. Cambridge Philos. Soc. **42** (1946), 15–23.
16. C. A. Rogers and S. J. Taylor, *Functions continuous and singular with respect to a Hausdorff measure*, Mathematika **8** (1961), 1–31.
17. W. Sierpiński, *Sur une courbe cantorienne qui contient une image biunivoque et continue de toute courbe donnée*, C. R. Acad. Sci. Paris **162** (1916), 626–631.
18. G. T. Whyburn, *Topological characterization of the Sierpiński curve*, Fund. Math. **45** (1958), 320–324.
19. U. Zähle, *Random fractals generated by random cutouts*, Math. Nachr. **116** (1984), 27–52.

DEPARTMENT OF MATHEMATICS, NORTH TEXAS STATE UNIVERSITY, DENTON, TEXAS
76203-5116

DEPARTMENT OF MATHEMATICS, UTAH STATE UNIVERSITY, LOGAN, UTAH 84322