

BOUNDARY BEHAVIOR OF A NONPARAMETRIC SURFACE OF PRESCRIBED MEAN CURVATURE NEAR A REENTRANT CORNER

ALAN R. ELCRAT AND KIRK E. LANCASTER

ABSTRACT. Let Ω be an open set in \mathbf{R}^2 which is locally convex at each point of its boundary except one, say $(0,0)$. Under certain mild assumptions, the solution of a prescribed mean curvature equation on Ω behaves as follows: All radial limits of the solution from directions in Ω exist at $(0,0)$, these limits are not identical, and the limits from a certain half-space (H) are identical. In particular, the restriction of the solution to $\Omega \cap H$ is the solution of an appropriate Dirichlet problem.

0. Introduction. We consider here the behavior of a generalized solution of the equation for surfaces of prescribed mean curvature at an inner corner of the boundary where the solution is discontinuous. This work is a generalization of the previous work of the second author [8], which dealt with the minimal surface equation. It was shown there that all radial limits exist and that they are constant in directions coming from a half-space. Here we find that the same result holds for (nonparametric) surfaces of prescribed mean curvature.

1. Preliminaries. Let Ω be a bounded, open, connected, simply connected subset of \mathbf{R}^2 with $N = (0,0) \in \partial\Omega$ such that Ω is locally convex at each point of $\partial\Omega \setminus \{N\}$. Let $H(x, y, t)$ be a continuous function on $\Omega \times \mathbf{R}$ and $\phi \in C^0(\partial\Omega)$. We will make a number of assumptions which will hold throughout this work.

ASSUMPTIONS. (A) The equation

$$(1) \quad (p/W)_x + (q/W)_y = 2H(x, y, z(x, y))$$

has a solution $z = f \in C^2(\Omega)$, where $p = z_x$, $q = z_y$, and $W^2 = 1 + p^2 + q^2$.

(B) $f \in C^0(\bar{\Omega} \setminus \{N\})$ and $f = \phi$ on $\partial\Omega \setminus \{N\}$.

(C) $F \notin C^0(\bar{\Omega})$.

(D) There exists $Q > 0$ such that $|H(x, y, f(x, y))| \leq Q$ for all $(x, y) \in \Omega$.

(E) The area of the graph of f over Ω is finite.

A function f satisfying the above conditions can be obtained as a generalized solution of the Dirichlet problem for (1) in Ω with data ϕ .

One method for obtaining such solutions is to minimize the functional

$$\iint_{\Omega} (1 + f_x + f_y)^{1/2} dx dy + \iint_{\Omega} \int_0^f 2H(x, y, t) dt dx dy + \int_{\partial\Omega} |f - \phi|$$

Received by the editors September 2, 1984.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 35J67, 35J65; Secondary 35J20.

in $BV(\Omega)$. Suppose that Ω has a locally Lipschitz boundary, H is Lipschitz and nondecreasing in t , and $H_0(x, y) = 2H(x, y, 0)$ satisfies

$$(2) \quad \left| \iint_A H_0(x, y) \, dx \, dy \right| \leq (1 - \varepsilon) \iint_A |D\chi_A|$$

for all Caccioppoli sets $A \subset \Omega$, where $0 < \varepsilon < 1$ is fixed and χ_A is the characteristic function of A . Then it is shown in [4] that there is a solution of (1) in Ω satisfying (E) (see also [5, Theorem 1.1]). Further, if $\partial\Omega \setminus \{N\}$ is smooth and its curvature $\kappa(x, y)$ satisfies

$$|2H(x, y, f(x, y))| \leq \kappa(x, y)$$

at each point, then (B) also holds.

On the other hand, if $H = H(x, y)$ and H has a bounded gradient, then the Perron method can be used as was done in [9] for H constant. (The essential ingredient necessary for the extension of Serrin’s work to variable H is an interior gradient estimate and this is available in [7 and 10].) In order to use Perron’s method, we need a bounded supersolution so that we know that the upper Perron class is nonempty. (A bounded subsolution would do just as well.) The results in [5] enable us to give sharp conditions on when such a supersolution can be found and also show that the two methods are closely related. If $H = H(x, y)$ is Lipschitz and Ω has a locally Lipschitz boundary, then Giusti shows that a necessary and sufficient condition for the existence of a solution of (1) in Ω is

$$\left| 2 \iint_A H(x, y) \, dx \, dy \right| < \iint_A |D\chi_A|$$

for all Caccioppoli sets A strictly contained in Ω with positive measure. If the strict inequality holds when $A = \Omega$, then it is shown that (2) holds. (In the boundary case $|2 \int \int_{\Omega} H| = \int \int_{\Omega} |D\chi_{\Omega}| = H_1(\partial\Omega)$, the unique (up to a constant) solution of (1) is bounded iff $\kappa(x, y) < 2H(x, y)$.) We see then that in many cases the Perron method can be applied if and only if the variational method can.

Finally, if $H = H(x, y)$, using results from [3, §15], we can give domains Ω and boundary data ϕ such that (C) must hold. In particular, if N is an inner boundary point of Ω [3], then a bound for the value of $z(N)$, $z \in C^0(\bar{\Omega})$, satisfying (1) can be given which depends only on values of z at points of $\partial\Omega$ bounded away from N .

2. A parametric representation of $z = f(x, y)$. Define

$$S = \{(x, y, f(x, y)) \mid (x, y) \in \bar{\Omega} \setminus \{N\}\},$$

$$\Gamma = \{(x, y, \phi(x, y)) \mid (x, y) \in \partial\Omega\},$$

and $S_0 = S \setminus \Gamma$. Let $T = \{(0, 0, z) \mid z \text{ is real}\}$ be the z -axis. We need some parameter domains, so set $E = \{(u, v) \mid u^2 + v^2 < 1\}$, $B = \{(u, v) \in E \mid v > 0\}$, $\partial' B = \{(u, v) \mid u^2 + v^2 = 1, v > 0\}$, $\partial'' B = \{(u, 0) \mid -1 < u < 1\}$, and $B' = B \cup \partial' B$. Let $P = (0, 0, \phi(0, 0))$ and $\Gamma_0 = \Gamma \setminus \{P\}$.

LEMMA 2.1. *There is a vector $X \in C^0(B' : R^3) \cap C^2(B : R^3)$ with the following properties:*

- (i) X is a homeomorphism of B onto S_0 .
- (ii) X maps $\partial' B$ strictly monotonically onto Γ_0 .
- (iii) X is conformal on B , i.e. $X_u \circ X_v = 0$, $X_u^2 = X_v^2$ on B .

- (iv) $X_{uu} + X_{vv} = 2H(X)X_u \times X_v$ on B .
- (v) If we write $X(u, v) = (x(u, v), y(u, v), z(u, v))$, then $x, y \in C^0(\bar{B})$ and $x(u, 0) = y(u, 0) = 0$ for $-1 \leq u \leq 1$.
- (vi) As $0 \leq \theta \leq \pi$ increases, $K(\cos(\theta), \sin(\theta))$ moves in a clockwise direction about $\partial\Omega$. Here $K(u, v) = (x(u, v), y(u, v))$.

PROOF. At each point Q of S_0 , there is a neighborhood U of Q in S_0 and a vector $Y \in C^2(E : \mathbf{R}^3)$ such that Y is a homeomorphism from E onto U , Y is conformal on E , and $\Delta Y = 2H(Y)Y_u \times Y_v$. Let us pick the vectors $\{Y\}$ so $\{Y_u \times Y_v\}$ gives a consistent orientation to S_0 . We may now regard S_0 as a Riemann surface with local uniformizing parameters $\{Y^{-1}\}$. By the uniformization theorem for simply connected Riemann surfaces, there is a global uniformizing parameter Φ mapping E onto S_0 . This means that Φ is a homeomorphism of E onto S_0 and that $Y^{-1} \circ \Phi$ is analytic for each local uniformizing parameter Y^{-1} when we regard the domains of Φ and Y as being subsets of the complex plane. Since $\Phi = Y \circ (Y^{-1} \circ \Phi)$ on an open subset of E , we see that Φ is conformal on E and that $\Delta\Phi = 2H(\Phi)\Phi_u \times \Phi_v$.

Set $\Phi(u, v) = (a(u, v), b(u, v), c(u, v))$ and consider the map $G(u, v) = (a(u, v), b(u, v))$. Notice that $\iint_E G_u^2 + G_v^2 \, du \, dv \leq 2 \iint_E |\Phi_u|^2 \, du \, dv = 2A(S) < \infty$, where $A(S)$ is the area of S . If we apply the proof of Theorem 2.4 of [1] to G , we see that $G \in C^0(\bar{E} : \mathbf{R}^2)$. If we define σ as the subset of ∂E which G maps onto $\partial\Omega \setminus \{N\}$, then $\Phi \in C^0(E \cup \sigma)$, since $f \in C^0(\bar{\Omega} \setminus \{N\})$.

Let $w_0 \in \partial\Omega$ and set $\Omega(\varepsilon) = \{w \in \Omega \mid |w - w_0| < \varepsilon\}$ and $E(\varepsilon) = G^{-1}(\Omega(\varepsilon))$ for all $\varepsilon > 0$. Since $G^{-1}(x, y) = \Phi^{-1}(x, y, f(x, y))$, G is a homeomorphism of E and Ω . For $\varepsilon > 0$ small enough, the open sets $\Omega(\varepsilon)$ and $\Omega \setminus \Omega(\varepsilon)$ are connected and simply connected and so $E(\varepsilon)$ and $E \setminus \overline{E(\varepsilon)}$ are connected and simply connected. Thus $\phi \neq \partial E \cap \overline{E(\varepsilon)}$ is a connected arc of ∂E . Since $\{E(\varepsilon)\}$ is a nested collection, $\overline{E(\varepsilon)}$ converges to a closed, connected arc $\tau(w_0) \subset \partial E$ as $\varepsilon \rightarrow 0$. Notice that $\tau(w_0) = G^{-1}(w_0)$ by construction and so G is weakly monotonic on ∂E . In particular σ is connected and Φ maps σ weakly monotonically on Γ_0 . Now if we use [6], we see that Φ maps σ strictly monotonically onto Γ_0 .

If $\partial E \setminus \sigma$ were a single point, we could use the proof of Lemma 2.2 to show that $\Phi \in C^0(\bar{E})$ and so $f \in C^0(\bar{\Omega})$, in contradiction to (C). To finish, we need only compose Φ with a suitable conformal map of B onto E and obtain X . Q.E.D.

The proof of the next lemma is a (minor) modification of the proof of Lemma 2.2 of [8]. Our proof is self-contained for the sake of clarity.

LEMMA 2.2. $X \in C^0(\bar{B})$.

PROOF. We need prove only that $z \in C^0(\bar{B})$. We will prove that z is uniformly continuous on B and so extends to a function in $C^0(\bar{B})$.

Let $\varepsilon > 0$. Define $g(x, y)$ as the function whose graph is the upper half-sphere of radius $1/2Q$ centered at $(0, 0, 0)$. Pick $d > 0$ so that $g(0, 0) - g(2d, 0) < \varepsilon/4$. For some $0 < \tau < \min(d, \varepsilon/4)$, the diameter of the shortest arc on $\Gamma \cup T$ joining two points on $\Gamma \cup T$ is less than $\min(d, \varepsilon/4)$ whenever the distance between the points is less than τ [1, p. 103]. Define $e(\delta) = 4A(S)/\ln(1/\delta)$ and pick $\delta > 0$ so that $2\pi e(\delta) < \tau^2$.

Let $w_0 = (u_0, v_0) \in B$. For $\delta \leq r \leq \sqrt{\delta}$, set $C_r = \{w \in \bar{B} \mid |w - w_0| = r\}$. Let (r, θ) be polar coordinates at w_0 and let $\xi(r, \theta) = X(w_0 + r(\cos(\theta), \sin(\theta)))$. If

$C_r \cap \partial B \neq \emptyset$, define $\alpha(r)$ and $\beta(r)$ by $0 \leq \alpha(r) < \beta(r) \leq 2\pi$ and $\{(r, \alpha(r)), (r, \beta(r))\} = C_r \cap \partial B$; otherwise $\alpha(r) = 0$ and $\beta(r) = 2\pi$. Set

$$p(r) = \int_{\alpha(r)}^{\beta(r)} |\xi_\theta(r, \theta)|^2 d\theta;$$

then

$$\int_\delta^{\sqrt{\delta}} p(r)/r dr \leq 2A(S).$$

For some $\rho \in [\delta, \sqrt{\delta}]$, $\ln(1/\delta)p(\rho)/2 \leq 2A(S)$ and so

$$p(\rho) \leq 4A(S)/\ln(1/\delta) = e(\delta).$$

If $C_\rho^* = X(C_\rho)$ and $L_\rho = \int_{\alpha(\rho)}^{\beta(\rho)} |\xi_\theta(\rho, \theta)| d\theta$ is the arclength of C'_ρ , then $L_\rho^2 \leq (\beta(\rho) - \alpha(\rho))p(\rho) \leq 2\pi e(\delta) < \tau^2$ and so $L_\rho < \tau < \min(d, \varepsilon/4)$. If $C_\rho \cap \partial B \neq \emptyset$, then the diameter of the shortest arc on $\Gamma \cup T$ joining the ends of C'_ρ is at most $\min(d, \varepsilon/4)$. For any $W \subset \bar{B}$, let $W^* = K(W)$.

Define J_ρ as the component of $B \setminus C_\rho$ which contains w_0 . Set

$$m = \inf\{z(u, v) | (u, v) \in C_\rho \text{ or } (u, v) \in \partial' B \cap \bar{J}_\rho\}$$

and

$$M = \sup\{z(u, v) | (u, v) \in C_\rho \text{ or } (u, v) \in \partial' B \cap \bar{J}_\rho\}.$$

Let $U = K(J_\rho)$ and $D_\rho = C_\rho \cup (\bar{J}_\rho \cap \partial' B)$. Notice that $X(J_\rho)$ is the graph of f over U and that $\partial U \setminus D_\rho^*$ is either empty or contains the single point $(0, 0)$. Now let us define $H'(x, y) = H(x, y, f(x, y))$ for $(x, y) \in \Omega$ and notice that $|H'(x, y)| \leq Q$ for all $(x, y) \in \Omega$. Notice that the diameter of U is less than $L_\rho + \text{diam}(D_\rho^* \setminus C_\rho^*)$, so less than $2d$. Now we apply Lemma 2.2 of [5] (with “ $\Omega = U$ ”, “ $u = M + g - g(2d, 0)$ ”, “ $v = f$ ”, and “ $\Gamma_1 = \partial U \setminus \{N\}$ ”) and see that

$$f(x, y) \leq M + g(x, y) - g(2d, 0) \leq M + \varepsilon/4 \quad \text{for } (x, y) \in U.$$

Since $z(u, v) = f(x(u, v), y(u, v))$, we get $z(u, v) < M + \varepsilon/4$ for $(u, v) \in J_\rho$. Similarly, $-z(u, v) \leq -m + \varepsilon/4$ for $(u, v) \in J_\rho$. Thus $m - \varepsilon/4 < z(u, v) < M + \varepsilon/4$ for $(u, v) \in J_\rho$.

Now the diameter of $X(\partial' B \cap \bar{J}_\rho)$ is less than $\varepsilon/4$ and $L_\rho < \varepsilon/4$, so $M - m < \varepsilon/2$. If $(u, v) \in B$ and $|(u, v) - (u_0, v_0)| < \delta$, then $(u, v) \in J_\rho$ and so $|z(u, v) - z(u_0, v_0)| < M - m + \varepsilon/2 < \varepsilon$. Since δ is independent of w_0 , z is uniformly continuous on B and so can be extended to \bar{B} as a continuous function. Q.E.D.

3. Boundary behavior. By [6], $X \in C^1(B \cup \partial'' B)$ and the branch points of X on $\partial'' B$ are isolated. We see that

$$X_u(u, 0) = (0, 0, z_u(u, 0)) \quad \text{and} \quad X_v(u, 0) = (x_v(u, 0), y_v(u, 0), 0).$$

Let (r, θ) be polar coordinates centered at $(0, 0)$. Then $\Omega = \{(r, \theta) | \alpha < \theta < \beta, 0 < r < r(\theta)\}$ for some $-\pi \leq \alpha < 0 < \beta \leq \pi$ (where we may need to rotate Ω about $(0, 0)$). For each $\alpha < \theta < \beta$, set $x(t) = x(t, \theta) = t \cdot \cos(\theta)$ and $y(t) = y(t, \theta) = t \cdot \sin(\theta)$. We denote the radial limit of f at $(0, 0)$ from the direction θ (if it exists) by

$$Rf(\theta) = \lim_{t \rightarrow 0^+} f(x(t, \theta), y(t, \theta)).$$

Set $\lambda(t) = (x(t), y(t), f(x(t)), y(t))$ and $\omega(t) = X^{-1}(\lambda(t))$. Notice that $\lambda(t) \rightarrow (0, 0, Rf(\theta))$ as $t \rightarrow 0+$ if $\lambda(t)$ converges to a point as $t \rightarrow 0+$. Finally define $Rf(\alpha) = Rf(\beta) = \phi(0, 0)$ and $u(\alpha) = -1, u(\beta) = 1$.

THEOREM 3.1. *For all $\alpha < \theta < \beta$, there is a unique $u(\theta) \in [-1, 1]$ such that $\omega(t) \rightarrow (u(\theta), 0)$ as $t \rightarrow 0+$. Also, $u \in C^0([\alpha, \beta])$. Thus $Rf(\theta) = z(u(\theta), 0)$ exists for all $\alpha \leq \theta \leq \beta$ and $Rf \in C^0([\alpha, \beta])$.*

The proof is the same as the proof of Lemma 3.1 of [8]. The only facts we use in this proof are that $X \in C^1(B \cup \partial''B)$, K is a homeomorphism of B and Ω , and the regular points of X on $\partial''B$ are dense in $\partial''B$. The fact that X maps $\partial'B$ (weakly or strongly) monotonically onto Γ_0 is not important and can be replaced by the following fact:

$$(x_v(u(\theta), 0), y_v(u(\theta), 0)) = |z_u(u(\theta), 0)|(\cos(\theta), \sin(\theta))$$

for $\alpha < \theta < \beta$ and so $u(\theta)$ is weakly monotonic on (α, β) .

THEOREM 3.2. *There exist $\lambda \in [\alpha, \beta - \pi]$ and $u_0 \in [-1, 1]$ such that $u(\theta) = u_0$ for all $\theta \in [\lambda, \lambda + \pi]$; also X is strictly monotonic on $[-1, u_0]$ and on $[u_0, 1]$.*

PROOF. Suppose $\theta_0 \in [\beta - \pi, \alpha + \pi]$ and $X(u, 0)$ is not weakly monotonic on $-1 \leq u \leq u(\theta_0)$. Then X has a branch point at $(b, 0)$ for some $-1 < b < u(\theta_0)$ and $z_u(u, 0)$ changes sign at $u = b$, say $z_u(u, 0) > 0$ on $(b - \varepsilon, b)$ and $z_u(u, 0) < 0$ on $(b, b + \varepsilon)$ for some $\varepsilon > 0$. Pick a and c so that $b - \varepsilon < a < b < c < b + \varepsilon$ and $z(a, 0) = z(c, 0)$. Then $a = u(\theta(a)), c = u(\theta(c))$, and $\alpha < \theta(a) < \theta(c) < \alpha + \pi$. Let ω be a smooth Jordan arc in B from $(a, 0)$ to $(c, 0)$ such that $\sigma = X(\omega)$ is a simple, closed Jordan curve, $\tau = K(\omega)$ is a convex Jordan curve which is smooth except at $(0, 0)$, and the curvature $\kappa(x, y)$ satisfies $\kappa(x, y) \geq 2Q$ for all $(0, 0) \neq (x, y) \in \tau$. Let U be the open region bounded by τ . We will use the function $H'(x, y) = H(x, y, f(x, y))$ mentioned earlier.

First, let $h \in C^0(\bar{U} \setminus \{N\})$ be the unique variational solution of $(p/W)_x + (q/W)_y = H'(x, y)$ in U with $h = f$ on $\tau \setminus \{N\}$ [4]. (We know that (2) is satisfied because of [5].) If we use a barrier argument of Serrin [9, pp. 375–376] together with Lemma 2.2 of [5], we see that $h \in C^0(\bar{U})$.

Second, by Lemma 2.2 of [5], we see that $f = h$ on U . This implies that $Rf(\theta) = h(0, 0)$ for all $\theta \in (\theta(a), \theta(c))$, a contradiction. Thus $X(u, 0)$ is weakly monotonic on $-1 \leq u \leq u(\theta_0)$. Using [6], we see that $X(u, 0)$ is strictly monotonic on $-1 \leq u \leq u(\theta_0)$. Similarly, $X(u, 0)$ is strictly monotonic on $u(\theta_0) \leq u \leq 1$. If θ_1 is another element of $[\beta - \pi, \alpha + \pi]$, the same argument proves that $X(u, 0)$ is strictly monotonic on $-1 \leq u \leq u(\theta_1)$ and on $u(\theta_1) \leq u \leq 1$. Since $X(-1, 0) = X(1, 0) = P$, we see that $u(\theta_0) = u(\theta_1)$. Thus $u(\theta)$ is constant on $[\beta - \pi, \alpha + \pi]$.

Suppose now that $\theta_2, \theta_3 \in [\alpha, \beta]$ with $0 < \theta_3 - \theta_2 < \pi$. The argument above shows that X is strictly monotonic on $[u(\theta_2), u(\theta_3)]$. Q.E.D.

Let $\theta_L = \inf\{\theta \in [\alpha, \beta] | u(\theta) = u_0\}$ and $\theta_R = \sup\{\theta \in [\alpha, \beta] | u(\theta) = u_0\}$. We know that $\theta_R - \theta_L \geq \pi$. Also $u(\cdot)$ is a homeomorphism of $[\alpha', \theta_L]$ onto $[-1, u_0]$ and of $[\theta_R, \beta']$ onto $[u_0, 1]$, for some $\alpha \leq \alpha' \leq \theta_L$ and $\theta_R < \beta' \leq \beta$. Thus

THEOREM 3.3. *The radial limits $Rf \in C^0([\alpha, \beta])$ behave as follows:*

(i) *The extreme values of $Rf(\theta)$ are $Rf(\alpha) = Rf(\beta) = \phi(0, 0)$ and $Rf(0) = z(u_0, 0)$.*

- (ii) $Rf(\theta)$ is monotonic on $[\alpha, \theta_L]$.
- (iii) $Rf(\theta) = z(u_0, 0)$ for all $\theta \in [\theta_L, \theta_R]$.
- (iv) $Rf(\theta)$ is monotonic on $[\theta_R, \beta]$.

Let U be open in \mathbf{R}^2 with $(0, 0) \in \partial U$, $H(x, y, t)$ be continuous on $U \times \mathbf{R}$, $\phi \in C^0(\partial U)$, and f be a C^2 solution of (1) in U . Suppose that for some $\varepsilon > 0$, the set $\Omega = \{(x, y) \in U \mid x^2 + y^2 < \varepsilon^2\}$ is locally convex at every point of $\partial\Omega$ except $(0, 0)$ and f satisfies assumptions (B), (D), (E) in Ω . Then f behaves at $(0, 0)$ as indicated in Theorem 3.3. As in [8], we make the following

CONJECTURE. If $\theta_R - \theta_L > \pi$, then $f \in C^0(\bar{\Omega})$.

ADDED IN PROOF. The conjecture mentioned above has been proven in *Non-parametric minimal surfaces in \mathbf{R}^3 whose boundaries have a jump discontinuity* (preprint) by the second author.

REFERENCES

1. R. Courant, *Dirichlet's principle, conformal mapping, and minimal surfaces*, Interscience, New York, 1950.
2. R. Courant and D. Hilbert, *Methods of mathematical physics*, Vol. 2, Interscience, New York, 1962.
3. R. Finn, *Remarks relevant to minimal surfaces, and to surfaces of prescribed mean curvature*, J. Analyse Math. **14** (1965), 139–160.
4. C. Gerhardt, *Existence, regularity, and boundary behavior of generalized surfaces of prescribed mean curvature*, Math. Z. **139** (1974), 173–198.
5. E. Giusti, *On the equation of surfaces of prescribed mean curvature*, Invent. Math. **46** (1978), 111–137.
6. E. Heinz, *Über das Randverhalten quasilinearer elliptischer Systeme mit isothermen Parametern*, Math. Z. **113** (1970), 99–105.
7. O. Ladyzhenskaya and N. Ural'tseva, *Local estimates for gradients of solutions of non-uniformly elliptic and parabolic equations*, Comm. Pure Appl. Math. **23** (1969), 677–703.
8. K. Lancaster, *Boundary behavior of a non-parametric minimal surface in \mathbf{R}^3 at a non-convex point*, Analysis **5** (1985), 61–69.
9. J. Serrin, *The Dirichlet problem for surfaces of constant mean curvature*, Proc. London Math. Soc. (3) **21** (1970), 361–384.
10. N. Trudinger, *Gradient estimates and mean curvature*, Math. Z. **131** (1973), 165–175.

DEPARTMENT OF MATHEMATICS AND STATISTICS, WICHITA STATE UNIVERSITY, WICHITA, KANSAS 67208