

## CONVERGENCE ACCELERATION FOR GENERALIZED CONTINUED FRACTIONS

PAUL LEVRIE AND LISA JACOBSEN

**ABSTRACT.** The main result in this paper is the proof of convergence acceleration for a suitable modification (as defined by de Bruin and Jacobsen) in the case of an  $n$ -fraction for which the underlying recurrence relation is of Perron-Kreuser type. It is assumed that the characteristic equations for this recurrence relation have only simple roots with differing absolute values.

**1. Introduction and notation.** Modifications of continued fractions have been studied extensively during the past few years by different authors [6, 10, 11, 12, 13]. Recently L. Jacobsen and M. G. de Bruin generalized this concept of modification to  $n$ -fractions or generalized continued fractions (GCF's). (For detail the reader is referred to [3]; the notation used will be that from [3].)

Let  $n$  be a fixed natural number ( $n \geq 1$ ). Consider a sequence of complex  $(n + 1)$ -tuples  $(b_k, a_k^{(1)}, \dots, a_k^{(n)})$ ,  $a_k^{(1)} \neq 0$ . Then the  $n$ -fraction associated with this sequence, written as

$$(1a) \quad K_{k=1}^{\infty} \left( \begin{array}{c} a_k^{(1)} \\ \vdots \\ a_k^{(n)} \\ b_k \end{array} \right)$$

is given by the sequence of approximants  $\{A_k^{(1)}/B_k, \dots, A_k^{(n)}/B_k\}_{k=1}^{\infty}$  (if they exist), where the numerators and denominators satisfy the recurrence relation

$$(1b) \quad X_k = b_k X_{k-1} + a_k^{(n)} X_{k-2} + \dots + a_k^{(1)} X_{k-n-1}, \quad k = 1, 2, \dots,$$

with initial values

$$(2) \quad A_{-j}^{(i)} = \delta_{i+j, n+1}, \quad B_{-j} = \delta_{n+1+j, n+1} \quad (j = 0, \dots, n)$$

for  $i = 1, \dots, n$ . The  $n$ -fraction is said to converge in  $\hat{\mathbb{C}}^n$  if the following limit exists in  $\hat{\mathbb{C}}^n$ :

$$(3) \quad \{\xi_0^{(1)}, \dots, \xi_0^{(n)}\} = \lim_{k \rightarrow \infty} \{A_k^{(1)}/B_k, \dots, A_k^{(n)}/B_k\}.$$

For  $k \geq 1$  we introduce the Moebius transforms

$$s_k^{(1)}(w^{(1)}, \dots, w^{(n)}) = \frac{a_k^{(1)}}{b_k + w^{(n)}},$$

$$s_k^{(i)}(w^{(1)}, \dots, w^{(n)}) = \frac{a_k^{(i)} + w^{(i-1)}}{b_k + w^{(n)}} \quad (i = 2, \dots, n)$$

Received by the editors September 1, 1986.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 65B99, 65Q05, 40A15.

©1988 American Mathematical Society  
 0002-9947/88 \$1.00 + \$.25 per page

and for  $i = 1, \dots, n$

$$(4) \quad \begin{aligned} S_1^{(i)}(w^{(1)}, \dots, w^{(n)}) &= s_1^{(i)}(w^{(1)}, \dots, w^{(n)}), \\ S_k^{(i)}(w^{(1)}, \dots, w^{(n)}) &= S_{k-1}^{(i)}(s_k^{(1)}, \dots, s_k^{(n)}) \quad (k \geq 2) \text{ (if they exist)}. \end{aligned}$$

Then we have the connection formula (cf. [1])

$$(5) \quad S_k^{(i)}(w^{(1)}, \dots, w^{(n)}) = \frac{A_k^{(i)} + A_{k-1}^{(i)}w^{(n)} + A_{k-2}^{(i)}w^{(n-1)} + \dots + A_{k-n}^{(i)}w^{(1)}}{B_k + B_{k-1}w^{(n)} + B_{k-2}w^{(n-1)} + \dots + B_{k-n}w^{(1)}}.$$

From this formula it follows that

$$(6) \quad A_k^{(i)} / B_k = S_k^{(i)}(0, \dots, 0) \quad \text{for } i = 1, \dots, n,$$

and we see that the sequence of approximants of the  $n$ -fraction (1) may be expressed by the Moebius transforms  $S_k^{(i)}$ . So as in the case of ordinary continued fractions, the approximants of the  $n$ -fraction (1) may be evaluated by replacing its tails, i.e.

$$(7) \quad K_{k=m+1}^\infty \begin{pmatrix} a_k^{(1)} \\ \vdots \\ a_k^{(n)} \\ b_k \end{pmatrix}$$

by zero for successive values of  $m$ . If the tail (7) converges, we let  $(\xi_m^{(1)}, \dots, \xi_m^{(n)})$  denote its value. Since the choice of zeros for the tails is a rather arbitrary one (the values of convergent tails converge to zero in exceptional cases only), the concept of modification of an  $n$ -fraction is introduced.

Given a sequence of  $n$ -tuples  $\{(w_k^{(1)}, \dots, w_k^{(n)})\}_{k=1}^\infty$  of numbers from  $\mathbf{C}$ , a modification of (1) is given by the sequence of  $n$ -tuples

$$(8) \quad \{S_k^{(1)}(w_k^{(1)}, \dots, w_k^{(n)}), \dots, S_k^{(n)}(w_k^{(1)}, \dots, w_k^{(n)})\}_{k=1}^\infty.$$

We assume  $(w_k^{(1)}, \dots, w_k^{(n)})$  to be chosen such that (8) is well defined. It can be seen from our main theorem that for convergent  $n$ -fractions and an appropriate choice of the  $w_k^{(i)}$  the sequence of modified approximants (8) converges to the value of the  $n$ -fraction faster than the sequence of ordinary approximants (6) (convergence acceleration is characterized by

$$(9) \quad \lim_{k \rightarrow \infty} \frac{\xi_0^{(i)} - S_k^{(i)}(w^{(1)}, \dots, w^{(n)})}{\xi_0^{(i)} - S_k^{(i)}(0, \dots, 0)} = 0 \quad (i = 1, \dots, n).$$

For the sequel we will restrict ourselves to the case that the  $n$ -fraction converges in  $\mathbf{C}^n$ . First we shall state a very important theorem in the theory of linear recurrence relations, i.e. the Perron-Kreuser theorem (cf. [4, 7, 8, 9]), which we shall use in the proof of Theorem I.

*The Perron-Kreuser theorem.* The recurrence relation

$$(1b) \quad X_k = b_k X_{k-1} + a_k^{(n)} X_{k-2} + \dots + a_k^{(1)} X_{k-n-1}, \quad k = 1, 2, \dots,$$

is said to be of Perron-Kreuser type if its coefficients satisfy

$$(10) \quad \begin{aligned} b_k &= -s_{n+1} k^{m_{n+1}} (1 + o(1)) \quad (k \rightarrow \infty), \\ a_k^{(i)} &= -s_i k^{m_i} (1 + o(1)) \quad (k \rightarrow \infty), \quad i = 1, \dots, n, \end{aligned}$$




FIGURE 1

where the  $s_i$  are real or complex numbers and  $s_{n+2}$  is defined to be one. The  $m_i$  are real or  $-\infty$  with  $m_1 > -\infty$ ;  $m_{n+2}$  is defined to be zero. With such a recurrence relation we can associate a uniquely defined Newton-Puiseux polygon in a rectangular  $(x, y)$ -coordinate system (see Figure 1).

Let the points  $p_0, p_1, \dots, p_{n+1}$  be defined by  $x = i, y = m_{n+2-i}$  ( $i = 0, \dots, n+1$ ). Then some of the points  $p_0, \dots, p_{n+1}$  are connected with linear segments in such a way that the resulting polygon is concave downwards, each point  $p_i$  ( $i = 1, \dots, n$ ) being on or below the resulting figure,  $p_0$  and  $p_{n+1}$  being the endpoints of the figure. If the polygon so constructed has  $\sigma$  distinct linear segments, their respective slopes are denoted by  $\alpha_1, \dots, \alpha_\sigma$  with  $\alpha_1 > \alpha_2 > \dots > \alpha_\sigma$ , and their abscissas are denoted by  $0 = e_0 < e_1 < \dots < e_\sigma = n + 1$ . It follows that

$$(11) \quad \alpha_\lambda = \frac{m_{n+2-e_\lambda} - m_{n+2-e_{\lambda-1}}}{e_\lambda - e_{\lambda-1}}, \quad \lambda = 1, \dots, \sigma,$$

and the Perron-Kreuser theorem states

**THEOREM.** *A linear recurrence relation with the above Newton-Puiseux polygon has a fundamental system of  $n + 1$  solutions which fall into  $\sigma$  classes. Each of these classes is further broken into subclasses, the  $\lambda$ th class ( $\lambda = 1, 2, \dots, \sigma$ ) containing  $\beta_\lambda$  subclasses. Let  $q_\gamma^{(\lambda)}$  ( $\gamma = 1, 2, \dots, \beta_\lambda$ ) denote the number of linearly independent solutions in the  $\gamma$ th subclass of the  $\lambda$ th class. Then each of the  $q_\gamma^{(\lambda)}$  solutions  $X_k$  and their nonzero linear combinations satisfy*

$$(12) \quad \limsup_{k \rightarrow \infty} (|X_k| / (k!)^{\alpha_\lambda})^{1/k} = t_\gamma^{(\lambda)}.$$

Here the  $t_\gamma^{(\lambda)}$  are distinct positive numbers which are the moduli of the roots of the following characteristic equation corresponding to the  $\lambda$ th class:

$$(13) \quad r_0^{(\lambda)} x^{e_\lambda - e_{\lambda-1}} + r_1^{(\lambda)} x^{e_\lambda - e_{\lambda-1} - 1} + \dots + r_{e_\lambda - e_{\lambda-1}}^{(\lambda)} = 0,$$

where

$$r_i^{(\lambda)} = s_{n+2-e_{\lambda-1}-i} \quad \text{or} \quad 0$$

depending on whether the point  $(e_{\lambda-1} + i, m_{n+2-e_{\lambda-1}-i})$  falls, respectively, on or below the  $\lambda$ th side of the Newton-Puiseux polygon. The number  $q_\gamma^{(\lambda)}$  is equal to

the number of roots, counting their multiplicities, of (13) with absolute value  $t_\gamma^{(\lambda)}$ . Thus, it follows that

$$(14) \quad q_1^{(\lambda)} + q_2^{(\lambda)} + \dots + q_{\beta_\gamma}^{(\lambda)} = e_\lambda - e_{\lambda-1}.$$

Further, to each simple root  $u$  of (13) whose absolute value is distinct from the absolute values of the other roots, there corresponds a solution  $X_k$  in the  $\lambda$ th class which satisfies

$$(15) \quad \lim_{k \rightarrow \infty} (X_{k+1}/k^{\alpha_\lambda} X_k) = u.$$

REMARKS. If  $e_\lambda > e_{\lambda-1} + 1$ , but none of the points  $p_i$ ,  $e_{\lambda-1} < i < e_\lambda$ , lies on the Newton-Puiseux polygon, then all the  $e_\lambda - e_{\lambda-1}$  roots of (13) are distinct but have coinciding absolute values

$$t_1^{(\lambda)} = |s_{n+2-e_\lambda}/s_{n+2-e_{\lambda-1}}|^{1/(e_\lambda-e_{\lambda-1})}.$$

(That is,  $\beta_\lambda = 1$  and  $q_1^{(\lambda)} = e_\lambda - e_{\lambda-1}$ .)

If  $e_\lambda = e_{\lambda-1} + 1$ , then there is only one linearly independent solution of (1b) from the  $\lambda$ th class. It behaves according to (15) with  $u = -s_{n+2-e_\lambda}/s_{n+2-e_{\lambda-1}}$ . If in addition  $\lambda = 1$ , then this is a dominant solution of (1b) and thus the GCF (1a) converges under mild conditions. Indeed, since then  $a_k^{(i)}/b_k b_{k-1} \dots b_{k-1-n+i} \rightarrow 0$  it follows that all the tails (7) converge from some  $m$  on ([1], see also Theorem 1).

From now on we will assume that for every  $\lambda$  the roots of (13) belonging to the  $\lambda$ th class are all different in modulus. Let  $u_i$  ( $i = e_{\lambda-1} + 1, \dots, e_\lambda$ ) be the roots of the characteristic equation for the  $\lambda$ th class, with

$$(16) \quad |u_{e_{\lambda-1}+1}| > \dots > |u_{e_\lambda}|.$$

Since  $r_{e_\lambda-e_{\lambda-1}}^{(\lambda)} = s_{n+2-e_\lambda} \neq 0$  we also have all  $|u_i| > 0$ .

It is easy to prove that in this case the recurrence relation (1b) has a basis ordered by domination: using (15) we are able to choose for each  $i = 1, \dots, n + 1$  a solution  $\{D_k^{(i)}\}$  of (1b) which satisfies

$$(17) \quad \lim_{k \rightarrow \infty} (D_{k+1}^{(i)}/k^{\alpha_\lambda} D_k^{(i)}) = u_i \quad \text{for } i = e_{\lambda-1} + 1, \dots, e_\lambda$$

and using (16) we then have

$$(18) \quad \lim_{k \rightarrow \infty} (D_k^{(i+1)}/D_k^{(i)}) = 0 \quad \text{for } i = 1, \dots, n.$$

Hence the  $\{D_k^{(i)}\}$  form a basis which is ordered by domination. From [2] follows therefore that the  $n$ -fraction connected with these recurrence relations, and all its tails, converge in  $\hat{C}^n$ . From (17) we have

$$(19) \quad \begin{aligned} D_{k+1}^{(i)}/D_k^{(i)} &= k^{\alpha_\lambda} u_i (1 + o(1)) && (k \rightarrow \infty) \\ &= v_i(k) (1 + o(1)) && (k \rightarrow \infty) \end{aligned}$$

where we have put  $v_i(k) = k^{\alpha_\lambda} u_i$ .

Later on we shall also need the following:

*Elementary symmetric functions.* The elementary symmetric functions for a set of  $m$  arbitrary complex numbers  $x_1, \dots, x_m$  are defined as

$$(20) \quad \begin{aligned} f_1(x_1, \dots, x_m) &= x_1 + \dots + x_m, \\ f_2(x_1, \dots, x_m) &= x_1x_2 + x_1x_3 + \dots + x_{m-1}x_m, \\ &\dots \\ f_m(x_1, \dots, x_m) &= x_1x_2 \dots x_m. \end{aligned}$$

If we construct from these  $f_i$  the polynomial

$$(21) \quad P_{x_1, \dots, x_m}(x) = x^m - f_1x^{m-1} + f_2x^{m-2} + \dots + (-1)^m f_m,$$

then this polynomial has the roots  $x_1, \dots, x_m$ , and so we have

$$(22) \quad P_{x_1, \dots, x_m}(x_i) = x_i^m - f_1x_i^{m-1} + f_2x_i^{m-2} + \dots + (-1)^m f_m = 0, \quad i = 1, \dots, m.$$

This can be seen as a system of  $m$  equations in the  $m$  unknowns  $f_i$  ( $i = 1, \dots, m$ ). Solving it using Cramer's rule gives us

$$(23) \quad f_{m-i+1} = \det V^{(i)} / \det V^{(m+1)}, \quad i = 1, \dots, m,$$

where  $V^{(i)}$  is the square matrix arising from the matrix  $V$  with

$$V = \begin{pmatrix} 1 & 1 & \dots & 1 \\ x_1 & x_2 & \dots & x_m \\ \vdots & \vdots & \vdots & \vdots \\ x_1^m & x_2^m & \dots & x_m^m \end{pmatrix}$$

by removing the  $i$ th row.

**2. Convergence of the values of the tails of an  $n$ -fraction.** In Theorem I we shall give a result concerning the limit of the tails of an  $n$ -fraction, and this for two special types of Perron-Kreuser recurrence relations:

$$(24) \quad \text{I. } e_1 > 1, \quad \alpha_1 = 0,$$

$$(25) \quad \text{II. } e_1 = 1, \quad \alpha_1 > \alpha_2 = 0.$$

(Note that using an equivalence transformation (see e.g. [1, 4]) every recurrence relation of Perron-Kreuser type can be brought into one of these two forms without changing the character of the Newton-Puiseux diagram or the equations (13).)

In our situation these two types of recurrence relations correspond to GCF's with the following properties.

*Type I.*  $m_i \leq 0$  for all  $i$ , where  $m_i = 0$  for  $i = n + 2 - e_1$  and at least one  $i > n + 2 - e_1$ , but for no  $i < n + 2 - e_1$ .

*Case (i).*  $m_{n+1} = 0$  (which is always the case if  $e_1 = 2$ ). Then

$$c_k^{(i)} = a_k^{(i)} / b_k b_{k-1} \dots b_{k-1-n+i} = (-1)^{n-i+1} (s_i / s_{n+1}^{n-i+2}) k^{m_i} (1 + o(1))$$

and thus  $a_k^{(i)} \rightarrow 0$  and  $c_k^{(i)} \rightarrow 0$  for all  $i < n + 2 - e_1$ . If in particular  $e_1 = 2$ , then (13) has the two solutions

$$u_{1,2} = -(s_{n+1}/2)(1 \pm \sqrt{1 - 4s_n/s_{n+1}^2})$$

for  $\lambda = 1$ , that is,  $-s_n/s_{n+1}^2 \notin (-\infty, -\frac{1}{4}]$ .

Case (ii).  $m_{n+1} < 0$  (which can only happen if  $e_1 > 2$  and thus  $n > 1$ ). Then

$$c_k^{(i)} = (-1)^{n-i+1} (s_i/s_{n+1}^{n-i+2}) k^{m_i - (n-i+2)m_{n+1}} (1 + o(1))$$

and thus  $c_k^{(i)} \rightarrow \infty$  for  $i = n + 2 - e_1$  and at least one  $i > n + 2 - e_1$ , but possibly also for other values of  $i$ . If in particular  $e_1 = 3$ , then (13) has the three solutions

$$u_{1,2,3} = (s_{n-1}/2)^{1/3} \left\{ \rho^p \left( -1 + \sqrt{1 + 4s_n^3/27s_{n-1}^2} \right)^{1/3} + \rho^q \left( -1 - \sqrt{1 + 4s_n^3/27s_{n-1}^2} \right)^{1/3} \right\}$$

for  $\lambda = 1$ , where  $\rho = e^{i2\pi/3}$  and  $(p, q) = (0, 0), (1, 2)$  and  $(2, 1)$ . That is  $-s_{n-1}^2/s_n^3 \notin (-\infty, 0] \cup [\frac{4}{27}, +\infty)$ .

Type II.  $m_{n+1} \geq m_i$  for all  $i$  with equality for  $i = n + 2 - e_2, m_{n+1} > 0$ . Then  $c_k^{(i)} \rightarrow 0$  for all  $i$ .

**THEOREM I.** *The values of the tails,  $\xi_k^{(i)}$  ( $i = 1, \dots, n$ ), of the GCF associated with a recurrence relation (1b) of Type I for which (16) holds for all  $\lambda$ , satisfy*

$$(26) \quad (a) \quad \lim_{k \rightarrow \infty} \xi_k^{(n-i+1)} = (-1)^i f_i(u_2, \dots, u_{e_1}) \quad \text{if } i = 1, \dots, e_1 - 1,$$

$$(27) \quad (b) \quad \lim_{k \rightarrow \infty} \frac{\xi_k^{(n-i+1)}}{k^{\alpha(i)}} = (-1)^{e_1-1} f_{e_1-1}(u_2, \dots, u_{e_1}) \frac{r_{i-e_{\lambda-1}+1}^{(\lambda)}}{r_0^{(2)}} \quad \text{if } i = e_{\lambda-1}, \dots, e_{\lambda} - 1,$$

for  $\lambda = 2, \dots, \sigma$  with

$$\alpha(i) = (e_2 - e_1)\alpha_2 + (e_3 - e_2)\alpha_3 + \dots + (i - e_{\lambda-1} + 1)\alpha_{\lambda}.$$

For a recurrence relation of Type II we have

$$(28) \quad \lim_{k \rightarrow \infty} \frac{\xi_k^{(n-i+1)}}{k^{\alpha(i)}} = \frac{r_{i-e_{\lambda-1}+1}^{(\lambda)}}{r_0^{(2)}} \quad \text{if } i = e_{\lambda-1}, \dots, e_{\lambda} - 1, \quad \lambda = 2, \dots, \sigma.$$

**REMARK.** In particular this means that

$$\lim_{k \rightarrow \infty} \xi_k^{(i)} = \begin{cases} 0 & \text{if } i \leq n + 1 - e_1 \\ (-1)^{n-i+1} f_{n-i+1}(u_2, \dots, u_{e_1}) & \text{if } i > n + 1 - e_1 \end{cases} \quad \text{for Type I}$$

and

$$\lim_{k \rightarrow \infty} \xi_k^{(i)} = \begin{cases} 0 & \text{if } i \leq n + 1 - e_2 \\ r_{n-i+1}^{(2)}/r_0^{(2)} & \text{if } i > n + 1 - e_2 \end{cases} \quad \text{for Type II.}$$

**PROOF.** Let us consider the shifted recurrence relation

$$(29) \quad X_m = b_{m+k} X_{m-1} + a_{m+k}^{(n)} X_{m-2} + \dots + a_{m+k}^{(1)} X_{m-n-1} \quad (m \geq 1).$$

The tails  $\xi_k^{(i)}$  then satisfy  $\xi_k^{(i)} = \lim_{m \rightarrow \infty} A_{k,m}^{(i)} / B_{k,m}$  where the  $A_{k,m}^{(i)}$  and  $B_{k,m}$  follow from (1b) and the initial values (2) as for the nonshifted case. Because the  $D_k^{(i)}$ 's form a basis for (1b) the shifted  $D_k^{(i)}$ 's form a basis for (29):

$$(30) \quad \begin{aligned} A_{k,m}^{(i)} &= \phi_1^{(k,i)} D_{m+k}^{(1)} + \phi_2^{(k,i)} D_{m+k}^{(2)} + \dots + \phi_{n+1}^{(k,i)} D_{m+k}^{(n+1)}, \quad i = 1, \dots, n, \\ B_{k,m} &= \psi_1^{(k)} D_{m+k}^{(1)} + \psi_2^{(k)} D_{m+k}^{(2)} + \dots + \psi_{n+1}^{(k)} D_{m+k}^{(n+1)}. \end{aligned}$$

If  $l^{(i)} = (l_1^{(i)}, \dots, l_{n+1}^{(i)})^T$  ( $i = 1, \dots, n+1$ ) with  $l_j^{(i)} = \delta_{i,j}$  ( $j = 1, \dots, n+1$ ) and if  $D_k$  is defined by

$$(31) \quad D_k = \begin{pmatrix} D_{k-n}^{(1)} & D_{k-n}^{(2)} & \dots & D_{k-n}^{(n+1)} \\ D_{k-n+1}^{(1)} & D_{k-n+1}^{(2)} & \dots & D_{k-n+1}^{(n+1)} \\ \vdots & \vdots & \ddots & \vdots \\ D_k^{(1)} & D_k^{(2)} & \dots & D_k^{(n+1)} \end{pmatrix}$$

then  $\phi_1^{(k,i)}$  and  $\psi_1^{(k)}$  may be calculated from

$$\begin{aligned} D_k (\phi_1^{(k,i)}, \dots, \phi_{n+1}^{(k,i)})^T &= l^{(i)} \quad \text{for } i = 1, \dots, n, \\ D_k (\psi_1^{(k)}, \dots, \psi_{n+1}^{(k)})^T &= l^{(n+1)}. \end{aligned}$$

Since the  $\{D_k^{(i)}\}$ 's form a basis, we have  $\det D_k \neq 0$  and Cramer's rule gives us

$$\phi_1^{(k,i)} = (-1)^{i+1} \frac{\det D_k^{(i,1)}}{\det D_k}, \quad \psi_1^{(k)} = (-1)^n \frac{\det D_k^{(n+1,1)}}{\det D_k}$$

where  $D_k^{(p,1)}$  is the matrix arising from  $D_k$  by removing the  $p$ th row and the first column. Since the basis of the  $\{D_k^{(i)}\}$ 's is ordered by domination, it follows that  $\det D_k^{(n+1,1)} \neq 0$  from some  $k$  on. Hence

$$\xi_k^{(n-i+1)} = \frac{\phi_1^{(k,n-i+1)}}{\psi_1^{(k)}} = (-1)^i \frac{\det D_k^{(n-i+1,1)}}{\det D_k^{(n+1,1)}}, \quad i = 1, \dots, n,$$

is well defined and finite. If we divide the  $l$ th column of  $D_k^{(n-i+1,1)}$  and  $D_k^{(n+1,1)}$  by  $D_{k-n}^{(l+1)}$  for  $l = 1, \dots, n$ , then, using (19) and (23) we find

$$\frac{\phi_1^{(k,n-i+1)}}{\psi_1^{(k)}} = (-1)^i f_i(v_2(k), \dots, v_{n+1}(k))(1 + o(1)) \quad (k \rightarrow \infty)$$

since

$$\frac{D_{k-n+j}^{(i)}}{D_{k-n}^{(i)}} = \frac{D_{k-n+j}^{(i)}}{D_{k-n+j-1}^{(i)}} \dots \frac{D_{k-n+1}^{(i)}}{D_{k-n}^{(i)}} = (v_i(k))^j (1 + o(1)) \quad (k \rightarrow \infty), j = 0, \dots, n.$$

Now if the recurrence relation (1b) is of Type I, we have

$$\begin{aligned} \phi_1^{(k,n-i+1)} / \psi_1^{(k)} &= (-1)^i f_i(u_2, \dots, u_{e_1}, k^{\alpha_2} u_{e_1+1}, \dots, k^{\alpha_2} u_{e_2}, \dots, \\ &\quad k^{\alpha_\sigma} u_{e_{\sigma-1}+1}, \dots, k^{\alpha_\sigma} u_{n+1})(1 + o(1)). \end{aligned}$$

If  $i = 1, \dots, e_1 - 1$  it is easy to see that

$$\phi_1^{(k, n-i+1)} / \psi_1^{(k)} = (-1)^i f_i(u_2, \dots, u_{e_1})(1 + o(1)) \quad (k \rightarrow \infty)$$

since  $0 > \alpha_2 > \alpha_3 > \dots > \alpha_\sigma$ . Hence

$$\begin{aligned} \lim_{k \rightarrow \infty} \xi_k^{(n-i+1)} &= \lim_{k \rightarrow \infty} \lim_{m \rightarrow \infty} \frac{A_{k,m}^{(n-i+1)}}{B_{k,m}} = \lim_{k \rightarrow \infty} \frac{\phi_1^{(k, n-i+1)}}{\psi_1^{(k)}} \\ &= (-1)^i f_i(u_2, \dots, u_{e_1}) \end{aligned}$$

using (30) and the fact that the basis of the  $\{D_k^{(i)}\}$ 's is ordered by domination. This proves part (a) of the theorem. If  $i = e_{\lambda-1}, \dots, e_\lambda - 1$  ( $\lambda = 2, \dots, \sigma$ ) we find, using the properties of the  $f_i$ :

$$\begin{aligned} \frac{\phi_1^{(k, n-i+1)}}{\psi_1^{(k)}} &= (-1)^i f_{e_1-1}(u_2, \dots, u_{e_1}) \left( \prod_{j=2}^{\lambda-1} k^{(e_j - e_{j-1})\alpha_j} f_{e_j - e_{j-1}}(u_{e_{j-1}+1}, \dots, u_{e_j}) \right) \\ &\quad \times k^{(i - e_{\lambda-1} + 1)\alpha_\lambda} f_{i - e_{\lambda-1} + 1}(u_{e_{\lambda-1}+1}, \dots, u_{e_\lambda})(1 + o(1)) \\ &= (-1)^i k^{\alpha(i)} f_{e_1-1}(u_2, \dots, u_{e_1}) \left( \prod_{j=2}^{\lambda-1} f_{e_j - e_{j-1}}(u_{e_{j-1}+1}, \dots, u_{e_j}) \right) \\ &\quad \times f_{i - e_{\lambda-1} + 1}(u_{e_{\lambda-1}+1}, \dots, u_{e_\lambda})(1 + o(1)) \\ &= (-1)^i k^{\alpha(i)} f_{e_1-1}(u_2, \dots, u_{e_1}) \left( \prod_{j=2}^{\lambda-1} (-1)^{e_j - e_{j-1}} \frac{r_{e_j - e_{j-1}}^{(j)}}{r_0^{(j)}} \right) \\ &\quad \times (-1)^{i - e_{\lambda-1} + 1} \frac{r_{i - e_{\lambda-1} + 1}^{(\lambda)}}{r_0^{(\lambda)}} (1 + o(1)) \quad \text{using (13), (21)} \\ &= (-1)^i k^{\alpha(i)} f_{e_1-1}(u_2, \dots, u_{e_1}) (-1)^{i - e_1 + 1} \left( \prod_{j=2}^{\lambda-1} \frac{r_{e_j - e_{j-1}}^{(j)}}{r_0^{(j)}} \right) \\ &\quad \times \frac{r_{i - e_{\lambda-1} + 1}^{(\lambda)}}{r_0^{(\lambda)}} (1 + o(1)) \\ &= (-1)^{e_1 - 1} k^{\alpha(i)} f_{e_1-1}(u_2, \dots, u_{e_1}) \frac{r_{i - e_{\lambda-1} + 1}^{(\lambda)}}{r_0^{(2)}} (1 + o(1)) \quad (k \rightarrow \infty) \end{aligned}$$

since  $r_0^{(j)} = r_{e_{j-1} - e_{j-2}}^{(j-1)}$  for  $j = 2, \dots, n + 1$ . Hence,

$$\begin{aligned} \lim_{k \rightarrow \infty} \frac{\xi_k^{(n-i+1)}}{k^{\alpha(i)}} &= \lim_{k \rightarrow \infty} \lim_{m \rightarrow \infty} \frac{A_{k,m}^{(n-i+1)}}{k^{\alpha(i)} B_{k,m}} = \lim_{k \rightarrow \infty} \frac{\phi_1^{(k, n-i+1)}}{k^{\alpha(i)} \psi_1^{(k)}} \\ &= (-1)^{e_1 - 1} f_{e_1-1}(u_2, \dots, u_{e_1}) \frac{r_{i - e_{\lambda-1} + 1}^{(\lambda)}}{r_0^{(2)}}. \end{aligned}$$

This proves part (b). To prove the second part of the theorem a similar argument may be used.

**3. The main theorem.** Let us define modifying factors  $w_k^{(i)} = w^{(i)}$  for  $i = 1, \dots, n; k \geq 1$  by

$$(32) \quad w^{(i)} = \lim_{k \rightarrow \infty} \xi_k^{(i)}$$

using Theorem I.

**THEOREM II.** *Given the  $n$ -fraction*

$$(1a) \quad K_{k=1}^\infty \begin{pmatrix} a_k^{(1)} \\ \vdots \\ a_k^{(n)} \\ b_k \end{pmatrix}$$

satisfying the assumptions (10), (16) and (24) or (25) and such that for  $j = 1, \dots, n$

$$(33) \quad \{A_k^{(j)}\} \notin \text{vect} \left\{ \{D_k^{(i)}\} | i = e_\rho + 1, \dots, n + 1 \right\} \quad \text{with } \rho = \begin{cases} 1 & \text{for Type I,} \\ 2 & \text{for Type II} \end{cases}$$

(for a set  $V$   $\text{vect } V$  denotes the vector space spanned by the elements of  $V$ ). If the  $n$ -fraction converges to a finite value  $\{\xi_0^{(1)}, \dots, \xi_0^{(n)}\} \in \mathbb{C}^n$ , we have

$$(34) \quad \lim_{k \rightarrow \infty} \frac{\xi_0^{(i)} - S_k^{(i)}(w^{(1)}, \dots, w^{(n)})}{\xi_0^{(i)} - S_k^{(i)}(0, \dots, 0)} = 0 \quad (i = 1, \dots, n).$$

**REMARK.** In [3] de Bruin and Jacobsen proved this theorem for limit-1-periodic  $n$ -fractions. As seen from our point of view limit-1-periodic  $n$ -fractions are  $n$ -fractions associated with a Perron-Kreuser recurrence relation of Type I with  $\rho = \sigma = 1$ .

**EXAMPLE.** Consider the 3-fraction

$$\begin{aligned} a_k^{(1)} &= 51k + 2500 + \frac{1}{k+1} + \frac{1}{k(k+1)}, \\ a_k^{(2)} &= -100k - 5051 + \frac{1}{k+1} + \frac{1}{k(k+1)}, \\ b_k &= k + 150 + \frac{1}{k+1} + \frac{1}{k(k+1)} \end{aligned}$$

with  $w^{(1)} = 51, w^{(2)} = -100$ . The exact value of the GCF (all digits correct) is

$$\xi_0^{(1)} = 25.849821113137, \quad \xi_0^{(2)} = -51.896713180952.$$

In the tables below values of  $S_k^{(i)}(0, 0)$  and  $S_k^{(i)}(w^{(1)}, w^{(2)})$  (Table 1) and of the acceleration coefficients

$$\eta_k^{(i)} = (\xi_0^{(i)} - S_k^{(i)}(w^{(1)}, w^{(2)})) / (\xi_0^{(i)} - S_k^{(i)}(0, 0))$$

(Table 2) are given. (The acceleration shows up rather late since the constant terms are so large.)

**PROOF OF THE MAIN THEOREM.** The proof given in [3] for the special case mentioned above is still valid, only a few changes have to be made. The proof consists of three steps:

(35) (a)  $S_{p+k}^{(i)}(0, \dots, 0) - S_k^{(i)}(0, \dots, 0) \neq 0$  for  $k \geq k_0, p \geq p_0$ .

(36) (b)  $\lim_{p \rightarrow \infty} \frac{S_{p+k}^{(i)}(0, \dots, 0) - S_k^{(i)}(w^{(1)}, \dots, w^{(n)})}{S_{p+k}^{(i)}(0, \dots, 0) - S_k^{(i)}(0, \dots, 0)} \in \mathbb{C}$ .

(c) The proof of (34).

Table 1

$\nu$	$S_\nu^{(1)}(0, 0)$	$S_\nu^{(1)}(w^{(1)}, w^{(2)})$	$S_\nu^{(2)}(0, 0)$	$S_\nu^{(2)}(w^{(1)}, w^{(2)})$
1	16.78947368	49.07692308	-33.88157895	-98.05769231
10	25.79789811	33.00217711	-51.79348166	-66.11677994
20	25.84765251	26.10655259	-51.89240164	-52.40713782
30	25.84878915	25.86690369	-51.89466146	-51.93067617
40	25.84890247	25.85223504	-51.89488677	-51.90151247
50	25.84894320	25.85029284	-51.89496774	-51.89765105
60	25.84898775	25.84990368	-51.89505631	-51.89687734
70	25.84910613	25.84983028	-51.89529167	-51.89673140
75	25.84925570	25.84982352	-51.89558906	-51.89671797
80	25.84948770	25.84982165	-51.89605030	-51.89671424
85	25.84969579	25.84982121	-51.89646401	-51.89671338
90	25.84979017	25.84982113	-51.89665165	-51.89671321
95	25.84981540	25.84982112	-51.89670181	-51.89671319
100	25.84982026	25.84982111	-51.89671148	-51.89671318
105	25.84982101	25.84982111	-51.89671297	-51.89671318
110	25.84982110	25.84982111	-51.89671316	-51.89671318
115	25.84982111	25.84982111	-51.89671318	-51.89671318

Table 2

$\nu$	$\eta_\nu^{(1)}$	$\eta_\nu^{(2)}$
1	-2.56359948	-2.56234444
10	-137.74927241	-137.74927241
20	-118.38567573	-118.38567573
30	-16.55343491	-16.55343491
40	-2.62770954	-2.62770954
50	-0.53732460	-0.53732460
60	-0.09907743	-0.09907743
70	-0.01281565	-0.01281565
75	-0.00425901	-0.00425901
80	-0.00159917	-0.00159917
85	-0.00079294	-0.00079294
90	-0.00050498	-0.00050498
95	-0.00036394	-0.00036394
100	-0.00027561	-0.00027561
105	-0.00021376	-0.00021376
110	-0.00016843	-0.00016843
115	-0.00013477	-0.00013468

The fact that all limits  $\xi_0^{(i)}$  are finite and the fact that the basis  $\{D_k^{(i)}\}$  is ordered by domination together imply that  $\{B_k, D_k^{(2)}, \dots, D_k^{(n+1)}\}$  is also a basis for (1b) with

$$(37) \quad \lim_{k \rightarrow \infty} \frac{B_k}{k^{\alpha_1} B_{k-1}} = u_1 \quad (\neq 0), \quad \lim_{k \rightarrow \infty} \frac{D_k^{(i)}}{B_k} = 0 \quad (i = 2, \dots, n).$$

Using the same argument as in [3], we have

$$(38) \quad A_k^{(i)} = \delta_1^{(i)} B_k + \sum_{q=2}^{n+1} \delta_q^{(i)} D_k^{(q)} \quad (i = 1, \dots, n; k \geq 1)$$

for some  $\delta_q^{(i)}$  with  $\delta_1^{(i)} = \xi_0^{(i)}$  and

$$(39) \quad \forall i \in \{1, \dots, n\} \exists t \in \{2, \dots, e_\rho\} (\delta_q^{(i)} = 0 \quad (q = 2, \dots, t-1) \& \delta_t^{(i)} \neq 0)$$

with  $\rho$  as in (33).

The proof of (a) is exactly the same as in [3]: we write

$$S_{p+k}^{(i)}(0, \dots, 0) - S_k^{(i)}(0, \dots, 0) = \frac{A_{p+k}^{(i)}}{B_{p+k}} - \frac{A_k^{(i)}}{B_k} = \frac{A_{p+k}^{(i)} B_k - A_k^{(i)} B_{p+k}}{B_{p+k} B_k}.$$

From (37) it follows that the denominator is different from zero for  $k \geq k_1, p \geq 1$ . Clearly it is also different from  $\infty$ . Using (38) the numerator becomes

$$A_{p+k}^{(i)} B_k - A_k^{(i)} B_{p+k} = B_{p+k} D_k^{(t)} \sum_{q=t}^{n+1} \delta_q^{(i)} \left( \frac{D_{p+k}^{(q)} B_k}{B_{p+k} D_k^{(t)}} - \frac{D_k^{(q)}}{D_k^{(t)}} \right)$$

where the factor before the summation sign is different from zero for  $k \geq k_2 \geq k_1, p \geq 1$  by (37) and (17) since  $u_i \neq 0$  for all  $i$ , and

$$\begin{aligned} \left| \sum_{q=t}^{n+1} \delta_q^{(i)} \left( \frac{D_{p+k}^{(q)} B_k}{B_{p+k} D_k^{(t)}} - \frac{D_k^{(q)}}{D_k^{(t)}} \right) \right| &= \left| \sum_{q=t}^{n+1} \delta_q^{(i)} \frac{D_k^{(q)}}{D_k^{(t)}} \left( \frac{D_{p+k}^{(q)} B_k}{D_k^{(q)} B_{p+k}} - 1 \right) \right| \\ &\geq \frac{1}{2} \left| \delta_t^{(i)} \left( \frac{D_{p+k}^{(t)} B_k}{D_k^{(t)} B_{p+k}} - 1 \right) \right| \geq \frac{1}{4} |\delta_t^{(i)}| > 0 \end{aligned}$$

for sufficiently large  $k$  and  $p$ . (The first inequality follows since  $D_k^{(q)}/D_k^{(t)} \rightarrow 0$  for all  $q > t$  and since  $D_{p+k}^{(q)} B_k / D_k^{(q)} B_{p+k} \rightarrow 0$  as  $p \rightarrow \infty$  and is thus bounded from some  $p$  and  $k$  on. The second inequality comes from using  $D_{p+k}^{(q)} B_k / D_k^{(q)} B_{p+k} \rightarrow 0$  again for  $q = t$ .) This proves (a).

Using (5) we rewrite the expression in (b) as

$$(40) \quad \frac{S_{p+k}^{(i)}(0, \dots, 0) - S_k^{(i)}(w^{(1)}, \dots, w^{(n)})}{S_{p+k}^{(i)}(0, \dots, 0) - S_k^{(i)}(0, \dots, 0)} = \left( 1 + \sum_{j=0}^{n-1} w^{(n-j)} \frac{B_{k-j-1}}{B_k} \right)^{-1} \sum_{j=-1}^{n-1} w^{(n-j)} q_{p,k,j}^{(i)}$$

where  $w^{(n+1)} = 1$  and

$$\begin{aligned}
 q_{p,k,j}^{(i)} &= \frac{A_{p+k}^{(i)} B_{k-j-1} - A_{k-j-1}^{(i)} B_{p+k}}{A_{p+k}^{(i)} B_k - A_k^{(i)} B_{p+k}} \\
 &= \sum_{q=2}^{n+1} \delta_q^{(i)} \left( \frac{D_{p+k}^{(q)}}{B_{p+k}} B_{k-j-1} - D_{k-j-1}^{(q)} \right) / \sum_{q=2}^{n+1} \delta_q^{(i)} \left( \frac{D_{p+k}^{(q)}}{B_{p+k}} B_k - D_k^{(q)} \right).
 \end{aligned}$$

The first factor in (40) satisfies (use (37))

(41)

$$\begin{aligned}
 \lim_{k \rightarrow \infty} \left( 1 + \sum_{j=0}^{n-1} w^{(n-j)} \frac{B_{k-j-1}}{B_k} \right) &= 1 \quad \text{if the recurrence relation is of Type II,} \\
 &= \frac{P_{u_2, \dots, u_{e_1}}(u_1)}{u_1^{e_1-1}} \neq 0 \quad \text{for Type I (use (21)).}
 \end{aligned}$$

It is therefore finite for sufficiently large  $k$ . For the second factor in (40) we have

$$\lim_{p \rightarrow \infty} q_{p,k,j}^{(i)} = \frac{\sum_{q=t}^{n+1} \delta_q^{(i)} D_{k-j-1}^{(q)}}{\sum_{q=t}^{n+1} \delta_q^{(i)} D_k^{(q)}} = \frac{\sum_{q=t}^{n+1} \delta_q^{(i)} D_{k-j-1}^{(q)} / D_k^{(t)}}{\sum_{q=t}^{n+1} \delta_q^{(i)} D_k^{(q)} / D_k^{(t)}}$$

where

$$\left| \sum_{q=t}^{n+1} \delta_q^{(i)} D_k^{(q)} / D_k^{(t)} \right| \geq \frac{1}{2} |\delta_t^{(i)}| > 0$$

for  $k \geq k_0$ . This proves (b).

(c) Since

$$\lim_{p \rightarrow \infty} S_{p+k}^{(i)}(0, \dots, 0) = \xi_0^{(i)} \quad (k \geq 1)$$

we have for the acceleration coefficients

$$\begin{aligned}
 &\lim_{k \rightarrow \infty} \frac{\xi_0^{(i)} - S_k^{(i)}(w^{(1)}, \dots, w^{(n)})}{\xi_0^{(i)} - S_k^{(i)}(0, \dots, 0)} \\
 &= \lim_{k \rightarrow \infty} \lim_{p \rightarrow \infty} \frac{S_{p+k}^{(i)}(0, \dots, 0) - S_k^{(i)}(w^{(1)}, \dots, w^{(n)})}{S_{p+k}^{(i)}(0, \dots, 0) - S_k^{(i)}(0, \dots, 0)} \\
 &= \lim_{k \rightarrow \infty} \left\{ \left( 1 + \sum_{j=0}^{n-1} w^{(n-j)} \frac{B_{k-j-1}}{B_k} \right)^{-1} \right. \\
 &\quad \left. \times \sum_{j=-1}^{n-1} w^{(n-j)} \frac{\sum_{q=t}^{n+1} \delta_q^{(i)} D_{k-j-1}^{(q)} / D_{k-j-1}^{(t)}}{\sum_{q=t}^{n+1} \delta_q^{(i)} D_k^{(q)} / D_{k-j-1}^{(t)}} \right\} \quad \text{with } t \leq e_\rho \\
 &= \frac{u_1^{e_1-1}}{P_{u_2, \dots, u_{e_1}}(u_1)} \sum_{j=-1}^{n-1} w^{(n-j)} \frac{\delta_t^{(i)}}{\delta_t^{(i)} u_t^{j+1}} = \frac{u_1^{e_1-1}}{P_{u_2, \dots, u_{e_1}}(u_1)} \frac{P_{u_2, \dots, u_{e_1}}(u_t)}{u_t^{e_1-1}} = 0
 \end{aligned}$$

if the recurrence relation is of Type I, and

$$\sum_{j=-1}^{n-1} w^{(n-j)} \frac{\delta_t^{(i)}}{\delta_t^{(i)} u_t^{j+1}} = \frac{P_{u_2, \dots, u_{e_2}}(u_t)}{u_t^{e_2-1}} = 0$$

if the recurrence relation is of Type II, where we have used (22) and (33).

#### REFERENCES

1. Marcel G. de Bruin, *Convergence of generalized C-fractions*, J. Approx. Theory **24** (1978), 177-207.
2. Marcel G. de Bruin and Lisa Jacobsen, *The dominance concept for linear recurrence relations with applications to continued fractions*, Nieuw Arch. Wisk. (4) **3** (1985), 253-266.
3. —, *Modification of generalised continued fractions*. I, Lecture Notes in Math., vol 1237 (J. Gilewicz, M. Pindor, W. Siemaszko, Eds.), Springer-Verlag, Berlin, 1987, pp. 161-176.
4. J. R. Cash, *A note on the numerical solution of linear recurrence relations*, Numer. Math. **34** (1980), 371-386.
5. P. Van der Cruyssen, *Linear difference equations and generalized continued fractions*, Computing **22** (1979), 269-287.
6. Lisa Jacobsen, *Modified approximants for continued fractions, construction and applications*, Norske Vid. Selsk. Skr., no. 3 (1983).
7. P. Kreuser, *Über das Verhalten der Integrale homogener linearer Differenzgleichungen im Unendlichen*, Thesis (Tubingen), Borna-Leipzig, 1914.
8. Oskar Perron, *Über Summengleichungen und Poincarésche Differenzgleichungen*, Math. Ann. **84** (1921), 1-15.
9. —, *Über lineare Differenzgleichungen und eine Anwendung auf lineare Differentialgleichungen mit Polynomkoeffizienten*, Math. Z. **72** (1959), 16-24.
10. Wolfgang J. Thron and Haakon Waadeland, *Accelerating convergence of limit-periodic continued fractions  $K(a_n/1)$* , Numer. Math. **34** (1980), 155-170.
11. —, *Analytic continuation of functions defined by means of continued fractions*, Math. Scand. **47** (1980), 72-90.
12. —, *Convergence questions for limit-periodic continued fractions*, Rocky Mountain J. Math. **11** (1981), 641-657.
13. W. B. Jones, W. J. Thron and H. Waadeland, Eds., *Analytic theory of continued fractions*, Proceedings Loen, Norway 1981, Lecture Notes in Math., vol. 932, Springer-Verlag, Berlin, 1982.

DEPARTMENT OF COMPUTER SCIENCE, UNIVERSITY OF LEUVEN, CELESTIJNENLAAN 200 A, B-3030 HEVERLEE, BELGIUM

INSTITUTT FOR MATEMATIKK OG STATISTIKK, UNIVERSITETET I TRONDHEIM, AVH, N-7055 DRAGVOLL, NORWAY