

## EQUIVALENT CONDITIONS TO THE SPECTRAL DECOMPOSITION PROPERTY FOR CLOSED OPERATORS

I. ERDELYI AND WANG SHENGWANG

**ABSTRACT.** The spectral decomposition property has been instrumental in developing a local spectral theory for closed operators acting on a complex Banach space. This paper gives some necessary and sufficient conditions for a closed operator to possess the spectral decomposition property.

In the monograph [3] and in a sequel of papers by the authors, a local spectral theory has been built for closed operators on the sole assumption of the spectral decomposition property. As an abstraction of Dunford's concept of "spectral reduction" [2, p. 1927] and that of Bishop's "duality theory of type 3" [1], an operator  $T$  endowed with the spectral decomposition property produces a spectral decomposition of the underlying space, pertinent to any finite open cover of the spectrum  $\sigma(T)$ . In this paper we obtain some conditions equivalent to the spectral decomposition property. Some of them generalize results from [4].

### 1. PRELIMINARIES

Given a Banach space  $X$  over the complex field  $\mathbb{C}$ , we denote by  $C(X)$  the class of closed operators with domain  $D_T$  and range in  $X$ , and we write  $C_d(X)$  for the subclass of the densely defined operators in  $C(X)$ . For a subset  $Y$  of  $X$ ,  $Y^\perp$  denotes the annihilator of  $Y$  in  $X^*$  and for  $Z \subset X^*$ , we use the symbol  ${}^\perp Z$  for the preannihilator of  $Z$  in  $X$ . For the rest, the terminology and notation conform to that employed in [3].

We shall adopt and adjust some concepts and ideas from Bishop's "duality theory of type 4" [1, §4]. A couple  $U_1$  and  $U_2$  of a bounded and an unbounded Cauchy domain, related by  $U_2 = (\overline{U_1})^c$ , are referred to as complementary simple sets.  $W_1$  and  $W_2$  are the sets of analytic functions from  $U_1$  to  $X$  and from  $U_2$  and  $X^*$ , respectively, which vanish at  $\infty$ . The seminorms

$$\|f\|_{K_1} = \max\{\|f(\lambda)\| : f \in W_1, \lambda \in K_1, K_1 (\subset U_1) \text{ is compact}\}$$

and

$$\|g\|_{K_2} = \max\{\|g(\lambda)\| : g \in W_2, \lambda \in K_2, K_2 (\subset U_2) \text{ is compact}\}$$

---

Received by the editors September 15, 1987.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 47B40, Secondary 47A10, 47A15, 47A30.

induce a locally convex topology on  $W_1$  and  $W_2$ , respectively. For  $i = 1, 2$  let  $V_i$  be the subset of  $W_i$  on which every function can be extended to be continuous on  $\overline{U}_i$ . For  $f \in V_1$ ,  $g \in V_2$ , the norms

$$\|f\|_{V_1} = \sup\{\|f(\lambda)\|: \lambda \in U_1\}, \quad \|g\|_{V_2} = \sup\{\|g(\lambda)\|: \lambda \in U_2\}$$

make  $(V_1, \|\cdot\|_{V_1})$  and  $(V_2, \|\cdot\|_{V_2})$  Banach spaces. For  $x \in X$ ,  $\mu \in U_1$  and  $\lambda \in U_2$ , define

$$(1.1) \quad \alpha(x, \mu, \lambda) = (\mu - \lambda)^{-1}x.$$

For fixed  $x \in X$  and  $\lambda \in U_2$ ,  $\alpha(x, \cdot, \lambda)$  is called an elementary element of  $V_1$ . Denote by  $V$  the subspace of  $V_1$  which is spanned by the elementary elements of  $V_1$ . For  $x^* \in X^*$ ,  $\mu \in U_1$  and  $\lambda \in U_2$ , define

$$(1.2) \quad \alpha(x^*, \mu, \lambda) = (\mu - \lambda)^{-1}x^*.$$

For fixed  $x^* \in X^*$  and  $\mu \in U_1$ , call  $\alpha(x^*, \mu, \cdot)$  and elementary element of  $V_2$ . For  $f \in V_1$  and  $g \in V_2$ , with continuous extensions to  $\Gamma = \partial U_1 = \partial U_2$ , endow  $\Gamma$  with the clockwise orientation and ascertain that the bilinear functional

$$(1.3) \quad \Phi(f) = \langle f, g \rangle = \frac{1}{2\pi i} \int_{\Gamma} \langle f(\lambda), g(\lambda) \rangle d\lambda$$

is jointly continuous.

For  $g \in V_2$ , (1.3) defines a bounded linear functional  $\Phi$  on  $V$ , i.e.  $\Phi \in V^*$ . For  $f = \alpha(x, \cdot, \lambda)$ , one obtains

$$(1.4) \quad \Phi(f) = \frac{1}{2\pi i} \int_{\Gamma} (\mu - \lambda)^{-1} \langle x, g(\mu) \rangle d\mu = \langle x, g(\lambda) \rangle.$$

The last equality holds because  $g(\infty) = 0$ .

The proof of the following lemma is similar to that of [3, Lemma 9.1].

**1.1. Lemma.** *Let  $U_1, U_2$  be complementary simple sets. With  $V, V_1$  and  $W_i$  ( $i = 1, 2$ ), as defined above, there exists a linear manifold  $Y$  in  $W_2$  and a norm on  $Y$  such that*

- (i)  $Y$  is a Banach space isometrically isomorphic to  $V^*$ ;
- (ii)  $V_2 \subset Y$ ;
- (iii) the mappings  $V_2 \rightarrow Y$  and  $Y \rightarrow W_2$  are continuous;
- (iv) the inner product between  $V$  and  $V_2$ , defined by (1.3), can be extended to an inner product between  $V$  and  $Y$  in conjunction with the isometric isomorphism between  $Y$  and  $V^*$ , as asserted by (i).

## 2. SOME DUAL PROPERTIES

For an operator  $T \in C_d(X)$ , define an operator  $H$  on  $V$  by

$$D_H = \{f \in V: Tf(\mu) \in V\}, \quad (Hf)(\mu) = (\mu - T)f(\mu).$$

**2.1. Lemma.** *The operator  $H$  is closed and densely defined on  $V$ .*

*Proof.* For  $f = \alpha(x, \cdot, \lambda)$  with  $x \in D_T$ , one has  $f \in D_H$  and

$$(2.1) \quad (Hf)(\mu) = (\mu - \lambda)^{-1}(\mu - T)x = x + (\mu - \lambda)^{-1}(\lambda - T)x.$$

The linear span of all elementary elements being dense in  $V$ , the operator  $H$  is densely defined.

Let  $\{f_n\}$  be a sequence in  $D_H$  such that  $f_n \rightarrow f$  and  $Hf_n \rightarrow g$ , for some functions  $f$  and  $g$ .  $T$  being closed, it follows from

$$(Hf_n)(\mu) = (\mu - T)f_n(\mu),$$

that  $f \in D_H$  and

$$(Hf)(\mu) = (\mu - T)f(\mu) = g(\mu), \quad \mu \in \overline{U}_1.$$

Thus  $H$  is closed.  $\square$

The next lemma defines the dual  $H^*$  of  $H$ . Henceforth,  $g$  will denote a typical element of  $V^* = Y$ .

**2.2. Lemma.** *The dual operator of  $H$  is defined by*

$$(2.2) \quad (H^*g)(\lambda) = - \lim_{\lambda \rightarrow \infty} \lambda g(\lambda) + (\lambda - T^*)g(\lambda), \quad g \in D_{H^*}.$$

*Proof.* For  $f(\mu) = \alpha(x, \mu, \lambda)$  with  $x \in D_T$  and  $\lambda \in U_2$  fixed, and  $g \in D_{H^*}$ , (2.1), (1.3) and (1.4) imply

$$(2.3) \quad \begin{aligned} \langle f, H^*g \rangle &= \langle Hf, g \rangle = \langle x, g \rangle + \langle \alpha((\lambda - T)x, \cdot, \lambda), g \rangle \\ &= \langle x, g \rangle + \langle (\lambda - T)x, g(\lambda) \rangle, \end{aligned}$$

where  $x$ , as a function of  $\mu \in \overline{U}_1$ , is an element of  $V_1$ . It follows from

$$x = \frac{1}{2\pi i} \int_{\Gamma'} (\mu - \lambda)^{-1} x d\lambda = \frac{1}{2\pi i} \int_{\Gamma'} \alpha(x, \mu, \lambda) d\lambda,$$

that  $x \in V$ , where  $\Gamma'$  is a closed  $C^2$ -Jordan curve with the clockwise orientation that contains  $\Gamma$  in its interior. Furthermore, with the help of (1.4), one obtains

$$(2.4) \quad \begin{aligned} \langle x, g \rangle &= \frac{1}{2\pi i} \int_{\Gamma'} \langle \alpha(x, \cdot, \lambda), g \rangle d\lambda = \frac{1}{2\pi i} \int_{\Gamma'} \langle x, g(\lambda) \rangle d\lambda \\ &= \lim_{z \rightarrow \infty} z \left( - \frac{1}{2\pi i} \int_{\Gamma'} (\lambda - z)^{-1} \langle x, g(\lambda) \rangle d\lambda \right) = - \lim_{z \rightarrow \infty} z \langle x, g(z) \rangle \\ &= - \lim_{\lambda \rightarrow \infty} \lambda \langle x, g(\lambda) \rangle = \left\langle x, - \lim_{\lambda \rightarrow \infty} \lambda g(\lambda) \right\rangle. \end{aligned}$$

It follows from (2.3) and (2.4) that

$$(2.5) \quad \begin{aligned} \langle f, H^*g \rangle &= \left\langle x, - \lim_{\lambda \rightarrow \infty} \lambda g(\lambda) \right\rangle + \langle (\lambda - T)x, g(\lambda) \rangle \\ &= \left\langle x, - \lim_{\lambda \rightarrow \infty} \lambda g(\lambda) + (\lambda - T)^*g(\lambda) \right\rangle. \end{aligned}$$

In fact,  $f = \alpha(x, \cdot, \lambda)$  and since  $\langle f, H^*g \rangle$  and  $\langle x, -\lim_{\lambda \rightarrow \infty} \lambda g(\lambda) \rangle$  are bounded linear functionals of  $x$ , so is  $\langle (\lambda - T)x, g(\lambda) \rangle$ . Thus  $g(\lambda) \in D_{T^*}$ , for every  $\lambda \in U_2$  and hence the last equality of (2.5) holds. Now (2.5) combined with (1.3) and (1.4), gives  $\langle f, H^*g \rangle = \langle x, (H^*g)(\lambda) \rangle$  and hence  $H^*$  is expressed by (2.2).  $\square$

Define the map  $\tau: V^* \rightarrow X^*$  by  $\tau g = \lim_{\lambda \rightarrow \infty} \lambda g(\lambda)$ . Then  $H^*$ , given by (2.2), is expressed by

$$(2.6) \quad (H^*g)(\lambda) = -\tau g + (\lambda - T^*)g(\lambda).$$

**2.3. Lemma.** *Let  $x^* \in X^*$ . Then  $x^* \in D_{T^*}$  iff there exists  $g \in D_{H^*}$  such that  $\tau g = x^*$ .*

*Proof.* First, assume that there is  $g \in D_{H^*}$  such that  $\tau g = x^*$ . Since  $H^*g \in V^*$ , the following limit exists

$$\lim_{\lambda \rightarrow \infty} T^* \lambda g(\lambda) = \lim_{\lambda \rightarrow \infty} \lambda T^* g(\lambda).$$

Furthermore,  $\lim_{\lambda \rightarrow \infty} \lambda g(\lambda)$  also exists and since  $T$  is closed, we have

$$x^* = \tau g = \lim_{\lambda \rightarrow \infty} \lambda g(\lambda) \in D_{T^*}.$$

Conversely, for every  $x^* \in D_{T^*}$ , the corresponding elementary element  $\alpha(x^*, \mu, \cdot)$  with  $\mu \in U_1$  fixed, is in  $D_{H^*}$ . It follows from (1.2), that

$$\tau(-\alpha) = -\lim \lambda \alpha(x^*, \mu, \lambda) = x^*$$

and the proof reaches its conclusion by setting  $g = -\alpha$ .  $\square$

### 3. NORMS ON THE DUAL SPACES

We introduce the norm

$$\|(f_1, f_2)\|_\eta = (\eta \|f_1\|^2 + \|f_2\|^2)^{1/2}, \quad \eta > 0,$$

in  $V \oplus V$ . This induces the norm

$$\|(g_1, g_2)\|_\eta = (\eta^{-1} \|g_1\|^2 + \|g_2\|^2)^{1/2}$$

in  $V^* \oplus V^*$ . Let  $G(H)$  and  $G(H^*)$  be the graphs of  $H$  and  $H^*$ , respectively.  $G(H)$ , as a subspace of  $V \oplus V$ , is endowed with the norm

$$(3.1) \quad \|(f, Hf)\|_\eta = (\eta \|f\|^2 + \|Hf\|^2)^{1/2}.$$

It follows from

$$(G(H))^\perp = \nu G(H^*), \quad \text{where } \nu(g_1, g_2) = (-g_2, g_1),$$

that  $\nu G(H^*)$  is the dual of  $(V \oplus V)/G(H)$ . The latter is equipped with the norm

$$(3.2) \quad \|(f_1, f_2)^\wedge\|_\eta = \inf\{(\|f_1 - f\|^2 + \|f_2 - Hf\|^2)^{1/2} : f \in D_H\},$$

where  $(f_1, f_2)^\wedge$  denotes a typical element of  $(V \oplus V)/G(H)$ . To the norm (3.2), there corresponds the following norm in  $\nu G(H^*)$ :

$$\|(-H^*g, g)\|_\eta = (\eta^{-1}\|H^*g\|^2 + \|g\|^2)^{1/2}.$$

**3.1. Lemma.** *The norm*

$$(3.3) \quad \|x^*\|_{T^*} = (\|x^*\|^2 + \|Tx^*\|^2)^{1/2}$$

in  $D_{T^*}$  is equivalent to the norm

$$(3.4) \quad \|x^*\|_\eta = \inf\{(\eta^{-1}\|H^*g\|^2 + \|g\|^2)^{1/2} : \tau g = x^*\}.$$

Furthermore,  $D_{T^*}$  equipped with the norm (3.3) or (3.4) is the dual of a Banach space.

*Proof.* First, we prove that  $D_{T^*}$  endowed with the norm (3.4) is a Banach space. Let  $\{x_n^*\}$  be a Cauchy sequence with respect to the norm (3.4). Without loss of generality, we may suppose that

$$(3.5) \quad \sum_{n=0}^{\infty} \|x_{n+1}^* - x_n^*\|_\eta < \infty, \quad x_0 = 0.$$

For each  $x_n$ , we may choose  $g_n \in D_{H^*}$  such that

$$(3.6) \quad (\eta^{-1}\|H^*(g_{n+1} - g_n)\|^2 + \|g_{n+1} - g_n\|^2)^{1/2} \leq 2\|x_{n+1}^* - x_n^*\|_\eta$$

and  $\tau g_n = x_n^*$ . Relations (3.5) and (3.6) imply that both  $\{g_n\}$  and  $\{H^*g_n\}$  converge. Put  $g = \lim_{n \rightarrow \infty} g_n$ .  $H^*$  being closed, one has  $g \in D_{H^*}$  and  $H^*g = \lim_{n \rightarrow \infty} H^*g_n$ . Then Lemma 2.3 implies that  $x^* = \tau g \in D_{T^*}$ . Since

$$\|x_n^* - x^*\|_\eta \leq (\eta^{-1}\|H^*(g_n - g)\|^2 + \|g_n - g\|^2)^{1/2} \rightarrow 0 \quad n \rightarrow \infty,$$

it follows that  $D_{T^*}$ , endowed with the norm (3.4), is a Banach space.

To show that the norms (3.3) and (3.4) are equivalent, let  $x^* \in D_{T^*}$  and  $g = \alpha(x^*, \mu, \cdot)$  with  $\mu \in U_1$  fixed. Since  $\tau g = x^*$ , one has

$$(3.7) \quad \|x^*\|_\eta \leq (\eta^{-1}\|H^*g\|^2 + \|g\|^2)^{1/2} \\ \leq \frac{1}{\delta}(\eta^{-1}(|\mu| \cdot \|x^*\| + \|T^*x^*\|)^2 + \|x^*\|^2)^{1/2},$$

where  $\delta = \text{dist}(\mu, U_2)$ . In view of (3.7), there exists  $K_\eta > 0$  such that

$$(3.8) \quad \|x^*\|_\eta \leq K_\eta(\|x^*\|^2 + \|T^*x^*\|^2)^{1/2} = K_\eta\|x^*\|_{T^*}.$$

$D_{T^*}$  being complete with respect to both  $\|\cdot\|_\eta$  and  $\|\cdot\|_{T^*}$ , (3.8) implies that the two norms are equivalent.  $D_{T^*}$  equipped with the norm (3.3) is isometrically isomorphic to  $\nu G(T^*) (= G(T)^\perp)$ . Since  $\nu G(T^*)$  is the dual of  $X \oplus X/G(T)$ , so is  $D_{T^*}$ .  $\square$

$D_{T^*}$  equipped with either of the two norms (3.3), (3.4), will be denoted by  $D$ . To obtain a further property of  $\tau$ , we need the following.

**3.2. Lemma.** Let  $Y, Z$  be Banach spaces and let  $S$  be a bounded surjective map of  $Y$  onto  $Z$ . In  $Z$  we define the norm

$$(3.9) \quad \|z\|_S = \inf\{\|y\|: y \in Y, Sy = z\}, \quad z \in Z.$$

Then, the corresponding norm in the dual space  $Z^*$  is

$$(3.10) \quad \|z^*\|_{S^*} = \|S^*z^*\|, \quad z^* \in Z^*.$$

*Proof.* Let  $N(S)$  be the null space of  $S$ . Then  $N(S)^\perp$  is the dual of  $Y/N(S)$ . Let  $y_0 \in Y$ ,  $z = Sy_0$  and let  $\hat{y}_0$  be the equivalence class of  $y_0$  in  $Y/N(S)$ . In terms of the norm (3.9), one has

$$\|\hat{y}_0\| = \inf\{\|y_0 - w\|: w \in N(S)\} = \inf\{\|y\|: Sy = z\} = \|z\|_S.$$

The dual norm of  $\|\hat{y}_0\|$  in  $N(S)^\perp$  is the usual norm in  $Y^*$ , restricted to  $N(S)^\perp$ . Note that  $S^*$  is a surjective map from  $Z^*$  onto  $N(S)^\perp$ . Therefore, the corresponding norm of  $\|\cdot\|_S$  in  $Z^*$  is the one expressed by (3.10).  $\square$

We define an operator  $K$  from  $\nu G(H^*)$  into  $D_T$  by  $K(-H^*g, g) = \tau g$ . Let  $D^\#$  be the dual of  $D$ . Then  $K^\#$ , the dual of  $K$ , is an operator from  $D^\#$  into the dual of  $\nu G(H^*)$ , i.e. from  $D^\#$  into  $(V^{**} \oplus V^{**})/(\nu G(H^*))^\perp$ .

For every  $x \in X$ , define a continuous linear functional  $\psi$  on  $D$ , by

$$(3.11) \quad \psi(x^*) = \langle x, x^* \rangle, \quad x^* \in D.$$

**3.3. Lemma.** The linear functional  $\psi$  (3.11) is a zero functional only if  $x = 0$ .

*Proof.* Assume that  $\psi = 0$ . Then  $\langle x, x^* \rangle = 0$  for every  $x^* \in D$ . Thus, we have  $\langle 0, -T^*x^* \rangle + \langle x, x^* \rangle = 0$ ,  $x^* \in D$ , equivalently,  $(0, x) \perp \nu G(T^*)$ , i.e.  $(0, x) \in G(T)$ . Consequently,  $x = 0$ .  $\square$

In view of Lemma 3.3, we may consider  $X$  as a subset of  $D^\#$ . In the following, we shall have a closer look at  $K^\#x$  for  $x \in X$ .

For  $x^* \in D$  and fixed  $\mu \in U_1$ , put  $g = \alpha(x^*, \mu, \cdot)$ . Then, one obtains

$$\begin{aligned} \langle K^\#x(-H^*g, g) \rangle &= \langle x, K(-H^*g, g) \rangle = \langle x, \tau g \rangle = \langle x, x^* \rangle \\ &= \frac{1}{2\pi i} \int_\Gamma (\mu - \lambda)^{-1} \langle x, x^* \rangle d\lambda = \langle x, g \rangle, \end{aligned}$$

where  $\Gamma$  has a clockwise orientation.

We know that  $(0, x)$  is an element of  $X \oplus X$ . We may also consider  $(0, x)$  as an element of  $V \oplus V$  and hence  $(0, x)$  can be assumed to be an element of  $V^{**} \oplus V^{**}$ .

Thus, we have

$$\langle x, g \rangle = \langle (0, x), (-H^*g, g) \rangle = \langle (0, x)^\sim, (-H^*g, g) \rangle,$$

where  $(0, x)^\sim$  is the equivalence class of  $(0, x)$  in  $(V^{**} \oplus V^{**})/(\nu G(H^*))^\perp$ . Consequently,  $K^\#x = (0, x)^\sim$ .

Denote by  $(0, x)^\wedge$  the equivalence class of  $(0, x)$  in  $(V \oplus V)/G(H)$ .

**3.4. Lemma.** *Let  $J$  be the natural embedding of  $(V \oplus V)/G(H)$  into*

$$(V^{**} \oplus V^{**})/(\nu G(H^*))^\perp.$$

*Then  $J(0, x)^\wedge = (0, x)^\sim$ .*

*Proof.* For any  $(-H^*g, g) \in \nu G(H^*)$ , one has

$$\begin{aligned} \langle (0, x)^\wedge, (-H^*g, g) \rangle &= \langle (0, x), (-H^*g, g) \rangle = \langle (-H^*g, g), (0, x) \rangle \\ &= \langle (-H^*g, g), (0, x)^\sim \rangle. \end{aligned}$$

Note that while in  $\langle (0, x), (-H^*g, g) \rangle$ ,  $(0, x) \in V \oplus V$ ; in  $\langle (-H^*g, g), (0, x) \rangle$ ,  $(0, x)$  is considered an element of  $V^{**} \oplus V^{**}$ . It follows from the above equalities that  $J(0, x)^\wedge = (0, x)^\sim$ .  $\square$

In particular, Lemma 3.4 implies

$$(3.12) \quad \|(0, x)^\wedge\| = \|(0, x)^\sim\|.$$

On the other hand,  $(0, x)^\wedge = 0$  implies  $(0, x) \in G(H)$  and the latter implies  $x = 0$ . Accordingly, we may define the following norm on  $X$ :

$$(3.13) \quad \|x\|_\eta = \|(0, x)^\wedge\| = \inf\{(\eta\|f\|^2 + \|x - Hf\|^2)^{1/2} : f \in D_H\}.$$

In view of Lemma 3.4, we may consider  $K^\#x = (0, x)^\sim$  as a point of  $(V \oplus V)/G(H)$ .

**3.5. Lemma.** *The norm  $\|\cdot\|_\eta$ , defined by (3.13), is the restriction of the norm on  $D^\#$ .*

*Proof.* The space  $(V^{**} \oplus V^{**})/(\nu G(H^*))^\perp$  is the conjugate of  $\nu G(H^*)$  and  $K$  is an operator from  $\nu G(H^*)$  into  $D$ . It follows from Lemma 3.2, that the dual norm on  $D^\#$  is given by

$$(3.14) \quad \|x^\#\| = \|K^\#x^\#\|_{**}$$

where  $x^\# \in D^\#$  and  $\|\cdot\|_{**}$  is the norm on  $V^{**} \oplus V^{**}/[\nu G(H^*)]^\perp$ . If  $x^\# = x \in X$ , then the norm (3.14) becomes  $\|x\|_\eta = \|K^\#x\| = \|(0, x)^\sim\|$  and it follows from (3.12) that the restriction of the norm (3.14) to  $X$  is that given by (3.13).  $\square$

#### 4. A DUALITY PROPERTY OF SOME SPECTRAL-TYPE MANIFOLDS

Define the following linear manifolds in  $X$ :

$$N = \{x \in X : \text{for every } \varepsilon > 0, \text{ there exists } f \in D_H \text{ with } \|x - Hf\| < \varepsilon\},$$

$$M = \{x^* \in D : \text{there exists } g \in D_H \text{ such that } H^*g = 0, \tau g = x^*\}.$$

**4.1. Lemma.** *The manifolds  $N$  and  $M$  have the following characterizations:*

$$(4.1) \quad N = \{x \in X : \|x\|_\eta \rightarrow 0 \text{ as } \eta \rightarrow 0\},$$

$$(4.2) \quad M = \{x^* \in D : \|x^*\|_\eta \leq R \text{ for } n > 0 \text{ and } R \text{ depends on } x^*\}.$$

*Proof.* First, we establish (4.1). Let  $x \in N$ . Since, for every  $\varepsilon > 0$ , there is  $f \in D_H$  such that  $\|x - Hf\| < \varepsilon$ , it follows from (3.13) that  $\overline{\lim}_{\eta \rightarrow 0} \|x\|_\eta \leq \varepsilon$ .  $\varepsilon$  being arbitrary, it follows that  $\lim_{\eta \rightarrow 0} \|x\|_\eta = 0$ .

Conversely, suppose that  $\|x\|_\eta \rightarrow 0$  as  $\eta \rightarrow 0$ . Then, for every  $\varepsilon > 0$ , there exists  $\eta > 0$  such that  $\|x\|_\eta < \varepsilon$  and hence  $\|x - Hf\| < \varepsilon$  for some  $f \in D_H$ .

Next, we prove (4.2). It is a straightforward consequence of (3.4) that

$$M \subset \{x^* \in D: \|x^*\|_\eta \text{ is bounded for } \eta > 0\}.$$

Conversely, suppose that  $x^* \in D$  and  $\|x^*\|_\eta$  is bounded for  $\eta > 0$ , i.e. there exists  $R > 0$  such that

$$\inf\{(n\|H^*g\|^2 + \|g\|^2)^{1/2}, \tau g = x^*\} < R, \quad n = 1, 2, \dots$$

Then, for every  $n$ , there exists  $g_n \in D_{H^*}$  satisfying conditions

$$(4.3) \quad n\|H^*g_n\|^2 + \|g_n\|^2 \leq R^2 \quad \text{and} \quad \tau g_n = x^*.$$

In view of (4.3), the sequences  $\{g_n\}$  and  $\{H^*g_n\}$  are bounded and hence the sequence  $\{(H^*g_n, g_n)\}$  is bounded. Consequently,  $\{(-H^*g_n, g_n)\}$  has a cluster point  $(h, g)$  in  $V^* \oplus V^*$ , with respect to the weak\* topology of  $V^* \oplus V^*$ . Since  $\nu G(H^*)$  is closed with respect to the same topology, one has  $\{h, g\} \in \nu G(H^*)$ , i.e.  $h = -H^*g$ . It follows from  $\|H^*g_n\| \leq R^2/n$  that  $\|H^*g\| = 0$ .

On the other hand, for every  $x \in X$ ,  $K^\#x = (0, x)^\sim \in (V \oplus V)/G(H)$ . Therefore,

$$\langle x, x^* \rangle = \langle x, \tau g_n \rangle = \langle x, K(-H^*g_n, g_n) \rangle = \langle K^\#x, (-H^*g_n, g_n) \rangle.$$

Since  $(-H^*g, g)$  is also a cluster point of  $\{(-H^*g_n, g_n)\}$  in the weak\* topology of  $\nu G(H^*)$ , the latter being the dual space of  $(V \oplus V)/G(H)$ , we have

$$\langle x, x^* \rangle = \langle K^\#x, (-H^*g, g) \rangle = \langle x, K(-H^*g, g) \rangle = \langle x, \tau g \rangle.$$

Thus  $\tau g = x^*$  and hence  $x^* \in M$ . Expression (4.2) is obtained.  $\square$

**4.2. Theorem.**  $N$  and  $M$ , as defined above, are related by

$$N^\perp = \overline{M}^w,$$

where  $^w$  denotes the weak\* closure in  $X^*$ .

*Proof.* Let  $x \in N$  and  $x^* \in M$ . It follows from Lemmas 3.5 and 4.1, that

$$|\langle x, x^* \rangle| \leq \|x\|_\eta \cdot \|x^*\|_\eta \rightarrow 0 \quad (\text{as } \eta \rightarrow 0).$$

Therefore,  $N^\perp \supset \overline{M}^w$ .

Next, we prove the opposite inclusion. For  $x \notin N$  ( $x \in X$ ), Lemma 4.1 implies that there exists  $\eta_n \downarrow 0$  such that

$$(4.4) \quad \|x\|_{\eta_n} > C > 0$$

for some constant  $C$ . In view of (4.4), we can find  $x_n^* \in D$  such that  $\|x_n^*\|_{\eta_n} \leq 1$  and  $|\langle x, x_n^* \rangle| > C$ . The sequence  $\{\eta_n\}$  being nonincreasing, so is the norm

(3.4), i.e.  $\|x_n^*\|_{\eta_n} \leq \|x_n^*\|_{\eta_n}$ . Consequently,  $\{x_n^*\}$  is bounded in the norm  $\|\cdot\|_{\eta_n}$ -topology. For every  $n$ , there exists  $g_n \in D_{H^*}$  such that

$$\eta_n^{-1} \|H^* g_n\|^2 + \|g_n\|^2 \leq 2\|x_n^*\|_{\eta_n}^2.$$

Thus  $\{(x_n^*, g_n, H^* g_n)\}$  is bounded in  $D \oplus G(H^*)$ . By Lemma 3.1 and the previous paragraph,  $D \oplus G(H^*)$  is the dual of a Banach space and  $\{(x_n^*, g_n, H^* g_n)\}$  has a cluster point  $(x^*, g, H^* g)$  in the weak\* topology of  $D \oplus G(H^*)$ . Since (3.11) defines a continuous linear functional on  $D$  for every  $x \in X$ , it follows that  $x^*$  is also a cluster point of  $\{x_n^*\}$  in the weak\* topology of  $X^*$ . Now it follows from the inequalities

$$\langle x, x_n^* \rangle = \langle x, \tau g_n \rangle = \langle x, K(-H^* g_n, g_n) \rangle = \langle K^* x, (-H^* g_n, g_n) \rangle$$

that, for  $x \in X$ , one has

$$\langle x, x^* \rangle = \langle K^* x, (-H^* g, g) \rangle = \langle x, K(-H^* g, g) \rangle = \langle x, \tau g \rangle.$$

Thus  $x^* = \tau g$ . By the definition of  $M$ ,  $x^* \in M$ . Hence  $N \supset^\perp M$ , or equivalently,  $N^\perp \subset \overline{M}^w$ .  $\square$

### 5. THE MAIN THEOREM

We recall the definition of the central topic of this paper.

**5.1. Definition.** An operator  $T \in C(X)$  is said to have the spectral decomposition property (SDP) if, for every finite open cover  $\{G_i\}_{i=0}^n$  of  $\mathbb{C}$  (or  $\sigma(T)$ ), where  $G_0$  is a neighborhood of infinity (i.e. its complement  $G_0^c$  is compact in  $\mathbb{C}$ ), there exists a system  $\{Y_i\}_{i=0}^n$  of invariant subspaces under  $T$  satisfying the following conditions:

- (I)  $X_i \subset D_T$  if  $G_i$  ( $1 \leq i \leq n$ ) is relatively compact;
- (II)  $\sigma(T|X_i) \subset G_i$  ( $0 \leq i \leq n$ );
- (III)  $X = \sum_{i=0}^n X_i$ .

The theory based on this property is greatly simplified by the fact [3, Corollary 6.3] that  $T$  has the SDP iff it has the two-summand spectral decomposition property that corresponds to  $n = 1$ . The theory also involves the concept of the spectral manifold  $X(T, H) = \{x \in X: \sigma(x, T) \subset H\}$ , where  $H \subset \mathbb{C}$  and  $\sigma(x, T)$  is the local spectrum at  $x \in X$ , and the concept of the  $T$ -bounded spectral maximal space  $\Xi(T, F)$  for  $F \subset \mathbb{C}$  compact. The  $T$ -bounded spectral maximal space  $\Xi(T, F)$  is associated to  $X(T, F)$  [3, Theorem 4.34] by

$$X(T, F) = \Xi(T, F) \oplus X(T, \emptyset) \quad \text{and} \quad \sigma(T|\Xi(T, F)) = \sigma(T|X(T, F)).$$

The given operator  $T$  may enjoy two specific properties:

$T$  is said to have property  $(\beta)$  [1, Definition 8 and 3, Definition 5.5] if, for any sequence  $\{f_n: G \rightarrow D_T\}$  of analytic functions, the condition  $(\lambda - T)f_n(\lambda) \rightarrow 0$  (as  $n \rightarrow \infty$ ) in the strong topology of  $X$  and uniformly on every compact

subset of  $G$  implies that  $f_n(\lambda) \rightarrow 0$  in the strong topology of  $X$  and uniformly on every compact subset of  $G$ .

$T$  is said to have property  $(\kappa)$  [3, Definition 5.4] if  $T$  has the single valued extension property and  $X(T, F)$  is closed for every closed  $F$ .

Property  $(\beta)$  implies property  $(\kappa)$ , as follows from [3, Proposition 5.6].

**5.2. Lemma.** *Suppose that  $S \in C(X^*)$ . Then  $S$  is the dual of a closed and densely defined operator  $T \in C_d(X)$  iff  $G(S)$  is closed in the weak\* topology of  $X^* \oplus X^*$  and  $D_S$  is total.*

*Proof.* Only if: Suppose that  $S$  is the dual of  $T \in C_d(X)$ , i.e.  $S = T^*$ . The equality

$$\nu G(S) = \nu G(T^*) = (G(T))^\perp$$

implies  $G(S)$  is closed in the weak\* topology of  $X^* \oplus X^*$ . To prove that  $D_S$  is total, let  $x \in X$  and  $\langle x, x^* \rangle = 0$  for all  $x^* \in D_S$ . Then

$$\langle x, x^* \rangle = 0 = \langle 0, Sx^* \rangle$$

is equivalent to

$$0 \oplus x \in {}^\perp(\nu G(S)) = G(T)$$

and hence  $x = T(0) = 0$ , so  $D_S$  is total.

If: Assume that  $G(S)$  is closed in the weak\* topology of  $X^* \oplus X^*$  and  $D_S$  is total. Letting  $W = {}^\perp(\nu G(S))$ , one has  $W = {}^\perp \nu G(S)$ . Let  $0 \oplus y \in W$ . For every  $x^* \in D_S$ , one has  $0 \oplus y \perp (-Sx^*) \oplus x^*$ , or equivalently,

$$(5.1) \quad 0 = \langle 0, Sx^* \rangle = \langle y, x^* \rangle \quad \text{for all } x^* \in D_S.$$

$D_S$  being total, (5.1) implies that  $y = 0$  and hence  $W$  is the graph of an operator  $T$ .  $W$  being closed,  $T$  is a closed operator.

To show that  $T$  is densely defined, let  $x^* \in X^*$  satisfy condition

$$\langle x, x^* \rangle = 0 \quad \text{for all } x \in D_T.$$

Then

$$x \oplus Tx \perp x^* \oplus 0 \quad \text{for all } x \in D_T$$

and hence  $x^* \oplus 0 \in (G(T))^\perp = W^\perp = \nu G(S)$ . Therefore  $x^* = -S(0) = 0$  and hence  $T$  is densely defined.  $\square$

**5.3. Lemma.** *Suppose that  $T \in C(X)$  and  $Y$  is invariant under  $T$ . Then  $T/Y$  is closed iff  $G(T/Y)$  is topologically isomorphic to  $G(T)/G(T|Y)$ .*

*Proof.* Only if: Assume that  $T/Y$  is closed. For  $x \in D_T$ , the following mapping  $x \oplus Tx + G(T|Y) \rightarrow (x \oplus Y) + (Tx + Y)$  is bijective from  $G(T)/G(T|Y)$  onto  $G(T/Y)$ . It follows from the inequalities

$$\begin{aligned} \|x \oplus Tx + G(T|Y)\| &= \inf\{\|x \oplus Tx + y \oplus Ty\|: y \in D_{T|Y}\} \\ &\geq \inf\{\|(x + y_1) \oplus (Tx + y_2)\|: y_1, y_2 \in Y\} \\ &= \|(x + Y) \oplus (Tx + Y)\| \end{aligned}$$

and from the open mapping theorem that  $G(T/Y)$  and  $G(T)/G(T|Y)$  are topologically isomorphic.

If: Assume that  $G(T/Y)$  and  $G(T)/G(T|Y)$  are topologically isomorphic. Then  $G(T/Y)$  is a Banach space and hence it is closed in  $X/Y \oplus X/Y$ . Thus  $T/Y$  is closed.  $\square$

**5.4. Lemma.** *Given  $T \in C_d(X)$ , let  $Z \subset D_T$  be an invariant subspace under  $T$ . Then*

- (i)  $Z^\perp$  is invariant under  $T^*$ ;
- (ii)  $T^*/Z^\perp$  is the dual of  $T|Z$  iff  $T^*/Z^\perp$  is closed.

*Proof.* (i) is evident.

(ii): If  $T^*/Z^\perp$  is the dual of  $T|Z$  then clearly  $T^*/Z^\perp$  is closed. Conversely, assume that  $T^*/Z^\perp$  is closed. Then, it follows from Lemma 5.3 that  $G(T^*/Z^\perp)$  is topologically isomorphic to  $G(T^*)/G(T^*|Z^\perp)$ . The following equalities

$$\nu G(T^*) = (G(T))^\perp; \quad G(T^*|Z^\perp) = G(T^*) \cap (Z^\perp \oplus Z^\perp)$$

imply that both  $G(T^*)$  and  $G(T^*|Z^\perp)$  are closed in the weak\* topology of  $X^* \oplus X^*$ . Then, it follows easily that  $G(T^*/Z^\perp)$  is closed in the weak\* topology of  $X^*/Z^\perp \oplus X^*/Z^\perp$ .

It follows from Lemma 5.2 that  $D_{T^*}$  is total and hence  $D_{T^*/Z^\perp}$  is total. Quoting again Lemma 5.2, it follows that  $T^*/Z^\perp$  is the dual of a densely defined closed operator  $U \in C_d(Z)$ .

The assumption  $Z \subset D_T$  implies that  $T|Z$  is bounded. Let  $(x^*)^\wedge$  be the equivalence class of  $x^* \in X^*$  in  $X^*/Z^\perp$ . Then, for every  $x^* \in D_{T^*}$  and  $x \in D_U$ , one has

$$\begin{aligned} \langle Tx, x^* \rangle &= \langle x, T^*x^* \rangle = \langle x, (T^*/Z^\perp)(x^*)^\wedge \rangle \\ &= \langle Ux, (x^*)^\wedge \rangle = \langle Ux, x^* \rangle. \end{aligned}$$

Since  $D_{T^*}$  is total, (5.2) implies that  $Tx = Ux$ , for each  $x \in D_U$ . Since  $T|Z$  is bounded and  $U$  is a densely defined closed operator, it follows that  $U = T|Z$  and hence  $T^*/Z^\perp$  is the dual of  $T|Z$ .  $\square$

Now we are in a position to prove our main theorem.

**5.5. Theorem.** *Given  $T \in C_d(X)$ , the following assertion are equivalent:*

- (i)  $T$  has the SDP;
- (ii) for every pair of open disks  $G, H$  with  $\overline{G} \subset H$ , there exist invariant subspaces  $X_G$  and  $X_H$  such that

$$(5.3) \quad X = X_G + X_H; \quad X_H \subset D_T;$$

$$(5.4) \quad \sigma(T|X_H) \subset H \quad \text{and} \quad \sigma(T|X_G) \subset G^c;$$

- (iii) for every pair of open disks  $G, H$  with  $\overline{G} \subset H$ , there exist invariant subspaces  $Y, Z$  such that

$$(a) \quad \sigma(T|Y) \subset G^c; \quad T/Y \text{ is bounded and } \sigma(T/Y) \subset H;$$

- (b)  $Z \subset D_T$ ,  $\sigma(T|Z) \subset H$ ,  $T/Z$  is closed and  $\sigma(T/Z) \subset G^c$ ;
- (c)  $T^*/Z^\perp$  is closed;
- (iv) both  $T$  and  $T^*$  have property  $(\beta)$ ;
- (v)  $T$  has property  $(\beta)$  and  $T^*$  has property  $(\kappa)$ .

*Proof.* The proof will be carried out through the following scheme of implications:

$$\begin{aligned} (i) \Rightarrow (ii) \Rightarrow (iv) & \Rightarrow (v) \rightarrow (i). \\ (i) \Rightarrow (iii) \Rightarrow (iv) & \end{aligned}$$

(i)  $\Rightarrow$  (ii) is evident.

(i)  $\Rightarrow$  (iii): Given  $T$  with the SDP, let  $G, H$  be open disks with  $\overline{G} \subset H$  and let  $L$  be an open set satisfying inclusions  $\overline{G} \subset L \subset \overline{L} \subset H$ . For  $Y = X(T, G^c)$  and  $Z = \overline{\Xi}(T, L)$ , we have  $X = Y + Z$ . Then, in view of [3, Proposition 3.4 and Corollary 3.3], conditions (a) and (b) of (iii) are satisfied. Furthermore, it follows from (i) and [3, Theorem 9.8 (II,ii)], that

$$Z^\perp = X^*(T^*, L^c).$$

Consequently, (iii,c) follows from [3, Proposition 3.4].

(ii)  $\Rightarrow$  (iv): Let  $G$  and  $H$  be a pair of open disks with  $\overline{G} \subset H$ . There exists invariant subspaces  $X_G$  and  $X_H$  satisfying conditions (5.3) and (5.4). It follows from [3, Proposition 3.4] that  $T/X_G$  is bounded and

$$\sigma(T/X_G) \subset \sigma(T/X_H) \cup \sigma(T|X_G \cap X_H) \subset H.$$

Then [3, Theorem 5.8] implies that  $T$  has property  $(\beta)$ .

To show that  $T^*$  has property  $(\beta)$ , let  $\{f_n^*\}$  be a sequence of  $D_{T^*}$ -valued analytic functions defined on an open set  $G \subset \mathbb{C}$  such that

$$(\lambda - T^*)f_n^*(\lambda) \rightarrow 0 \quad (\text{as } n \rightarrow \infty)$$

uniformly on every compact subset of  $G$  in the strong topology of  $X^*$ . Without loss of generality, we may suppose that  $G = \{\lambda: |\lambda| < r\}$  for some  $r > 0$  and that  $K \subset G$  is compact. Let  $G_0$  and  $H_0$  be open disks satisfying inclusions

$$K \subset G_0 \subset \overline{G_0} \subset H_0 \subset \overline{H_0} \subset G.$$

Since  $T$  has property  $(\beta)$ , the subspaces  $X(T, G_0^c)$ ,  $\Xi(T, \overline{H_0})$  are defined. In view of conditions (5.3) and (5.4) applied to the open disks  $G_0, H_0$ , one obtains

$$(5.5) \quad X = X(T, G_0^c) + \Xi(T, \overline{H_0}).$$

Since  $K \subset \rho(T|X(T, G_0^c))$ , for  $\lambda \in K$  and  $x \in X(T, G_0^c)$ , one has

$$|\langle x, f_n^*(\lambda) \rangle| = |\langle R(\lambda; T|X(T, G_0^c))x, (\lambda - T^*)f_n^*(\lambda) \rangle| \leq M_0 \|(\lambda - T^*)f_n^*(\lambda)\| \cdot \|x\|,$$

where  $M_0 > 0$  is a constant independent of  $\lambda \in K$ . Then for every  $\varepsilon > 0$ , there exists  $N_0 > 0$  such that

$$(5.6) \quad |\langle x, f_n^*(\lambda) \rangle| \leq \varepsilon \|x\|, \quad \text{for all } \lambda \in K \text{ as } n > N_0.$$

Let  $C_0 = \{\lambda: |\lambda| = r_0\} \subset G$  with  $\overline{H}_0$  in the interior of the disk bounded by  $C_0$ , for some  $r_0 > 0$ . Then  $C_0 \subset \rho(T|\Xi(T, \overline{H}_0))$  and hence for  $\lambda \in C_0$  and  $x \in \Xi(T, \overline{H}_0)$  one has

$$|\langle x, f_n^*(\lambda) \rangle| = |\langle R(\lambda; T|\Xi(T, \overline{H}_0))x, (\lambda - T^*)f_n^*(\lambda) \rangle| \leq M_1 \|(\lambda - T^*)f_n^*(\lambda)\|,$$

where  $M_1 > 0$  is a constant independent of  $\lambda \in C_0$ . Then there is  $N_1$  such that

$$|\langle x, f_n^*(\lambda) \rangle| \leq \varepsilon \frac{\text{dist}(K, C_0)}{r_0} \|x\| \quad \text{for all } \lambda \in C_0 \text{ as } n > N_1.$$

It follows from the Cauchy integral formula that

$$(5.7) \quad |\langle x, f_n^*(\lambda) \rangle| \leq \frac{1}{2\pi} \int_{|\xi|=r_0} \frac{|\langle x, f_n^*(\lambda) \rangle|}{|\xi - \lambda|} |d\xi| \leq \varepsilon \|x\|,$$

for all  $\lambda \in K$  as  $n > N_1$ .

The decomposition (5.5) and the inequalities (5.6), (5.7) imply that there is a constant  $M > 0$  such that

$$|\langle x, f_n^*(\lambda) \rangle| \leq \varepsilon M \|x\| \quad \text{for all } x \in X, \lambda \in K \text{ as } n > \max\{N_0, N_1\}.$$

Thus it follows that  $\{f_n^*(\lambda)\}$  converges to zero uniformly on  $K$  in the strong topology of  $X^*$  and hence  $T^*$  has property  $(\beta)$ .

(iii)  $\Rightarrow$  (iv): Condition (iii,a) and [3, Theorem 5.8] imply that  $T$  has property  $(\beta)$ . By Lemma 5.4,  $Z^* = Z^\perp$  is invariant under  $T^*$  and then

$$\sigma(T^*|Z^\perp) = \sigma(T/Z) \subset G^c.$$

Again, by Lemma 5.4,  $T^*/Z^\perp$  is bounded and hence so is  $T|Z$ . We have

$$\sigma(T^*/Z^\perp) = \sigma(T|Z) \subset H.$$

Thus [3, Theorem 5.8] applies again and states that  $T^*$  has property  $(\beta)$ .

(iv)  $\Rightarrow$  (v) is evident.

(v)  $\Rightarrow$  (i): Let  $\{G_0, G_1\}$  be an open cover of  $\mathbb{C}$ , where  $G_0$  is a neighborhood of infinity and  $G_1$  is relatively compact. Let  $U_1, U_2$  be a couple of Cauchy domains with  $U_1$  bounded,  $U_2$  unbounded such that  $U_2 = (\overline{U_1})^c$ . Furthermore, we request that  $U_2$  verify inclusions

$$G_1^c \subset U_2 \subset \overline{U_2} \subset G_0.$$

Next, we define the linear manifolds  $N$  and  $M$  as in §4. We claim that the following inclusions hold:

$$(5.8) \quad \text{(a) } N \subset \overline{X(T, G_0)}, \quad \text{(b) } \overline{M}^w \subset \Xi^*(T^*, \overline{G_1}).$$

To prove (5.8a), let  $x \in N$ . For  $n = 1, 2, 3, \dots$  choose  $f_n \in D_H$  such that  $\|x - Hf_n\| < 1/n$ . Since  $T$  has property  $(\beta)$ ,  $\{f_n\}$  converges uniformly on compact sets in  $U_1$ . Put  $f(\lambda) = \lim_{n \rightarrow \infty} f_n(\lambda)$ , for  $\lambda \in U_1$ . Then  $f(\lambda) \in D_T$  and  $(\lambda - T)f(\lambda) = x, \lambda \in U_1$ . Consequently,

$$\sigma(x, T) \subset U_1^c = \overline{U_2} \subset G_0$$

and (5.8a) follows.

To prove (5.8b), let  $x^* \in M$ . There exists  $g \in D_{H^*}$  such that  $H^*g = 0$  and  $\tau g = x^*$ , or equivalently,

$$(\lambda - T^*)g(\lambda) = \tau g = x^*, \quad \lambda \in U_2.$$

Thus it follows that

$$\sigma(x^*, T^*) \subset U_2^c \subset \overline{G}_1$$

and hence  $x^* \in X^*(T^*, \overline{G}_1)$ . Since  $g(\lambda) \in V^*$  implies  $\lim_{\lambda \rightarrow \infty} \|g(\lambda)\| = 0$ , it follows from [3, Lemma 5.11] that  $x^* \in \Xi^*(T^*, \overline{G}_1)$ . Therefore,  $M \subset \Xi^*(T^*, \overline{G}_1)$ . Now [3, Theorem 9.4] implies that  $\Xi^*(T^*, \overline{G}_1)$  is weak\* closed and hence  $\overline{M}^w \subset \Xi^*(T^*, \overline{G}_1)$ . Now (5.8) and Theorem 4.2 imply

$$(5.9) \quad (X(T, G_0))^\perp \subset N^\perp = \overline{M}^w \subset \Xi^*(T^*, \overline{G}_1).$$

With  $G_0$  fixed, we may choose a sequence of open sets  $\{G_n\}$  such that  $\bigcap_{n=1}^\infty \overline{G}_n = G_0^c = F_0$  and  $\{G_0, G_n\}$  covers  $\mathbf{C}$  for every  $n$ . Then (5.9) implies that

$$(X(T, G_0))^\perp \subset \Xi^*(T^*, \overline{G}_n) \quad \text{for every } n.$$

Consequently,

$$(5.10) \quad (X(T, \overline{G}_0))^\perp \subset \bigcap_{n=1}^\infty \Xi^*(T^*, \overline{G}_n) = \Xi^*(T^*, F_0).$$

Combining (5.10) with the evident inclusion  $(X(T, G_0))^\perp \supset \Xi^*(T^*, F_0)$ , one finds

$$(5.11) \quad (X(T, G_0))^\perp = \Xi^*(T^*, F_0).$$

Since  $\Xi^*(T^*, F_0)$  is invariant under  $T^*$ , (5.11) implies that  $\overline{X(T, G_0)}$  is invariant under  $T$ . In fact, for every  $x \in X(T, \overline{G}_0) \cap D_T$  and  $x^* \in \Xi^*(T^*, F_0)$ , one has  $\langle Tx, x^* \rangle = \langle x, T^*x^* \rangle = 0$  so that  $\overline{X(T, G_0)}$  is invariant under  $T$ . Furthermore, we shall show that

$$(5.12) \quad \sigma(T|_{\overline{X(T, G_0)}}) \subset \overline{G}_0.$$

Let  $x \in \overline{X(T, G_0)}$  and choose a sequence  $\{x_n\} \subset X(T, G_0)$  such that  $x_n \rightarrow x$ . Let  $x_n(\cdot)$  denote the local resolvent of  $T$  at  $x_n$ . By property  $(\beta)$ , the convergence

$$(\lambda - T)x_n(\lambda) = x_n \rightarrow x, \quad \lambda \in (\overline{G}_0)^c$$

implies  $x_n(\lambda) \rightarrow f(\lambda)$  and  $(\lambda - T)f(\lambda) = x$ .

Therefore  $\sigma(x, T) \subset \overline{G}_0$ . On the other hand, for every  $\lambda \in (\overline{G}_0)^c$ , we have

$$\sigma(x_n(\lambda), T) = \sigma(x_n, T) \subset G_0,$$

so  $x_n(\lambda) \in X(T, G_0)$  and hence  $x(\lambda) \in \overline{X(T, G_0)}$  for  $\lambda \in (\overline{G}_0)^c$ . Then, by a known property [5, see also 3, Proposition 2.7], inclusion (5.12) follows.

Now we are in a position to show that  $T$  has the SDP. Let  $\{G_0, G_1\}$  be an open cover of  $\mathbf{C}$  with  $G_0$  a neighborhood of infinity and  $G_1$  relatively compact.

Let  $H_0$  be another open neighborhood of infinity such that  $\overline{G}_1 \cap \overline{H}_0 = \emptyset$  and  $\overline{H}_0 \subset G_0$ . Then  $\tilde{G}_0 = G_1 \cup H_0$  is a neighborhood of infinity and in virtue of (5.11) one writes

$$X(T, \tilde{G}_0)^\perp = \Xi^*(T^*, \tilde{F}_0),$$

where  $\tilde{F}_0 = (\tilde{G}_0)^c$  and both  $\Xi^*(T^*, F_0)$ ,  $\Xi^*(T^*, \tilde{F}_0)$  are closed in the weak\* topology of  $X^*$ . Similarly,  $\Xi^*(T^*, F_0 \cup \tilde{F}_0)$  is closed in the weak\* topology. Since  $F_0 \cap \tilde{F}_0 = \emptyset$  ( $F_0 = G_0^c$ ), we have

$$(5.13) \quad \Xi^*(T^*, F_0 \cup \tilde{F}_0) = \Xi^*(T^*, F_0) \oplus \Xi^*(T^*, \tilde{F}_0).$$

Set  $Z^* = \Xi^*(T^*, F_0 \cup \tilde{F}_0)$ .

Let  $x \in X$ ,  $x^* \in Z^*$  and denote by  $x_0^*$  the projection of  $x^*$  onto  $\Xi^*(T^*, F_0)$ , in conjunction with (5.13). The linear functional  $x_0$  on  $Z^*$ , defined by

$$(5.14) \quad \langle x_0, x^* \rangle = \langle x, x_0^* \rangle$$

is continuous in the weak\* topology. Use the Hahn-Banach theorem on locally convex spaces to extend  $x_0$  to a linear functional on  $X^*$ , that is continuous in the weak\* topology. Therefore  $x_0 \in X$ . Since the projection  $x_0^*$  of  $x^* \in \Xi^*(T^*, \tilde{F}_0)$  onto  $\Xi^*(T^*, F_0)$  is zero, it follows from (5.14) that  $\langle x_0, x^* \rangle = 0$  for  $x^* \in \Xi^*(T^*, \tilde{F}_0)$ . Thus,  $x_0 \in {}^\perp \Xi^*(T^*, \tilde{F}_0) = \overline{X(T, \tilde{G}_0)}$ . Put  $x_1 = x - x_0$  and for  $x^* \in \Xi^*(T^*, F_0)$ , use (5.14) to obtain  $\langle x_1, x^* \rangle = 0$ . Then  $x_1 \in {}^\perp \Xi^*(T^*, F_0) = \overline{X(T, G_0)}$ . Since  $x \in X$  is arbitrary, the representation  $x = x_0 + x_1$  with  $x_0 \in \overline{X(T, \tilde{G}_0)}$ ,  $x_1 \in \overline{X(T, G_0)}$  implies

$$(5.15) \quad X = \overline{X(T, G_0)} + \overline{X(T, \tilde{G}_0)}.$$

As regarding  $\overline{X(T, \tilde{G}_0)}$ , it follows from (5.12) that

$$\sigma(T|_{\overline{X(T, \tilde{G}_0)}}) \subset \overline{\tilde{G}_0} = \overline{G}_1 \cup \overline{H}_0.$$

Having  $\overline{G}_1 \cap \overline{H}_0 = \emptyset$  and  $G_1$  relatively compact, the functional calculus for closed operators produces the following decomposition

$$(5.16) \quad (a) \quad \overline{X(T, \tilde{G}_0)} = Y_1 \oplus Y_2, \quad (b) \quad Y_1 \subset D_T;$$

$$(5.17) \quad (a) \quad \sigma(T|_{Y_1}) \subset \overline{G}_1, \quad (b) \quad \sigma(T|_{Y_2}) \subset \overline{H}_0.$$

Since  $\overline{H}_0 \subset G_0$ ,  $Y_2 \subset \overline{X(T, G_0)}$ , (5.15) and (5.16) imply

$$(5.18) \quad X = Y_1 + \overline{X(T, G_0)}.$$

In view of (5.16b), (5.12), (5.17a) and (5.18),  $T$  has the SDP.

*Remark.* A more restrictive version of property  $(\beta)$  is used in [6, Lemma 4.6]. Given  $T \in C(X)$ , a function  $f: G \rightarrow D_T$  defined on an open subset  $G$  of the compactified complex plane  $C_\infty$ , is said to be  $T$ -analytic if both  $f$  and  $Tf$  are

analytic on  $G$ .  $T$  has property  $(\beta)$ , in this stronger version, if for any sequence of  $T$ -analytic functions  $\{f_n: G \rightarrow D_T\}$ , the condition  $(\lambda - T)f_n(\lambda) \rightarrow 0$  (as  $n \rightarrow \infty$ ) in the strong topology of  $X$  and uniformly on every compact subset of  $G$  implies that  $f_n(\lambda) \rightarrow 0$  in the strong topology of  $X$  and uniformly on every compact subset of  $G$ .

It follows from the definition of the operator  $H$  and Lemma 2.2 in §2 that both  $Tf(\mu)$  and  $T^*g(\lambda)$  are analytic. Consequently, Theorem 5.5 holds if we use the above-mentioned stronger version of property  $(\beta)$  in (iv) and (v).

#### REFERENCES

1. E. Bishop, *A duality theorem for an arbitrary operator*, Pacific J. Math. **9** (1959), 379–397.
2. N. Dunford and J. T. Schwartz, *Linear operators*, Part III, Wiley, New York, 1971.
3. I. Erdelyi and Wang Shengwang, *A local spectral theory for closed operators*, London Math. Soc. Lecture Note Ser. 105, Cambridge Univ. Press, 1985.
4. R. Lange and Wang Shengwang, *New criteria for a decomposable operator*, Illinois J. Math. (in print).
5. R. C. Sine, *Spectral decomposition of a class of operators*, Pacific J. Math. **14** (1964), 333–352.
6. Florian-Horia Vasilescu, *Analytic functional calculus and spectral decompositions*, Reidel, Dordrecht, Boston and London 1982.

DEPARTMENT OF MATHEMATICS, TEMPLE UNIVERSITY, PHILADELPHIA, PENNSYLVANIA 19122

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF PITTSBURGH, PITTSBURGH, PENNSYLVANIA 15260