

## THE MINIMAL NORMAL EXTENSION FOR $M_z$ ON THE HARDY SPACE OF A PLANAR REGION

JOHN SPRAKER

ABSTRACT. Multiplication by the independent variable on  $H^2(R)$  for  $R$  a bounded open region in the complex plane  $\mathbb{C}$  is a subnormal operator. This paper characterizes its minimal normal extension  $N$ . Any normal operator is determined by a scalar-valued spectral measure and a multiplicity function. It is a consequence of some standard operator theory that a scalar-valued spectral measure for  $N$  is harmonic measure for  $R$ ,  $\omega$ . This paper investigates the multiplicity function  $m$  for  $N$ . It is shown that  $m$  is bounded above by two  $\omega$ -a.e., and necessary and sufficient conditions are given for  $m$  to attain this upper bound on a set of positive harmonic measure. Examples are given which indicate the relationship between  $N$  and the boundary of  $R$ .

### 1. INTRODUCTION AND PRELIMINARIES

One of the major difficulties encountered in the study of subnormal operators is the shortage of examples. This paper examines the subnormal operator defined by multiplication by the independent variable on  $H^2(R)$ , the space of analytic functions  $f$  on a bounded region  $R$  such that  $|f|^2$  has a harmonic majorant. Recall that there exists a finite Borel measure  $\omega_a$  carried by  $\partial R$  with the property that for all  $f \in C_{\mathbb{R}}(\partial R)$ ,  $\hat{f}(a) = \int_{\partial R} \hat{f} d\omega_a$  where  $\hat{f}$  denotes the solution of the Dirichlet problem for  $R$  associated with  $f$ . This measure is known as *harmonic measure* for  $R$  at  $a$ . For more on this see [11, p. 332].

The following result is probably known, but seems to be documented only for the case of Dirichlet regions. Let  $\sigma$  be a normalized arc-length measure for the unit disk  $D$ .

**1.1. Theorem.** *Let  $R$  be a bounded region and let  $\phi: D \rightarrow R$  be a uniformization map from  $D$  onto  $R$  which takes 0 to  $a$ . Let  $\tilde{\phi}$  be a Borel function on  $\partial D$  which agrees  $\sigma$ -a.e. with the radial limits of  $\phi$ . Then  $\omega_a(E) = \sigma \circ \tilde{\phi}^{-1}(E)$  for any Borel set  $E$ .*

*Proof.* For  $f$  in  $C_{\mathbb{R}}(\partial R)$ , define  $\hat{f}$  by  $\hat{f}(x) = \int_{\partial R} f d\omega_x$ , where  $\hat{f}$  is a bounded harmonic function on  $R$ . Let  $A \subset \partial D$  have the following properties: (i)  $\sigma(A) = 1$ , (ii) the radial limits of  $\phi$  and  $\hat{f} \circ \phi$  exist on  $A$ , and (iii)

---

Received by the editors January 8, 1988. The contents of this paper were presented at the January 1988 meeting of the AMS in Atlanta.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 47B20; Secondary 31A15.

*Key words and phrases.* Harmonic measure, multiplication operator, minimal normal extension.

the radial limits of  $\phi$  agree with  $\tilde{\phi}$  on  $\widetilde{A}$ . The radial limit function associated with  $\hat{f} \circ \phi$  will be denoted by  $\hat{f} \circ \tilde{\phi}$ . Choose  $\lambda \in A$  such that  $\tilde{\phi}(\lambda)$  is a regular boundary point. Then as  $r \rightarrow 1$ , we have that  $\phi(r\lambda) \rightarrow \tilde{\phi}(\lambda)$ , which implies that  $\hat{f}(\phi(r\lambda)) \rightarrow f(\tilde{\phi}(\lambda))$ . By [27, p. 42] the set  $I$  of irregular points is a countable union of compact sets of capacity zero; and, by [14, p. 119]  $\tilde{\phi}^{-1}(I)$  has measure zero. Thus, we have that  $\hat{f}(\phi) = f \circ \tilde{\phi}$ , which holds  $\sigma$ -a.e. This allows us to define a bounded harmonic function  $g$  on  $D$  by

$$g(\alpha) = \int_{\partial D} f \circ \tilde{\phi}(\lambda) d\tau_{\alpha(\lambda)},$$

where  $\tau_{\alpha}$  is harmonic measure for  $D$  at  $\alpha$ . Since the radial limits of  $g$  and  $\hat{f} \circ \phi$  agree, then  $g = f \circ \phi$  and so  $g(0) = \hat{f}(\phi(0)) = \hat{f}(a)$ . Thus,

$$\int_{\partial R} f d\omega_a = \int_{\partial D} f \circ \tilde{\phi}(\lambda) d\sigma(\lambda). \quad \square$$

Hereafter let  $D$  be a region defined by the removal from the open unit disk of the closure of a countable pairwise disjoint collection of closed subdisks. Furthermore, we will assume that the set of accumulation points of the subdisks, those points with the property that every neighborhood of them contains one of the subdisks, is countable and that 0 lies in  $D$ . Let  $R$  be a bounded region in  $\mathbb{C}$  which is conformally equivalent to  $D$ . If  $R$  is simply connected, let  $D$  be the unit disk. Let  $\phi: D \rightarrow R$  be a one-to-one conformal map such that  $a = \phi(0)$ . Harmonic measure for  $D$  at 0 will be denoted by  $\tau_0$  and  $\omega_a$  will denote harmonic measure for  $R$  at  $a$ .

The operator obtained by composition with  $\phi$  is an isomorphism between  $H^2(R)$  and  $H^2(D)$ , and  $H^2(D)$  embeds onto a subspace  $H^2(\partial D)$  of  $L^2(\partial D, \tau_0)$  via nontangential limits. Let  $\tilde{\phi}$  be a Borel function on  $\partial D$  equal  $\tau_0$ -a.e. to the nontangential limits of  $\phi$ . It is easy to show that  $\omega_a = \tau_0 \circ \tilde{\phi}^{-1}$ . Hereafter  $S$  will denote the operator on  $H^2(R)$  defined by multiplication by the independent variable.  $S$  is unitarily equivalent to  $M_{\tilde{\phi}}$  on  $H^2(D)$  which has a normal extension, namely  $M_{\tilde{\phi}}$  on  $L^2(\partial D, \tau_0)$ . Thus,  $S$  is subnormal. In this paper we will characterize its minimal normal extension, denoted by  $N$ , and study the relationship between  $N$  and  $R$ . To do this we will make use of the fact that every normal operator is characterized by a scalar-valued spectral measure and a multiplicity function. It will be shown later (Theorem 2.4) that  $M_{\tilde{\phi}}$  on  $L^2(\partial D, \tau_0)$  is not only a normal extension of  $S$ , but, in fact, the minimal one  $N$ . Since  $\omega_a = \tau_0 \circ \tilde{\phi}^{-1}$ , Proposition 8.12 in [8, p. 296] shows that harmonic measure for  $R$  is a scalar-valued spectral measure for  $N$ .

## 2. THE MULTIPLICITY FUNCTION

What is the multiplicity function for  $N$ ? We will discover that the structure of  $\partial R$  determines the answer.

Abrahamse and Kriete have done some pioneering work on multiplicity theory for general multiplication operators. They have established a connection between the multiplicity function  $m$  and  $\tilde{\phi}^{-1}$ . In order to understand this, a definition is required.

**2.1. Definition.** A *disintegration* of  $\tau_0$  with respect to  $\tilde{\phi}$  is a function  $y \rightarrow \tau_0^y$  from the essential range of  $\tilde{\phi}$  into the set of regular Borel probability measures on  $D$  such that

- (a)  $\tau_0^y(\partial D \setminus \tilde{\phi}^{-1}(y)) = 0$ ,  $\omega_a$ -a.e., and
- (b) for every Borel set  $E$  in  $\partial D$ , the function  $y \rightarrow \tau_0^y(E)$  is  $\omega_a$ -measurable and  $\tau_0(E) = \int_{\partial R} \tau_0^y(E) d\omega_a(y)$ .

It may be shown (see [1]) that the disintegration is unique and depends only on the equivalence class of  $\tilde{\phi}$  in  $L^\infty(\partial D, \tau_0)$ . The following result, proven in [1, 3], shows the importance of this concept.

**2.2. Theorem.** Let  $X$  be a locally compact separable metric space, and let  $\mu$  be a Borel probability measure on  $X$ . If  $\psi$  is in  $L^\infty(\mu)$ ,  $\mu_y$  is a disintegration of  $\mu$  with respect to  $\psi$ , and  $m$  is the multiplicity function for the operator  $M_\psi$  on  $L^2(\mu)$ , then

$$m(y) = \# \text{supp}(\mu_y) \quad \text{for } \mu\text{-almost every } y$$

and

$$\text{supp}(\mu_y) \subset \text{cl}\{\psi^{-1}(y)\} \quad \text{for } \mu\text{-almost every } y.$$

The difficulty with using this theorem lies in the fact that it is often hard to obtain a disintegration of the measure. It does tell us, however, that  $m(y) \leq \#\psi^{-1}(y)$ ,  $\mu$ -a.e. This can be used to obtain an upper bound on the multiplicity function for  $N$  provided that we can show that  $N \cong M_{\tilde{\phi}}$  on  $L^2(\partial D, \tau_0)$  and provided that we can gain some knowledge of  $\tilde{\phi}^{-1}$ . To accomplish this we resurrect an old topological result of R. L. Moore [24]. Professor N. K. Nikolskii drew the author's attention to this result. It states that any pairwise disjoint collection of triods (homeomorphs of the letter "T"), in the complex plane can be at most countable. Now let us define a Borel set  $B$  contained in  $\partial R$  of full  $\omega_a$ -measure such that  $B$  is contained in the set of nontangential limits of  $\phi$  and  $\tilde{\phi}$  is equal to the nontangential limit function on  $\tilde{\phi}^{-1}(B)$ . The argument below is basically Nikolskii's and is a very interesting application of Moore's theorem.

**2.3. Lemma.**  $\{x \in B: \tilde{\phi}^{-1}(x) \geq 3\}$  is countable.

*Proof.* Suppose  $\#\tilde{\phi}^{-1}(x) \geq 3$ . It may be shown via a small topological argument that  $\tilde{\phi}^{-1}(x)$  lies in exactly one of the components of  $\partial D$ . Since the collection of accumulation points of the subdisks removed in the construction of  $D$  is countable, we may consider only those  $x$  for which  $\tilde{\phi}^{-1}(x)$  contains

no such accumulation point. Let  $\partial B(z, \delta)$  be the component of  $\partial D$  containing  $\tilde{\phi}^{-1}(x)$ , and choose  $a_1, a_2, a_3$  in  $\tilde{\phi}^{-1}(x)$ . Assume for the moment that  $B(z, \delta)$  is not the unit disk. One can always choose  $\varepsilon_x$  so that

$$\bigcup_{i=1}^3 \phi(\{a_i + t(a_i - z) : 0 < t \leq \varepsilon_x\}) \cup \{x\}$$

will form a triod. For  $y \neq x$  with  $\#\tilde{\phi}^{-1}(y) \geq 3$  and  $\tilde{\phi}^{-1}(y) \subset \partial B(z, \delta)$  any corresponding triod would be disjoint from the one above. This is true because  $\phi$  is one-to-one, and these triods are formed by taking the images under  $\phi$  of line segments extending radially from  $\partial B(z, \delta)$ . Such line segments are, of course, disjoint. Hence it is possible to choose a pairwise disjoint collection of triods which lies in a one-to-one correspondence with the set of points  $x$  such that  $\#\tilde{\phi}^{-1}(x) \geq 3$  and  $\tilde{\phi}^{-1}(x) \subset \partial B(z, \delta)$ . By Moore's result this set must be countable. For those  $x$  such that  $\tilde{\phi}^{-1}(x)$  is contained in the unit circle form triods of the form

$$\bigcup_{i=1}^3 \phi(\{ta_i : 1 - \varepsilon_x \leq t < 1\}) \cup \{x\}$$

and follow the same argument. Since the number of components of  $\partial D$  is countable, the result follows.  $\square$

The following theorem will tell us that  $N$ , the minimal normal extension of  $S$ , actually is  $M_{\tilde{\phi}}$  on  $L^2(\partial D, \tau)$ . The technique used in the proof was discovered by Professor John B. Conway [10].

**2.4. Theorem.**  $N \cong M_{\tilde{\phi}}$  on  $L^2(\partial D, \tau_0)$ .

*Proof.* Let  $A_0$  be the closed linear span of  $\{\tilde{\phi}^k f : f \in H^2(D), k \geq 0\}$  in  $L^2(\partial D, \tau_0)$ . It must be shown that  $A_0 = L^2(\partial D, \tau_0)$ . If  $h \in L^2(\partial D, \tau_0) \ominus A_0$ , then for all  $k \geq 0$  and for all  $f$  in  $H^2(D)$ ,

$$\begin{aligned} 0 &= \langle \tilde{\phi}^k f, h \rangle = \int_{\partial D} \tilde{\phi}^k f \bar{h} d\tau_0 \\ &= \int_{\partial R} \left[ \int_{\tilde{\phi}^{-1}(x)} \tilde{\phi}^k f \bar{h} d\lambda_x \right] d\omega_a(x) \end{aligned}$$

where  $\omega_a$  is harmonic measure for  $R$  and  $\lambda_x$  is a disintegration of  $\tau_0$  with respect to  $\tilde{\phi}$ . Clearly,  $f$  may be replaced by  $\tilde{\phi}^j f$  for all  $j \geq 0$ . Therefore,

$$\begin{aligned} 0 &= \int_{\partial R} \left[ \int_{\tilde{\phi}^{-1}(x)} \tilde{\phi}^k \tilde{\phi}^j f \bar{h} d\lambda_x \right] d\omega_a(x) \\ &= \int_{\partial R} \bar{x}^k x^j \int_{\tilde{\phi}^{-1}(x)} f \bar{h} d\lambda_x d\omega_a(x). \end{aligned}$$

This tells us that  $\int_{\tilde{\phi}^{-1}(x)} f \bar{h} d\lambda_x = 0$  for  $\omega_a$ -a.e.  $x$ , by the Weierstrass approximation theorem. Thus for each  $f$  in  $H^2(R)$  there exists  $G_r \subset R$  such

that  $x \notin G_f$  implies  $\int_{\tilde{\phi}^{-1}(x)} f \bar{h} d\lambda_x = 0$  and  $\omega_a(G_f) = 0$ . Let  $A = \{x \in \partial R : \#\tilde{\phi}^{-1}(x) \leq 2\}$ . A previous lemma shows us that  $\omega_a(A) = 1$ . Denote by  $P$  the collection of polynomials with complex rational coefficients, and let  $G = (\bigcup_{p \in P} G_p) \cup (\partial R \setminus A)$ . Clearly  $\omega_a(G) = 0$  since  $P$  is a countable set. By taking uniform limits, we find that for any polynomial  $p$  it follows that  $\int_{\tilde{\phi}^{-1}(x)} p \bar{h} d\lambda_x = 0$  for  $x \in G$ . Choose  $\alpha \in \tilde{\phi}^{-1}(\partial R \setminus G)$  and  $x \notin G$  such that  $\alpha \in \tilde{\phi}^{-1}(x)$ . There are now two cases to consider.

(1)  $\#\tilde{\phi}^{-1}(x) = 1$ . Here  $\lambda_x$  is a point mass, and since 1 is a polynomial  $0 = \int_{\alpha} \bar{h} d\lambda_x = \bar{h}(\alpha)$ .

(2)  $\#\tilde{\phi}^{-1}(x) = 2$ . Let  $\beta \in \tilde{\phi}^{-1}(x)$ ,  $\beta \neq \alpha$ . Then  $\int_{\tilde{\phi}^{-1}(x)} (z - \beta) h d\lambda_x = 0$ , which implies that  $\bar{h}(\alpha) = 0$  or that  $\lambda_x(\alpha) = 0$ . Let

$$B = \{\alpha \in \tilde{\phi}^{-1}(\partial R \setminus G) : h(\alpha) \neq 0\}.$$

$B$  is contained in

$$\{\alpha \in \tilde{\phi}^{-1}(\partial R \setminus G) : \text{if } \alpha \in \tilde{\phi}^{-1}(x) \text{ then } \lambda_x(\alpha) = 0\}$$

which implies that

$$\tau_0(B) = \int_{\partial D} \chi_B(x) d\tau_0(x) = \int_{\partial R} \left[ \int_{\tilde{\phi}^{-1}(x)} \chi_B d\tau_0 \right] d\omega_a(x) = 0$$

since for any  $x \notin G$ ,  $\lambda_x(B \cap \tilde{\phi}^{-1}(x)) = 0$  by definition of  $B$ . Therefore,  $\tau_0$ -a.e.  $h = 0$ . So  $A_0 = L^2(\partial D, \tau_0)$ .  $\square$

Since countable sets have harmonic measure zero, we have a nice upper bound on the multiplicity function.

**2.5. Theorem.**  $m(y) \leq 2 \omega_a$ -a.e.

*Proof.* The result follows from the previous theorem, Lemma 2.3, and the fact that  $m(y) \leq \#\tilde{\phi}^{-1}(y)$   $\omega_a$ -a.e.  $\square$

The technique in the proof of Theorem 2.5 immediately gives the next result, though there are certainly other methods of arriving at the same conclusion.

**2.6. Theorem.** *If  $R$  is bounded by a Jordan curve, then  $N$  is of multiplicity one.*

*Proof.* The hypothesis implies that all the prime ends of  $R$  are of the first kind and that every point of  $\partial R$  lies in the impression of exactly one prime end. This means that  $\phi$  is continuous to the boundary of  $D$  and is one-to-one there. Clearly,  $m(y) \leq \#\tilde{\phi}^{-1}(y) = 1$   $\omega_a$ -a.e.  $\square$

Hereafter we will let  $B \subset \partial R$  be a Borel set of full  $\omega_a$  measure contained in the nontangential limits of  $\phi$  such that  $\tilde{\phi}$  is equal to the nontangential limit function on  $\tilde{\phi}^{-1}(B)$  as before, and

- (a) for every  $x \in B$ ,  $\#\tilde{\phi}^{-1}(x) \leq 2$ ,
- (b) for every  $x \in B$ ,  $m(x) = \#\text{supp}(\tau_0^x)$  where  $\tau_0^x$  is a fixed disintegration of  $\tau_0$  with respect to  $\tilde{\phi}$ , and
- (c)  $B \subset \text{essran}(\tilde{\phi})$ .

Our next goal is to determine whether it is possible for the multiplicity function to attain its upper bound on a set of positive harmonic measure. With this in mind let  $A \subset \partial R$  be a Borel set with the following properties:

- (1)  $\omega_a(A) > 0$ ,
- (2) for every  $x \in A$ ,  $\#\tilde{\phi}^{-1}(x) = 2$ .

It may be shown using a measurable selection function  $s: A \rightarrow \tilde{\phi}^{-1}(A)$  [25, p. 515] that there exist disjoint Borel sets  $A_1 = s(A)$  and  $A_2 = \tilde{\phi}^{-1}(A) \setminus A_1$  contained in  $\partial D$  such that for every  $x \in A$  one element of  $\tilde{\phi}^{-1}(x)$  lies in each  $A_i$ . Define Borel measures  $\omega_i$ ,  $i = 1, 2$  on  $A$  by  $\omega_i(E) = \tau_0(A_i \cap \tilde{\phi}^{-1}(E))$ ,  $i = 1, 2$ .

**2.7. Theorem.** *The multiplicity function is equal to 2 almost everywhere on  $A$  if and only if  $\omega_a$ ,  $\omega_1$ , and  $\omega_2$  are pairwise mutually absolutely continuous.*

*Proof.* Suppose  $m = 2$  a.e. on  $A$ . Assume that  $\omega_1$  is not absolutely continuous with respect to  $\omega_2$ . Then there exists a Borel set  $A' \subset A$  such that  $\omega_1(A') \neq 0$  and  $\omega_2(A') = 0$ . Let us relabel and let  $A = A'$ ,  $\omega_i = \omega_i|_{A'}$ ,  $A_1 = s(A')$ , and let  $A_2 = \tilde{\phi}^{-1}(A') \setminus A_1$ . Clearly,

$$(2.7.1) \quad N \cong M_{\tilde{\phi}}|_{L^2(\partial D/(A_1 \cup A_2), \tau_0)} \oplus M_{\tilde{\phi}}|_{L^2(A_1, \tau_0)} \oplus M_{\tilde{\phi}}|_{L^2(A_2, \tau_0)}.$$

Theorem 1.1 tells us that  $\tau_0(A_2) = \omega_2(A) = 0$ , so

$$N \cong M_{\tilde{\phi}}|_{L^2(\partial D \setminus (A_1 \cup A_2), \tau_0)} \oplus M_{\tilde{\phi}}|_{L^2(A_1, \tau_0)}.$$

Since  $\tilde{\phi}$  is one-to-one on  $A_1$  and  $m(y) \leq \#\tilde{\phi}^{-1}(y)$  a.e., it is clear that  $M_{\tilde{\phi}}|_{L^2(A_1, \tau_0)}$  is a multiplicity one normal operator. Since  $\text{supp}(\tau_0^y) \subset \text{cl}(\phi^{-1}(y))$  almost everywhere and  $\#\tilde{\phi}^{-1}(y) \leq 2$  a.e. the multiplicity function of  $M_{\tilde{\phi}}|_{L^2(\partial D \setminus (A_1 \cup A_2), \tau_0)}$  is zero a.e. on  $A$ . This means that for almost all  $x \in A$ ,  $m(x) = 1$  contradicting the hypothesis.

Let us assume that  $\omega_1$  and  $\omega_2$  are m.a.c. Of course, it follows from the fact that  $\omega_a|_A = \omega_1 + \omega_2$  that  $\omega_i$  and  $\omega_a$  are m.a.c.,  $i = 1, 2$ . Define  $k_i(x)$ ,  $i = 1, 2$ , by  $k_i(x) = (d\omega_a/d\omega_i)^{1/2}(x)$  and define  $U: L^2(A, \omega_a)^{(2)} \rightarrow L^2(A_1 \cup A_2, \tau_0)$  by

$$U(f_1 \oplus f_2) = \begin{cases} k_1(\tilde{\phi}(x))f_1(\tilde{\phi}(x)), & x \in A_1, \\ k_2(\tilde{\phi}(x))f_2(\tilde{\phi}(x)), & x \in A_2. \end{cases}$$

Note that

$$\begin{aligned}
 \|U(f_1 \oplus f_2)\|^2 &= \int_{A_1 \cup A_2} |U(f_1 \oplus f_2)|^2 d\tau_0 \\
 &= \sum_{i=1}^2 \int_{A_i} (d\omega_a/d\omega_i(\tilde{\phi}(x))) |f_i(\tilde{\phi}(x))|^2 d\tau_0(x) \\
 &= \sum_{i=1}^2 \int_{A_i} |f_i(x)|^2 d\omega_a(x) \\
 &= \|f_1\|^2 + \|f_2\|^2 = \|(f_1 \oplus f_2)\|^2.
 \end{aligned}$$

Therefore,  $U$  is an isometry. We need to show that it is also surjective and therefore unitary.

First observe that since  $\tilde{\phi}$  is one-to-one on  $A_i$ ,  $i = 1, 2$ , then  $h_i(x) = \tilde{\phi}^{-1}(x) \cap A_i$  defines a function for  $i = 1, 2$ . Also, a one-to-one Borel image of a Borel set is a Borel set (see [19, p. 489]). This means that  $h_i(x)$  is a Borel function. Let  $f \in L^2(A_1 \cup A_2, \tau_0)$  be a Borel function. Since  $h_i(x)$  is Borel, the following holds:

$$f(x) = U(1/k_1(x))f(h_1(x)) \oplus 1/k_2(x)f(h_2(x)).$$

Thus  $U$  is unitary. Let  $T = M_z$  on  $L^2(A, \omega_a)$ . Then

$$\begin{aligned}
 UT^{(2)}(f \oplus g) &= U(zf \oplus zg) \\
 &= \begin{cases} k_1(\tilde{\phi}(x))\tilde{\phi}(x)f(\tilde{\phi}(x)), & x \in A_1, \\ k_2(\tilde{\phi}(x))\tilde{\phi}(x)g(\tilde{\phi}(x)), & x \in A_2, \end{cases} \\
 &= M_{\tilde{\phi}}U(f \oplus g).
 \end{aligned}$$

Hence  $T^{(2)} \cong M_{\tilde{\phi}}|_{L^2(A_1 \cup A_2, \tau_0)}$ . Clearly  $T^{(2)}$  is a multiplicity two normal operator. By the decomposition in (2.7.1) we find that

$$N \cong T^{(2)} \oplus M_{\tilde{\phi}}|_{L^2(\partial D \setminus (A_1 \cup A_2), \tau_0)}.$$

Again the second term contributes no multiplicity to  $A$ . Therefore the result follows.  $\square$

The preceding theorem gives a necessary and sufficient measure-theoretic condition for the multiplicity function to attain its upper bound on a set of positive harmonic measure. This condition may in fact be used in a large number of examples. If  $R$  is the slit disk and  $A$  is some subinterval of the slit, it is clear that  $A_1$  and  $A_2$  may be taken to be subarcs of  $\partial D$ . A well-known result of F. and M. Riesz [30] may be used to show that  $\omega_i$  and arclength measure are mutually absolutely continuous, for  $i = 1, 2$ , so  $m$  is 2 a.e. on  $A$ . This example may be generalized to other simply connected regions which contain a subset  $V$  of their boundaries with the property that for all  $x \in V$  there exists  $\varepsilon_x > 0$

such that

- (1)  $\overline{B}(x, \varepsilon_x) \cap \partial R$  is a rectifiable Jordan crosscut of  $B(x, \varepsilon_x)$  and
- (2)  $B(x, \varepsilon_x) \setminus \partial R \subset R$ .

Such a set  $V$  will be called *slitlike*.

The condition in Theorem 2.7 is not always satisfied, however. Consider Figure 1. Begin with an open rectangle slit along the interval from 0 to 1. As in the construction of a Cantor set of positive linear measure, remove a middle interval of length smaller than  $\frac{1}{3}$ . Above the endpoints of this interval draw two vertical line segments of length  $\frac{1}{2}$  starting at  $a_1$  and  $a_2$  as shown. For the time being, connect  $a_1$  to  $a_2$  with a horizontal segment. Let  $\lambda_2$  be harmonic measure at the point  $a$  for the rectangle with vertices  $a_1, a_2, \alpha_1,$  and  $\alpha_2$ . Construct horizontal segments  $\Gamma_1$  and  $\Gamma_2$  of equal length so that

$$\lambda_2\{\overline{\alpha_1\alpha_2} \setminus (\Gamma_1 \cup \Gamma_2)\} < \frac{1}{2}.$$




FIGURE 1

Now remove the next two intervals in the construction of the aforementioned Cantor set and draw “branches” as shown. The four points  $b_1, b_2, b_3,$  and  $b_4$  illustrated lie above the endpoints of the intervals removed as before. Let  $\lambda_3$  be harmonic measure at the point  $a$  for the region enclosed by drawing line segments from  $b_1$  to  $b_2$  and from  $b_3$  and  $b_4$ . Construct horizontal segments  $\Gamma'_i, i = 1, 2, 3, 4,$  at the ends of the branches so that

$$\lambda_3\{\overline{b_1b_2} \setminus (\Gamma'_1 \cup \Gamma'_2) \cup \overline{b_3b_4} \setminus (\Gamma'_3 \cup \Gamma'_4)\} < \frac{1}{3}.$$

Continue in this manner. The branches will “converge” down to the Cantor set. Call the union of the branches the tree. The tree forms a simply connected domain, since any loop in it being compact must lie above the real axis, and if one slices the tree horizontally anywhere above the real line, then the subregion above the cut is a contractible space. It may also be shown that the tree is a

Jordan domain. This may be accomplished by either showing that all the prime ends are of the first kind and every point on the boundary of the tree lies in the impression of exactly one prime end or by observing that the connection of the boundary of the tree is destroyed by the removal of any two arbitrary points. That the latter statement is equivalent to being a Jordan domain is a result found in [26, p. 93]. Both of these arguments are fairly straightforward and are left to the reader. Let  $C$  denote the Cantor set. The harmonic measure of  $C$  with respect to the tree is majorized by each of

$$\lambda_2(\overline{\alpha_1\alpha_2}\setminus(\Gamma_1 \cup \Gamma_2)), \lambda_3\{\overline{b_1b_2}\setminus(\Gamma'_1 \cup \Gamma'_2) \cup \overline{b_3b_4}\setminus(\Gamma'_3 \cup \Gamma'_4)\}, \dots$$

Therefore it is zero. Now remove the segment from  $a_1$  to  $a_2$ , and let  $R$  be the open region illustrated in Figure 1. Since the segment from  $a_1$  to  $a_2$ , a subarc of the boundary of the tree, was removed,  $R$  is an open rectangle minus two Jordan arcs which intersect only at 0. Thus  $R$  is simply connected. Define  $\phi$ ,  $\tilde{\phi}$ , and  $\omega_a$  as before. By constructing a rectifiable Jordan subdomain of  $R$  below the real line whose boundary contains  $[0, 1]$  it may be shown that  $\omega_a(C) > 0$ , because  $C$  has positive arc-length measure. The result of F. and M. Riesz in [30] states that arclength and harmonic measure are mutually absolutely continuous for rectifiable Jordan domains. Also for almost every  $x$  in  $C$ ,  $\#\tilde{\phi}^{-1}(x) = 2$  since  $x$  corresponds to a prime end defined by a chain in the tree and one defined by a chain below the real line. Let  $A_1$  be those points in  $\partial D$  corresponding to prime ends of  $C$  defined by chains in the tree and let  $A_2 = \tilde{\phi}^{-1}(C)\setminus A_1$ . These sets are easily seen to be Borel. Define  $\omega_1$  and  $\omega_2$  as before. Since harmonic measure is carried between Jordan domains such as the tree and  $\tilde{\phi}^{-1}$  (tree) by conformal maps, and  $\omega_1$  and harmonic measure for the tree are certainly mutually absolutely continuous,  $\omega_1(C) = 0$ . Since  $\omega_a(C) > 0$  and  $\omega_a|_C = \omega_1 + \omega_2$ , it is clear that  $\omega_2(C) > 0$ . Therefore,  $\omega_1$  and  $\omega_2$  are not mutually absolutely continuous; so, there must exist  $V \subset C$  of positive harmonic measure such that the multiplicity function is equal to one almost everywhere on  $V$ . For another example see [20].

It is well known that a normal operator has multiplicity identically one if and only if it has a star-cyclic vector. This fact along with some of our previous examples may be used to answer some approximation theory questions.

**2.8. Theorem.** *If  $R$  is the slit disk, then the polynomials are not dense in  $H^2(R)$ . In fact,  $S = M_{\tilde{\phi}}$  on  $H^2(R)$  has no cyclic vector.*

*Proof.* It is clear by the minimality of  $N$  that

$$\begin{aligned} L^2(\partial D, \tau) &= \text{cls}\{N^{*n}f: n \geq 0, f \in H^2(D)\} \\ &= \text{cls}\{\tilde{\phi}^n f: n \geq 0, f \in H^2(D)\} \end{aligned}$$

where cls denotes the closed linear span. If the polynomials were dense in  $H^2(R)$ , then polynomials in  $\phi$  would be dense in  $H^2(D)$  since the map  $T: H^2(R) \rightarrow H^2(D)$  which takes  $f$  to  $f \circ \phi$  is an isometric isomorphism. This

means that

$$L^2(\partial D, \tau) = \text{cls}\{\bar{\phi}^n \bar{\phi}^m : n, m \geq 0\} = \text{cls}\{R1 : R \in C^*(N)\}.$$

In other words, one is a star-cyclic vector for  $N$ . This, of course, cannot be since the multiplicity function is two on the slit. If  $f$  is a cyclic vector for  $S$ , repeat the above argument for polynomials in  $(\phi)(f \circ \phi)$  to obtain a contradiction.  $\square$

Again we can generalize this to any region with a set of multiplicity two which has positive harmonic measure. Slitlike sets come to mind for instance. The advantage of this approach lies in the fact that one can tell by inspection that the polynomials are not dense. It is easy to recognize slits.

*Acknowledgements.* I wish to thank Professor John B. Conway for his constant guidance and excellent suggestions during the development of this paper, and Professor James B. Barksdale for the use of his time and word-processing facilities.

#### BIBLIOGRAPHY

1. M. B. Abrahamse, *Multiplication operators*, Lecture Notes in Math., vol. 693, Springer-Verlag, Berlin, 1978, pp. 17–36.
2. M. B. Abrahamse and R. G. Douglas, *A class of subnormal operators related to multiply-connected domains*, Adv. in Math. **19** (1976), 106–148.
3. M. B. Abrahamse and T. Kriete, *The spectral multiplicity of a multiplication operator*, Indiana Univ. Math. J. **22** (1973), 845–857.
4. L. Ahlfors, *Conformal invariants, topics in geometric function theory*, McGraw-Hill, New York, 1973.
5. M. Brelot, *On topologies and boundaries in potential theory*, Lecture Notes in Math., vol. 175, Springer-Verlag, Berlin, 1971.
6. C. Caratheodory, *Über die Begrenzung einfach Zusammenhängerder Gebiete*, Math. Ann. **73** (1913), 323–370.
7. E. F. Collingwood and A. J. Lohwater, *The theory of cluster sets*, Cambridge Univ. Press, New York, 1966.
8. J. B. Conway, *A course in functional analysis*, Springer-Verlag, New York, 1985.
9. ———, *Functions of one complex variable* (2nd ed.), Springer-Verlag, New York, 1986.
10. ———, *The minimal normal extension of a function of a subnormal operator*, Preprint, 1987.
11. ———, *Subnormal operators*, Pitman, 1981.
12. J. Dudziak, *The minimal normal extension problem for subnormal operators*, J. Funct. Anal. **65** (1986), 314–338.
13. P. L. Duren, *Theory of  $H^p$  space*, Academic Press, New York, 1970.
14. S. D. Fisher, *Function theory on planar domains*, Wiley, New York, 1983.
15. T. W. Gamelin, *Uniform algebras*, (2nd ed.), Chelsea, New York, 1984.
16. M. Hasumi, *Hardy classes on infinitely connected Riemann surfaces*, Lecture Notes in Math., vol. 1027, Springer-Verlag, Berlin, 1983.
17. L. L. Helms, *Introduction to potential theory*, Wiley-Interscience, New York, 1969.
18. T. L. Kriete, *An elementary approach to the multiplicity theory of multiplication operators*, Rocky Mountain J. Math. **16** (1986), 23–33.
19. K. Kuratowski, *Topology*, Vol. 1, Academic Press, New York, 1966.

20. A. J. Lohwater and W. Seidel, *An example in conformal mapping*, Duke Math J. **15** (1948), 137–143.
21. A. I. Markushevitch, *Theory of functions of a complex variable*, (2nd ed.), Chelsea, New York, 1985.
22. R. S. Martin, *Minimal positive harmonic functions*, Trans. Amer. Math. Soc. **49** (1941), 137–172.
23. J. McMillan and G. Piranian, *Compression and expansion of boundary sets*, Duke Math. J. **40** (1973), 599–605.
24. R. L. Moore, *Concerning triods in the plane and the junction points of plane continua*, Proc. Nat. Acad. Sci. U.S.A. **14** (1928), 85–88.
25. M. A. Naimark, *Normed rings*, Noordhoff, Groningen, 1964, pp. 515–519.
26. M. H. A. Newman, *Elements of the topology of plane sets of points*, (2nd ed.), Cambridge Univ. Press, New York, 1954.
27. M. Ohtsuka, *Dirichlet problem, extremal length and prime ends*, Van Nostrand Reinhold, New York, 1970.
28. A. Pfluger, *Lectures on conformal mapping*, Dept. of Math., Indiana University, 1969.
29. C. Pommerenke, *Univalent functions*, Vanderhoeck & Ruprecht, Gottingen, 1975.
30. F. Riesz and M. Riesz, *Über die Randwerte einer analytischen Funktion 4*, Congr. Scand. Math., Stockholm, 1916, pp. 27–44.
31. W. Rudin, *Analytic functions of class  $HP$* , Trans. Amer. Math. Soc. **78** (1955), 46–66.
32. M. Tsuji, *Potential theory in modern function theory*, (2nd ed.), Chelsea, New York, 1975.

DEPARTMENT OF MATHEMATICS, WESTERN KENTUCKY UNIVERSITY, BOWLING GREEN,  
KENTUCKY 42101