

ROOTS OF UNITY AND THE ADAMS-NOVIKOV SPECTRAL SEQUENCE FOR FORMAL A -MODULES

KEITH JOHNSON

ABSTRACT. The cohomology of a Hopf algebroid related to the Adams-Novikov spectral sequence for formal A -modules is studied in the special case in which A is the ring of integers in the field obtained by adjoining p th roots of unity to $\widehat{\mathbb{Q}}_p$, the p -adic numbers. Information about these cohomology groups is used to give new proofs of results about the E_2 term of the Adams spectral sequence based on 2-local complex K -theory, and about the odd primary Kervaire invariant elements in the usual Adams-Novikov spectral sequence.

One of the most powerful tools used in the computation of stable homotopy groups is the Adams-Novikov spectral sequence. The E_2 term of this spectral sequence is a certain Ext group derived from a universal formal group law. In [R3] the corresponding Ext group for a universal formal A -module, for A the ring of algebraic integers in an algebraic number field, K , or its p -adic completion, was introduced and certain conjectures about these groups were formulated. One of these conjectures (concerning the value of $\text{Ext}^{1,*}$) was confirmed in [J] using a Hopf algebroid (i.e., a generalized Hopf algebra in which the left and right units need not agree), $E_A T$, which generalizes the Hopf algebroid $K_* K$ of stable cooperations for complex K -theory. The present paper is concerned with the cohomology of $E_A T$ in the special case of $A = \widehat{\mathbb{Z}}_p[\zeta]$ where ζ is a p th root of unity and $\widehat{\mathbb{Z}}_p$ denotes the p -adic integers. We will show that in this case $E_A T$ is contained in an extension of Hopf algebroids

$$\widehat{E}_A T \xrightarrow{j} E_A T \xrightarrow{\rho} \overline{E}_A T$$

and that the cohomology of $\overline{E}_A T$ can be completely described. This provides us with information about the cohomology of $E_A T$ via the Cartan-Eilenberg spectral sequence associated to this extension.

Two applications of this result are presented. In the case $p = 2$, $E_A T$ can be identified with the 2-adic completion of the Hopf algebroid $K_* K_{(2)}$ of stable cooperations for 2-primary complex K -theory. In this case the cohomology

Received by the editors February 1, 1989.
1980 *Mathematics Subject Classification* (1985 Revision). Primary 55T25; Secondary 55N22, 14L05.

of $\widetilde{E}_A T$ can also be described, so that we can completely describe the Cartan-Eilenberg spectral sequence (there are no nontrivial differentials for dimensional reasons). We therefore obtain a new proof of the results in [R1, K], computing $H^*(K_* K_{(2)})$.

A second application is the construction of nontrivial elements in the classical Adams-Novikov spectral sequence based on BP , the Brown-Peterson spectrum, which is a summand of MU , the spectrum representing complex cobordism. In [R3] a map of Hopf algebroids

$$\Psi: VT = BP_* BP \rightarrow V_A T$$

was described (here $V_A T$ is the Hopf algebroid generalizing VT to the category of formal A -modules, and is constructed using isomorphisms of A -typical formal A -modules). Composing Ψ with the Conner-Floyd map

$$\Phi: V_A T \rightarrow E_A T$$

constructed in [J] and the map ρ , we have a map

$$\chi: VT \rightarrow \overline{E}_A T$$

from VT to a Hopf algebroid whose cohomology is known. We thus have a tool for identifying nonzero elements of $H^{**}VT$, the E_2 term of the classical Adams-Novikov spectral sequence. We apply this to give a new proof of Theorem 4 of [R1] concerning the odd primary Kervaire invariant elements.

1. AN EXTENSION CONTAINING $E_A T$

In this section, we recall the definition and some of the structure of $E_A T$. We describe the homogeneous components of $E_A T$ and construct two related Hopf algebroids, \widehat{C}_n and \overline{C}_n , with which we construct the extension described in the introduction. We conclude by computing the cohomology of \overline{C}_n .

The ring $A = \widehat{\mathbb{Z}}_p[\zeta]$ is the ring of integers in the field $K = \widehat{\mathbb{Q}}_p[\zeta]$, which is an extension of $\widehat{\mathbb{Q}}_p$ of degree $p - 1$. A has a unique prime ideal (π) whose generator may be taken to be $\pi = \zeta - 1$, and the residue field of A , i.e., $A/(\pi)$, is $\mathbb{Z}/p\mathbb{Z}$. p is totally ramified in A , with $(\pi)^{p-1} = (p)$.

Recall from [J] that the Hopf algebroid $(E_A, E_A T)$ is defined by

$$E_A = A[t, t^{-1}],$$

$$E_A T = \{f \in K[u, u^{-1}, v, v^{-1}] \mid f(at, bt) \in E_A, \text{ if } a, b \in A, a, b \equiv 1 \pmod{(\pi)}\}$$

and that $E_A, E_A T$ are graded with $\deg(t) = \deg(u) = \deg(v) = 2(p - 1)$. The structure maps for $(E_A, E_A T)$ are:

$$\begin{aligned} \eta_L(t) &= u, & \eta_R(t) &= v, \\ \psi(u) &= u \otimes 1, & \psi(v) &= 1 \otimes v, \\ c(u) &= v, & c(v) &= u, \\ \varepsilon(u) &= t, & \varepsilon(v) &= t. \end{aligned}$$

If we denote the homogeneous component of $E_A T$ of degree $2 \cdot n \cdot (p - 1)$ by $(E_A T)_n$ then we obtain a Hopf algebroid $(A, (E_A T)_n)$. Let us also define

$$C_n = \{f \in K[w, w^{-1}] \mid f(a) \in A \text{ if } a \in A, a \equiv 1 \pmod{\pi}\}.$$

(A, C_n) can be given the structure of a Hopf algebroid via the maps

$$\begin{aligned} \eta_L(1) &= 1, & \eta_R(1) &= w^n, \\ \psi(w) &= w \otimes w, & c(w) &= w^{-1}, & \varepsilon(w) &= 1. \end{aligned}$$

We may define a map $C_n \rightarrow (E_A T)_n$ by $f \mapsto u^n \cdot f(v/u)$ and it is straightforward to check that this defines an isomorphism of Hopf algebroids. Thus, in particular we have

$$H^{s, 2n \cdot (p-1)}(E_A T) \simeq H^s(C_n).$$

We will do most of our computations using C_n rather than $E_A T$, and write C in place of C_n if the choice of right unit is not relevant.

Let us also write $B = C \cap K[w]$. We may define a sequence of polynomials in B inductively by

$$q_0 = (w - 1)/\pi, \quad q_{i+1} = (q_i^q - q_i)/\pi.$$

Also, let us denote

$$q^I(w) = q_0^{i_0} \cdots q_m^{i_m}$$

if $I = (i_0, \dots, i_m)$ is a multi-index.

Lemma 1. *The polynomials $\{q^I \mid 0 \leq i_j < p, m = 0, 1, 2, \dots\}$ form a basis for B as an A -module.*

Proof. This is Proposition 7 of [J] (note that these polynomials are denoted there by f_i).

Corollary 2. *The polynomials $\{q_i \mid i = 0, 1, \dots\}$ generate C over $A[w, w^{-1}]$.*

It will be useful for us to have a slightly different generating set for C available in addition to this one. Define inductively

$$\tilde{q}_0 = (w - 1)/\pi, \quad \tilde{q}_1 = (w^p - 1)/\pi^{p+1}, \quad \tilde{q}_{i+1} = (\tilde{q}_i^p - \tilde{q}_i)/\pi$$

and

$$\tilde{q}^I = \tilde{q}_0^{i_0} \cdots \tilde{q}_m^{i_m} \quad \text{if } I = (i_0, \dots, i_m).$$

Lemma 3. *The polynomials $\{\tilde{q}^I \mid 0 \leq i_j < p, m = 0, 1, 2, \dots\}$ form a basis for B as an A -module.*

Proof. Part of this lemma is, of course, that $\tilde{q}_i(w) \in B$. Since for any $a \in A$, $a^p - a \equiv 0 \pmod{\pi}$, it is sufficient for us to show that $\tilde{q}_1(w) \in B$. This, however, follows from [J, Lemma 17].

To see that this set forms a basis, note that the $(n + 1) \times (n + 1)$ matrix that expresses the polynomials \tilde{q}^I with $\sum i_j p^j \leq n$ as a linear combination of the polynomials q^I is triangular, with diagonal entries equal to 1. Thus, it is

invertible over A , and so the polynomials \tilde{q}^I span B . They are clearly linearly independent.

Corollary 4. *The polynomials $\{\tilde{q}_i | i = 0, 1, \dots\}$ generate C over $A[w, w^{-1}]$.*

Our interest in this second generating set is motivated by the fact, easily proved by induction, that $\tilde{q}_i(w)$ for $i \geq 1$ is a polynomial in w^p . If we denote $\tilde{C} = C \cap K[w^p, w^{-p}]$ and $\tilde{B} = \tilde{C} \cap K[w^p]$, then we have

Corollary 5. *The polynomials $\{\tilde{q}^I | i_0 = 0, 0 \leq i_j < p, m = 1, 2, \dots\}$ form a basis for \tilde{B} as an A -module.*

Corollary 6. *The polynomials $\{\tilde{q}_i | i = 1, 2, \dots\}$ generate \tilde{C} over $A[w^p, w^{-p}]$.*

A third algebra related to C and \tilde{C} is

$$\hat{C} = \{f \in K[x, x^{-1}] | f(a) \in A \text{ if } a \equiv 1 \pmod{\pi^{p+1}}\}.$$

We make (A, \hat{C}_n) into a Hopf algebroid by defining

$$\begin{aligned} \eta_L(1) &= 1, & \eta_R(1) &= x^n, \\ \psi(x) &= x \otimes x, & c(x) &= x^{-1}, & \varepsilon(x) &= 1. \end{aligned}$$

We also define $\hat{B} = \hat{C} \cap K[x]$.

The analogs of the polynomials q_i and \tilde{q}_i in this case are the polynomials defined by

$$\hat{q}_1 = (x - 1)/\pi^{p+1}, \quad \hat{q}_{i+1} = (\hat{q}_i^p - \hat{q}_i)/\pi.$$

We also use the notation $\hat{q}^I = \hat{q}_1^{i_1} \cdots \hat{q}_m^{i_m}$ if $I = (i_1, \dots, i_m)$. The analog of Lemmas 1 and 3 is

Lemma 7. *The polynomials $\{\hat{q}^I | 0 \leq i_j < p, m = 1, 2, \dots\}$ form a basis for \hat{B} as an A -module.*

Proof. The map $K[x] \rightarrow K[x]$ defined by $g(x) \mapsto g((x - 1)/\pi^{p+1})$ maps the algebra of polynomials with the property that $g(a) \in A$ if $a \in A$ isomorphically to \hat{B} . Since it also maps the basis for this former algebra constructed in [J, Proposition 7], onto the set $\{\hat{q}^I\}$, the latter must be a basis for \hat{B} .

Corollary 8. *The polynomials $\{\hat{q}_i | i = 1, 2, \dots\}$ generate \hat{C} over $A[x, x^{-1}]$.*

The connection between \hat{C} and the previous two Hopf algebras we have considered is given by

Proposition 9. *The map from \hat{C} to C that sends x to w^p is an injection of Hopf algebroids whose image is \tilde{C} .*

Proof. Since this map sends \hat{q}_i to \tilde{q}_i , the result is clear.

We next describe the Hopf algebroid $(\bar{E}_A, \bar{E}_A \bar{T})$, or rather we describe its homogeneous, degree $n \cdot 2 \cdot (p - 1)$ component, \bar{C}_n . Let \bar{C}_n denote the dual of the group algebra of the cyclic group of order p :

$$\bar{C}_n = A[\mathbb{Z}/p\mathbb{Z}]^* = \text{Hom}_A(A[\mathbb{Z}/p\mathbb{Z}], A).$$

The structure maps for \overline{C}_n are, using δ to denote a generator for $\mathbb{Z}/p\mathbb{Z}$,

$$\begin{aligned} \psi(f)(\delta^i \otimes \delta^j) &= f(\delta^{i+j}), \\ \eta_L(1)(\delta^i) &= 1, \quad \eta_R(1)(\delta^i) = \zeta^{ni}, \\ c(f)(\delta^i) &= f(\delta^{-i}), \quad \varepsilon(f) = f(1). \end{aligned}$$

Let us also define a map of Hopf algebroids $\rho: (A, C_n) \rightarrow (A, \overline{C}_n)$ by $\rho(f)(\delta^i) = f(\zeta^i)$.

The critical fact about ρ is

Lemma 10. ρ is a normal map of Hopf algebroids.

Proof. It is straightforward that ρ preserves the Hopf algebroid structure maps and so defines a map of Hopf algebroids; the question is whether it is normal. Referring to [R4, A1.1.10] we must verify that

$$C_n \square_{\overline{C}_n'} A = A \square_{\overline{C}_n'} C_n$$

where \square denotes the cotensor product and, for (A, Γ) a Hopf algebroid, Γ' is the associated Hopf algebra, defined by

$$\Gamma' = \Gamma / (\eta_R(a) - \eta_L(a) | a \in A).$$

In the case $\Gamma = \overline{C}_n$, this becomes

$$\overline{C}_n' = \begin{cases} \overline{C}_n & \text{if } p|n, \\ A & \text{if } (p, n) = 1. \end{cases}$$

To see this, note that if $p|n$, then

$$\begin{aligned} (\eta_R(a) - \eta_L(a))(\delta^i) &= a \cdot (\eta_R(1) - \eta_L(1))(\delta^i) \\ &= a \cdot (\zeta^{ni} - 1) = a \cdot (1 - 1) = 0 \end{aligned}$$

while if $(n, p) = 1$, then the ideal generated by $\eta_R(a) - \eta_L(a)$ is

$$I = \{\phi \in \overline{C}_n | \phi(1) = 0\}.$$

Thus, the map $\overline{C}_n' = \overline{C}_n / I \rightarrow A$ that sends ϕ to $\phi(1)$ is an isomorphism.

Since $C_n \square_A A = A \square_A C_n = C_n$, we may assume that $p|n$. The cotensor product $C_n \square_{\overline{C}_n'} A$ is defined to be the kernel of the map

$$C_n \simeq C_n \otimes_A A \rightarrow C_n \otimes_A \overline{C}_n \otimes_A A$$

which sends f to $(1 \otimes \rho)(\psi f) \otimes 1 - f \otimes \eta_L(1) \otimes 1$. This kernel consists of those elements $f \in \overline{C}_n$ for which

$$(1 \otimes \rho)(f(w \otimes w)) = f(w) \otimes 1$$

in $C_n \otimes_A \overline{C}_n$. These are precisely those elements $f \in \overline{C}_n$ of the form $f(w) = g(w^p)$. Similarly, $A \square_{\overline{C}_n'} C_n$ consists of those $f \in C_n$ for which

$$(\rho \otimes 1)(f(w \otimes w)) = \eta_R(1) \otimes f(w)$$

in $\overline{C}_n \otimes C_n$. Since $\eta_R(1) = 1$ in \overline{C}_n when $p|n$, we see that this also consists of those $f \in \overline{C}_n$ of the form $f(w) = g(w^p)$.

If we define the sub-Hopf algebroid (\tilde{A}, \tilde{C}_n) of (A, C_n) by

$$\tilde{A} = A \square_{\overline{C}_n} A, \quad \tilde{C}_n = A \square_{\overline{C}_n} C_n \square_{\overline{C}_n} A,$$

then, following [R4, A1.1.15], we have

Corollary 11. $(\tilde{A}, \tilde{C}_n) \xrightarrow{i} (A, C_n) \xrightarrow{\rho} (A, \overline{C}_n)$ is an extension of Hopf algebroids. (The fact that i is an inclusion is [R4, A1.1.14].)

For this to be useful we must describe (\tilde{A}, \tilde{C}_n) . As noted in the proof of [R4, A1.1.14], we have

$$\begin{aligned} \tilde{A} &= \{a \in A \mid \eta_L(a) = \eta_R(a) \text{ in } \overline{C}_n\}, \\ \tilde{C}_n &= \{f \in C_n \mid (\rho \otimes 1 \otimes \rho)\psi^2 f = \eta_L(1) \otimes f \otimes \eta_R(1)\}. \end{aligned}$$

and so

$$(\tilde{A}, \tilde{C}_n) = \begin{cases} 0 & \text{if } (n, p) = 1, \\ (A, \tilde{C}_n) \simeq (A, \widehat{C}_{n/p}) & \text{if } n|p. \end{cases}$$

The applications we have in mind for this extension involve the cohomology of C_n , which we approach via that of \tilde{C}_n and \overline{C}_n . We conclude this section, therefore, by recalling the cohomology of \overline{C}_n . Let us define two homomorphisms $S, T: \overline{C}_n \rightarrow \overline{C}_n$ by

$$S(f)(x) = f(\delta x) - f(x) \quad \text{and} \quad T(f)(x) = \sum_{i=0}^{p-1} f(\delta^i x).$$

A straightforward computation yields

Lemma 12. $0 \rightarrow A \xrightarrow{\eta_L} \overline{C}_n \xrightarrow{S} \overline{C}_n \xrightarrow{T} \overline{C}_n \xrightarrow{S} \dots$ is an injective resolution of A considered as a left C_n comodule.

Corollary 13. The cohomology of \overline{C}_n is given by

$$H^s(\overline{C}_n) = \begin{cases} A/\pi A, & s \text{ odd,} \\ 0, & s \text{ even,} \end{cases}$$

if $(n, p) = 1$, and by

$$H^s(\overline{C}_n) = \begin{cases} A, & s = 0, \\ A/pA, & s > 0, \text{ } s \text{ even,} \\ 0, & s \text{ odd,} \end{cases}$$

if $p|n$.

Proof. Applying the functor $A \square_{C_n} ()$ to the resolution of A and using the identification $A \square_{C_n} C_n = A$ gives the complexes

$$A \xrightarrow{\zeta^n - 1} A \xrightarrow{0} A \xrightarrow{\zeta^n - 1} A \xrightarrow{0} \dots$$

if $(n, p) = 1$, and

$$A \xrightarrow{0} A \xrightarrow{p} A \xrightarrow{0} A \xrightarrow{p} A \rightarrow \dots$$

if $p|n$.

2. APPLICATIONS

2.1. The cohomology of $K_*K_{(2)}$. If the prime p is chosen to be 2, then $A = \widehat{\mathbb{Z}}_2$ and C_n can be described as

$$C_n = \{f \in \widehat{\mathbb{Q}}_2[w, w^{-1}] \mid f(a) \in \widehat{\mathbb{Z}}_2 \text{ if } a \equiv 1 \pmod{2}\}.$$

The description of K_*K given in [AHS]

$$K_*K = \{f \in \mathbb{Q}[u, u^{-1}, v, v^{-1}] \mid f(at, bt) \in \mathbb{Z}[t, t^{-1}, 1/a, 1/b] \text{ if } a, b \in \mathbb{Z}, a, b \neq 0\}$$

shows that C_n can be identified with $(K_*K_{(2)})_n$ so that the E_2 term of the Adams spectral sequence based on 2-local complex K -theory has as its completion

$$E_2^{*,n} = H^*(C_n) = \text{Ext}_{C_n}^*(\widehat{\mathbb{Z}}_2, \widehat{\mathbb{Z}}_2).$$

The Cartan-Eilenberg spectral sequence, [R4, A1.3.14], allows us to describe these groups in terms of the cohomology of \widehat{C}_n and \overline{C}_n :

Proposition 14. *There is a spectral sequence converging to $\text{Ext}_{C_n}^*(\widehat{\mathbb{Z}}_2, \widehat{\mathbb{Z}}_2)$ whose E_2 term is $E_2^{s,t} = \text{Ext}_{\widehat{C}_n}^s(\widehat{\mathbb{Z}}_2, \text{Ext}_{\overline{C}_n}^t(\widehat{\mathbb{Z}}_2, \widehat{\mathbb{Z}}_2))$.*

Since $\text{Ext}_{\widehat{C}_n}^*(\widehat{\mathbb{Z}}_2, \widehat{\mathbb{Z}}_2)$ is described at the end of §1, we turn to describing $\text{Ext}_{\overline{C}_n}^*(\widehat{\mathbb{Z}}_2, \widehat{\mathbb{Z}}_2)$. The key to this description is the following injective resolution, which is the analog at the prime 2 of a resolution constructed for odd primes in [B, §7].

Lemma 15. *The sequence*

$$0 \rightarrow \widehat{\mathbb{Z}}_2 \xrightarrow{p_0} \widehat{C}_n \xrightarrow{p_1} \widehat{C}_n \xrightarrow{p_2} \widehat{\mathbb{Q}}_2 \rightarrow 0$$

defined by $p_1(f) = f(9w) - f(w)$ and $p_2(\sum a_i w^i) = a_0$ is an injective resolution of $\widehat{\mathbb{Z}}_2$.

(The left \widehat{C}_n comodule structure of $\widehat{\mathbb{Z}}_2$ and $\widehat{\mathbb{Q}}_2$ is that defined by $\eta_{L \cdot}$)

(The factor $9 = 2^3 + 1$ occurs here because it is a generator of

$$(1 + 2^3 \widehat{\mathbb{Z}}_2) / (1 + 2^n \widehat{\mathbb{Z}}_2)$$

for $n \geq 4$.)

Proof. If $p_1(\sum a_i w^i) = \sum_i a_i (9^i - 1) \cdot w^i = 0$ then $a_i = 0$ for $i \neq 0$ and the integrality condition for \widehat{C}_n shows that $a_0 \in \widehat{\mathbb{Z}}_2$. Thus, $\ker(p_1) = \text{Im}(p_0)$.

The fact that the polynomials $(w^{2^k} - 1)/2^{k+3}$ are in \widehat{C}_n shows that p_2 is surjective.

It remains to verify that $\ker(p_2) = \text{Im}(p_1)$. Suppose that

$$f = \sum_{i \neq 0} a_i w^i \in \ker(p_2).$$

For any $a \in \widehat{Q}_2$, the polynomial

$$g(w) = a + \sum_{i \neq 0} \frac{a_i w^i}{9^i - 1}$$

is mapped to f by p_1 . The question is whether a can be chosen so that $g \in \widehat{C}_n$. Choose a so that $g(1) = 0$. Since $p_1(g) = g(9w) - g(w) \in \widehat{C}_n$ it follows by induction on k that $g(9^k) \in A$ for any k , and this is enough to imply $g \in \widehat{C}_n$. To see this, first note that there exists m such that $2^m \cdot g \in \widehat{Z}_2[w, w^{-1}]$, and such that if $a, b \in \widehat{Z}_2$, then $g(b) \in \widehat{Z}_2$. However, $(1 + 2^3 \widehat{Z}_2)/(1 + 2^m \widehat{Z}_2)$ is cyclic, generated by 9. Thus, if $a \in 1 + 2^3 \widehat{Z}_2$, then $a \equiv 9^k \pmod{2^m}$ for some k and so $g(a) \in \widehat{Z}_2$.

Corollary 16.

(a)

$$\text{Ext}_{\widehat{C}_n}^s(\widehat{Z}_2, \widehat{Z}_2) = \begin{cases} \widehat{Z}_2, & s = 0, \\ \widehat{Q}_2/\widehat{Z}_2, & s = 2, \\ 0, & \text{otherwise}; \end{cases}$$

(b) for $n \neq 0$

$$\text{Ext}_{\widehat{C}_n}^s(\widehat{Z}_2, \widehat{Z}_2) = \begin{cases} \mathbb{Z}/2^{d(n)}\mathbb{Z}, & s = 1, \\ 0, & \text{otherwise}; \end{cases}$$

(c)

$$\text{Ext}_{\widehat{C}_n}^s(\widehat{Z}_2, \mathbb{Z}/2\mathbb{Z}) = \begin{cases} \mathbb{Z}/2\mathbb{Z}, & s = 0, 1, \\ 0, & \text{otherwise}. \end{cases}$$

Here $d(n)$ is the largest integer such that $2^{d(n)}$ divides $2^3 \cdot n$.

Proof. $\text{Ext}_{\widehat{C}_n}(\widehat{Z}_2, \widehat{Z}_2)$ is the cohomology of the complex

$$\widehat{Z}_2 \square_{\widehat{C}_n} \widehat{C}_n \rightarrow \widehat{Z}_2 \square_{\widehat{C}_n} \widehat{C}_n \rightarrow \widehat{Z}_2 \square_{\widehat{C}_n} \widehat{Q}_2 \rightarrow 0$$

If $n = 0$ this complex is

$$\widehat{Z}_2 \xrightarrow{0} \widehat{Z}_2 \xrightarrow{1} \widehat{Q}_2 \rightarrow 0$$

and, if $n \neq 0$

$$\widehat{Z}_2 \xrightarrow{9^n - 1} \widehat{Z}_2 \rightarrow 0 \rightarrow 0.$$

These account for (a) and (b), since the highest power of 2 dividing $9^n - 1$ is $2^{d(n)}$. For (c), we are interested in the cohomology of

$$\widehat{\mathbb{Z}}_2 \square_{\widehat{C}_n} \widehat{C}_n \otimes_A \mathbb{Z}/2\mathbb{Z} \rightarrow \widehat{\mathbb{Z}}_2 \square_{\widehat{C}_n} \widehat{C}_n \otimes_A \widehat{\mathbb{Z}}/2\mathbb{Z} \rightarrow \widehat{\mathbb{Z}}_2 \square_{\widehat{C}_n} \otimes_A \mathbb{Z}/2\mathbb{Z} \rightarrow 0.$$

This complex is, for any n ,

$$\mathbb{Z}/2\mathbb{Z} \xrightarrow{0} \mathbb{Z}/2\mathbb{Z} \rightarrow 0.$$

Combining these results with Proposition 14 and Corollary 13, we obtain

Corollary 17. *The E_2 term of the spectral sequence of Proposition 14 is*

$$E_2^{s,t} = \begin{cases} \widehat{\mathbb{Z}}_2, & \text{if } (s, t) = (0, 0), \ n = 0, \\ \widehat{\mathbb{Q}}_2/\widehat{\mathbb{Z}}_2, & \text{if } (s, t) = (2, 0), \ n = 0, \\ \mathbb{Z}/2^{d(m)}\mathbb{Z}, & \text{if } (s, t) = (1, 0), \ n = 2m = 0, \\ \mathbb{Z}/2\mathbb{Z}, & \text{if } (s, t) = (0, 2t'), \ n = 2m, \text{ or } (1, 2t'), \\ 0, & \text{otherwise.} \end{cases}$$

Corollary 18.

$$\text{Ext}_{K_*K_{(2)}}^{s,t}(\pi_*K, \pi_*K) = \begin{cases} \mathbb{Z}_{(2)}, & \text{if } (s, t) = (0, 0), \\ \mathbb{Z}/2^\infty\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}, & \text{if } (s, t) = (2, 0), \\ \mathbb{Z}/2^{d(m)}\mathbb{Z}, & \text{if } (s, t) = (1, 2m) \neq (1, 0), \\ \mathbb{Z}/2\mathbb{Z}, & \text{if } (s, t) = (s, 2t') \neq (2, 0), \ s \geq 2, \\ 0, & \text{otherwise.} \end{cases}$$

2.2. The odd primary Kervaire invariant elements. The Hopf algebroid $(V_A, V_A T)$ is constructed using isomorphisms of A -typical formal A -modules. If $A = \mathbb{Z}_{(p)}$, then one obtains $(V, VT) = (BP_*, BP_*BP)$, the Hopf algebroid of Brown-Peterson homology. If A is a $\mathbb{Z}_{(p)}$ algebra as in the case $A = \widehat{\mathbb{Z}}_p[\zeta]$ with which we are concerned, then a formal A -module is also a formal $\mathbb{Z}_{(p)}$ module. Thus, we obtain, as in [R3, 3.11], as map of Hopf algebroids

$$\Psi: (V, VT) \rightarrow (V_A, V_A T).$$

Composing this with the generalized Conner-Floyd map

$$\Phi: (V_A, V_A T) \rightarrow (E_A, E_A T)$$

of [J] and with $\rho: (E_A, E_A T) \rightarrow (E_A, \overline{E_A T})$ we obtain a map

$$\chi: (V, VT) \rightarrow (E_A, \overline{E_A T})$$

and so a map in cohomology

$$\chi^*: H^*VT \rightarrow H^*(E_A T).$$

We will show that a family of interesting elements in H^*VT , the odd primary Kervaire invariant elements, have nonzero image under this map.

Recall that (V, VT) has the description

$$V = \mathbb{Z}_{(p)}[v_1, v_2, \dots], \quad VT = V[t_1, t_2, \dots],$$

and that V, VT are graded with $\deg(v_i) = \deg(t_i) = 2(p^i - 1)$. The elements $h_0, b_i \in H^{1, 2(p-1)}(VT), H^{2, 2(p-1)p^{i+1}}(VT)$, respectively, are represented in the cobar complex of VT by $h_0 = [t_1]$ and

$$b_i = \frac{1}{p} \sum_{j=1}^{p^{i+1}-1} (p_j^{i+1}) [t_1^j \otimes t_1^{p^{i+1}-j}].$$

Our result is

Proposition 19. *All monomials in $h_0, b_i, i = 0, 1, 2, \dots$, have nonzero image in $H^*(E_A T)$ under χ^* .*

Proof. It is straightforward to describe the map of cobar complexes induced by χ . We also need, however, a method of identifying cohomologically nontrivial elements in the cobar complex of $E_A T$ or \overline{C}_n . For this we define a chain map from the cobar complex of \overline{C}_n to the complex described in §1, Lemma 12.

Recall from [R4, A1.2.11] that the cobar resolution of A as a \overline{C}_n comodule has as its s th term $\overline{C}_n \otimes (\ker(\varepsilon))^{\otimes s}$ and that the differential is given by

$$d(\gamma_0 \otimes \dots \otimes \gamma_s) = \sum_{i=0}^s (-1)^i \gamma_0 \otimes \dots \otimes \psi(\gamma_i) \otimes \dots \otimes \gamma_s + (-1)^{s+1} \gamma_0 \otimes \dots \otimes \gamma_s.$$

If we identify elements of

$$\overline{C}_n \otimes (\ker(\varepsilon))^{\otimes s} \subseteq \overline{C}_n^{\otimes s+1} = \text{Hom}_A(A[\mathbb{Z}/p\mathbb{Z}], A)^{\otimes s+1}$$

with multilinear maps from $A[\mathbb{Z}/p\mathbb{Z}]^{s+1}$ to A , then the differential becomes

$$df(w_0, \dots, w_{s+1}) = \sum_{i=0}^s (-1)^i f(w_0, \dots, w_i \cdot w_{i+1}, \dots, w_{s+1}) + (-1)^{s+1} f(w_0, \dots, w_s).$$

Using this identification we define a chain map, R , from the cobar resolution of A over \overline{C}_n to the resolution described in Lemma 12.

$$R(f)(w) = \begin{cases} f(w, \zeta) & \text{if } s = 1, \\ \sum_{i_1, \dots, i_{s-1/2}=1}^{p-1} f(w, \zeta^{i_1}, \zeta, \zeta^{i_2}, \dots, \zeta), & s \text{ odd}, \\ \sum_{i_1, \dots, i_{s/2}=1}^{p-1} f(w, \zeta, \zeta^{i_1}, \zeta, \zeta^{i_2}, \dots, \zeta^{i_{s/2}}), & s \text{ even}. \end{cases}$$

Applying $A \square_{\overline{C}_n} ()$ we obtain a map from the cobar complex of \overline{C}_n to the complex of Corollary 13. We denote this map by R as well. It is given by

$$R(f) = \begin{cases} f(\zeta), & s = 1, \\ \sum f(\zeta^{i_1}, \zeta, \zeta^{i_2}, \dots, \zeta), & s \text{ odd}, \\ \sum f(\zeta^{i_1}, \zeta, \zeta^{i_2}, \dots, \zeta^{i_{s/2}}), & s \text{ even}. \end{cases}$$

Under the composition $\Phi \circ \Psi$, the elements h_0 and b_i are mapped to $(w - 1)/\pi$ and

$$\frac{1}{p} \sum_{j=1}^{p^{i+1}-1} \binom{p^{i+1}}{j} \left(\frac{w-1}{\pi}\right)^j \otimes \left(\frac{w-1}{p}\right)^{p^{i+1}-j}$$

in the cobar complex of C_n . Under the composition $R \circ \rho$, these are mapped to 1 and

$$\frac{1}{p} \sum_{j=1}^{p-1} \left(\left(\frac{\zeta^j - 1}{\pi} + 1\right)^{p^{i+1}} - \left(\frac{\zeta^j - 1}{\pi}\right)^{p^{i+1}} - 1 \right),$$

respectively. We denote the latter element of A by k_i . This series of maps will send the monomial $h_0^e b_1^{i_1} \dots b_m^{i_m}$ to $k_1^{i_1} \dots k_m^{i_m}$. Showing that $k_i \not\equiv 0 \pmod{\pi}$ will, therefore, complete the proof of Proposition 19.

$$\begin{aligned} & \frac{1}{p} \sum_{j=1}^{p-1} \left(\left(\frac{\zeta^j - 1}{\pi} + 1\right)^{p^{i+1}} - \left(\frac{\zeta^j - 1}{\pi}\right)^{p^{i+1}} - 1 \right) \\ &= \frac{1}{p} \sum_{j=1}^{p-1} \sum_{k=1}^{p-1} \binom{p^{i+1}}{k} \left(\frac{\zeta^j - 1}{\pi}\right)^k \\ &\equiv \frac{1}{p} \sum_{j=1}^{p-1} \sum_{k=1}^{p-1} \binom{p^{m+1}}{k \cdot p^m} \left(\frac{\gamma^{j-1}}{\pi}\right)^{k \cdot p^m} \pmod{\pi} \\ &\equiv \frac{1}{p} \sum_{j=1}^{p-1} \sum_{k=1}^{p-1} \binom{p}{k} \left(\frac{\gamma^{j-1}}{\pi}\right)^k \pmod{\pi} \\ &\equiv \frac{1}{p} \sum_{j=1}^{p-1} \sum_{k=1}^{p-1} \binom{p}{k} j^k \pmod{\pi} \\ &= \frac{1}{p} \sum_{k=1}^{p-1} \binom{p}{k} \sum_{j=1}^{p-1} j^k \equiv \frac{-1}{p} \binom{p}{p-1} \pmod{\pi} \\ &\equiv -1 \pmod{\pi}. \end{aligned}$$

In these congruences, we have used the fact that $\binom{p^{i+1}}{k}$ is divisible by p^2 unless k is divisible by p^i , that $\binom{p^{i+1}}{kp^i} \equiv \binom{p}{k} \pmod{p}$, that since $\zeta = \pi + 1$, $((\zeta^i - 1)/\pi) \equiv i \pmod{\pi}$, and, finally, that $\sum_{j=1}^{p-1} j^k \equiv 0 \pmod{p}$ if $k < p - 1$ and that $\sum_{j=1}^{p-1} j^{p-1} \equiv -1 \pmod{p}$.

REFERENCES

- [AHS] J. F. Adams, A. S. Harris, and R. M. Switzer, *Hopf algebras of cooperations for real and complex K-theory*, Proc. London Math. Soc. (3) **23** (1971), 385–408.
- [B] A. K. Bousfield, *On the homotopy theory of K-local spectra at an odd prime*, Amer. J. Math. **107** (1985), 895–932.
- [J] K. Johnson, *The Conner-Floyd map for formal A-modules*, Trans. Amer. Math. Soc. **302** (1987), 319–332.
- [K] W. Kreuger, *The 2-primary K-theory Adams spectral sequence*, J. Pure Appl. Algebra **36** (1985), 143–158.
- [R1] D. C. Ravenel, *The non-existence of odd primary Arf invariant elements in stable homotopy theory*, Proc. Cambridge Philos. Soc. **83** (1978), 429–443.
- [R2] ———, *Localization with respect to certain periodic homology theories*, Amer. J. Math. **106** (1984), 351–414.
- [R3] ———, *Formal A-modules and the Adams Novikov spectral sequence*, J. Pure Appl. Algebra **32** (1984), 327–345.
- [R4] ———, *Complex cobordism and stable homotopy groups of spheres*, Academic Press, New York, 1986.

DEPARTMENT OF MATHEMATICS, STATISTICS AND COMPUTING SCIENCE, DALHOUSIE UNIVERSITY, HALIFAX, NOVA SCOTIA, CANADA