

## ON WEAK CONVERGENCE IN DYNAMICAL SYSTEMS TO SELF-SIMILAR PROCESSES WITH SPECTRAL REPRESENTATION

MICHAEL T. LACEY

**ABSTRACT.** Let  $(X, \mu, T)$  be an aperiodic dynamical system. Set  $S_m f = f + \dots + f \circ T^{m-1}$ , where  $f$  is a measurable function on  $X$ . Let  $Y(t)$  be one of a class of self-similar process with a "nice" spectral representation, for instance, either a fractional Brownian motion, a Hermite process, or a harmonizable fractional stable motion. We show that there is an  $f$  on  $X$ , and constants  $A_m \rightarrow +\infty$  so that

$$A_m^{-1} S_{[mt]} f \xrightarrow{d} Y(t),$$

the convergence being understood in the sense of weak convergence of all finite dimensional distributions in  $t$ .

### 1. INTRODUCTION

Let  $(X, \mathcal{A}, \mu, T)$  be a dynamical system, which is to say that  $(X, \mathcal{A}, \mu)$  is a Lebesgue space with  $\mu(X) = 1$ , and  $T: X \rightarrow X$  preserves  $\mu$ -measure. For a measurable function  $f$  on  $X$ , set  $S_m f = f + \dots + f \circ T^{m-1}$ . We are interested in the weak convergence of the sums  $S_m f$ , as a function of  $m$ . To be more precise, the problem we consider is this: Start with a stochastic process  $Y(t)$ . To avoid some trivial cases, assume that for all  $t > 0$ ,  $Y(\cdot)$  is continuous in probability at  $t$ , and the distribution of  $Y(t)$  is nondegenerate. We then ask: is there a measurable function  $f$  on  $X$ , and constants  $A_m \rightarrow +\infty$  so that

$$(1.1) \quad A_m^{-1} S_{[mt]} f \xrightarrow{d} Y(t)?$$

Here, convergence is in the sense of weak convergence of all finite dimensional distributions in  $t$ .

Previous results of this type have been in specific kinds of dynamical systems, for instance ergodic Markov shifts in [GL] and certain flows on manifolds in [S], to cite two well-known references. Moreover, the limiting process  $Y(t)$  has been a Brownian motion. The goal of this paper is to show that in any aperiodic dynamical system, (1.1) holds for a reasonably broad class of stochastic

Received by the editors October 12, 1989.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 60F17; Secondary 60F15, 28D05.

*Key words and phrases.* Aperiodic dynamical system; weak convergence; almost sure invariance principle; spectral representation; self-similar; fractional Brownian motion; Hermite processes; harmonizable fractional stable motion.

Research supported in part by an NSF Grant.

processes, which include the Brownian motion. These results generalize (and were motivated by) the interesting Central Limit Theorem of [BD].

Before stating our results, we note that (1.1) already imposes some necessary conditions on the stochastic process  $Y(t)$ . Namely,  $Y(0) = 0$ ;  $Y(t)$  has stationary increments; and [Lam]  $Y(t)$  enjoys the following time rescaling property: for some  $H > 0$ ,

$$(1.2) \quad a^{-H} Y(at) \stackrel{d}{=} Y(t), \quad a > 0.$$

Similar to (1.1), equality here is understood to be with respect to all finite dimensional distributions. Such a process is referred to in the literature as a self-similar of index  $H$ , stationary increment process. We shall refer to this briefly as  $H$ -ssi. The interest in these processes is due in part to their usefulness in modeling various physical phenomena, especially those with long-range dependence. We refer the reader to the surveys [T2 and M].

A last necessary condition for (1.1) is  $A_m = m^H L(m)$ , where  $L(m)$  is slowly varying at infinity. [Lam] again. Also, if  $Y(t)$  is square integrable, normalize it so that  $EY^2(1) = 1$ . Then  $EY^2(t) = t^{2H}$ , so that one has  $0 < H \leq 1$ . The case  $H = 1$  is trivial, as  $Y(t)$  can be realized as  $t \cdot Y(1)$ , and so is excluded from consideration in this paper.

We show that several kinds of processes can arise in (1.1). Three examples are the fractional Brownian motions, briefly fBm [MvN]; the Hermite processes, of index  $1/2 < H < 1$  [Dob, Maj]; and the harmonizable fractional stable motions [CM, T1]. The unifying element behind these three classes of processes is an especially nice spectral representation. In this section, we shall only state a result in the easiest case of the fBm as the other two cases are almost corollaries of the proof in that case. The modifications needed are sketched in §4 below.

A process  $B_H(t)$  is a standard fBm of index  $H$  if  $B_H(t)$  is  $H$ -ssi; is a real-valued Gaussian process; and  $EB_H(t)^2 = 1$ . Note that  $B_{1/2}(t)$  is a standard Brownian motion.

Recall that  $(X, \mathcal{A}, \mu, T)$  is aperiodic if for all  $A \in \mathcal{A}$  with  $\mu(A) > 0$ , and integers  $m$ , there is a measurable  $B \subset A$  so that  $\mu(T^m B \cap B) < \mu(B)$ .

**Theorem 1.1.** *Let  $(X, \mathcal{A}, \mu, T)$  be aperiodic, and  $0 < H < 1$ . Then there is an  $f \in L^2(\mu)$  so that*

$$m^{-H} S_{[m]} f \stackrel{d}{\Rightarrow} B_H(t).$$

In fact, we can prove a stronger almost everywhere result, see §3. By taking  $t = 1$  in this theorem, we obtain the Central Limit Theorem of [BD] mentioned above. We mention that the techniques of [BD] do *not* allow the deduction of a functional limit theorem, like this one.

One would like to apply the weak convergence results of this paper to dynamical systems, just as previous results, with the Brownian motion as a limit, have found applications (See [DK, PT, PUZ, and the survey D].) First however, information about the kinds of functions satisfying our theorems must be developed.

2. SPECTRAL REPRESENTATION

Fix  $0 < H < 1$ , and consider the stationary sequence  $\xi_k = B_H(k + 1) - B_H(k)$ , for  $k = 0, 1, 2, \dots$ . This sequence admits a representation

$$\xi_k = \int_{-\pi}^{\pi} e^{ik\lambda} W(d\lambda),$$

where  $W$  is a Gaussian random spectral measure, with control measure  $G$ .  $W(\cdot)$  can be defined this way: For all  $A, B \subset (0, \pi)$

- (i)  $W(A)$  is a complex Gaussian random variable of mean zero and variance  $G(A)$ ;
- (ii)  $EW(A)\overline{W(B)} = G(A \cap B)$ ;
- (iii)  $W(-A) = \overline{W(A)}$ ;
- (iv)  $W(\bigcup_1^n A_i) = \sum_{i=1}^n W(A_i)$  for mutually disjoint  $A_1, \dots, A_n$ .

Further, for any  $f \in L^2(G)$ ,

$$(2.1) \quad \left\| \int_{-\pi}^{\pi} f(\lambda) W(d\lambda) \right\|_2 = \left( \int_{-\pi}^{\pi} |f(\lambda)|^2 G(d\lambda) \right)^{1/2}.$$

We also note that as  $H < 1$ ,  $\frac{1}{m} B_H(m) \rightarrow 0$  a.s. so that in particular  $G(\{0\}) = 0$ .

Set  $K_m(\lambda) = \sum_{k=0}^{m-1} e^{ik\lambda} = (e^{im\lambda} - 1)/(e^{i\lambda} - 1)$ . Then

$$B_H(m) = \int_{-\pi}^{\pi} K_m(\lambda) W(d\lambda).$$

For the proof of Theorem 1.1, it is convenient to work with a discrete spectral measure, whose properties are collected together in the next lemma.

Let  $\{a_j: j \geq 0\}$  be decreasing to 0, with  $a_0 = \pi$ . Then the intervals  $I_j = [a_j, a_{j-1})$ ,  $j \geq 1$ , partition  $(0, \pi)$ . Let  $\lambda_j \in I_j$  be rational with respect to  $\pi$ . Define a new Gaussian random spectral measure  $V(\cdot)$  on  $(0, \pi)$  by  $V(\{\lambda_j\}) = W(I_j)$ . This measure is defined on the same probability space as  $W$ . Set  $l(m) = \lambda_{[\log m]}$ , for  $m \geq 4$ .

**Lemma 2.1.** *The intervals  $I_j$  above can be chosen so that the following three conditions hold.*

$$(2.2) \quad l(m) \geq (\log m)^{-1}, \quad m \geq 4,$$

$$(2.3) \quad \left| \int_{l(m)}^{\pi} K_m(\lambda) W(d\lambda) \right|, \quad \left| \int_{l(m)}^{\pi} K_m(\lambda) V(d\lambda) \right| = O((\log m)^2) \quad \text{a.s.},$$

and

$$(2.4) \quad \left| \int_0^{l(m)} K_m(\lambda) (W - V)(d\lambda) \right| = O((\log m)^2) \quad \text{a.s.}$$

*Proof.* To satisfy (2.2), choose the intervals  $I_j$  so that

$$I_{[\log m]} \subset ((\log m)^{-1}, \pi), \quad m \geq 4,$$

which can clearly be accomplished. In addition, require that

$$(2.5) \quad \sup_{j > \log m} \text{length}(I_j) \leq m^{-2}.$$

To see (2.3) for  $W(\cdot)$ , observe that by (2.1),

$$(2.6) \quad \left\| \int_{l(m)}^{\pi} K_m(\lambda) W(d\lambda) \right\|_2 \leq C \sup_{l(m) \leq \lambda < \pi} |K_m(\lambda)| \leq C(\log m)$$

Here, we have used (2.2) and the following inequality for  $K_m$ .

$$(2.7) \quad |K_m(\lambda)| \leq C/|\lambda|, \quad 0 < |\lambda| \leq \pi.$$

( $C$  is a positive constant, which might change from line to line.) Using the exponential-squared tail of Gaussian random variables, one easily sees that (2.6) implies (2.3), for  $W(\cdot)$ . The proof for  $V(\cdot)$  is completely similar.

To see (2.4), use (2.5) and (2.1) to see that

$$\begin{aligned} \left\| \int_0^{l(m)} K_m(\lambda)(W - V) d\lambda \right\|_2 &\leq C \sup_{j \geq \log m} \sup_{\lambda, \lambda' \in I_j} |K_m(\lambda) - K_m(\lambda')| \\ &\leq C \|K'_m\|_{\infty} \sup_{j \geq \log m} \text{length}(I_j) \leq C. \end{aligned}$$

This is more than enough to prove (2.4).  $\square$

### 3. ASIP

We can in fact prove a stronger form of Theorem 1.1 by suitably enlarging the dynamical system  $(X, \mathcal{A}, \mu, T)$ . For  $0 < H < 1$ , let  $B_H(t)$  be a standard fBm, with underlying probability space  $(\Omega, \mathcal{F}, P)$ . Define a new dynamical system  $(\bar{X}, \bar{\mathcal{A}}, \bar{\mu}, \bar{T})$  by

$$\bar{X} = X \otimes \Omega, \quad \bar{\mathcal{A}} = \mathcal{A} \otimes \mathcal{F}, \quad \bar{\mu} = \mu \otimes P, \quad \bar{T} = T \otimes \text{Id}_{\Omega}.$$

For a measurable function from  $\bar{X}$ , set  $\bar{S}_m f = \sum_{k=0}^{m-1} f \circ \bar{T}^k$ .

**Theorem 3.1.** *Let  $(X, \mathcal{A}, \mu, T)$  be aperiodic, and fix  $0 < H < 1$ . Then there is a standard fBm  $\bar{B}_H$  defined on  $(\bar{X}, \bar{\mathcal{A}}, \bar{\mu})$ , and an  $\bar{f} \in L^2(\bar{\mu})$  so that*

$$|\bar{S}_m \bar{f} - \bar{B}_H(m)| = O((\log m)^2) \quad \text{a.e. } (\bar{X})$$

This result is known in probability theory as an Almost Sure Invariance Principle (ASIP). Its strength lies in the fact that it not only proves Theorem 1.1 (on  $\bar{X}$ ), it also shows that a wide range of almost sure fluctuation results for  $B_H$  are inherited by the sums  $\bar{S}_m \bar{f}$ . This point is explained in the introduction to [PS], in the case of the Brownian motion. (Yet another almost sure result inherited by  $\bar{S}_m \bar{f}$  is seen from Theorem 1 of [LP].) The only known results of this type for fBm concern generalizations of the law of the iterated logarithm. See [TC, O, and Or].

Also, it appears that Theorem 3.1 is the first ASIP with the fBm as a limit. Every similar result known to the author approximates partial sums by a process with independent increments.

Noting that (1.1) is an isomorphic invariant of the dynamical system, we see that in many cases there is no loss of generality in working with  $\bar{X}$ . Moreover, it will be clear that much of our proof could be carried out on  $X$  alone, at the cost of obscuring the proof with more detail. See comments at the end of this section.

*Proof.* We divide the proof into three steps.

*Step 1.* Let  $J$  be an integer, and  $\varepsilon > 0$ . A set  $F \in \mathcal{A}$  is called an  $(J, \varepsilon)$  tower if  $F, TF, \dots, T^{J-1}F$  are disjoint, and  $\mu(\bigcup_{k=0}^{J-1} T^k F) > 1 - \varepsilon$ . Rohlin's lemma states that there is an  $(J, \varepsilon)$  tower in  $X$  for all pairs  $(J, \varepsilon)$ .

Let  $\lambda$  be rational with respect to  $\pi$ . With towers we will be able to construct a special function  $g$  so that  $S_m g(x)$  "acts like"  $K_m(\lambda)$  for an arbitrarily large number of  $m$ 's. (This is sufficient in view of (2.3) above.) It is done this way: let  $J$  and  $N$  be integers, and  $\varepsilon > 0$ . The second integer  $N$  should further satisfy  $e^{iN\lambda} = 1$ . Let  $F$  be a  $(JN, \varepsilon)$  tower. Call the following function special for  $(\lambda, F, J, N, \varepsilon)$ .

$$g(x) = \begin{cases} e^{ik\lambda}, & x \in T^k F, \quad 0 \leq k < JN, \\ 0, & \text{otherwise.} \end{cases}$$

The properties of  $g$  which we need are

$$(3.1) \quad S_m(g)(x) = g(x)K_m(\lambda), \quad x \in \bigcup_0^{(J-1)N-1} T^k F, \quad m \leq N,$$

$$S_N(g)(x) = 0, \quad x \in \bigcup_0^{(J-1)N-1} T^k F;$$

and

$$(3.2) \quad |S_m(g)(x)| \leq C/|\lambda|, \quad \text{all } m \text{ and } x.$$

The first property follows immediately from the definition of  $g$ ; the second follows from the choice of  $N$ ; and the last follows from the first two, and (2.7).

*Step 2.* We will now construct the function  $\bar{f}$ , and the process  $\bar{B}_H$ . Let  $\{\lambda_j\}$  be as in Lemma 2.1. We will need a special function for each  $\lambda_j$ , the parameters of which are determined in this way. Choose integers  $N_j$  so that

$$e^{iN_j\lambda_j} = 1, \quad \text{all } j$$

and

$$(3.3) \quad N_{[\log m]} > m, \quad m \geq 4.$$

Choose integers  $J_j$  so that  $\sum_j J_j^{-1} < +\infty$ . Let  $\varepsilon_j = J_j^{-1}$ , let  $F_j$  be a  $(J_j N_j, \varepsilon_j)$  tower, and  $g_j$  be special for  $(\lambda_j, F_j, J_j, N_j, \varepsilon_j)$ . Let

$$R_j = X \setminus \bigcup_0^{(J_j-1)N_j-1} T^k F_j.$$

Then

$$(3.4) \quad \sum_j \mu(R_j) < +\infty.$$

Let  $\bar{f} = \sum_j g_j \otimes V_j + \overline{g_j \otimes V_j}$ . Here we have set  $V_j = V(\{\lambda_j\})$ . ( $V(\cdot)$  is the spectral measure of Lemma 2.1.) To see that  $\bar{f} \in L^2(\bar{\mu})$ , note that the  $V_j$  are independent random variables, moreover,  $e^{i\lambda} V_j \stackrel{d}{=} V_j$  for  $0 \leq \lambda < 2\pi$ . Thus for each  $x \in X$ ,  $\bar{f}(x, \cdot)$  is a Gaussian series, with

$$(3.5) \quad E(\bar{f}(x, \cdot))^2 \leq 2 \sum_j E V_j^2 = E \left( \int_0^\pi V(d\lambda) \right)^2 < +\infty.$$

Thus, the sum  $\sum g_j(x) V_j$  converges a.s. for all  $x$ . Hence  $\bar{f}$  is well-defined a.e.  $(\bar{X})$  and is in  $L^2(\bar{X})$ .

We now define the process  $\bar{B}_H$ . It is enough to define a version of the spectral measure  $W$ , introduced in §2. In turn, it is enough to define  $W$  on  $(0, \pi)$ . Recall the intervals  $I_j$  of Lemma 2.1 partition  $(0, \pi)$ . Then for a measurable  $A \subset (0, \pi)$ , set

$$\bar{W}(A \cap I_j)(x, \omega) = \begin{cases} g_j(x) W(A \cap I_j)(\omega), & g_j(x) \neq 0, \\ W(A \cap I_j)(\omega), & g_j(x) = 0. \end{cases}$$

As  $W$  is a  $\sigma$ -finite measure almost surely, this defines  $\bar{W}$ . Also note that applying Lemma 2.1 to  $\bar{W}$ , we see that

$$\bar{V}_j = \bar{V}(\{\lambda_j\})(x, \omega) = \begin{cases} g_j(x) W(I_j)(\omega), & g_j(x) \neq 0, \\ W(I_j)(\omega), & g_j(x) = 0. \end{cases}$$

*Step 3.* We can now conclude the proof of Theorem 3.1. Set

$$h_m = \sum_{j \leq \log m} g_j \otimes V_j \quad \text{and} \quad h^m = \sum_{j > \log m} g_j \otimes V_j.$$

By Lemma 2.1 it suffices to prove the following:

$$(3.6) \quad \bar{S}_m(h_m) = O((\log m)^2) \quad \text{a.e. } (\bar{X}),$$

and

$$(3.7) \quad \bar{S}_m(h^m) - \sum_{j \geq \log m} K_m(\lambda_j) \bar{V}_j = O(1) \quad \text{a.e. } (\bar{X}).$$

For the first result, fix  $x \in X$ . Then  $\bar{S}_m h_m(x, \cdot)$  is a Gaussian random variable. Moreover,

$$(E|\bar{S}_m h_m(x, \cdot)|^2)^{1/2} \leq c \sup_{j \leq \log m} |S_m g_j(x)| \leq c(\log m),$$

by (2.2) and (3.2). Then (3.6) follows from this fact just as (2.3) follows from (2.6) above.

To see (3.7), for almost all  $x \in X$  there is an  $m$  so that  $x \in X \setminus \bigcup_{j \geq \log m} R_j$  by (3.4). Then, by (3.1) and (3.3), we have

$$\begin{aligned} \bar{S}_m h^m(x, \cdot) &= \sum_{j \geq \log m} S_m g_j(x) \otimes V_j \\ &= \sum_{j \geq \log m} g_j(x) K_m(\lambda_j) \otimes V_j \\ &= \sum_{j \geq \log m} K_m(\lambda_j) \bar{V}_j(x, \cdot), \end{aligned}$$

thus (3.7) holds.  $\square$

The proof above can be carried out on  $X$ , instead of  $\bar{X}$ . The only new step needed is to discretize the random variables  $V_j$ , to get random variables  $V'_j$ . Do this in such a way that

$$e^{ik\lambda_j} V'_j \stackrel{d}{=} V'_j, \quad k = 1, 2, \dots$$

As the  $V_j$  are independent random variables, the  $V'_j$  will be independent. Moreover, if the discretization is fine enough, one can work with the  $V'_j$ , up to a  $O((\log m)^2)$  a.e. error term. Then one constructs special functions  $g_j$  so that

$$\begin{aligned} g_j|_{F_j} &\stackrel{d}{=} V'_j, \\ g_j(T^k x) &= e^{ik\lambda_j} g_j(x), \quad x \in F_j, \quad 0 \leq k < J_j N_j, \end{aligned}$$

and  $g_j(x) = 0$  for  $x \in X \setminus \bigcup_0^{J_j N_j - 1} T^k F_j$ . Lemma 4 of [BD] should be used to define the  $g_j$ . The rest of the details are left to the interested reader.

#### 4. FURTHER PROCESSES

We discuss some extensions of the proof of Theorem 3.1 mentioned in the Introduction. These examples below are not complete, but suggest the range of processes that can arise as a limit. Other examples can be found in §3 of [GS].

**4.a. Hermite processes.** The Hermite processes are square-integrable ss si processes, which can be represented as multiple Wiener-Itô integrals. We follow [Dob and Maj] in our presentation below. Let  $M$  be a complex Gaussian random spectral measure, with control measure  $G$  satisfying

$$G(cA) = c^q G(A), \quad A \subset \mathbf{R},$$

for some  $q$ . Set  $\phi_t(x) = (e^{itx} - 1)/(ix)$ , fix an integer  $k$ , and let  $0 < q < 1/k$ . Then define

$$(4.1) \quad Y(t) = C_{qk} \int_{\mathbf{R}^k}' \phi_t(x_1 + \dots + x_k) |x_1|^q \dots |x_k|^q M(dx_1) \dots M(dx_k).$$

This is a multiple Wiener-Itô integral, where the prime on the integral sign indicates that integration is not performed over the hyperplanes  $|x_i| = |x_j|$ ,  $1 \leq i < j \leq k$ .  $Y(t)$  is  $H$ -ss si, with  $\frac{1}{2} < H = 1 - kq/2 < 1$ . For  $k = 1$ , it is a fBm, and for  $k \neq 1$ , it has a nonclassical distribution. (For  $k = 2$ , the characteristic function can be found in [R, T3].) The constant  $C_{qk}$  is chosen so that  $EZ^2(1) = 1$ .

**Theorem 4.1.** *For each  $k$  and  $0 < q < 1/k$  there is an  $f \in L^2(X)$  so that*

$$(4.2) \quad m^{1-kq/2} S_{[mt]} f \xrightarrow{d} Y(t).$$

We indicate how the proof would go. First, fix a stationary Gaussian sequence  $\xi_j$  so that

$$E\xi_j \xi_0 \sim j^{-q}, \quad j \rightarrow +\infty,$$

and write  $\xi_j = \int_{-\pi}^{\pi} e^{ijx} W(dx)$ . (The  $\xi_j$  can be taken to be  $B_H(j+1) - B_H(j)$ , where  $H = 1 - q/2$ .) It is then known that the “sums”

$$(4.3) \quad Y_m = C_{qk} \int_{(-\pi, \pi)^k} K_m(x_1 + \dots + x_k) W(dx_1) \dots W(dx_k)$$

satisfy (4.2). (See §4 of [Mae] for a nice summary of this; §8 of [Maj] for a proof.) A critical point in the proof of this fact, which we also use, is that the representations (4.1) and (4.3) are increasing concentrated near the origin. (This is how (2.2) and (2.3) are used in the proof of Theorem 3.1.)

The methods of §3 allow one to easily approximate (4.3): Using the notation of that section, set

$$\bar{f}_k = \sum'_{(j_1, \dots, j_k)} g_{j_1} \dots g_{j_k} \otimes V_{j_1} \dots V_{j_k} + \overline{g_{j_1} \dots g_{j_k} \otimes V_{j_1} \dots V_{j_k}}$$

where the prime on the sum indicates that the sum is *not* performed over the hyperdiagonals  $|j_u| = |j_v|$ ,  $1 \leq u < v \leq k$ .

Observe that by modifying the definition of the special functions  $g_j$ , one can insure that the difference

$$S_m(g_{j_1} \dots g_{j_k})(x) - g_{j_1}(x) \dots g_{j_k}(x) K_m(\lambda_{j_1} + \dots + \lambda_{j_k})$$

can be made arbitrarily small, in measure and in  $L^2(X)$ , for arbitrarily many  $m$ . (The only change is that the “junk sets”  $R_j$ , in (3.4) above, have to be made smaller in measure.)

The analogue of (2.1) is also needed. For any function  $f: \mathbf{R}^k \rightarrow \mathbf{C}$ , which is symmetric with respect to permutations of its arguments,

$$\begin{aligned} & \left\| \int_{\mathbf{R}^k} f(x_1, \dots, x_k) W(dx_1) \cdots W(dx_k) \right\|_2 \\ &= C_k \left( \int_{\mathbf{R}^k} |f(x_1, \dots, x_k)|^2 dx_1 \cdots dx_k \right)^{1/2}. \end{aligned}$$

With these observations, the proof of Theorem 3.1 is easily modified to yield Theorem 4.1.

**4.b. Harmonizable fractional stable motion.** We consider a class of ss si processes without second moment, obtained by replacing the Gaussian spectral measure used thus far, by a stable spectral measure. The resulting processes are special in that they do have a spectral representation, for such a representation need not exist for a stable process [H]. To define these processes, let  $l(\cdot)$  denote Lebesgue measure on  $\mathbf{R}$ , fix  $0 < \alpha \leq 2$ , and  $0 < H < 1$ . Let  $M(dx)$  be a complex-valued, isotropic (i.e. rotationally invariant) independently scattered  $\alpha$ -stable random measure on  $\mathbf{R}_+$ , with control measure  $l$ . A more detailed description of  $M$  is this: for any two Borel sets  $A, B \subset \mathbf{R}_+$ , with  $l(A), l(B) < +\infty$ ,  $M(A)$  and  $M(B)$  are complex-valued random variables satisfying

$$\begin{aligned} e^{it} M(A) &\stackrel{d}{=} M(A), \quad t \in \mathbf{R}, \\ E \exp\{it(\theta_1 \operatorname{Re}(M(A) - M(B)) + \theta_2 \operatorname{Im}(M(A) - M(B)))\} \\ &= \exp(-l(A\Delta B)|t(\theta_1, \theta_2)|^\alpha), \quad t \in \mathbf{R}, (\theta_1, \theta_2) \in \mathbf{R}^2, \end{aligned}$$

and  $M(A)$  and  $M(B)$  are independent if  $A \cap B = 0$ . Extended  $M$  to all of  $\mathbf{R}$  by  $M(-A) = \overline{M(A)}$ .

For  $0 < H < 1$ , set

$$Y_{\alpha, H}(t) = \int_{-\infty}^{\infty} \frac{e^{itx} - 1}{ix} |x|^{1-H-1/\alpha} M(dx).$$

This is an  $H$ -ss si process, and for  $\alpha = 2$  it is a fractional Brownian motion. For  $0 < \alpha < 2$ ,  $Y_{\alpha, H}(t)$  is a symmetric  $\alpha$ -stable random variable, so in particular  $P(|Y_{\alpha, H}(t)| > x) \sim C_{\alpha, H} t^{\alpha H} x^{-\alpha}$ , as  $x \rightarrow +\infty$ . Thus  $Y_{\alpha, H} \in \text{weak-}L^\alpha = L^{\alpha\infty}$ , and for  $0 < \alpha \leq 1$  is not integrable. Define the  $L^{\alpha\infty}$  "norm" by

$$\|Z\|_{\alpha\infty} = \sup_{z>0} z(\mu(|z| > z))^{1/\alpha}.$$

For  $\alpha \geq 1$ ,  $\|\cdot\|_{\alpha\infty}$  is equivalent to a norm; for  $0 < \alpha < 1$ , this fails.

**Theorem 4.2.** Let  $(X, \mathcal{A}, \mu, T)$  be aperiodic. Fix  $1 \leq \alpha < 2$ , and  $0 < H < 1$ . Then there is an  $f \in L_{\alpha\infty}(\mu)$  so that

$$m^{-H} S_{[m]} f \stackrel{d}{\Rightarrow} Y_{\alpha, H}(t).$$

The proof involves only trivial modifications of the proof of Theorem 3.1. First observe that  $Y_{\alpha, H}(m)$ ,  $m = 1, 2, \dots$ , admits a spectral representation

on  $(-\pi, \pi)$ :

$$Y_{\alpha, H}(m) = \int_{-\pi}^{\pi} K_m(\lambda)N(d\lambda),$$

with control measure  $\eta$ . Then  $\eta(\{0\}) = 0$ , and for all  $f \in L^\alpha(\eta)$ ,

$$\left\| \int_{-\pi}^{\pi} f(\lambda)N(d\lambda) \right\|_{\alpha\infty} = C_\alpha \left( \int_{-\pi}^{\pi} |f(\lambda)|^\alpha d\lambda \right)^{1/\alpha}.$$

With these observations, the remainder of the proof is easily checked.

**4.3. Stationary sequences.** One has the feeling that the range of limit behavior present in an aperiodic dynamical system should be rich. A rigorous result in this direction would be to show (1.1) for any square-integrable ss si process  $Y(t)$ . In fact, a result can be stated without any reference to self-similarity, which we do now. Let

$$\xi_k = \int_{-\pi}^{\pi} e^{ik\lambda} W(d\lambda)$$

be a square-integrable stationary sequence. Set  $S_m = \xi_0 + \dots + \xi_{m-1}$ , and  $s_m^2 = ES_m^2$ . Assume that  $W(\{0\}) = 0$  a.s., and  $m^\varepsilon \leq s_m^2 \leq m^{2-\varepsilon}$ , for some  $\varepsilon > 0$  and all large  $m$ . These two conditions imply that  $\frac{1}{m}S_m \rightarrow 0$  a.s. Further assume that  $W$  satisfies the following ‘‘isotropic’’ property: for any finite number of pairwise disjoint Borel measurable sets  $A_1, A_2, \dots, A_n \subset (0, \pi)$ , and  $\lambda_1, \lambda_2, \dots, \lambda_n \subset (0, 2\pi)$ ,

$$\{e^{i\lambda_i} W(A_i)\} \stackrel{d}{=} \{W(A_i)\}.$$

**Theorem 4.3.** *Let  $(X, \mathcal{A}, \mu, T)$  be aperiodic, and  $\xi_k$  as above. Then there is an  $f \in L^2(\mu)$  so that*

$$\sup_{a \in \mathbb{R}} |\mu(S_m f > a) - P(S_m > a)| \rightarrow 0, \quad m \rightarrow +\infty.$$

This appears to be the most general result available from the method of proof used in this paper. Note that it contains Theorem 1.1 as an immediate consequence, but this appears to be the only corollary of Theorem 4.3 of importance. We have stated this result to advertise the problem of generalizing it.

The only part of our proof that needs modification is Lemma 2.1. Instead, with the notation of that lemma, one must prove.

**Lemma 4.1.** *The intervals  $I_j$  can be chosen so that the following four conditions hold:*

$$\begin{aligned} l(m) &\geq (\log m)^{-1}; \\ \left\| \int_{I(m)}^{\pi} K_m(\lambda)W(dx) \right\|_2, \left\| \int_{I(m)}^{\pi} K_m(\lambda)V(d\lambda) \right\|_2 &= O(\log m); \\ \left\| \int_0^{l(m)} K_m(\lambda)(W - V)(d\lambda) \right\|_2 &= O(\log m) \end{aligned}$$

and for all complex numbers  $z_i$ , of modulus 1, the sum  $\sum z_i V(I_i)$  converges a.e.

Notice that the middle two properties are *not* almost sure statements. The last condition is needed to make sure that  $\bar{f}$  is well defined. (See the argument concerning (3.5) above.) The Rademacher-Menchoff Theorem [Z, vol. II, p. 193] is ideally suited to obtain this last condition. Use it, together with the following observation, to make a judicious choice of the intervals  $I_j$ . For all large  $m$ ,

$$m^2 EW([0, 1/m])^2 \leq CE \left| \int_{-\pi}^{\pi} K_m(\lambda) W(d\lambda) \right|^2 \leq Cm^{2-\varepsilon}$$

so that  $EW([0, 1/m])^2 \leq Cm^{-\varepsilon}$ .

#### ACKNOWLEDGMENTS

We express our thanks to three people who played important roles in the development of these results. They are Manfred Denker, Karl Petersen and Murad Taqqu. Also, an earlier version of this paper was completed in at the Department of Statistics, University of North Carolina, Chapel Hill. We are grateful for its hospitality.

#### REFERENCES

- [BD] R. Burton and M. Denker, *On the central limit theorem for dynamical systems*, Trans. Amer. Math. Soc. **302** (1987), 715–726.
- [CM] S. Cambanis and M. Maejima, *Two classes of self-similar stable processes with stationary increments*, Stochastic Process. Appl. **32** (1989), 305–329.
- [D] M. Denker, *The Central Limit Theorem for dynamical systems*, Dynamical Systems and Ergodic Theory, Banach Center Publications, vol. 23, Warsaw, 1989.
- [DK] M. Denker and G. Keller, *Rigorous statistical procedures for data from dynamical systems*, J. Statist. Phys. **44** (1986), 67–93.
- [Dob] R. L. Dobrushin, *Gaussian and their subordinated self-similar random generalized fields*, Ann. Probab. **7** (1979), 1–28.
- [GS] L. Giraitis and D. Surgailis, *CLT and other limit theorems for functionals of Gaussian processes*, Z. Wahrsch. Verw. Gebiete **70** (1985), 191–212.
- [GL] M. I. Gordin and B. A. Lifsic, *The central limit theorem for stationary Markov processes*, J. Soviet Math. **19** (1978), 392–394.
- [H] C. D. Hardin, *On the spectral representation of symmetric stable processes*, J. Multivariate Anal. **12** (1982), 385–401.
- [LP] M. Lacey and W. Philipp, *A note on the almost sure central limit theorem*, Statist. Probab. Lett. **9** (1990), 201–205.
- [Lam] J. W. Lamperti, *Semistable stochastic processes*, Trans. Amer. Math. Soc. **104** (1962), 62–78.
- [M] M. Maejima, *Self-similar processes and limit theorems*, Sugaku Expositions **2** (1989), 103–123.
- [Maj] P. Major, *Multiple Wiener-Itô Integrals*, Lecture Notes in Math., vol. 849, Springer-Verlag, Berlin and New York, 1981.
- [MvN] B. B. Mandelbrot and J. W. van Ness, *Fractional Brownian motion, fractional noises and applications*, SIAM Rev. **10** (1968), 422–437.

- [O] H. Oodaira, *On Strassen's version of the law of the iterated logarithm for Gaussian processes*, Z. Wahrsch. Verw. Gebiete **21** (1972), 289–299.
- [Or] J. Ortega, *Upper classes for the increments of fractional Wiener processes*, Probab. Theory Related Fields **80** (1989), 365–379.
- [PS] W. Philipp and W. Stout, *Almost sure invariance principles for partial sums of weakly dependent random variables*, Mem. Amer. Math. Soc. no. 161, 1975.
- [PUZ] F. Przytycki, M. Urbanski, and A. Zdunik, *Harmonic, Gibbs and Hausdorff measures on repellers for holomorphic maps*, Ann. of Math. (2) **130** (1989), 1–40.
- [R] M. Rosenblatt, *Independence and dependence*, Proc. 4th Berkeley Sympos. Math. Statist. Probab., Univ. of California Press, Berkeley, Calif., 1961, pp. 411–443.
- [PT] W. Parry and S. Tuncel, *Classification problems in ergodic theory*, Cambridge Univ. Press, London and New York, 1982.
- [S] Ya. G. Sinai, *The central limit theorem for geodesic flows on manifolds of constant negative curvature*, Soviet Math. **1** (1960), 983–986.
- [T1] M. S. Taqqu, *Random processes with long range dependence and high variability*, J. Geophys. Res. **92** (1987), 9683–9686.
- [T2] ———, *Self-similar processes and related ultraviolet and infrared catastrophes*, Random Fields: Rigorous Results in Statistical Mechanics and Quantum Field Theory, Colloquia Mathematica Societatis Janos Bolyai, vol. 27, Book 2, North-Holland, Amsterdam, 1981, pp. 1057–1096.
- [T3] ———, *Weak convergence to fractional Brownian motion and to the Rosenblatt process*, Z. Wahrsch. Verw. Gebiete **31** (1975), 287–302.
- [TC] ———, *A survey of functional laws of the iterated logarithm for self-similar process*, Stochastic Models **1** (1985), 77–115.
- [Z] A. Zygmund, *Trigonometric series*, (2nd ed.), Cambridge Univ. Press, London and New York, 1959.

DEPARTMENT OF MATHEMATICS, INDIANA UNIVERSITY, BLOOMINGTON, INDIANA 47405  
E-mail address: mlacey@iubacs