

CLASSIFICATION OF THE TOR-ALGEBRAS OF CODIMENSION FOUR ALMOST COMPLETE INTERSECTIONS

ANDREW R. KUSTIN

ABSTRACT. Let (R, \mathfrak{m}, k) be a local ring in which 2 is a unit. Assume that every element of k has a square root in k . We classify the algebras $\mathrm{Tor}_\bullet^R(R/J, k)$ as J varies over all grade four almost complete intersection ideals in R . The analogous classification has already been found when J varies over all grade four Gorenstein ideals [21], and when J varies over all ideals of grade at most three [5, 30]. The present paper makes use of the classification, in [21], of the Tor-algebras of codimension four Gorenstein rings, as well as the (usually nonminimal) DG-algebra resolution of a codimension four almost complete intersection which is produced in [25 and 26].

Fix, for the time being, a regular local ring (R, \mathfrak{m}, k) . For each Cohen-Macaulay ring A of the form $A = R/I$, we consider the Tor-algebra $T_\bullet = T_\bullet(A) = \mathrm{Tor}_\bullet^R(A, k)$. A great deal of information about A is encoded in $T_\bullet(A)$. Some of the classical results along these lines are: A is regular if and only if $T_\bullet = T_0$ [27]; A is Gorenstein if and only if T_\bullet is a Poincaré duality algebra [4]; A is a complete intersection if and only if T_\bullet is the exterior algebra on T_1 [29, 1]. There are at least three types of modern applications of theorems which classify Tor-algebras. The major impetus for studying T_\bullet is Avramov's machine for converting questions about the local ring A into questions about the algebra T_\bullet , provided the minimal R -free resolution of A is a DG-algebra. The algebra T_\bullet is graded-commutative, instead of commutative; nonetheless, it is a much simpler object than the original ring A . In particular, T_\bullet is always a finite dimensional vector space over k . Avramov's machine has been successfully applied when the codimension of A is at most three; or A is Gorenstein of codimension four; or A is one link from a complete intersection; or A is Gorenstein and two links from a complete intersection. In each case the minimal R -resolution of A is a DG-algebra [6, 17, 19, 16, 5] and the Tor-algebra $T_\bullet(A)$ has been classified [21, 30, 5]. Once the key hypotheses are established, then one is able to prove [12, 5] that the Poincaré series

$$P_A^M(z) = \sum_{i=0}^{\infty} \dim_k \mathrm{Tor}_i^A(M, k) z^i$$

is a rational function for all finitely generated A -modules M . One is also able to prove [2] that all of these rings A satisfy the Eisenbud Conjecture [8]; that

Received by the editors May 1, 1991.

1991 *Mathematics Subject Classification.* Primary 13D03, 13C40, 14M07.

Key words and phrases. Almost complete intersection, Betti numbers, DG-algebra, Gorenstein ideal, linkage, perfect ideal, Poincaré series, Tor-algebra.

is, if M is a finitely generated A -module whose Betti numbers are bounded, then the minimal resolution of M is eventually periodic of period two. See [3] for further results and problems along these lines.

Avramov's machine has been applied to Gorenstein rings of codimension four and to rings which are a "small" number of links from other "nice" rings. It is our hope that these techniques may also be applied to rings which are one link from a Gorenstein ring of codimension four, in other words to almost complete intersections of codimension four. The first step in this direction was taken in Palmer's thesis [25, 26]. Let A be a codimension four almost complete intersection. Palmer produced a DG-algebra resolution of A . Palmer's resolution is close to, but not always equal to, the minimal resolution of A . Palmer's work provides evidence that the minimal resolution of A is a DG-algebra and it is very useful in the present paper where the second step—the classification of $T_*(A)$ —takes place. Palmer's work is summarized in §3, and is applied to $T_*(A)$ in §4. (It is noteworthy that the present paper represents the first time that $T_*(A)$ has been classified before the minimal resolution of A was known to be a DG-algebra; indeed, it is likely that the present work will help complete the project initiated in [25].)

A second application of theorems which classify Tor-algebras is to the Buchsbaum-Eisenbud conjecture [6] about lower bounds for Betti numbers. Charalambous, Evans, and Miller [7] have proved that if the dimension, d , of R is at most four, and M is an R -module of finite length, with M not equal to R modulo a regular sequence, then the Betti numbers of M satisfy $\binom{d}{i} \leq \beta_i(M)$ for $0 < i < d$, and $2^d + 2^{d-1} \leq \sum_{i=0}^d \beta_i(M)$. One of the key ingredients in their proof is the classification in [21] of $T_*(A)$ for codimension four Gorenstein rings A . The classification of Tor-algebras contained in the present paper should lead to further progress on establishing lower bounds for Betti numbers.

Multiplicative operations in Tor-algebras also play some role in determining the generating set of a residual intersection. This theme is initiated in [23]. Further results along these lines will appear in subsequent papers.

The algebra $T_*(A)$ has been classified when A is a codimension four Gorenstein ring [21]; and when A is a codimension three ring [30, 5]. In each case, there are at most five different families of Tor-algebras. Furthermore, each family is discrete, in the sense that the family members are parameterized by integers. The proofs in [21] and [5] are based on the theory of linkage. The proof in [30] comes from invariant theory. The proofs look quite different, but the ultimate linear algebra calculations are roughly equivalent. The linkage theory proof is like an induction; one must know the answer before one can prove it. For rings of codimension three, the proof in [30] preceded proof in [5]; indeed, the authors of [5] took Weyman's answer and reproved it using their linkage technique. Some further details may be found in [24]. The classification in the present paper uses the linkage style of argument. Once again the answer consists of a small number of discrete families of Tor-algebras; see Theorem 1.5.

The main result is stated in §1 and proved in §4. Palmer's DG-algebra resolution \mathbb{M} of a codimension four almost complete intersection is recorded in §3. The multiplication in \mathbb{M} uses the multiplication on a resolution of a codi-

mension four Gorenstein ring. In §2 we recall the classification of $T_\bullet(A)$ for codimension four Gorenstein rings A . In §5 we give examples and ask questions. The remainder of the present section is a discussion of the conventions that are used throughout the paper.

In this paper “ring” means commutative noetherian ring with one. The *grade* of a proper ideal I in a ring R is the length of the longest regular sequence on R in I . The ideal I of R is called *perfect* if the grade of I is equal to the projective dimension, $\text{pd}_R(R/I)$, of the R -module R/I . A grade g ideal I is called a *complete intersection* if it can be generated by g generators. Complete intersection ideals are necessarily perfect. The grade g ideal I is called an *almost complete intersection* if it is a *perfect* ideal which is *not* a complete intersection and which can be generated by $g + 1$ generators. The grade g ideal I is called *Gorenstein* if it is perfect and $\text{Ext}_R^g(R/I, R) \cong R/I$.

Let k be a fixed field. Throughout this paper, we write

$$(0.1) \quad \text{“} S_\bullet \text{ is a graded } k\text{-algebra”}$$

to mean that S_\bullet is a finite dimensional graded-commutative associative k -algebra of the form $S_\bullet = \bigoplus_{i=0}^n S_i$ with $S_0 = k$. In particular,

$$s_i s_j = (-1)^{ij} s_j s_i \quad \text{for all } s_i \in S_i \text{ and } s_j \in S_j$$

and

$$s_i s_i = 0 \quad \text{if } s_i \in S_i \text{ and } i \text{ is odd.}$$

For example, if (R, \mathfrak{m}, k) is a local ring and I is an R -ideal of finite projective dimension, then $\text{Tor}_\bullet^R(R/I, k)$ is a graded k -algebra in the sense of (0.1). For a more concrete example, let V be a vector space of dimension d over k . The exterior algebra

$$\bigwedge^\bullet V = \bigwedge_k^\bullet V = k \oplus V \oplus \bigwedge^2 V \oplus \bigwedge^3 V \oplus \dots \oplus \bigwedge^d V,$$

with multiplication given by exterior product, is a graded k -algebra in the sense of (0.1). We use the usual conventions regarding grading. If $M = \bigoplus M_j$ is a graded S_\bullet -module, then $M(a)$ is the graded S_\bullet -module with the property that $M(a)_j = M_{a+j}$ and $\text{Hom}_{S_\bullet}(S_\bullet(-a), M) = M(a)$. In particular, there is an isomorphism of graded k -vector spaces from $k(-1)^d$ to the subspace $V = \bigwedge^1 V$ of $\bigwedge^\bullet V$.

In this paper the word “trivial” is given two distinct meanings. Suppose that S_\bullet is a graded k -algebra and W is a positively graded S_\bullet -module. Then the *trivial extension of S_\bullet by W* , $S_\bullet \times W$, is the graded k -algebra whose graded vector space structure is given by $S_\bullet \oplus W$ and whose multiplication is given by

$$(s_i, w_j)(s_k, w_l) = (s_i s_k, s_i w_l + (-1)^{jk} s_k w_j)$$

for all $s_a \in S_a$ and all $w_b \in W_b$. On the other hand, we say that W is a *trivial S_\bullet -module* if $S_+ W = 0$. In particular, if k is viewed as an S_\bullet -module by way of the natural quotient map $S_\bullet \rightarrow S_\bullet/S_+ = k$, then $\bigoplus_{i=1}^n k(-i)^{m_i}$ is a trivial S_\bullet -module.

Elementary results about linkage and DG-algebras may be found in [6 and 17]. In this paper, “DG-algebra” always means *associative* DG-algebra.

1. THE STATEMENT OF THE MAIN THEOREM

Let k be a fixed field. In Table 1.3 we define the graded k -algebras (in the sense of (0.1)) which appear in Theorem 1.5, the main theorem of the

paper. Each of these algebras has the form $S_\bullet = \bigoplus_{i=0}^4 S_i$ with $S_0 = k$ and $d_i = \dim_k S_i$. Select bases $\{x_i\}$ for S_1 , $\{y_i\}$ for S_2 , $\{z_i\}$ for S_3 , and $\{w_i\}$ for S_4 . View S_2 as the direct sum $S'_2 \oplus S_1^2$. Numerical information about these algebras is collected in Table 1.4. One may combine Lemma 1.2 with Table 1.4 in order to conclude that each of the algebras of Table 1.3 represents a distinct isomorphism class of k -algebras, provided the parameters p , q , and r satisfy

$$(1.1) \quad 0 \leq p, \quad 2 \leq q \leq 3, \quad \text{and} \quad 2 \leq r \leq 5.$$

(If we had allowed q and r to take the value 1, then $E^{(1)}$ would equal $E[1]$ and $F^{(1)}$ would equal $F[1]$.)

Lemma 1.2. *If S_\bullet is one of the algebras from Table 1.3, then there is a four dimensional subspace V of S_1 with the property that $\dim V^2 = 6$ if and only if S_\bullet is not equal to $C^{(2)}$, C^* , or $C[p]$ for any p .*

Proof. If S_\bullet is not equal to $C^{(2)}$, C^* , or $C[p]$ for any p , then the subspace V of S_1 spanned by x_1, x_2, x_3 , and x_4 has $\dim V^2 = 6$. On the other hand, we now suppose that S_\bullet is equal to $C^{(2)}$, C^* , or $C[p]$ for some p . Let x'_1, x'_2, x'_3, x'_4 be a basis for V . Select α_{ij} in k with $x'_j = \sum_{i=1}^5 \alpha_{ij} x_i$; let $\Delta(i, j; a, b)$ and $D(a, b, c, d)$ represent the following determinants:

$$\Delta(i, j; a, b) = \begin{vmatrix} \alpha_{ia} & \alpha_{ib} \\ \alpha_{ja} & \alpha_{jb} \end{vmatrix} \quad \text{and} \quad D(a, b, c, d) = \begin{vmatrix} \alpha_{a1} & \alpha_{a2} & \alpha_{a3} & \alpha_{a4} \\ \alpha_{b1} & \alpha_{b2} & \alpha_{b3} & \alpha_{b4} \\ \alpha_{c1} & \alpha_{c2} & \alpha_{c3} & \alpha_{c4} \\ \alpha_{d1} & \alpha_{d2} & \alpha_{d3} & \alpha_{d4} \end{vmatrix}.$$

Recall that $x_3x_4 = x_3x_5 = x_4x_5 = 0$ in S_\bullet . It follows that

$$x'_a x'_b = \sum_{i=1}^2 \sum_{j=i+1}^5 \Delta(i, j; a, b) x_i x_j \quad \text{in } S_\bullet.$$

Observe that

$$\begin{aligned} & \Delta(1, 2; 3, 4)x'_1 x'_2 - \Delta(1, 2; 2, 4)x'_1 x'_3 + \Delta(1, 2; 2, 3)x'_1 x'_4 + \Delta(1, 2; 1, 2)x'_3 x'_4 \\ & - \Delta(1, 2; 1, 3)x'_2 x'_4 + \Delta(1, 2; 1, 4)x'_2 x'_3 = \sum_{i=1}^2 \sum_{j=i+1}^5 D(1, 2, i, j) x_i x_j = 0. \end{aligned}$$

There are two possibilities. If $\Delta(1, 2; a, b) \neq 0$ for some pair (a, b) , then $\dim V^2 \leq 5$. If $\Delta(1, 2; a, b) = 0$ for all (a, b) , then the rank of

$$\begin{bmatrix} \alpha_{11} & \alpha_{12} & \alpha_{13} & \alpha_{14} \\ \alpha_{21} & \alpha_{22} & \alpha_{23} & \alpha_{24} \end{bmatrix}$$

is at most one, and V is contained in $U = (\lambda x_1 + \mu x_2, x_3, x_4, x_5)$ for some λ and μ in k . It follows that $\dim V^2 \leq \dim U^2 \leq 3$. \square

Key to Table 1.3.

- (a) $x_1x_2 = y_1, x_1x_3 = y_2, x_1x_4 = y_3, x_2x_3 = y_4, x_2x_4 = y_5, x_3x_4 = y_6$
- (a') $x_1x_2x_3 = z_1, x_1x_2x_4 = z_2, x_1x_3x_4 = z_3, x_2x_3x_4 = z_4$
- (b) $x_1x_2 = y_{l+1}, x_1x_3 = y_{l+2}, x_1x_4 = y_{l+3}, x_1x_5 = y_{l+4}, x_2x_3 = y_{l+5}, x_2x_4 = y_{l+6}, x_3x_4 = y_{l+7}$
- (b') $x_1x_2x_3 = z_{l+1}, x_1x_2x_4 = z_{l+2}, x_1x_3x_4 = z_{l+3}$
- (c) $x_1x_2 = y_{l+1}, x_1x_3 = y_{l+2}, x_1x_4 = y_{l+3}, x_1x_5 = y_{l+4}, x_2x_3 = y_{l+5}, x_2x_4 = y_{l+6}, x_2x_5 = y_{l+7}$

TABLE 1.3. The definition of the algebras A–F*

S_\bullet	d_1	d_2	d_3	d_4	$S_1 \times S_1$	$S_1 \times S_1 \times S_1$	$S_1 \times S'_2$	$S_1 \times S_3$	$S_2 \times S_2$
A	4	6	4	0	(a)	(a')	0	0	0
B[p]	5	$p+7$	$2p+3$	p	(b) with $l=p$	(b') with $l=2p$	(g)	(g')	0
C[p]	5	$p+7$	$2p+3$	p	(c) with $l=p$	(c') with $l=2p$	(g)	(g')	0
C ⁽²⁾	5	8	7	1	(c) with $l=1$	(c') with $l=4$	(h) with $j=2$	(h') with $j=2$	0
C*	5	9	7	2	(c) with $l=2$	(c') with $l=4$	(i)	(i')	(i')
D[p]	5	$p+8$	$2p+2$	p	(d) with $l=p$	(d') with $l=2p$	(g)	(g')	0
D ⁽²⁾	5	9	6	1	(d) with $l=1$	(d') with $l=4$	(h) with $j=2$	(h') with $j=2$	0
E[p]	5	$p+9$	$2p+1$	p	(e) with $l=p$	(e') with $l=2p$	(g)	(g')	0
E ^(q)	5	10	$2q+1$	1	(e) with $l=1$	(e') with $l=2q$	(h) with $j=q$	(h') with $j=q$	0
F[p]	5	$p+10$	$2p$	p	(f) with $l=p$	0	(g)	(g')	0
F ^(r)	5	11	$2r$	1	(f) with $l=1$	0	(h) with $j=r$	(h') with $j=r$	0
F*	5	12	10	2	(f) with $l=2$	0	(h) with $j=5$	(h') with $j=5$	(j)

The k -algebra S_\bullet is graded-commutative and associated; every product of basis vectors which is not listed has been set equal to zero. The notation is explained at the beginning of the section. The parameters p , q , and r satisfy (1.1).

- (c') $x_1x_2x_3 = z_{l+1}$, $x_1x_2x_4 = z_{l+2}$, $x_1x_2x_5 = z_{l+3}$
 (d) $x_1x_2 = y_{l+1}$, $x_1x_3 = y_{l+2}$, $x_1x_4 = y_{l+3}$, $x_1x_5 = y_{l+4}$, $x_2x_3 = y_{l+5}$,
 $x_2x_4 = y_{l+6}$, $x_2x_5 = y_{l+7}$, $x_3x_4 = y_{l+8}$
 (d') $x_1x_2x_3 = z_{l+1}$, $x_1x_2x_4 = z_{l+2}$
 (e) $x_1x_2 = y_{l+1}$, $x_1x_3 = y_{l+2}$, $x_1x_4 = y_{l+3}$, $x_1x_5 = y_{l+4}$, $x_2x_3 = y_{l+5}$,
 $x_2x_4 = y_{l+6}$, $x_2x_5 = y_{l+7}$, $x_3x_4 = y_{l+8}$, $x_3x_5 = y_{l+9}$
 (e') $x_1x_2x_3 = z_{l+1}$,
 (f) $x_1x_2 = y_{l+1}$, $x_1x_3 = y_{l+2}$, $x_1x_4 = y_{l+3}$, $x_1x_5 = y_{l+4}$, $x_2x_3 = y_{l+5}$,
 $x_2x_4 = y_{l+6}$, $x_2x_5 = y_{l+7}$, $x_3x_4 = y_{l+8}$, $x_3x_5 = y_{l+9}$, $x_4x_5 = y_{l+10}$
 (g) $x_1y_i = z_i$ for $1 \leq i \leq p$,
 (g') $x_1z_{p+i} = w_i$ for $1 \leq i \leq p$,
 (h) $x_iy_1 = z_i$ for $1 \leq i \leq j$,
 (h') $x_iz_{j+i} = w_1$ for $1 \leq i \leq j$,
 (i) $x_1y_1 = z_1$, $x_1y_2 = z_2$, $x_2y_1 = z_3$, $x_2y_2 = z_4$
 (i') $x_1x_2y_1 = w_1$, $x_1x_2y_2 = w_2$,
 (j) $y_1y_2 = w_1$, $y_1^2 = w_2$.

TABLE 1.4. Numerical information about the algebra $\mathbf{A-F}^*$

S_\bullet	$\dim S_1^2$	$\dim S_1^3$	$\dim S_1S_2 - \dim S_1^3$	$\dim S_1S_3$	$\dim S_2^2$
A	6	4	0	0	0
B [p]	7	3	p	p	0
C [p]	7	3	p	p	0
C ⁽²⁾	7	3	2	1	0
C [*]	7	3	4	2	2
D [p]	8	2	p	p	0
D ⁽²⁾	8	2	2	1	0
E [p]	9	1	p	p	0
E ^(q)	9	1	q	1	0
F [p]	10	0	p	p	0
F ^(r)	10	0	r	1	0
F [*]	10	0	5	1	2

Theorem 1.5. *Let (R, \mathfrak{m}, k) be a local ring in which 2 is a unit. Assume that every element of k has a square root in k . Let J be a grade four almost complete intersection ideal in R , and let T_\bullet be the graded k -algebra $\text{Tor}_\bullet^R(R/J, k)$. Then there is a parameter p , q , or r which satisfies (1.1), an algebra S_\bullet from the list **A**, **B**[p], **C**[p], **C**⁽²⁾, **C**^{*}, **D**[p], **D**⁽²⁾, **E**[p], **E**^(q), **F**[p], **F**^(r), **F**^{*}, and a positively graded vector space W such that, T_\bullet is isomorphic (as a graded k -algebra) to the trivial extension $S_\bullet \ltimes W$ of S_\bullet by the trivial S_\bullet -module W .*

Note. In the above theorem, the vector space W has the form $\bigoplus_{i=1}^4 k(-i)^{m_i}$ where $m_1 = 1$ if $S_\bullet = \mathbf{A}$, and $m_1 = 0$ in all other cases.

The proof of Theorem 1.5 is contained in §4. We next record a few consequences of Theorem 1.5. If one is interested only in the subalgebra of T_\bullet which is generated by T_1 , then the classification of Theorem 1.5 can be made even cleaner.

Corollary 1.6. *If the notation of Theorem 1.5 is adopted, then the subalgebra $k[T_1]$ of T_\bullet is isomorphic to one of the algebras $\mathbf{A} \times k(-1)$, $\mathbf{B}[0]$, $\mathbf{C}[0]$, $\mathbf{D}[0]$, $\mathbf{E}[0]$, $\mathbf{F}[0]$. In particular, the following numerical statements hold:*

- (a) $6 \leq \dim T_1^2 \leq 10$, and
- (b) $\dim T_1^2 + \dim T_1^3 = 10$.

Proof. It is easy to see that if T_\bullet has the form $S_\bullet \times W$ (as described in Theorem 1.5) where S_\bullet is $\mathbf{C}^{(2)}$, or \mathbf{C}^* , or $\mathbf{C}[p]$ (for some p), then the subalgebra $k[T_1]$ of T_\bullet is $\mathbf{C}[0]$. An analogous statement holds for all of the other algebras of Table 1.3. The numerical assertions follow from Table 1.4. \square

The next corollary follows from Lemma 1.2 by way of a prime avoidance argument.

Corollary 1.7. *Adopt the notation of Theorem 1.5. Exactly one of the following statements holds:*

- (a) *the subalgebra $k[T_1]$ of T_\bullet is $\mathbf{C}[0]$; or*
- (b) *there is a minimal presentation*

$$(1.8) \quad R^n \xrightarrow{d_2} R^5 [a_1, \dots, a_5] J$$

for J with the property that a_1, a_2, a_3, a_4 is a regular R -sequence and the first six columns of d_2 are

$$\begin{bmatrix} -a_2 & -a_3 & -a_4 & 0 & 0 & 0 \\ a_1 & 0 & 0 & -a_3 & -a_4 & 0 \\ 0 & a_1 & 0 & a_2 & 0 & -a_4 \\ 0 & 0 & a_1 & 0 & a_2 & a_3 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

Remarks 1.9. Some of the algebras of Table 1.3 have a compact coordinate-free representation:

(a) If V is the graded vector space $k(-1)^4$, then $\mathbf{A} \cong \bigwedge^\bullet V / \bigwedge^4 V$. In the notation of Theorem 1.5, one can show (see, for example, [25, Proposition 3.2] or [26, Proposition 4.2]) that $T_\bullet \cong \mathbf{A} \times W$ if and only if there is a grade four Gorenstein ideal I and a grade four complete intersection ideal K with

$$(1.10) \quad K \subseteq \mathfrak{m}I$$

such that $J = K : I$. (The significant hypothesis in the last sentence is the one we have isolated as (1.10).)

(b) If V is the graded vector space $k(-1)^3$, then $\mathbf{B}[p]$ is isomorphic to

$$\left(\frac{\bigwedge^\bullet V}{\bigwedge^3 V} \times [k(-1) \oplus k(-2)^p \oplus k(-3)^p] \right) \otimes_k \bigwedge^\bullet k(-1).$$

(c) The algebra $\mathbf{C}[p]$ is isomorphic to

$$[[k \times k(-1)^3] \otimes_k \bigwedge^\bullet k(-1)] \times (k(-2)^p \oplus k(-3)^p) \otimes_k \bigwedge^\bullet k(-1).$$

The algebra \mathbf{C}^* is isomorphic to

$$[k \times (k(-1)^3 \oplus k(-2)^2)] \otimes_k \bigwedge^\bullet k(-1)^2.$$

If J' is a grade two almost complete intersection (in other words, J' is a determinantal ideal generated by the 2×2 minors of a 3×2 matrix), and J

is the ideal (J', a, b) for some elements a and b of R , with a, b a regular sequence on R/J' , then $\text{Tor}_\bullet^R(R/J, k)$ is isomorphic to \mathbf{C}^* . (See the proof of case one in §4.)

(d) Let $V = k(-1)^2$ and $V' = k(-1)^2$ be graded vector spaces and S_\bullet be the graded k -algebra $\bigwedge^\bullet(V \oplus V')$. Let \overline{S}_\bullet be the graded k -algebra and W be the \overline{S}_\bullet -module defined by

$$\overline{S}_\bullet = S_\bullet / (\bigwedge^2 V') S_+ \quad \text{and} \quad W = S_\bullet / (V' + S_2) S_\bullet.$$

If W^* is the \overline{S}_\bullet -module $\text{Hom}_k(W, k)$, then

$$\mathbf{D}[0] = \overline{S}_\bullet \times W(-1), \quad \text{and}$$

$$\mathbf{D}^{(2)} = \overline{S}_\bullet \times (W(-1) \oplus W(-2) \oplus W^*(-4)).$$

(e) Let V be the graded vector space $k(-1)^3$, W be the $\bigwedge^\bullet V$ -module $\bigwedge^\bullet V / \bigwedge^2 V$, and W^* be the $\bigwedge^\bullet V$ -module $\text{Hom}_k(W, k)$. It is not difficult to see that

$$\mathbf{E}[0] = \bigwedge^\bullet V \times W(-1)^2,$$

$$\mathbf{E}^{(3)} = \bigwedge^\bullet V \times (W(-1)^2 \oplus W(-2) \oplus W^*(-4)).$$

(f) If V is the graded vector space $k(-1)^5$, then $\mathbf{F}[0] \cong \bigwedge^\bullet V / \bigwedge^3 V$. Suppose that J is an ideal from Theorem 1.5 with the property that the subalgebra $k[T_1]$ of T_\bullet is isomorphic to $\mathbf{F}[0]$. Let (1.8) be a *minimal* presentation of J . It follows that the basis for R^n can be chosen so that the first 10 columns of d_2 are

$$\begin{bmatrix} -a_2 & -a_3 & -a_4 & -a_5 & 0 & 0 & 0 & 0 & 0 & 0 \\ a_1 & 0 & 0 & 0 & -a_3 & -a_4 & -a_5 & 0 & 0 & 0 \\ 0 & a_1 & 0 & 0 & a_2 & 0 & 0 & -a_4 & -a_5 & 0 \\ 0 & 0 & a_1 & 0 & 0 & a_2 & 0 & a_3 & 0 & -a_5 \\ 0 & 0 & 0 & a_1 & 0 & 0 & a_2 & 0 & a_3 & a_4 \end{bmatrix}.$$

Let W be the $\mathbf{F}[0]$ -module $\mathbf{F}[0]/\mathbf{F}_+[0]^2$, and let W^* be the $\mathbf{F}[0]$ -module $\text{Hom}_k(W, k)$. It is not difficult to see that

$$\mathbf{F}^{(5)} \cong \mathbf{F}[0] \times [W(-2) \oplus W^*(-4)].$$

2. THE TOR-ALGEBRA OF A CODIMENSION FOUR GORENSTEIN RING

The classification of Tor-algebras for rings defined by grade four Gorenstein ideals plays a crucial role in the proof of Theorem 1.5. The following result is proved in [21] (when $\text{char } k \neq 2$) and [16]. (The results in [17, 21], and [16] are stated for Gorenstein ideals in Gorenstein local rings; however, it is not difficult to check that the proofs hold for Gorenstein ideals in arbitrary local rings.) The Tor-algebra $\text{Tor}_\bullet^R(R/I, k)$ may be described intrinsically without any mention of the minimal resolution

$$(2.1) \quad \mathbb{L}: 0 \rightarrow L_4 \xrightarrow{l_4} L_3 \xrightarrow{l_3} L_2 \xrightarrow{l_2} L_1 \xrightarrow{l_1} L_0$$

of R/I . We have chosen to introduce \mathbb{L} in Theorem 2.2 so that the notation in the present section coincides with the notation in §4. We know from [17] and [16] that \mathbb{L} is a DG-algebra; so, the graded k -algebras $\overline{\mathbb{L}}$ and $\text{Tor}_\bullet^R(R/I, k)$ are equal. (Throughout the paper we write $\bar{}$ to mean $\otimes_R k$ and $a \equiv b$ to mean $\bar{a} = \bar{b}$.)

Theorem 2.2. *Let (R, \mathfrak{m}, k) be a local ring. Assume that either every element in k has a square root in k , or else that the characteristic of k is equal to two. Let I be a grade four Gorenstein ideal in R , \mathbb{L} be the minimal R -resolution of R/I , and $\overline{\mathbb{L}}$ be the graded k -algebra $\text{Tor}_\bullet^R(R/I, k)$. If I is not a complete intersection, then there are bases e_1, \dots, e_n for L_1 ; $f_1, \dots, f_{n-1}, f'_1, \dots, f'_{n-1}$ for L_2 ; g_1, \dots, g_n for L_3 ; and h for L_4 such that the multiplication $\overline{L}_i \times \overline{L}_{4-i} \rightarrow \overline{L}_4$ is given by*

$$(2.3) \quad e_i g_j = \delta_{ij} h, \quad f_i f'_j \equiv \delta_{ij} h, \quad f_i f_j \equiv f'_i f'_j \equiv 0,$$

and the other products in $\overline{\mathbb{L}}$ are given by one of the following cases:

- (a) All products in $\overline{L}_1 \overline{L}_1$ and $\overline{L}_1 \overline{L}_2$ are zero.
- (b) All products in $\overline{L}_1 \overline{L}_1$ and $\overline{L}_1 \overline{L}_2$ are zero except:

$$(2.4) \quad \begin{aligned} e_1 e_2 = f_1, \quad e_1 e_3 = f_2, \quad e_2 e_3 = f_3, \\ e_2 f'_1 \equiv e_3 f'_2 \equiv g_1, \quad -e_1 f'_1 \equiv e_3 f'_3 \equiv g_2, \quad \text{and} \quad e_1 f'_2 \equiv e_2 f'_3 \equiv -g_3. \end{aligned}$$

- (c) There is an integer p such that $e_{p+1} e_i = f_i$, $e_i f'_i \equiv g_{p+1}$, and $e_{p+1} f'_i \equiv -g_i$ for $1 \leq i \leq p$ and all other products in $\overline{L}_1 \overline{L}_1$ and $\overline{L}_1 \overline{L}_2$ are zero.

Note. It is possible to choose the basis for \mathbb{L} so that the multiplication is correct “on the nose” for $L_1 \otimes L_1 \rightarrow L_2$ and $L_1 \otimes L_3 \rightarrow L_4$, and is also correct for $\overline{L}_1 \otimes \overline{L}_2 \rightarrow \overline{L}_3$ and $\overline{L}_2 \otimes \overline{L}_2 \rightarrow \overline{L}_4$.

Remark 2.5. One consequence of the above classification is the well-known fact that $\overline{L}_1^3 = 0$ when I is a grade four Gorenstein ideal which is not a complete intersection.

The proof of Theorem 1.5 requires that we understand the multiplication $V \otimes \overline{\mathbb{L}} \rightarrow \overline{\mathbb{L}}$, where V is an arbitrary subspace of \overline{L}_1 . It is not difficult to guess all of the possibilities. For example, if the multiplication of $\overline{\mathbb{L}}$ is described in Theorem 2.2(c), then the distinguished element \overline{e}_{p+1} “may be taken” to be either in V (case (iii) below) or not in V (case (iv)). A complete proof of Corollary 2.7 (in contrast to the above heuristic argument) has two parts. We use linear algebra to find an appropriate basis of \overline{L}_1 , and then we use the fact that $\overline{\mathbb{L}}$ is a Poincaré duality algebra to determine the rest of the multiplication in $\overline{\mathbb{L}}$. The second part of the argument is summarized in the following lemma, which appears as [21, Lemma 2.3]. (The characteristic two version of the lemma may be found at the end of [16].) The proof of Lemma 2.6, which is due to Avramov, is the only place in the present paper that the square roots of elements of k are used.

Lemma 2.6. *Let \mathbb{L} be as in Theorem 2.2. If e_1, \dots, e_n is any basis for L_1 , h is any basis for L_4 , and f_1, \dots, f_m is the beginning of a basis for L_2 with $m \leq n - 1$ and $f_i f_j \equiv 0$ for all i and j , then there is a basis g_1, \dots, g_n for L_3 and an extension of f_1, \dots, f_m to a basis $f_1, \dots, f_{n-1}, f'_1, \dots, f'_{n-1}$ for L_2 such that (2.3) holds.*

Corollary 2.7. *Adopt the notation and hypotheses of Theorem 2.2. If V is a nonzero subspace of \overline{L}_1 of dimension t , then there are bases $\{e_i\}$ for L_1 , $\{f_i, f'_i\}$ for L_2 , $\{g_i\}$ for L_3 and h for L_4 such that (2.3) holds, $\overline{e}_1, \dots, \overline{e}_t$ is a basis for V , and the multiplication $V \otimes \overline{L}_1 \rightarrow \overline{L}_2$ and $V \otimes \overline{L}_2 \rightarrow \overline{L}_3$ is given by one of the following cases:*

(i) The integer t is at least 3 and the only nonzero products in $V\bar{L}_1$ and $V\bar{L}_2$ are given in (2.4).

(ii) The integer t is at least 2 and all products in $V\bar{L}_1$ and $V\bar{L}_2$ are zero, except

$$e_1e_2 = f_1, \quad e_1e_{t+1} = f_2, \quad e_2e_{t+1} = f_3, \\ e_2f'_1 \equiv g_1, \quad -e_1f'_1 \equiv g_2, \quad \text{and} \quad e_1f'_2 \equiv e_2f'_3 \equiv -g_{t+1}.$$

(iii) There are integers a and b , with $0 \leq a \leq t-1$ and $0 \leq b$, such that the only nonzero products of basis vectors in $V\bar{L}_1$ and $V\bar{L}_2$ are

$$e_1e_{1+i} = f_i, \quad e_1f'_i \equiv -g_{1+i}, \quad e_{1+i}f'_i \equiv g_1, \quad \text{for } 1 \leq i \leq a, \\ e_1e_{t+i} = f_{a+i}, \quad e_1f'_{a+i} \equiv -g_{t+i}, \quad \text{for } 1 \leq i \leq b.$$

(iv) There is an integer j , with $2 \leq j \leq t$, such that the only nonzero products of basis vectors in $V\bar{L}_1$ and $V\bar{L}_2$ are $e_{t+1}e_i = f_i$ and $e_if'_i \equiv g_{t+1}$ for $1 \leq i \leq j$.

Proof. If $\bar{\mathbb{L}}$ is described in Theorem 2.2(a), then it is clear that $V\bar{\mathbb{L}}$ is given by (iii) with $a = b = 0$. We next suppose that $\bar{\mathbb{L}}$ is described by Theorem 2.2(c). In this case \bar{L}_1 decomposes as $k\bar{e} \oplus U$ for some $e \in L_1$ and some $U \subseteq \bar{L}_1$ with $U^2 = 0$. There are two possibilities: either $V \subseteq U$, or else there is an element u of U such that $\bar{e} + u \in V$. If $V \subseteq U$, then we let e_{t+1} be the element e of L_1 . Select elements e_1, \dots, e_t of L_1 such that $\bar{e}_1, \dots, \bar{e}_t$ is a basis for V , $\bar{e}_{t+1}\bar{e}_1, \dots, \bar{e}_{t+1}\bar{e}_j$ is a basis for $\bar{e}_{t+1}V$, and $e_{t+1}e_i \equiv 0$ for $j+1 \leq i \leq t$. Define $f_i = e_{t+1}e_i$ in L_2 for $1 \leq i \leq j$. Observe that $(f_1, \dots, f_j)^2 = 0$. Complete the basis for $\bar{\mathbb{L}}$ using Lemma 2.6. Observe that the multiplication $V\bar{\mathbb{L}}$ is described in (iv) (if $2 \leq j$) or (iii) (with $a = 0$ and $b = j$ if $0 \leq j \leq 1$). If $\bar{e} + u \in V$, then let $e_1 \in L_1$ be a preimage of this element. Observe that $\bar{L}_1 = k\bar{e}_1 \oplus U$. Select $e_2, \dots, e_n \in L_1$ such that $\bar{e}_2, \dots, \bar{e}_n \in U$, $\bar{e}_1, \dots, \bar{e}_t$ is a basis for V , e_1, \dots, e_n is a basis for L_1 , $\bar{e}_1\bar{e}_2, \dots, \bar{e}_1\bar{e}_{a+1}$ is a basis for \bar{e}_1V , $\bar{e}_1\bar{e}_2, \dots, \bar{e}_1\bar{e}_{a+1}, \bar{e}_1\bar{e}_{t+1}, \dots, \bar{e}_1\bar{e}_{t+b}$ is a basis for $\bar{e}_1\bar{L}_1$, and $e_1e_i \equiv 0$ whenever $a+2 \leq i \leq t$ or $t+b+1 \leq i \leq n$. Define the elements f_1, \dots, f_{a+b} in L_2 in the obvious manner and proceed, as in the case $V \subseteq U$, to show that $V\bar{\mathbb{L}}$ is described by (iii).

Finally, suppose that $\bar{\mathbb{L}}$ is described in Theorem 2.2(b). In other words, we are given a decomposition $\bar{L}_1 = E \oplus U$ with $\dim E = \dim E^2 = 3$ and $U \cdot \bar{L}_1 = 0$. Consider the map $\pi: V \rightarrow E$ which is the composition

$$V \xrightarrow{\text{incl.}} \bar{L}_1 = E \oplus U \xrightarrow{\text{proj.}} E.$$

Let r be the rank of π . It is clear that the kernel of π is $V \cap U$; consequently, we may select e_1, \dots, e_t in L_1 such that $\bar{e}_1, \dots, \bar{e}_t$ is a basis for V , and $\bar{e}_{r+1}, \dots, \bar{e}_t$ is a basis for $V \cap U$. It follows that $\pi(\bar{e}_1), \dots, \pi(\bar{e}_r)$ is a basis for $\text{im } \pi$. Let $s = 3 - r$ and let e_{t+1}, \dots, e_{t+s} be elements of L_1 such that $\bar{e}_{t+1}, \dots, \bar{e}_{t+s}$ are in E , and $\pi(\bar{e}_1), \dots, \pi(\bar{e}_r), \bar{e}_{t+1}, \dots, \bar{e}_{t+s}$ is a basis for E . If E' is the subspace $(\bar{e}_1, \dots, \bar{e}_r, \bar{e}_{t+1}, \dots, \bar{e}_{t+s})$ of \bar{L}_1 , then it is clear that $\dim E' = \dim (E')^2 = 3$ and that $E' \oplus U = \bar{L}_1$. It follows that we can find e_{t+s+1}, \dots, e_n in L_1 such that $\bar{e}_{t+s+1}, \dots, \bar{e}_n$ are in U , and e_1, \dots, e_n is a basis of L_1 . This basis has been chosen so that $\bar{e}_1, \dots, \bar{e}_r \in E' \cap V$, $\bar{e}_{r+1}, \dots, \bar{e}_t \in V \cap U$, $\bar{e}_{t+1}, \dots, \bar{e}_{t+s} \in E' \setminus V$, $\bar{e}_{t+s+1}, \dots, \bar{e}_n \in U \setminus V$. Complete the basis for $\bar{\mathbb{L}}$ by using the technique of the preceding paragraph. It is

now clear that $V\overline{\mathbb{L}}$ is described by (i), if $r = 3$; by (ii), if $r = 2$; by (iii) with $a = 0$ and $b = 2$, if $r = 1$; and by (iii) with $a = b = 0$, if $r = 0$. \square

3. A DG-RESOLUTION OF ALMOST COMPLETE INTERSECTIONS

Let J be a grade four almost complete intersection in the local ring (R, \mathfrak{m}, k) . In this section we describe Palmer's DG-algebra resolution \mathbb{M} of R/J . This resolution, in general, is not the minimal resolution of R/J ; nonetheless, we are able to use it in §4 to compute the multiplication in $\text{Tor}_\bullet^R(R/J, k)$.

Let K be a grade four complete intersection ideal with $K \subseteq J$ and $\mu(J/K) = 1$. (We use $\mu(M)$ to mean the minimal number of generators of the R -module M .) The ideal $I = K : J$ is known to be a grade four Gorenstein ideal. It is shown in [17] and [16] (the results in these references hold for Gorenstein ideals in arbitrary local rings) that the minimal resolution \mathbb{L} of R/I is a DG-algebra. Let \mathbb{K} be a Koszul complex which is the minimal resolution of R/K and let $\alpha_\bullet : \mathbb{K} \rightarrow \mathbb{L}$

$$(3.1) \quad \begin{array}{ccccccccccc} 0 & \rightarrow & K_4 & \xrightarrow{k_4} & K_3 & \xrightarrow{k_3} & K_2 & \xrightarrow{k_2} & K_1 & \xrightarrow{k_1} & K_0 \\ & & \alpha_4 \downarrow & & \alpha_3 \downarrow & & \alpha_2 \downarrow & & \alpha_1 \downarrow & & \alpha_0 \downarrow \\ 0 & \rightarrow & L_4 & \xrightarrow{l_4} & L_3 & \xrightarrow{l_3} & L_2 & \xrightarrow{l_2} & L_1 & \xrightarrow{l_1} & L_0 \end{array}$$

be a map of DG-algebras which extends the identity map $\alpha_0 : R \rightarrow R$. Fix orientation isomorphisms $[\] : K_4 \rightarrow R$ and $[\] : L_4 \rightarrow R$. A routine mapping cone argument establishes the following result.

Proposition 3.2. *Let J be a grade four almost complete intersection in the local ring (R, \mathfrak{m}, k) and let K be a grade four complete intersection ideal with $K \subseteq J$ and $\mu(J/K) = 1$. Let \mathbb{K} be the minimal resolution of R/K , \mathbb{L} be the minimal resolution of R/I for $I = K : J$, and $\alpha_\bullet : \mathbb{K} \rightarrow \mathbb{L}$, as in (3.1), be a map of oriented DG-algebras. If $\beta_i : L_i \rightarrow K_i$ is the map defined by*

$$(3.3) \quad [\beta_i(v_i)u_{4-i}] = [v_i\alpha_{4-i}(u_{4-i})]$$

for all $u_j \in K_j$ and all $v_i \in L_i$, then

$$\mathbb{M} = \mathbb{M}(\alpha_\bullet) : 0 \rightarrow M_4 \xrightarrow{m_4} M_3 \xrightarrow{m_3} M_2 \xrightarrow{m_2} M_1 \xrightarrow{m_1} M_0$$

is a resolution of R/J , where $M_0 = R$, $M_1 = K_1 \oplus L_0$, $M_2 = K_2 \oplus L_1$, $M_3 = K_3 \oplus L_2$, $M_4 = L_3$, $m_1 = [k_1 \ \beta_0]$,

$$m_2 = \begin{bmatrix} k_2 & -\beta_1 \\ 0 & l_1 \end{bmatrix}, \quad m_3 = \begin{bmatrix} k_3 & \beta_2 \\ 0 & l_2 \end{bmatrix}, \quad \text{and} \quad m_4 = \begin{bmatrix} -\beta_3 \\ l_3 \end{bmatrix}.$$

Note. The definition of β_i makes use of the well-known perfect pairings $K_i \otimes K_{4-i} \rightarrow R$ and $L_i \otimes L_{4-i} \rightarrow R$, which are given by $u_i \otimes u_{4-i} \mapsto [u_i u_{4-i}]$ and $v_i \otimes v_{4-i} \mapsto [v_i v_{4-i}]$. The orientation on the left side of (3.3) is the orientation on \mathbb{K} , whereas the orientation on the right side of (3.3) is the orientation on \mathbb{L} .

The next result asserts that \mathbb{M} has the structure of a DG-algebra, provided 2 is a unit in R . A small amount of notation is needed in order to describe the multiplication in \mathbb{M} . Let h be the element of L_4 with $[h] = 1$ and let $\varepsilon_1, \varepsilon_2, \varepsilon_3, \varepsilon_4$ be a basis for K_1 with $[\varepsilon_1 \wedge \varepsilon_2 \wedge \varepsilon_3 \wedge \varepsilon_4] = 1$. The result claims the existence of an R -module homomorphism $P : \bigwedge^5 L_1 \rightarrow L_2$ which satisfies

a long list of properties. Two homomorphisms, $p: L_1 \rightarrow L_2$ and $q: L_2 \rightarrow L_3$, are defined in terms of P by

$$(3.4) \quad p(v_1) = P(v_1 \wedge \alpha_1(\varepsilon_1) \wedge \alpha_1(\varepsilon_2) \wedge \alpha_1(\varepsilon_3) \wedge \alpha_1(\varepsilon_4)),$$

and $v_1q(v_2) = v_2p(v_1)$ for all $v_i \in L_i$.

Theorem 3.5 [25]. *Adopt the notation of the preceding paragraph together with the notation and hypotheses of Proposition 3.2. If 2 is a unit in R , then there is a map $P: \wedge^3 L_1 \rightarrow L_2$ such that the following maps give \mathbb{M} the structure of a DG-algebra:*

$$\begin{aligned} M_1 \otimes M_1 &\rightarrow M_2: \begin{bmatrix} u_1 \\ v_0 \end{bmatrix} \begin{bmatrix} u'_1 \\ v'_0 \end{bmatrix} = \begin{bmatrix} u_1 u'_1 \\ v'_0 \alpha_1(u_1) - v_0 \alpha_1(u'_1) \end{bmatrix}, \\ M_1 \otimes M_2 &\rightarrow M_3: \begin{bmatrix} u_1 \\ v_0 \end{bmatrix} \begin{bmatrix} u_2 \\ v_1 \end{bmatrix} = \begin{bmatrix} u_1 u_2 \\ v_0 \alpha_2(u_2) + \alpha_1(u_1) v_1 + v_0 p(v_1) \end{bmatrix}, \\ M_1 \otimes M_3 &\rightarrow M_4: \begin{bmatrix} u_1 \\ v_0 \end{bmatrix} \begin{bmatrix} u_3 \\ v_2 \end{bmatrix} = -[u_1 u_3] l_4(h) - v_0 \alpha_3(u_3) + \alpha_1(u_1) v_2 - v_0 q(v_2), \\ M_2 \times M_2 &\rightarrow M_4: \begin{bmatrix} u_2 \\ v_1 \end{bmatrix} \begin{bmatrix} u'_2 \\ v'_1 \end{bmatrix} = -[u_2 u'_2] l_4(h) + \alpha_2(u_2) v'_1 + v_1 \alpha_2(u'_2) \\ &\quad + v_1 p(v'_1) + v'_1 p(v_1) \end{aligned}$$

for all $u_i, u'_i \in K_i$ and $v_i, v'_i \in L_i$. Furthermore, the map P also has the property that

$$v_1 v'_1 P(v_1 \wedge v'_1 \wedge _): \wedge^3 L_1 \rightarrow L_4$$

is the zero map for all $v_1, v'_1 \in L_1$.

Note. There are two parts to the proof in [25, 26]. In the first part, a long list of properties for P is compiled such that whenever a map P satisfies all of these properties, then the above indicated multiplication gives \mathbb{M} the structure of a DG-algebra. The one property for P that is highlighted in Theorem 3.5 is just one of the many properties from this list; however, it happens to be the only property of P that we use explicitly in §4. The second, and much more difficult, part of the proof in [25, 26] is to prove that the desired P (a “higher order multiplication” on the resolution \mathbb{L} of a codimension four Gorenstein ring) does exist.

4. THE PROOF OF THE MAIN THEOREM

Fix the notation and hypotheses of Theorem 1.5. If K is a grade four complete intersection ideal with $K \subseteq J$ and $\mu(J/K) = 1$, then we say that the grade four Gorenstein ideal $I = K : J$ is (directly) linked to J by K . For each such K , let

$$t(K) = \dim_k \left(\frac{K + \mathfrak{m}I}{\mathfrak{m}I} \right).$$

In other words, $t(K)$ is the cardinality of the largest subset of K which begins a minimal generating set for the ideal $K : J$. It is clear that $0 \leq t(K) \leq 4$. Our proof of Theorem 1.5 is divided into three cases:

Case 1. The ideal J is directly linked to a complete intersection.

Case 2. The ideal J is not directly linked to a complete intersection; and there exists a grade four complete intersection ideal K with $K \subseteq J$, $\mu(J/K) = 1$, and $t(K) \leq 3$.

Case 3. The ideal J is not directly linked to a complete intersection; and $t(K) = 4$ for every grade four complete intersection ideal K with $K \subseteq J$ and $\mu(J/K) = 1$.

The proof of Theorem 1.5 in Case 1. According to the hypothesis, there are complete intersection ideals I and K with $K \subseteq J$, $\mu(J/K) = 1$, and I linked to J by K . Let $t = t(K)$ and $s = 4 - t$. It is known (see, for example, [5, Theorem 3.2]) that there are matrices $\mathbf{a}_{1 \times s}$, $\mathbf{b}_{1 \times t}$, and $X_{s \times s}$ with entries in \mathfrak{m} such that $J = J' + I_1(\mathbf{b})$ and the entries b_1, \dots, b_t of \mathbf{b} form a regular sequence on both R and R/J' where $J' = I_1(\mathbf{a}X) + I_s(X)$. (If M is a matrix with entries in R , then we use $I_l(M)$ to denote the ideal in R generated by the $l \times l$ minors of M .) Let \mathbb{L}' be the minimal resolution of R/J' and \mathbb{K} be the Koszul complex which is the minimal resolution of $R/I_1(\mathbf{b})$. Both of these resolutions are DG-algebras. (See [5, Proposition 4.4] for the multiplication on \mathbb{L}' .) It follows that the resolution $\mathbb{L}' \otimes_R \mathbb{K}$ of R/J is a DG-algebra; and therefore,

$$T_\bullet \cong \text{Tor}_\bullet^R(R/J', k) \otimes_k \text{Tor}_\bullet^R(R/I_1(\mathbf{b}), k).$$

We know that $\text{Tor}_\bullet^R(R/I_1(\mathbf{b}), k)$ is the exterior algebra $\bigwedge^\bullet k(-1)^t$. Proposition 4.4 of [5] shows that

$$\text{Tor}_\bullet^R(R/J', k) \cong S_\bullet \ltimes W \quad \text{where } V = k(-1)^s, S_\bullet = \bigwedge^\bullet V / \bigwedge^s V,$$

and W is the trivial S_\bullet -module

$$\sum_{i=1}^s k(-i)^{\binom{s}{i-1}}.$$

The hypothesis that J is a proper ideal which is not a complete intersection ensures that $0 \leq t \leq 2$. It is now clear that

$$T_\bullet = \begin{cases} \mathbf{C}^*, & \text{if } t = 2, \\ \mathbf{B}[3], & \text{if } t = 1, \\ \mathbf{A} \ltimes W, & \text{if } t = 0. \end{cases}$$

The proof of Theorem 1.5 in Case 1 is complete.

For each choice of a grade four complete intersection ideal K with $K \subseteq J$ and J/K cyclic, we are able to use the information of §§2 and 3 in order to calculate part of the multiplication in T_\bullet . To prove Theorem 1.5 in Cases 2 and 3, we piece together this incomplete information in order to produce the entire multiplication table for T_\bullet . For the time being, let K be a fixed grade four complete intersection with $K \subseteq J$ and J/K cyclic. Let t denote $t(K)$, and let I be the Gorenstein ideal $K : J$. (We are finished with Case 1; so we may assume that the ideal I is not a complete intersection.) Define \mathbb{K} , \mathbb{L} , and α_\bullet as in (3.1); β_i and $\mathbb{M} = \mathbb{M}(\alpha_\bullet)$ as in Proposition 3.2; and an algebra structure on \mathbb{M} as in Theorem 3.5. We calculate multiplication in T_\bullet by using the fact that T_\bullet is equal to the homology algebra $H_\bullet(\overline{\mathbb{M}})$. A quick look at Proposition 3.2 shows that $T_\bullet = \bigoplus_{i=0}^4 T_i$, where $T_0 = k$, $T_1 = \overline{K}_1 \oplus \overline{L}_0$,

$$(4.1) \quad T_2 = \frac{\overline{K}_2}{\text{im } \beta_2} \oplus \overline{L}_1, \quad T_3 = \frac{\overline{K}_3}{\text{im } \beta_3} \oplus \ker \overline{\beta}_2, \quad \text{and} \quad T_4 = \ker \overline{\beta}_3.$$

Much of the multiplication on \mathbb{M} becomes zero in $\overline{\mathbb{M}}$. The resolution \mathbb{L} is minimal; and therefore, $l_4 \equiv 0$. We know from Remark 2.5 that $L_1^3 \subseteq \mathfrak{m}L_3$; thus,

$$\text{im } \alpha_3 = (\text{im } \alpha_1)^3 \subseteq L_1^3 \subseteq \mathfrak{m}L_3 \quad \text{and} \quad L_1 \cdot (\text{im } \alpha_2) \subseteq \mathfrak{m}L_3.$$

It follows that the multiplication of Theorem 3.5 induces the following multiplication on T_\bullet :

(4.2)

$$\begin{aligned} T_1 \otimes T_1 &\rightarrow T_2: \begin{bmatrix} \overline{u}_1 \\ \overline{v}_0 \end{bmatrix} \begin{bmatrix} \overline{u}'_1 \\ \overline{v}'_0 \end{bmatrix} = \begin{bmatrix} \overline{u}_1 \overline{u}'_1 \pmod{\text{im } \overline{\beta}_2} \\ \overline{v}'_0 \overline{\alpha}_1(\overline{u}_1) - \overline{v}_0 \overline{\alpha}_1(\overline{u}'_1) \end{bmatrix}, \\ T_1 \otimes T_2 &\rightarrow T_3: \begin{bmatrix} \overline{u}_1 \\ \overline{v}_0 \end{bmatrix} \begin{bmatrix} \overline{u}_2 \pmod{\text{im } \overline{\beta}_2} \\ \overline{v}_1 \end{bmatrix} = \begin{bmatrix} \overline{u}_1 \overline{u}_2 \pmod{\text{im } \overline{\beta}_3} \\ \overline{v}_0 \overline{\alpha}_2(\overline{u}_2) + \overline{\alpha}_1(\overline{u}_1) \overline{v}_1 + \overline{v}_0 \overline{p}(\overline{v}_1) \end{bmatrix}, \\ T_1 \otimes T_3 &\rightarrow T_4: \begin{bmatrix} \overline{u}_1 \\ \overline{v}_0 \end{bmatrix} \begin{bmatrix} \overline{u}_3 \pmod{\text{im } \overline{\beta}_3} \\ \overline{v}_2 \end{bmatrix} = \overline{\alpha}_1(\overline{u}_1) \overline{v}_2 - \overline{v}_0 \overline{q}(\overline{v}_2), \\ T_2 \otimes T_2 &\rightarrow T_4: \begin{bmatrix} \overline{u}_2 \pmod{\text{im } \overline{\beta}_2} \\ \overline{v}_1 \end{bmatrix} \begin{bmatrix} \overline{u}'_2 \pmod{\text{im } \overline{\beta}_2} \\ \overline{v}'_1 \end{bmatrix} = \overline{v}_1 \overline{p}(\overline{v}'_1) + \overline{v}'_1 \overline{p}(\overline{v}_1) \end{aligned}$$

for $u_i, u'_i \in K_i$ and $v_i, v'_i \in L_i$.

Apply Corollary 2.7 to the subspace $\text{im } \overline{\alpha}_1$ of \overline{L}_1 in the Tor-algebra $\overline{\mathbb{L}} = \text{Tor}_\bullet^R(R/I, k)$ in order to find bases e_1, \dots, e_n for L_1 ; $f_1, \dots, f_{n-1}, f'_1, \dots, f'_{n-1}$ for L_2 ; g_1, \dots, g_n for L_3 ; and h for L_4 such that $\overline{e}_1, \dots, \overline{e}_t$ is a basis for $\text{im } \overline{\alpha}$, and the multiplication $(\text{im } \overline{\alpha}_1) \cdot \overline{\mathbb{L}}$ is described by one of (i)–(iv). In particular, there are five possibilities for the multiplication $(\text{im } \overline{\alpha}_1) \cdot (\text{im } \overline{\alpha}_1)$:

- (A) all products are zero; or
- (B) $\overline{e}_1 \overline{e}_2 = \overline{f}_1$; or
- (C) $\overline{e}_1 \overline{e}_2 = \overline{f}_1$, and $\overline{e}_1 \overline{e}_3 = \overline{f}_2$; or
- (D) $\overline{e}_1 \overline{e}_2 = \overline{f}_1$, $\overline{e}_1 \overline{e}_3 = \overline{f}_2$, and $\overline{e}_2 \overline{e}_3 = \overline{f}_3$; or
- (E) $\overline{e}_1 \overline{e}_2 = \overline{f}_1$, $\overline{e}_1 \overline{e}_3 = \overline{f}_2$, and $\overline{e}_1 \overline{e}_4 = \overline{f}_3$.

For each possibility we have listed the nonzero products; all other products of basis vectors are zero. In Case 2 of our proof of Theorem 1.5, we have $t \leq 3$, so possibility (E) does not occur in this case. Furthermore, Lemma 4.14(b) shows that in Case 3 the multiplication $(\text{im } \overline{\alpha}_1)^2$ is described by (A); consequently there is no loss of generality if we set up our notation under the hypothesis that

$$(4.3) \quad \text{the multiplication } (\text{im } \overline{\alpha}_1)^2 \text{ is described by one of (A)–(D).}$$

Choose a basis $\varepsilon_1, \varepsilon_2, \varepsilon_3, \varepsilon_4$ for K_1 such that

$$(4.4) \quad \alpha_1(\varepsilon_i) = \varepsilon_i \quad \text{for } 1 \leq i \leq t, \quad \alpha_1(\varepsilon_i) \equiv 0 \quad \text{for } t+1 \leq i \leq 4, \quad \text{and} \\ [\varepsilon_1 \wedge \varepsilon_2 \wedge \varepsilon_3 \wedge \varepsilon_4] = 1.$$

(Notice that the definition of p in (3.4) appears to use a particular basis for K_1 ; however, every basis $\varepsilon_1, \varepsilon_2, \varepsilon_3, \varepsilon_4$ of K_1 which satisfies (4.4) gives rise to the exact same function p .) Now that the basis for K_1 is set, we give names to the corresponding basis elements of K_2 and K_3 :

$$\begin{aligned} \varphi_1 = \varepsilon_1 \varepsilon_2, \quad \varphi_2 = \varepsilon_1 \varepsilon_3, \quad \varphi_3 = \varepsilon_2 \varepsilon_3, \quad \varphi'_1 = \varepsilon_3 \varepsilon_4, \quad \varphi'_2 = -\varepsilon_2 \varepsilon_4, \quad \varphi'_3 = \varepsilon_1 \varepsilon_4, \\ \gamma_1 = \varepsilon_2 \varepsilon_3 \varepsilon_4, \quad \gamma_2 = -\varepsilon_1 \varepsilon_3 \varepsilon_4, \quad \gamma_3 = \varepsilon_1 \varepsilon_2 \varepsilon_4, \quad \text{and} \quad \gamma_4 = -\varepsilon_1 \varepsilon_2 \varepsilon_3. \end{aligned}$$

Let $d = \text{rank } \bar{\alpha}_2$. It is clear that $0 \leq d \leq \binom{t}{2}$. The notation has been set up, thanks to (4.3), so that

$$\begin{aligned} \alpha_2(\varphi_i) &= f_i \quad \text{for } 1 \leq i \leq d, \quad \text{and} \\ \alpha_2(\varphi_i) &\equiv \alpha_2(\varphi'_j) \equiv 0 \quad \text{for } d+1 \leq i \leq 3 \text{ and } 1 \leq j \leq 3. \end{aligned}$$

A straightforward application of (3.3) yields

$$\beta_3(g_i) \equiv \begin{cases} \gamma_i, & \text{if } 1 \leq i \leq t, \\ 0, & \text{if } t+1 \leq i \leq n, \end{cases} \quad \beta_2(f'_i) \equiv \begin{cases} \varphi'_i, & \text{if } 1 \leq i \leq d, \\ 0, & \text{if } d+1 \leq i \leq n-1, \end{cases}$$

and $\beta_2(f_i) \equiv 0$ for $1 \leq i \leq n-1$. Thus,

$$(4.5) \quad \begin{aligned} \ker \bar{\beta}_3 &= (\bar{g}_{t+1}, \dots, \bar{g}_n) \subseteq \bar{L}_3, \\ \ker \bar{\beta}_2 &= (\bar{f}'_{d+1}, \dots, \bar{f}'_{n-1}, \bar{f}_1, \dots, \bar{f}_{n-1}) \subseteq \bar{L}_3, \\ \text{im } \bar{\beta}_3 &= (\bar{\gamma}_1, \dots, \bar{\gamma}_t) \subseteq \bar{K}_3, \quad \text{and} \quad \text{im } \bar{\beta}_2 = (\bar{\varphi}'_1, \dots, \bar{\varphi}'_d) \subseteq \bar{K}_2. \end{aligned}$$

Label the following elements of T_\bullet :

$$\begin{aligned} x_i &= \begin{bmatrix} \bar{e}_i \\ 0 \end{bmatrix}, \quad x_5 = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \in T_1 \quad \text{for } 1 \leq i \leq 4; \\ y_i &= \begin{bmatrix} \bar{\varphi}'_i \pmod{\text{im } \bar{\beta}_2} \\ 0 \end{bmatrix}, \quad y_{3+i} = \begin{bmatrix} \bar{\varphi}_i \pmod{\text{im } \bar{\beta}_2} \\ 0 \end{bmatrix}, \quad y_{6+j} = \begin{bmatrix} 0 \\ \bar{e}_j \end{bmatrix} \in T_2 \end{aligned}$$

for $1 \leq i \leq 3$ and $1 \leq j \leq n$;

$$z_i = \begin{bmatrix} \bar{\gamma}_i \pmod{\text{im } \bar{\beta}_3} \\ 0 \end{bmatrix}, \quad z_{4+j} = \begin{bmatrix} 0 \\ \bar{f}_j \end{bmatrix}, \quad z_{n-d+3+l} = \begin{bmatrix} 0 \\ \bar{f}'_l \end{bmatrix} \in T_3$$

for $1 \leq i \leq 4$, $1 \leq j \leq n-1$, and $d+1 \leq l \leq n-1$; and

$$w_i = \bar{g}_{t+i} \in T_4 \quad \text{for } 1 \leq i \leq n-t.$$

(Notice that the above labeling depends on the choice of K .) We see from (4.1) and (4.5) that $y_1 = \dots = y_d = 0$, $z_1 = \dots = z_t = 0$. Furthermore,

$$(4.6) \quad \begin{aligned} x_1, \dots, x_5 &\text{ is a basis for } T_1; \quad y_{d+1}, \dots, y_{6+n} \text{ is a basis for } T_2; \\ & z_{t+1}, \dots, z_{2n+2-d} \end{aligned}$$

is a basis for T_3 ; and w_1, \dots, w_{n-t} is a basis for T_4 .

It is easy to see, using (4.2), that the multiplication $T_1 \otimes T_1 \rightarrow T_2$ is given by

$$(4.7) \quad \begin{aligned} x_1 x_2 &= y_4, \quad x_1 x_3 = y_5, \quad x_2 x_3 = y_6, \quad x_3 x_4 = y_1, \quad x_2 x_4 = -y_2, \\ x_1 x_4 &= y_3, \quad \text{and} \quad x_i x_5 = \begin{cases} y_{6+i}, & \text{for } 1 \leq i \leq t, \\ 0, & \text{for } t+1 \leq i \leq 4; \end{cases} \end{aligned}$$

and that the multiplication $T_1 \otimes T_1 \otimes T_1 \rightarrow T_3$ is given by

$$(4.8) \quad \begin{aligned} x_2 x_3 x_4 &= z_1, \quad x_1 x_3 x_4 = -z_2, \quad x_1 x_2 x_4 = z_3, \quad x_1 x_2 x_3 = -z_4, \\ x_1 x_2 x_5 &= \begin{cases} z_5, & \text{if } 1 \leq d, \\ 0, & \text{if } d = 0, \end{cases} \quad x_1 x_3 x_5 = \begin{cases} z_6, & \text{if } 2 \leq d, \\ 0, & \text{if } d \leq 1, \end{cases} \\ x_2 x_3 x_5 &= \begin{cases} z_7, & \text{if } 3 \leq d, \\ 0, & \text{if } d \leq 2, \end{cases} \end{aligned}$$

and $x_i x_4 x_5 = 0$ for all i . Furthermore, all of the products of basis vectors from

$$(4.9) \quad (x_1, \dots, x_4) \cdot (y_{t+7}, \dots, y_{n+6}) \quad \text{and} \quad (x_1, \dots, x_4) \cdot T_3$$

are zero except

$$x_1 y_{t+7} = z_6, \quad x_2 y_{t+7} = z_7, \quad \text{and} \quad x_1 z_{n+4} = x_2 z_{n+5} = -w_1$$

when the multiplication $(\text{im } \bar{\alpha}_1) \cdot \bar{\mathbb{L}}$ is described by Corollary 2.7(ii);

$$(4.10) \quad x_1 y_{6+t+i} = z_{4+d+i} \quad \text{and} \quad x_1 z_{n+3+i} = -w_i \quad \text{for } 1 \leq i \leq b$$

when the multiplication $(\text{im } \bar{\alpha}_1) \cdot \bar{\mathbb{L}}$ is described by Corollary 2.7(iii); and

$$(4.11) \quad x_i y_{t+7} = -z_{4+i} \quad \text{and} \quad x_i z_{n+3+i} = w_1 \quad \text{for } 1 \leq i \leq j$$

when the multiplication $(\text{im } \bar{\alpha}_1) \cdot \bar{\mathbb{L}}$ is described by Corollary 2.7(iv). It is not possible to determine

$$(4.12) \quad x_5 \cdot (y_{t+7}, \dots, y_{n+6}), \quad x_5 \cdot T_3, \quad \text{or} \quad T_2 \cdot T_2$$

at the present level of generality.

The proof of Theorem 1.5 in Case 2. Fix a complete intersection ideal K with $K \subseteq J$, $\mu(J/K) = 1$, and $t(K) \leq 3$. Use K to calculate multiplication in T_\bullet as described in (4.1) and (4.2). The map p of (3.4) satisfies $p \equiv 0$ because $\text{rank } \bar{\alpha}_1 = t \leq 3$. The map q is defined in terms of p ; hence, $q \equiv 0$. It follows that all of the products of (4.12) are zero. Combine the basis for T_\bullet given in (4.6) with the multiplication from (4.7), (4.8), and (4.9) in order to see that Table 4.13 is correct and complete, where $T_\bullet = S_\bullet \ltimes W$ for some trivial S_\bullet -module W . Recall that the algebras $\mathbf{A}\text{-F}^*$ are defined in Table 1.3. If the multiplication $(\text{im } \bar{\alpha}_1) \cdot \bar{\mathbb{L}}$ is described in part (iii) of Corollary 2.7, then the parameter a must equal d . The multiplications in part (ii) and part (iv) each require that $2 \leq t$; but (ii) must have $d = 1$, whereas (iv) requires $d = 0$.

TABLE 4.13. The conclusion of the proof of Theorem 1.5 in Case 2

t	d	$(\text{im } \bar{\alpha}_1) \cdot \bar{\mathbb{L}}$	$k[T_1]$	S_\bullet
0	0	$(\text{im } \bar{\alpha}_1) = 0$	$\mathbf{A} \ltimes k(-1)$	\mathbf{A}
1	0	(iii) with $a = 0$ and $b \geq 0$	$\mathbf{B}[0]$	$\mathbf{B}[b]$
2	0	(iii) with $a = 0$ and $b \geq 0$	$\mathbf{D}[0]$	$\mathbf{D}[b]$
2	0	(iv) with $j = 2$	$\mathbf{D}[0]$	$\mathbf{D}^{(2)}$
2	1	(ii)	$\mathbf{C}[0]$	$\mathbf{C}^{(2)}$
2	1	(iii) with $a = 1$ and $b \geq 0$	$\mathbf{C}[0]$	$\mathbf{C}[b]$
3	0	(iii) with $a = 0$ and $b \geq 0$	$\mathbf{E}[0]$	$\mathbf{E}[b]$
3	0	(iv) with $2 \leq j \leq 3$	$\mathbf{E}[0]$	$\mathbf{E}^{(j)}$
3	1	(ii)	$\mathbf{D}[0]$	$\mathbf{D}^{(2)}$
3	1	(iii) with $a = 1$ and $b \geq 0$	$\mathbf{D}[0]$	$\mathbf{D}[b]$
3	2	(iii) with $a = 2$ and $b \geq 0$	$\mathbf{B}[0]$	$\mathbf{B}[b]$
3	3	(i)	$\mathbf{A} \ltimes k(-1)$	\mathbf{A}

The proof of Theorem 1.5 in Case 2 is complete.

Without any further ado, we are able to identify the subalgebra $k[T_1]$ of T_\bullet in Case 3 of Theorem 1.5. Part (b) of the following lemma appears to be technical; but, in particular, it yields a complete description of the minimal resolution of R/J .

Lemma 4.14. *If the notation and hypotheses for Case 3 (from the beginning of the present section) are adopted, then the following statements hold:*

(a) *The algebra $k[T_1]$ is isomorphic to $\mathbf{F}[0]$.*

(b) *Let K be any grade four complete intersection ideal with $K \subseteq J$ and $\mu(J/K) = 1$. If \mathbf{M} from Proposition 3.2 is the corresponding resolution of R/J , then $\beta_2 \equiv 0$, $\alpha_2 \equiv 0$, and $\text{im } \beta_3 = K_3$.*

Proof. We first prove that $\dim_k T_1^2 = 10$. Let $\mathbf{a} = \{a_1, \dots, a_5\}$ be a minimal generating set of J with the property that every four element subset of \mathbf{a} is a regular sequence; and let x'_i be the image of \bar{a}_i under the natural isomorphism

$$(4.15) \quad J/\mathfrak{m}J \xrightarrow{\cong} \text{Tor}_1^R(R/J, k).$$

It suffices to show that

$$(4.16) \quad \dim_k(T_1^2/(x'_1, \dots, \hat{x}'_i, \dots, x'_5)^2) = 4$$

for $i = 1, \dots, 5$. We establish (4.16) for $i = 5$; the other four cases follow from the symmetry of the situation. Let K be the complete intersection ideal (a_1, \dots, a_4) . Consider T_\bullet as described in (4.1). If $\varepsilon'_1, \dots, \varepsilon'_4$ is a basis for K_1 with $k_1(\varepsilon'_i) = a_i$, then it follows that

$$x' = \begin{bmatrix} \bar{\varepsilon}'_i \\ 0 \end{bmatrix} \quad \text{for } 1 \leq i \leq 4.$$

Let

$$x_5 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

It is not necessarily true that $x'_5 = x_5$; but we do know that $x'_5 = \lambda x_5 + x'$ for some unit $\lambda \in k$ and some $x' \in (x'_1, \dots, x'_4)$. The multiplication in T_\bullet can be read from (4.2):

$$x'_i x'_j = \begin{bmatrix} \overline{\varepsilon'_i \varepsilon'_j} \pmod{\text{im } \bar{\beta}_2} \\ 0 \end{bmatrix} \quad \text{and} \quad x'_i x_5 = \begin{bmatrix} 0 \\ \bar{\alpha}_1(\varepsilon'_i) \end{bmatrix}$$

for $1 \leq i, j \leq 4$. The hypothesis ensures that $t = 4$; so $\alpha_1(\varepsilon'_1), \dots, \alpha_1(\varepsilon'_4)$ is the beginning of a basis for L_1 . We have established that $x'_1 x'_5, x'_2 x'_5, x'_3 x'_5$, and $x'_4 x'_5$ generate a four dimensional subspace of $T_1^2/(x'_1, \dots, x'_4)^2$; therefore, (4.16) holds and $\dim T_1^2 = 10$.

Furthermore, now that we know that $\dim T_1^2 = 10$, we may read the preceding paragraph from bottom to top in order to conclude that $\text{im } \bar{\beta}_2 = 0$ for every resolution \mathbf{M} from Proposition 3.2. It is clear that $\text{rank } \bar{\alpha}_2 = \text{rank } \bar{\beta}_2 = 0$, and that $\text{rank } \bar{\beta}_3 = \text{rank } \bar{\alpha}_1 = t = 4$; consequently, (b) has been established.

To finish the proof of (a) we must show that $T_1^3 = 0$. Once again, we use (4.2) to see that

$$x'_i x'_j x'_l = \begin{bmatrix} \overline{\varepsilon'_i \varepsilon'_j \varepsilon'_l} \pmod{\text{im } \bar{\beta}_3} \\ 0 \end{bmatrix} \quad \text{and} \quad x'_i x'_j x_5 = \begin{bmatrix} 0 \\ \bar{\alpha}_2(\varepsilon'_i \varepsilon'_j) \end{bmatrix}$$

for $1 \leq i, j, l \leq 4$. The product $x'_i x'_j x'_l$ is equal to 0 because $\bar{\beta}_3$ is surjective; and $x'_i x'_j x_5 = 0$ because $\bar{\alpha}_2 = 0$. \square

We now subdivide Case 3 into two subcases:

Case 3A. There is a nonzero element $x \in T_1$ such that $xT_2 = 0$ and $xT_3 = 0$.

Case 3B. If $x \in T_1$ with $x \neq 0$, then either $xT_2 \neq 0$ or $xT_3 \neq 0$.

The proof of Theorem 1.5 in Case 3A. Let a be an element of J with the property that \bar{a} is sent to x under the isomorphism of (4.15), and K be a grade four complete intersection ideal such that $J = (K, a)$. Adopt the notation of the paragraph preceding (4.1) and apply Corollary 2.7 in order to pick a basis for \mathbb{L} so that the multiplication in $(\text{im } \bar{\alpha}_1) \cdot \bar{\mathbb{L}}$ is described by one of the cases (i)–(iv). Recall from part (b) of Lemma 4.14 that $\alpha_2 \equiv 0$; hence, the multiplication $(\text{im } \bar{\alpha}_1) \cdot \bar{\mathbb{L}}$ is actually described by either (iii) with $a = 0$ or (iv). Label the elements x_i, y_i, z_i , and w_i of T_\bullet exactly as was done in (4.6). (Keep in mind that $t = 4$ and $d = 0$.) Notice that $x_5 = \lambda x + x'$ for some unit $\lambda \in k$ and some $x' \in (x_1, \dots, x_4)$. We will know all of the multiplication in T_\bullet once we show that $T_2 \cdot T_2 = 0$. According to (4.2) it suffices to prove that $v'_1 p(v_1) \equiv 0$ for all $v_1, v'_1 \in L_1$; and therefore, by Remark 2.5, it suffices to show that $\bar{p}(\bar{v}_1) \in \bar{L}_1^2$. Since $x' \in (x_1, \dots, x_4)$, there is an element $\bar{\varepsilon} \in K_1$ such that

$$x' = \begin{bmatrix} \bar{\varepsilon} \\ 0 \end{bmatrix} \quad \text{in } T_1.$$

Recall that $xT_2 = 0$. Use (4.2) to compute that

$$\begin{aligned} \begin{bmatrix} 0 \\ \bar{\alpha}_1(\bar{\varepsilon})\bar{v}_1 \end{bmatrix} &= \begin{bmatrix} \bar{\varepsilon} \\ 0 \end{bmatrix} \begin{bmatrix} 0 \\ \bar{v}_1 \end{bmatrix} = x' \begin{bmatrix} 0 \\ \bar{v}_1 \end{bmatrix} = x_5 \begin{bmatrix} 0 \\ \bar{v}_1 \end{bmatrix} \\ &= \begin{bmatrix} 0 \\ \bar{1} \end{bmatrix} \begin{bmatrix} 0 \\ \bar{v}_1 \end{bmatrix} = \begin{bmatrix} 0 \\ \bar{p}(\bar{v}_1) \end{bmatrix} \in T_1 \cdot T_2. \end{aligned}$$

We conclude that $\bar{p}(\bar{v}_1) = \bar{\alpha}_1(\bar{\varepsilon})\bar{v}_1 \in \bar{L}_1^2$ and $T_2^2 = 0$.

Combine Lemma 4.14(a), together with the hypothesis $xT_2 = xT_3 = 0$ and the fact $T_2^2 = 0$, in order to see that $T_\bullet = S_\bullet \rtimes W$ for some trivial S_\bullet -module W where

$$S_\bullet = \begin{cases} \mathbf{F}[b] & \text{with } 0 \leq b, & \text{if (4.9) is described by (4.10), and} \\ \mathbf{F}^{(j)} & \text{with } 2 \leq j \leq 4, & \text{if (4.9) is described by (4.11).} \end{cases}$$

The proof of Theorem 1.5 in Case 3A is complete.

Case 3B is the most interesting case. In Lemma 4.17 we record the consequences in T_\bullet of the observation that the multiplication $(\text{im } \bar{\alpha}_1) \cdot \bar{\mathbb{L}}$ must be described by part (iv) of Corollary 2.7. This result gives many incomplete multiplication tables for T_\bullet . In Lemma 4.18 we paste the incomplete multiplication tables of Lemma 4.17 together to learn all of the multiplication in T_\bullet , except the multiplication T_2^2 . The proof of Lemma 4.20 is where the hard work takes place in Case 3B with $T_2^2 \neq 0$.

Lemma 4.17. *Adopt the notation and hypotheses of Case 3B. If X_1 is a four dimensional subspace of T_1 , then there are elements $y_1 \in T_2$ and $w_1 \in T_4$, and there are subspaces $Y_1 \subseteq T_2$, $Z_1 \subseteq T_3$, and $Z'_1 \subseteq T_3$ such that $T_2 = ky_1 \oplus Y_1$, $T_3 = Z_1 \oplus Z'_1$, and*

- (a) $\dim(y_1 \cdot X_1) = 4$,
- (b) $X_1 \cdot Y_1 = 0$,
- (c) $X_1 \cdot T_3 \subseteq kw_1$,
- (d) *the multiplication map $X_1 \otimes Z_1 \rightarrow kw_1$ is a perfect pairing*,
- (e) $X_1 \cdot Z'_1 = 0$, and
- (f) $X_1 \cdot T_2 \subseteq Z'_1$.

Proof. Select a grade four complete intersection K with the property that the image of \bar{K} under (4.15) is X_1 . Use K to calculate multiplication in T_\bullet as described in (4.1) and (4.2). Observe that the elements x_1, \dots, x_4 , which are defined above (4.6), form a basis for X_1 . We know from Lemma 4.14(a) that $X_1 \cdot T_1^2 = 0$; consequently, all of the multiplication in $X_1 \cdot T_2$ and $X_1 \cdot T_3$ is given in (4.9). Recall the hypothesis that if x is a nonzero element of X_1 , then either $xT_2 \neq 0$ or $xT_3 \neq 0$. It follows that the multiplication in $X_1 \cdot T_2$ and $X_1 \cdot T_3$ is described by (4.11) with $j = 4$. There is no difficulty seeing that the multiplication of (4.11), with $j = 4$, is the same as the coordinate-free description which is given in the statement of the result. \square

Lemma 4.18. *If the notation and hypotheses of Case 3B are adopted, then there are elements $y \in T_2$ and $w \in T_4$, and there are subspaces $Y \subseteq T_2$, $Z \subseteq T_3$, and $Z' \subseteq T_3$ such that $T_2 = ky \oplus Y$, $T_3 = Z \oplus Z'$, and*

- (a) $\dim(y \cdot T_1) = 5$,
- (b) $T_1 \cdot Y = 0$,
- (c) $T_1 \cdot T_3 \subseteq kw$,
- (d) *the multiplication map $T_1 \otimes Z \rightarrow kw$ is a perfect pairing*,
- (e) $T_1 \cdot Z' = 0$, and
- (f) $T_1 \cdot T_2 \subseteq Z'$.

Before proving the above result, we notice that Lemmas 4.14 and 4.18 complete the proof in Case 3B when $T_2^2 = 0$.

Corollary 4.19. *If the notation and hypotheses of Case 3B are adopted and $T_2^2 = 0$, then T_\bullet has the form $\mathbb{F}^{(5)} \ltimes W$ for some trivial $\mathbb{F}^{(5)}$ -module W .*

Proof of Lemma 4.18. Let X_1 and X_2 be four dimensional subspaces of T_1 with $X_1 = X_2$. Apply Lemma 4.17 to find $y_i \in T_2$, $w_i \in T_4$, $Y_i \subseteq T_2$, $Z_i \subseteq T_3$, and $Z'_i \subseteq T_3$ with $\dim(y_i \cdot X_i) = 4$, $X_i \cdot Y_i = 0$, $X_i \cdot T_3 \subseteq kw_i$, the multiplication map $X_i \otimes Z_i \rightarrow kw_i$ a perfect pairing, and $X_i \cdot Z'_i = 0$, for $i = 1$ and $i = 2$. Let $y = y_1$, $w = w_1$, and $Y = Y_1$.

(b) Let x be a nonzero element of $X_1 \cap X_2$ and let $(x)^\perp = \{y_0 \in T_2 \mid xy_0 = 0\}$. It is clear that $Y_1 = (x)^\perp = Y_2$. Furthermore, we know that $X_1 + X_2 = T_1$; therefore, $Y \cdot T_1 = 0$.

(a) It suffices to show that $\dim(y \cdot X) = 4$ for every four dimensional subspace X of T_1 . The choice of X_2 is independent of our definition of y ; consequently, it suffices to show that $\dim(y \cdot X_2) = 4$. But, this fact follows from the following observations which we have already established: $ky \oplus Y = ky_2 \oplus Y$, $\dim(y_2 \cdot X_2) = 4$ and $Y \cdot X_2 = 0$.

(c) Take x from the proof of (b). The hypothesis ensures that $x \cdot T_3$ is a nonzero subspace of $(w_1) \cap (w_2)$. It follows that the one dimensional subspaces (w_1) and (w_2) of T_4 are equal. Use $X_1 + X_2 = T_1$ in order to conclude that $T_1 \cdot T_3 \subseteq (w)$.

(d) and (e) Let φ be the name of the map $T_3 \rightarrow \text{Hom}_k(T_1, kw)$ which is induced by the multiplication map $T_1 \otimes T_3 \rightarrow kw$, let x_1, \dots, x_5 be a fixed basis for T_1 and let x_1^*, \dots, x_5^* be the corresponding dual basis for $\text{Hom}_k(T_1, kw)$. Apply parts (d) and (e) of Lemma 4.17 to the subspace (x_1, \dots, x_4) of T_1 in order to find a basis for T_3 for which the matrix of φ is

$$\left[\begin{array}{c|c} I & 0 \\ \hline \lambda_1 \cdots \lambda_4 & \lambda_5 \cdots \lambda_n \end{array} \right]$$

for some $\lambda_i \in k$. If $\lambda_5 = \dots = \lambda_n = 0$, then $x \cdot T_3 = 0$ for $x = x_5 - \sum_{i=1}^4 \lambda_i x_i$ and this contradicts Lemma 4.17(d). Thus, $\lambda_i \neq 0$ for some i with $5 \leq i \leq n$ and a basis z_1, \dots, z_n for T_3 may be found for which the matrix of φ is $[I \ 0]$. Let $Z = (z_1, \dots, z_5)$ and $Z' = (z_6, \dots, z_n)$.

(f) It is immediate from Lemma 4.17(f) that $T_1 \cdot T_1 \cdot T_2 = 0$; hence, $T_1 \cdot T_2 \subseteq Z'$. \square

Lemma 4.20. *Adopt the notation and hypotheses of Case 3B with $T_2^2 \neq 0$. Let K be any complete intersection ideal with $K \subseteq J$ and J/K cyclic, \mathbb{L} be the minimal resolution of $R/(K : J)$ which is shown in (2.1), and $p: L_1 \rightarrow L_2$ be the map of (3.4). Then there exists an integer b , with $b \geq 6$, and there exists bases e_1, \dots, e_n for L_1 ; $f_1, \dots, f_{n-1}, f'_1, \dots, f'_{n-1}$ for L_2 ; g_1, \dots, g_n for L_3 ; and h for L_4 such that*

- (a) $K = (l_1(e_1), \dots, l_1(e_4))$,
- (b) (2.3) holds,
- (c) all products of basis vectors in $\bar{L}_1 \cdot \bar{L}_1$ and $\bar{L}_1 \cdot \bar{L}_2$ are zero except $e_b e_i = f_i$, $e_i f'_i \equiv g_b$, and $e_b f'_i \equiv -g_i$ for $1 \leq i \leq b-1$, and
- (d) $p(e_b) \equiv f'_3$ and $p(e_i) \equiv 0$ for all $i \neq 5$.

Proof. Let h be any generator for L_4 . We have two ways to view the multiplication in T_\bullet . On the one hand, we can use the multiplication in $\bar{\mathbb{L}}$ to compute $T_\bullet \cdot T_\bullet$ as described in (4.1) and (4.2). On the other hand, Lemma 4.18 gives a complete description of all of the multiplication in T_\bullet , except the multiplication $T_2 \cdot T_2$. In the present proof we use the interplay between these two descriptions of $T_\bullet \cdot T_\bullet$ in order to learn about the multiplication in $\bar{\mathbb{L}}$.

Let e_1, \dots, e_4 be elements in L_1 with $(l_1(e_1), \dots, l_1(e_4)) = K$. The hypothesis $t = 4$, ensures that e_1, \dots, e_4 is the beginning of a basis for L_1 . Let $\varepsilon_1, \dots, \varepsilon_4$ be the basis for K_1 which is defined by $\alpha_1(\varepsilon_i) = e_i$ for $1 \leq i \leq 4$, and let x_1, \dots, x_5 be the basis for T_1 which is given above (4.6). According to Lemma 4.18, we may decompose T_2 into $ky \oplus Y$ with

$$(4.21) \quad \dim(y \cdot T_1) = 5 \quad \text{and} \quad T_1 \cdot Y = 0.$$

We know from Lemma 4.14(a) that $T_1^2 \subseteq Y$; consequently,

$$\begin{bmatrix} \bar{\varphi} \\ 0 \end{bmatrix} \in Y \quad \text{and} \quad \begin{bmatrix} 0 \\ \bar{e}_i \end{bmatrix} \in Y$$

for all $\varphi \in K_2$ and for all i with $1 \leq i \leq 4$. It follows that we may modify y in order to assume that

$$y = \begin{bmatrix} 0 \\ \bar{e}_0 \end{bmatrix}$$

for some $e_0 \in L_1$. It also follows that L_1 decomposes into $Re_0 \oplus E$ where $(e_1, \dots, e_4) \subseteq E$ and E has the property that

$$\begin{bmatrix} 0 \\ \bar{e} \end{bmatrix} \in Y$$

for all $e \in E$. When the products of (4.21) are interpreted using (4.2), we see that $e_0e_1, \dots, e_0e_4, p(e_0)$ is the beginning of a basis for L_2 , $(\bar{e}_1, \dots, \bar{e}_4) \cdot \bar{E} = 0$, and $p(e) \equiv 0$ for all $e \in E$.

We next show that $E \cdot E \equiv 0$. We have observed that $\dim \bar{L}_1^2 \geq 4$; consequently, a quick look at Theorem 2.2 shows that the multiplication in $\bar{\mathbb{L}}$ is given in multiplication table (c). In other words, there is a decomposition $\bar{L}_1 = kv \oplus V$ with $V^2 = 0$. The fact that $\dim \bar{e}_0 \cdot \bar{L}_1 \geq 4$ ensures that $\bar{e}_0 \notin V$; and therefore, $\bar{L}_1 = k\bar{e}_0 \oplus V$. It is easy to select a nonzero element \bar{e} of $(\bar{e}_1, \dots, \bar{e}_4) \cap V$. Indeed, if we write $\bar{e}_i = \lambda_i \bar{e}_0 + v_i$ with $\lambda_i \in k$ and $v_i \in V$, then either $\lambda_1 = 0$ (in which case we take $\bar{e} = \bar{e}_1$) or $\lambda_1 \neq 0$ (in which case we take $\bar{e} = \lambda_1 \bar{e}_2 - \lambda_2 \bar{e}_1$). Let $\bar{e}' = \lambda \bar{e}_0 + v$ be an arbitrary element of \bar{E} . We know that $(\bar{e}_1, \dots, \bar{e}_4) \cdot \bar{E} = 0$, $V^2 = 0$ and $\dim \bar{e}_0(\bar{e}_1, \dots, \bar{e}_4) = 4$. It follows from

$$0 = \bar{e}'\bar{e} = (\lambda \bar{e}_0 + v)\bar{e} = \lambda \bar{e}_0 \bar{e},$$

that $\lambda = 0$; thus $\bar{E} \subseteq V$ and $E \cdot E \equiv 0$.

We may decompose E as $(e_1, \dots, e_4) \oplus E' \oplus E''$, where

$$(4.22) \quad \dim \bar{e}_0((\bar{e}_1, \dots, \bar{e}_4) \oplus \bar{E}') = \dim((\bar{e}_1, \dots, \bar{e}_4) \oplus \bar{E}')$$

and $e_0E'' \equiv 0$. Let $b - 1$ denote the dimension of the vector spaces on line (4.22). Rename e_0 by calling it e_b . Pick any basis e_{b+1}, \dots, e_n for E'' .

The hypothesis $T_2^2 \neq 0$ guarantees that there are elements v_1 and v'_1 in L_1 with $\bar{v}_1 \bar{p}(v'_1) \neq 0$. We have seen that $\bar{L}_1 = k\bar{e}_b \oplus \ker \bar{p}$; thus, $\bar{v}_1 \bar{p}(\bar{e}_b)$ is a nonzero element of \bar{L}_3 for some $v_1 \in L_1$. The multiplication $\bar{L}_1 \otimes \bar{L}_3 \rightarrow \bar{L}_4$ is a perfect pairing; consequently, $\bar{p}(\bar{e}_b) \cdot \bar{L}_1^2 \neq 0$. On the other hand, we have seen that $\bar{L}_1^2 = \bar{e}_b \bar{L}_1$. Thus, $\bar{p}(\bar{e}_b) \bar{e}_b$ is a nonzero element of \bar{L}_3 . The very last assertion in Theorem 3.5 shows that $\bar{p}(\bar{e}_b) \bar{e}_b (\bar{e}_1, \dots, \bar{e}_4) = 0$. Thus, we may select a basis e_5, \dots, e_{b-1} for E' with $\bar{p}(\bar{e}_b) \bar{e}_b \bar{e}_5 = \bar{h}$ and $\bar{p}(\bar{e}_b) \bar{e}_b \bar{e}_i = 0$ for $6 \leq i \leq b - 1$. Select the basis g_1, \dots, g_n for L_3 with the property $e_i g_j = \delta_{ij} h$. Observe that $\bar{p}(\bar{e}_b) \bar{e}_b = -\bar{g}_5$. Label $f_i = e_b e_i$ for $1 \leq i \leq b - 1$ and $f'_5 = p(e_b)$ in L_2 . Observe that $(f_1, \dots, f_{b-1})^2 = 0$ and $f_i f'_5 \equiv \delta_{i5} h$ for $1 \leq i \leq b - 1$. The proof of Lemma 2.6 (see [21] for details) allows us to extend $f_1, \dots, f_{b-1}, f'_5$ to be a basis $f_1, \dots, f_{b-1}, f'_1, \dots, f'_{b-1}$ of L_2 which satisfies (2.3). It is now clear that the basis we have constructed for \mathbb{L} satisfies conditions (a)–(d). \square

Corollary 4.23. *If the notation and hypotheses of Case 3B are adopted and $T_2^2 \neq 0$, then T_\bullet has the form $\mathbf{F}^* \ltimes W$ for some trivial \mathbf{F}^* -module W .*

Proof. Let K be any grade four complete intersection with $K \subseteq J$ and J/K cyclic. Let \mathbb{L} be the minimal resolution of $R/(K : J)$. Fix a basis for \mathbb{L} as described in Lemma 4.20. Compute multiplication in T_\bullet as described in (4.1) and (4.2). Consider the basis for T_\bullet which is given in (4.6). We know from Lemma 4.14 that $\dim T_1^2 = 10$ and $T_1^3 = 0$; furthermore, the individual

products in T_1^2 are given in (4.7). Use (4.2) and Lemma 4.20 to compute that all products in $T_1 \cdot T_2$ are zero except

$$x_i y_{b+6} = -z_{4+i} \quad \text{for } 1 \leq i \leq 4 \quad \text{and} \quad x_5 y_{b+6} = z_{n+8}.$$

The map $q: L_2 \rightarrow L_3$ is defined below (3.4). It follows from Lemma 4.20(d) that $q(f_5) \equiv g_b$, but $q(f_i) \equiv q(f'_j) \equiv 0$ for all $i \neq 5$ and all j . It is now clear that all products in $T_1 \cdot T_3$ are zero except

$$x_i z_{n+3+i} = -w_{b-4} \quad \text{for } 1 \leq i \leq 4 \quad \text{and} \quad x_5 z_9 = -w_{b-4}.$$

Finally, we use (4.2) and Lemma 4.20(d) once again to see that all products in $T_2 \cdot T_2$ are zero except

$$y_{11} y_{b+6} = w_{b-4} \quad \text{and} \quad y_{b+6} y_{b+6} = -2w_1.$$

Recall that 2 is a unit in k . There is no difficulty in verifying that $T_\bullet = \mathbf{F}^* \ltimes W$ for some trivial \mathbf{F}^* -module W . \square

The proof of Theorem 1.5 is complete.

5. EXAMPLES AND QUESTIONS

We begin this section by commenting on the hypotheses of Theorem 1.5. The hypothesis that k have square roots is used only in the proof of Lemma 2.6 and it is not a particularly annoying hypothesis. Indeed, if (R', m', k') is an arbitrary local ring, then the technique of residue field inflation (see, for example, [9, 0_{III} 10.3.1]) yields a faithfully flat extension (R, m, k) of R' for which k is closed under the square root operation. Many of the consequences of Theorem 1.5, applied to R , will descend back to R' ; however, we do not know if the conclusion of Theorem 1.5 will descend to R' . The hypothesis that 2 is a unit in R is also used only sporadically. There is a very trivial division by 2 at the end of the proof of Corollary 4.23; however, if the characteristic of k had been two, then we would have calculated the second divided power $y^{(2)}$ of each element y of T_2 and in particular, we would have written $y_{b+6}^{(2)} = -w_1$ instead of $y_{b+6}^2 = -2w_1$, thereby avoiding the division by 2. The more serious use of $\text{char } k \neq 2$ occurs when we appeal to Theorem 3.5. The proof of this result in [25] and [26] involves many divisions by 2. We presume (but have not proved) that these divisions can be circumvented.

We next consider the question of the existence of grade four almost complete intersection ideals with prescribed Tor-algebras.

Question 5.1. Let S_\bullet be a graded k -algebra from the list in Theorem 1.5. Does there exist a grade four almost complete intersection ideal J such that

$$(5.2) \quad \text{Tor}_\bullet^R(R/J, k) \cong S_\bullet \ltimes W$$

for some trivial S_\bullet -module W ?

We are able to answer most of Question 5.1. All of the potential Tor-algebras which are listed in Theorem 2.2 for grade four Gorenstein ideals actually do exist (see [14] for Gorenstein rings whose Tor-algebras are described in Theorem 2.2(c)); consequently, the proof in Cases 1 and 2 (especially Table 4.13) can be read as an algorithm for producing an ideal J for which (5.2) holds, provided S_\bullet is from the list \mathbf{A} , $\mathbf{B}[p]$, $\mathbf{C}[p]$, $\mathbf{C}^{(2)}$, \mathbf{C}^* , $\mathbf{D}[p]$, $\mathbf{D}^{(2)}$, $\mathbf{E}[p]$, $\mathbf{E}^{(2)}$, and $\mathbf{E}^{(3)}$

with $0 \leq p$. In Examples 5.6, 5.7, and 5.8 we exhibit ideals J for which (5.2) holds with $S_\bullet = \mathbf{F}[0], \mathbf{F}[1], \mathbf{F}[2], \mathbf{F}[3], \mathbf{F}^{(2)}, \mathbf{F}^{(4)},$ and \mathbf{F}^* .

A more complete classification of Tor-algebras remains elusive.

Question 5.3. Let S_\bullet be a graded k -algebra from the list in Theorem 1.5. What are necessary and sufficient conditions on the vector space dimensions $\dim W_i$ in order that (5.2) hold with $W = \bigoplus_{i=0}^4 W_i$ for some grade four almost complete intersection J ?

For example, the proof of Theorem 1.5 shows that if (5.2) holds with $S_\bullet = \mathbf{C}^*$ for some grade four almost complete intersection J , then $W = 0$. (In fact, the entire resolution of R/J is known in this case.) On the other hand, every example that we have considered for which

$$(5.4) \quad \text{Tor}_\bullet^R(R/J, k) \cong \mathbf{F}^* \times W,$$

also has $W = 0$. We wonder if (5.4) implies that $W = 0$; we also wonder if a structure theorem exists for the minimal resolution of R/J for those J which satisfy (5.4).

Finally, the variable of linkage class should also be thrown into the question about the classification of Tor-algebras. A number of years ago, Matthew Miller and the present author knew many Gorenstein rings of projective dimension four with $T_1^2 = 0$. None of these rings were in the linkage class of a complete intersection (licci). We conjectured that if A is a licci Gorenstein ring of projective dimension four, then $T_1^2 \neq 0$, and we deduced a number of consequences assuming that the conjecture held. Most of the consequences of the conjecture [20] have since been proved [11]; furthermore, various attempts to gather evidence for the conjecture have netted results which are interesting in their own right [22]. In the meantime, we have shown that the conjecture itself is false. The following question remains unanswered.

Question 5.5. Suppose T_\bullet is the Tor-algebra of some Cohen-Macaulay ring. Does there exist a licci ring A with $T_\bullet(A) \cong T_\bullet$?

Example 5.6. Let $Y_{1 \times 5}$ be a generic matrix, $X_{5 \times 5}$ be a generic alternating matrix, and R be the local ring $k[X, Y]_{(X, Y)}$. Huneke and Ulrich [10, Proposition 5.8] introduced the grade four almost complete intersection $J = I_1(YX)$. One can compute that $\text{Tor}_\bullet^R(R/J, k) = \mathbf{F}^*$. The Huneke-Ulrich almost complete intersection ideals are closely related to the Huneke-Ulrich deviation two Gorenstein ideals which have been studied rather extensively; see [15, 13, 28].

Example 5.7. Let $Y_{1 \times 4}$ and $X_{4 \times 3}$ be generic matrices and v be an indeterminate. Consider the local ring $R = k[X, Y, v]_{(X, Y, v)}$. Let $I = (a_1, \dots, a_7)$ be the grade four Gorenstein ideal with $a_j = \sum_{i=1}^4 y_i x_{ij}$ for $1 \leq j \leq 3$ and $a_{4+j} = c_j + v y_j$ for $1 \leq j \leq 4$, where c_j is equal to $(-1)^{j+1}$ times the determinant of X with row j removed. (The ideal I is known as a Herzog ideal; see, for example, [19], [5, §3], or [23, Example 7.16].) If $J = (a_1, a_4, a_5, a_6) : I$, then (5.2) holds with $S_\bullet = \mathbf{F}[2]$ and W equal to $k(-2) \oplus k(-3)^8 \oplus k(-4)$. If $J = (y_1 a_3 + a_4, a_5, a_6, a_7) : I$, then (5.2) holds with $S_\bullet = \mathbf{F}[0]$ and $W = k(-2)^3 \oplus k(-3)^{12} \oplus k(-4)^3$.

Example 5.8. Let $I = (a_1, \dots, a_9)$ be the grade four Gorenstein ideal defined in [18] with $\tau = 5$, $x_{11} = 1$, and $x_{21} = x_{31} = x_{41} = x_{51} = x_{12} = x_{13} = 0$. If

$J = (a_i, a_j, a_k, a_l): I$, then

$$\begin{aligned}
 \mathrm{Tor}_\bullet^R(R/J, k) &= \mathbf{F}[1] \times (k(-2)^3 \oplus k(-3)^{12} \oplus k(-4)^3), \\
 &\quad \text{if } \{i, j, k, l\} = \{3, 5, 6, 7\}, \\
 &= \mathbf{F}[3] \times (k(-2)^1 \oplus k(-3)^8 \oplus k(-4)^1), \\
 &\quad \text{if } \{i, j, k, l\} = \{1, 5, 6, 7\}, \\
 &= \mathbf{F}^{(2)} \times (k(-2)^3 \oplus k(-3)^{10} \oplus k(-4)^3), \\
 &\quad \text{if } \{i, j, k, l\} = \{3, 5, 6, 9\}, \\
 &= \mathbf{F}^{(4)} \times (k(-2)^3 \oplus k(-3)^6 \oplus k(-4)^3), \\
 &\quad \text{if } \{i, j, k, l\} = \{2, 3, 6, 9\}.
 \end{aligned}$$

ACKNOWLEDGMENT

A large number of examples were calculated using the “script” capability of version 3.0 of the MACAULAY program. The author sends his thanks to Bayer and Stillman for creating and sharing this wonderful program.

REFERENCES

1. E. Assmus, *On the homology of local rings*, Illinois J. Math. **3** (1959), 187–199.
2. L. Avramov, *Homological asymptotics of modules over local rings*, Commutative Algebra, Mathematical Sciences Research Institute Publications, no. 15, Springer-Verlag, Berlin, Heidelberg, New York, 1989, pp. 33–62.
3. —, *Problems on infinite free resolutions*, Free Resolutions in Commutative Algebra and Algebraic Geometry, papers from the conference held in Sundance, Utah, May 1990, (D. Eisenbud and C. Huneke, eds.), Jones and Bartlett, Boston, Mass., 1992.
4. L. Avramov and E. Golod, *Homology algebra of the Koszul complex of a local Gorenstein ring*, Mat. Zametki **9** (1971), 53–58; English transl., Math. Notes **9** (1971), 30–32.
5. L. Avramov, A. Kustin, and M. Miller, *Poincaré series of modules over local rings of small embedding codimension or small linking number*, J. Algebra **118** (1988), 162–204.
6. D. Buchsbaum and D. Eisenbud, *Algebra structures for finite free resolutions, and some structure theorems for ideals of codimension 3*, Amer. J. Math. **99** (1977), 447–485.
7. H. Charalambous, E. G. Evans, and M. Miller, *Betti numbers for modules of finite length*, Proc. Amer. Math. Soc. **109** (1990), 63–70.
8. D. Eisenbud, *Homological algebra on a complete intersection, with an application to group representations*, Trans. Amer. Math. Soc. **260** (1980), 35–64.
9. A. Grothendieck, *Eléments de géométrie algébrique III*, Inst. Hautes. Études Sci. Publ. Math. **11** (1961).
10. C. Huneke and B. Ulrich, *Divisor class groups and deformations*, Amer. J. Math. **107** (1985), 1265–1303.
11. —, *The structure of linkage*, Ann. of Math. (2) **126** (1987), 277–334.
12. C. Jacobsson, A. Kustin, and M. Miller, *The Poincaré series of a codimension four Gorenstein ring is rational*, J. Pure Appl. Algebra **38** (1985), 255–275.
13. S. Kim, *Projective resolutions of generic order ideals*, Ph.D. thesis, University of Illinois, Urbana, 1988.
14. A. Kustin, *New examples of rigid Gorenstein unique factorization domains*, Comm. Algebra **12** (1984), 2409–2439.
15. —, *The minimal free resolutions of the Huneke-Ulrich deviation two Gorenstein ideals*, J. Algebra **100** (1986), 265–304.

16. —, *Gorenstein algebras of codimension four and characteristic two*, *Comm. Algebra* **15** (1987), 2417–2429.
17. A. Kustin and M. Miller, *Algebra structures on minimal resolutions of Gorenstein rings of embedding codimension four*, *Math. Z.* **173** (1980), 171–184.
18. —, *Structure theory for a class of grade four Gorenstein ideals*, *Trans. Amer. Math. Soc.* **270** (1982), 287–307.
19. —, *Multiplicative structure on resolutions of algebras defined by Herzog ideals*, *J. London Math. Soc.* (2) **28** (1983), 247–260.
20. —, *Tight double linkage of Gorenstein algebras*, *J. Algebra* **95** (1985), 384–397.
21. —, *Classification of the Tor-algebras of codimension four Gorenstein local rings*, *Math. Z.* **190** (1985), 341–355.
22. A. Kustin, M. Miller, and B. Ulrich, *Linkage theory for algebras with pure resolutions*, *J. Algebra* **102** (1986), 199–228.
23. A. Kustin, M. Miller, and B. Ulrich, *Generating a residual intersection*, *J. Algebra* **146** (1992), 335–384.
24. M. Miller, *Multiplicative structures on finite free resolutions*, *Free Resolutions in Commutative Algebra and Algebraic Geometry*, papers from the conference held in Sundance, Utah, May 1990, (D. Eisenbud and C. Huneke, eds.), Jones and Bartlett, Boston, Mass., 1992.
25. S. Palmer, *Algebra structures on resolutions of rings defined by grade four almost complete intersections*, Ph.D. thesis, University of South Carolina, 1990.
26. —, *Algebra structures on resolutions of rings defined by grade four almost complete intersections*, *J. Algebra* (to appear).
27. J.-P. Serre, *Sur la dimension homologique des anneaux et des modules noetheriens*, *Proc. Internat. Sympos. Algebraic Number Theory*, 1955, Science Council of Japan, Tokyo, 1956, pp. 175–189.
28. H. Srinivasan, *Minimal algebra resolutions for cyclic modules defined by Huneke-Ulrich ideals*, *J. Algebra* **137** (1991), 433–472.
29. J. Tate, *Homology of noetherian rings and local rings*, *Illinois J. Math.* **1** (1957), 14–27.
30. J. Weyman, *On the structure of free resolutions of length 3*, *J. Algebra* **126** (1989), 1–33.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTH CAROLINA, COLUMBIA, SOUTH CAROLINA 29208

E-mail address: kustin@milo.math.sc Carolina.edu