

## CLASSIFICATIONS OF BAIRE-1 FUNCTIONS AND $c_0$ -SPREADING MODELS

V. FARMAKI

**ABSTRACT.** We prove that if for a bounded function  $f$  defined on a compact space  $K$  there exists a bounded sequence  $(f_n)$  of continuous functions, with spreading model of order  $\xi$ ,  $1 \leq \xi < \omega_1$ , equivalent to the summing basis of  $c_0$ , converging pointwise to  $f$ , then  $r_{\text{ND}}(f) > \omega^\xi$  (the index  $r_{\text{ND}}$  as defined by A. Kechris and A. Louveau). As a corollary of this result we have that the Banach spaces  $V_\xi(K)$ ,  $1 \leq \xi < \omega_1$ , which previously defined by the author, consist of functions with rank greater than  $\omega^\xi$ . For the case  $\xi = 1$  we have the equality of these classes. For every countable ordinal number  $\xi$  we construct a function  $S$  with  $r_{\text{ND}}(S) > \omega^\xi$ . Defining the notion of null-coefficient sequences of order  $\xi$ ,  $1 \leq \xi < \omega_1$ , we prove that every bounded sequence  $(f_n)$  of continuous functions converging pointwise to a function  $f$  with  $r_{\text{ND}}(f) \leq \omega^\xi$  is a null-coefficient sequence of order  $\xi$ . As a corollary to this we have the following  $c_0$ -spreading model theorem: Every nontrivial, weak-Cauchy sequence in a Banach space either has a convex block subsequence generating a spreading model equivalent to the summing basis of  $c_0$  or is a null-coefficient sequence of order 1.

### INTRODUCTION

In the last few years various classifications of the class  $B_1(K)$  of bounded Baire-1 functions on a compact metric space  $K$  were given by many authors (see [1, 7, 8]). Recently in [5] the class  $B_1(K)$  was classified into a transfinite, decreasing hierarchy  $V_\xi(K)$ ,  $1 \leq \xi < \omega_1$ , of Banach spaces. The first space coincides with  $B_{1/4}(K)$ , which was first defined in [7]; and the intersection of all  $V_\xi(K)$  is equal to the space  $\text{DBSC}(K)$  of differences of bounded semicontinuous functions on  $K$ . As proved in [7] and [5],  $f \in B_{1/4}(K)$  if and only if there exists a sequence  $(f_n)$  of continuous functions on  $K$  converging pointwise to  $f$  and generating a spreading model equivalent to the summing basis of  $c_0$ . Extending the notion of spreading models in [5], it is proved that the functions in  $V_\xi(K)$  have a stronger property, namely, that there exists a sequence of continuous functions on  $K$  with spreading model of order  $\xi$  equivalent to the summing basis of  $c_0$ , converging pointwise to  $f$ .

A. Kechris and A. Louveau in [8] defined a natural rank  $r_{\text{ND}}$  on every bounded function  $f$  defined on a compact metric space  $K$  not in  $\text{DBSC}(K)$ , which has values of the form  $\omega^\xi$  for countable ordinals  $\xi$  [6] (by [8] all such ordinals are obtained). With a different terminology but equivalent formulation

---

Received by the editors April 5, 1993 and, in revised form, January 26, 1994.  
1991 *Mathematics Subject Classification.* Primary 26A21; Secondary 46B25.

©1994 American Mathematical Society

this index is used by H. Rosenthal in [9] to prove the important result: that every bounded sequence  $(f_n)$  of continuous functions on  $K$  converging pointwise to a function  $f$  not in  $\text{DBSC}(K)$  has a strongly summing subsequence. From this result and the characterization of functions in  $\text{DBSC}(K)$  given by C. Bessaga and A. Pelczynski [4], there follows the  $c_0$ -theorem of Rosenthal, namely, that every nontrivial, weak-Cauchy sequence in a Banach space has either a strongly summing subsequence or a convex block basis equivalent to the summing basis of  $c_0$ .

In this paper we give a relation between the rank  $r_{\text{ND}}$  and the functions which are pointwise limits of sequences of continuous functions with spreading model of order  $\xi$ ,  $1 \leq \xi < \omega_1$ , equivalent to the summing basis of  $c_0$ . Namely, we prove (Theorem 9) that if for a bounded function  $f$  defined on a compact metric space  $K$  there exists a bounded sequence  $(f_n)$  of continuous functions on  $K$ , with spreading model of order  $\xi$  ( $1 \leq \xi < \omega_1$ ), equivalent to the summing basis of  $c_0$ , converging pointwise to  $f$ , then  $r_{\text{ND}}(f) > \omega^\xi$ . As a corollary of this result we have that for every  $1 \leq \xi < \omega_1$

$$V_\xi(K) \subseteq \{f \in B_1(K) : r_{\text{ND}}(f) > \omega^\xi\}.$$

For the case  $\xi = 1$  we have the equality of these classes. Finally, for every countable ordinal number  $\xi$  we construct a linear, Baire-1 function  $S$  on a compact metric space  $K$  which is not in  $\text{DBSC}(K)$  and prove that  $r_{\text{ND}}(S) > \omega^\xi$  using Theorem 9.

Defining the notion of null-coefficient sequences of order  $\xi$ ,  $1 \leq \xi < \omega_1$ , we prove a result similar to Rosenthal's for the case of functions with rank less or equal to  $\omega^\xi$ . Namely, we prove that every bounded sequence  $(f_n)$  of continuous functions converging pointwise to a function  $f$  with  $r_{\text{ND}}(f) \leq \omega^\xi$  ( $1 \leq \xi < \omega$ ) is null-coefficient of order  $\xi$  (Theorem 14). In particular (case  $\xi = 1$ ) it is proved that  $f \notin B_{1/4}(K)$  if and only if every bounded sequence of continuous functions converging pointwise to  $f$  is null-coefficient of order 1. As a corollary to this and the characterization of functions in  $B_{1/4}(K) \setminus C(K)$  given in [5] we have the following  $c_0$ -spreading model theorem: Every nontrivial, weak-Cauchy sequence in a Banach space either has a convex block subsequence generating a spreading model equivalent to the summing basis of  $c_0$  or is a null-coefficient sequence of order 1 (Theorem 18).

We will use standard terminology and notation. For completeness we will give some definitions and notation which we will use in the following.

Let  $K$  be a compact, metrizable space. The class of continuous functions on  $K$  is denoted by  $C(K)$  and the class of Baire-1 functions on  $K$  (i.e., the pointwise limits of uniformly bounded sequences of continuous functions on  $K$ ) by  $B_1(K)$ .  $\text{DBSC}(K)$  denotes the subclass of  $B_1(K)$  consisting of differences of bounded semicontinuous functions. It is easy to see that

$$\text{DBSC}(K) = \left\{ f \in B_1(K) : \text{there exists } (f_n) \subseteq C(K) \right. \\ \left. \text{so that } f = \sum f_n \text{ and } \sum |f_n| \text{ is bounded} \right\}.$$

The class  $\text{DBSC}(K)$  is a Banach space with respect to the norm

$$\|f\|_D = \inf \left\{ \left\| \sum |f_n| \right\|_\infty : (f_n) \subseteq C(K) \text{ and } \sum f_n = f \right\}.$$

It is not hard to check that  $\|f\|_\infty \leq \|f\|_D$ , but the two norms are not equivalent in general. The norm-closure of  $\text{DBSC}(K)$  is denoted by  $B_{1/2}(K)$  in [7]. In the same paper the authors define the subclass  $B_{1/4}(K)$  by

$$B_{1/4}(K) = \left\{ f \in B_1(K) : \text{there exists } (f_n) \subseteq \text{DBSC}(K) \right. \\ \left. \text{such that } \|f_n - f\|_\infty \rightarrow 0 \text{ and } \sup_n \|f_n\|_D < \infty \right\}.$$

The space  $B_{1/4}(K)$  is complete with respect to the norm

$$\|f\|_{1/4} = \inf \left\{ \sup_n \|f_n\|_D : (f_n) \subseteq \text{DBSC}(K) \text{ and } \|f_n - f\|_\infty \rightarrow 0 \right\}.$$

In [5] this definition was extended in the transfinite as follows: Let

$$V_1(K) = B_{1/4}(K) \text{ and } \| \cdot \|_1 = \| \cdot \|_{1/4}.$$

If the normed space  $(V_\xi(K), \| \cdot \|_\xi)$  has been defined, then

$$V_{\xi+1}(K) = \left\{ f \in B_1(K) : \text{there exists } (f_n) \subseteq \text{DBSC}(K) \right. \\ \left. \text{with } \|f_n - f\|_\xi \rightarrow 0 \text{ and } \sup \|f_n\|_D < \infty \right\}$$

and

$$\|f\|_{\xi+1} = \inf \left\{ \sup_n \|f_n\|_D : (f_n) \subseteq \text{DBSC}(K) \text{ and } \|f_n - f\|_\xi \rightarrow 0 \right\}.$$

Finally, for a limit ordinal  $\xi$

$$\|f\|_\xi = \sup \{ \|f\|_\beta : 1 \leq \beta < \xi \} \text{ for every } f \in \bigcap_{\beta < \xi} V_\beta(K)$$

and

$$V_\xi(K) = \{ f \in B_1(K) : \|f\|_\xi < \infty \}.$$

The spaces  $(V_\xi(K), \| \cdot \|_\xi)$ ,  $1 \leq \xi < \omega_1$ , are complete, and their intersection coincides with  $\text{DBSC}(K)$  [5]. It is easy to see that  $V_\xi(K) \subseteq V_\beta(K)$  and  $\|f\|_\infty \leq \|f\|_\beta \leq \|f\|_\xi$  for every  $f \in V_\xi(K)$  and  $\beta < \xi < \omega_1$ . According to [7] and [5], the functions in  $B_{1/4}(K) \setminus C(K)$  are characterized in terms of  $c_0$ -spreading models and the functions in  $V_\xi(K) \setminus C(K)$  have an analogous stronger property. As we will need these results, we include a precise statement:

Let  $(x_n)$  be a seminormalized basic sequence in a Banach space  $X$ . A basic sequence  $(e_n)$  is said to be a spreading model of  $(x_n)$  if for every  $k \in \mathbb{N}$  and  $\varepsilon > 0$  there exists  $m \in \mathbb{N}$  so that if  $m < n_1 < n_2 < \dots < n_k$ , then

$$\left\| \sum_{i=1}^k \lambda_i x_{n_i} \right\| - \left\| \sum_{i=1}^k \lambda_i e_i \right\| < \varepsilon \text{ for all scalars } \lambda_1, \dots, \lambda_k \text{ with } \max_{1 \leq i < k} |\lambda_i| \leq 1.$$

Every seminormalized basic sequence has a subsequence generating a spreading model.

If  $H, F$  are two finite subsets of  $\mathbb{N}$ , we denote  $H < F$  iff  $\max H < \min F$ .

The summing basis  $(s_n)$  of  $c_0$  is characterized by

$$\left\| \sum_{i=1}^\infty \lambda_i s_i \right\| = \sup_n \left| \sum_{i=1}^n \lambda_i \right|.$$

**Definition 1** [1]. For every limit ordinal  $\xi$ , let  $(\xi_n)$  be a sequence of ordinal numbers strictly increasing to  $\xi$ . We define:

$$\mathcal{F}_0 = \{\{n\} : n \in \mathbb{N}\},$$

$$\mathcal{F}_{\xi+1} = \{F \subseteq \mathbb{N} : F \subseteq F_1 \cup \dots \cup F_n \text{ with } \{n\} \leq F_1 < \dots < F_n \text{ and } F_i \in \mathcal{F}_\xi, \\ i = 1, \dots, n\},$$

and if  $\xi$  is a limit ordinal

$$\mathcal{F}_\xi = \{F \subseteq \mathbb{N} : F \in \mathcal{F}_{\xi_n} \text{ and } n \leq \min F\}.$$

**Definition 2.** Let  $X$  be a Banach space and  $(x_n)$  a sequence in  $X$ . We say that  $(x_n)$  has a spreading model of order  $\xi$  equivalent (or  $\delta$ -equivalent) to the summing basis of  $c_0$  if there exists  $\delta > 0$  such that

$$(1/\delta) \left\| \sum_{i=1}^k \lambda_i s_i \right\|_\infty \leq \left\| \sum_{i=1}^k \lambda_i x_{n_i} \right\| \leq \delta \left\| \sum_{i=1}^k \lambda_i s_i \right\|_\infty,$$

for every  $(n_1, \dots, n_k) \in \mathcal{F}_\xi$  and scalars  $\lambda_1, \dots, \lambda_k$ .

It is easy to see that a sequence  $(y_n)$  in  $X$  has a subsequence generating a spreading model equivalent to the summing basis of  $c_0$  if and only if it has a subsequence with spreading model of order 1 equivalent to the summing basis of  $c_0$ .

**Theorem 3** [5, 7]. Let  $K$  be a compact metric space,  $f$  a real bounded function on  $K$ , and  $\xi$  a countable ordinal number. If  $f \in V_\xi(K) \setminus C(K)$ , then there exists a sequence  $(f_n) \subseteq C(K)$ , with spreading model of order  $\xi$  (for every choice of  $(\mathcal{F}_\xi)$ ) equivalent to the summing basis of  $c_0$ , converging pointwise to  $f$ . Moreover,  $f \in B_{1/4}(K) \setminus C(K)$  if and only if there exists  $(f_n) \subseteq C(K)$ , with spreading model (or order 1) equivalent to the summing basis of  $c_0$ , converging pointwise to  $f$ .

In [8] the authors define a natural rank  $r_{ND}$  on every bounded function defined on a compact metric space  $K$ , as follows:

Let  $f$  be a bounded function on  $K$ . One defines the upper regularization of  $f$ ,  $ur(f)$  (usually denoted by  $\hat{f}$ ), by

$$ur(f) = \inf\{g : g \in C(K) \text{ and } g \geq f\}.$$

The function  $ur(f)$  is upper semicontinuous, and one has

$$ur(f)(x) = \overline{\lim}_{y \rightarrow x} f(y) = \max\{L \in [-\infty, \infty] : \exists x_n \rightarrow x, f(x_n) \rightarrow L\}$$

$$= \inf \left\{ \sup_{y \in U} f(y) : U \text{ is a neighbourhood of } x \right\}.$$

In [8] the authors associate with each bounded function  $f$  an increasing sequence  $(f_\xi)_{1 \leq \xi < \omega_1}$  of upper semicontinuous functions. In a different formulation (but equivalently) in [9] the author defines an increasing sequence  $(u_\xi(f))_{1 \leq \xi < \omega_1}$  as

$$u_1(f) = ur(ur(f) - f).$$

If  $u_\xi(f)$  is defined,

$$u_{\xi+1}(f) = ur(ur(u_\xi(f) + f) - f).$$

For a limit  $\xi$ ,  $u_\xi(f)$  is defined if and only if  $u_\beta(f)$  is defined for all  $\beta < \xi$  and  $\sup_{\beta < \xi} u_\beta(f)$  is bounded, and then

$$u_\xi(f) = \text{ur} \left( \sup_{\beta < \xi} u_\beta(f) \right).$$

According to [8],  $f$  is in  $\text{DBSC}(K)$  if and only if  $u_\xi(f)$  is defined for all  $\xi < \omega_1$  or, equivalently, if there exists a  $\xi < \omega_1$  such that  $u_\xi(f)$  is defined and  $u_{\xi+1}(f) = u_\xi(f)$ . Hence, to every bounded function  $f$  on  $K$  there corresponds a rank:

$$r_{\text{ND}}(f) = \inf\{1 \leq \xi < \omega_1 : u_\xi(f) \text{ is undefined}\}, \quad \text{if such a } \xi \text{ exists}$$

and  $r_{\text{ND}}(f) = \omega_1$  otherwise.

Note that the values of this rank are always limit ordinals. It is proved in [6] that if  $f \notin \text{DBSC}(K)$ , then  $r_{\text{ND}}(f) = \omega^\xi$  for some  $1 \leq \xi < \omega_1$  (by [8] all such ordinals are obtained) according to the following lemma.

**Lemma 4** [6]. *Let  $f$  be a bounded function on  $K$ , and suppose that  $u_\xi(f)$  is defined. Then  $u_{\xi \cdot n}(f)$  is defined and  $\|u_{\xi \cdot n}(f)\|_\infty \leq n \|u_\xi(f)\|_\infty$  for all  $n \in \mathbb{N}$ .*

*Proof.* Let  $M = \|u_\xi(f)\|_\infty$ . By induction  $u_{\xi+\beta}(f)$  is defined and  $M + u_\beta(f) \geq u_{\xi+\beta}(f)$  for every  $\beta \leq \xi$ . Finally,  $u_{\xi \cdot 2}(f)$  is defined and  $\|u_{\xi \cdot 2}(f)\|_\infty \leq 2 \|u_\xi(f)\|_\infty$ . The result then follows by induction on  $n$ .

In the proof of the main theorem we will use two lemmas which are proved in [9]. For completeness we give them below.

**Lemma 5** [9]. *Let  $f$  be a bounded real function defined on a compact metric space  $K$ ,  $\xi$  a countable ordinal number, and  $x \in K$ . Assume that  $0 < u_\xi(f)(x) < u_{\xi+1}(f)(x) = M < \infty$ . If  $U$  is an open neighborhood of  $x$  and  $0 < \varepsilon < 1$ , then there exist positive numbers  $\lambda, \delta$ , and  $x_1 \in U$  such that:*

- (i)  $(1 - \varepsilon_1)M < \lambda + \delta < (1 - \varepsilon_1)M$ ,
- (\*\*) (ii)  $x_1 \in \text{cl}(L)$ , where  $L = \{y \in K : \lambda \leq u_\xi(f)(y) < (1 - \varepsilon_1)M - \delta\}$ ,
- (iii)  $\overline{\lim}_{y \in L, y \rightarrow x_1} (f(y) - f(x_1)) = \delta$ .

**Lemma 6** [9]. *Let  $K$  be a compact metric space and  $(f_n) \subseteq C(K)$  converging pointwise to a bounded function  $f$ . If  $x_1 \in K$ ,  $L$  is a subset of  $K$  with  $x_1 \in \text{cl}(L)$ ,  $\delta = \overline{\lim}_{y \in L, y \rightarrow x_1} (f(y) - f(x_1)) > 0$ ,  $0 < \varepsilon < 1$ , and  $U$  is an open neighborhood of  $x_1$ , then there exists a subsequence  $(f_{n_i})$  of  $(f_n)$  such that given  $t > 1$  there exists an  $x_2 \in U \cap L$  satisfying:*

- (i)  $f(x_2) - f(x_1) > (1 - \varepsilon)\delta$ ,
- (ii)  $\sum_{1 \leq i < t} |f_{n_i}(x_2) - f(x_1)| < \varepsilon\delta$ ,
- (\*) (iii)  $\sum_{i \geq t} |f_{n_i}(x_2) - f(x_2)| < \varepsilon\delta$ .

We will define for every countable ordinal number  $\xi$  a family  $\mathcal{A}_\xi$  of finite subsets of  $\mathbb{N}$  such that  $\mathcal{A}_{\omega^\beta} = \mathcal{F}_\beta$  for every  $1 \leq \beta < \omega_1$ .

**Definition 7.** Let  $(\mathcal{F}_\xi)_{1 \leq \xi < \omega_1}$  be a family of finite subsets of  $\mathbb{N}$  as described in Definition 1. We define:

$$\mathcal{A}_1 = \{F \subseteq \mathbb{N} : \#F = 2\},$$

$$\mathcal{A}_{\xi+1} = \{F \subseteq \mathbb{N} : F \subseteq F_1 \cup F_2 \text{ where } F_1 < F_2, F_1 \in \mathcal{A}_1, \text{ and } F_2 \in \mathcal{A}_\xi\}.$$

If  $\xi$  is a limit ordinal, then  $\xi = \sum_{i=1}^m \rho_i \omega^{\beta_i}$ , where  $m, \rho_1, \dots, \rho_m \in \mathbb{N}$  and  $\beta_1, \dots, \beta_m$  are ordinal numbers with  $\beta_1 > \dots > \beta_m > 0$ . We define

$$\mathcal{A}_{\rho \omega^\beta} = \{F \subseteq \mathbb{N} : F \subseteq F_1 \cup \dots \cup F_\rho$$

$$\text{with } F_1 < \dots < F_\rho \text{ and } F_i \in \mathcal{F}_\beta \text{ for } i = 1, \dots, \rho\}$$

and in general

$$\mathcal{A}_\xi = \left\{ F \subseteq \mathbb{N} : F \subseteq F_1 \cup F_2 \text{ with } F_1 < F_2, F_1 \in \mathcal{A}_\gamma, \text{ and } F_2 \in \mathcal{A}_\beta \right.$$

$$\left. \text{where } \gamma = \rho_m \omega^{\beta_m}, \beta = \sum_{i=1}^{m-1} \rho_i \omega^{\beta_i} \right\}.$$

The following theorem is inspired by Theorem 4.1 of Rosenthal in [9].

**Theorem 8.** Let  $f$  be a real function defined on a compact metric space  $K$  and  $(f_n)$  a uniformly bounded sequence of continuous functions converging pointwise to  $f$ . Let also  $\xi$  be a countable ordinal and  $x \in K$  with  $0 < u_\xi(f)(x) < \infty$ . For every open neighborhood  $U$  of  $x$  and  $0 < \varepsilon < 1$  there exists a subsequence  $(f_{n_i})$  of  $(f_n)$  with the following properties: Given an infinite sequence of integers  $1 \leq t_1 < t_2 < \dots$  there exists  $F \in \mathcal{A}_\xi$ , where  $F = \{n_{t_1}, \dots, n_{t_k}\}$  ( $k \in \mathbb{N}$ ), and  $y \in U$  such that:

- (i)  $f_{n_{t_i}} - f_{n_{t_{i-1}}}(y) > 0$  for  $i = 1, \dots, k$  and
- (ii)  $\sum_{i=1}^k f_{n_{t_i}} - f_{n_{t_{i-1}}}(y) > (1 - \varepsilon)u_\xi(f)(x)$ .

*Proof.* The argument is similar to the proof of Theorem 4.1 in [9], except that additional work is required to locate  $F$  in  $\mathcal{A}_\xi$ .

Let  $1 < \varepsilon < 0$  and  $U$  be an open neighborhood of  $x$ .

Case  $\xi = 1$ . Let  $0 < \varepsilon_1 < 1$  with  $(1 - \varepsilon_1)(1 - 3\varepsilon_1) > 1 - \varepsilon$  and  $M = u_1(f)(x)$ . According to the definition there exists  $x_1 \in U$  with

$$(1 - \varepsilon_1)M < \text{ur}(f)(x_1) - f(x_1) = \delta < (1 - \varepsilon_1)M.$$

From Lemma 6 there exists a subsequence  $(f_{n_t})$  of  $(f_n)$  such that given  $t > 1$  there exists  $x_2 \in U$  satisfying (\*) (i)–(iii):

- (i)  $f(x_2) - f(x_1) > (1 - \varepsilon_1)\delta$ ,
  - (ii)  $\sum_{1 \leq i < t} |f_{n_i}(x_2) - f(x_1)| < \varepsilon_1\delta$ ,
  - (iii)  $\sum_{i \geq t} |f_{n_i}(x_2) - f(x_2)| < \varepsilon_1\delta$ .
- (\*)

Then given  $1 \leq t_1 < t_2$  there exists  $x_2 \in U$  satisfying (\*) for  $t = t_2$ . Thus  $F = \{n_{t_1}, n_{t_2}\} \in \mathcal{A}_1$  and

$$f_{n_{t_2}}(x_2) - f_{n_{t_1}}(x_2) > f(x_2) - f(x_1) - 2\varepsilon_1\delta$$

$$> (1 - \varepsilon_1)\delta - 2\varepsilon_1\delta > (1 - 3\varepsilon_1)(1 - \varepsilon_1)M > (1 - \varepsilon)M.$$

Case  $\xi + 1$ . Suppose the result is established for  $\xi$ . Let  $0 < u_{\xi+1}(f)(x) = M < \infty$  and  $0 < \varepsilon_1 < 1$  with  $(1 - \varepsilon_1)(1 - 3\varepsilon_1) > 1 - \varepsilon$ . We may assume that  $0 < u_\xi(f)(x) < u_{\xi+1}(f)(x)$ . Otherwise, if  $0 < u_\xi(f)(x) = u_{\xi+1}(f)(x)$ , the result follows by hypothesis and  $u_\xi(f)(x) = 0$  is impossible.

According to Lemma 5 there exist  $\lambda > 0$ ,  $\delta > 0$ , and  $x_1 \in U$  satisfying (\*\*)  
(i)–(iii):

- (i)  $(1 - \varepsilon_1)M < \lambda + \delta < (1 - \varepsilon_1)M$ ,
- (\*\*) (ii)  $x_1 \in \text{cl}(L)$ , where  $L = \{y \in K : \lambda \leq u_\xi(f)(y) < (1 - \varepsilon_1)M - \delta\}$ ,
- (iii)  $\overline{\lim}_{y \in L, y \rightarrow x_1} (f(y) - f(x_1)) = \delta$ .

From Lemma 6 there exists a subsequence  $(f_{n_t})$  of  $(f_n)$  such that given  $t > 1$  there exists  $x_2 \in U \cap L$  satisfying (\*) (i)–(iii). Without loss of generality we may assume that  $(f_n)$  itself has this property.

We will construct positive integers  $n_s$ ,  $s \in \mathbb{N}$ , and infinite subsets  $M_s$ ,  $s \in \mathbb{N}$ , of  $\mathbb{N}$  satisfying (\*\*\*) (i)–(viii):

- (i)  $n_1 < \dots < n_s < M_s$ ,
- (ii)  $M_s \subseteq M_{s-1}$ ,
- (iii)  $n_s = \min M_{s-1}$ .

Given  $r \in \mathbb{N}$  with  $1 < r \leq s$  there exist an open set  $V \subseteq U$  and  $x_2 \in V$  so that:

- (\*\*\*) (iv)  $f(x_2) - f(x_1) > (1 - \varepsilon_1)\delta$ ,
- (v)  $\sum_{1 \leq i < r} |f_{n_i}(y) - f(x_1)| < \varepsilon_1\delta$  for every  $y \in V$ ,
- (vi)  $\sum_{r \leq i \leq s} |f_{n_i}(y) - f(x_2)| < \varepsilon_1\delta$  for every  $y \in V$ ,
- (vii)  $\lambda \leq u_\xi(f)(x_2) < (1 + \varepsilon_1)M - \delta$ ,
- (viii) given  $\{m_1, m_2, \dots\} \subseteq M_s$  with  $1 \leq m_1 < m_2 < \dots$  there exists  $y \in V$  and  $F = \{m_1, m_2, \dots, m_{2k}\} \in \mathcal{A}_\xi$  ( $k \in \mathbb{N}$ ) such that  $f_{m_{2i}} - f_{m_{2i-1}}(y) > 0$ ,  $i = 1, \dots, k$ , and

$$\sum_{i=1}^k f_{m_{2i}} - f_{m_{2i-1}}(y) > (1 - \varepsilon_1)u_\xi(f)(x_2).$$

Let  $M_1 = \mathbb{N} \setminus \{1\}$ ,  $n_1 = 1$ , and  $n_2 = 2$ . We set  $s = 2 = r$ . As we assumed previously, there exists  $x_2 \in U \cap L$  such that

$$f(x_2) - f(x_1) > (1 - \varepsilon_1)\delta, \quad |f_1(x_2) - f(x_1)| < \varepsilon_1\delta, \quad \sum_{i \geq 2} |f_i(x_2) - f(x_2)| < \varepsilon_1\delta.$$

Using the continuity of  $f_1$  and  $f_2$  we can choose an open subset  $V$  of  $U$  with  $x_2 \in V$  such that  $|f_1(y) - f(x_1)| < \varepsilon_1\delta$  and  $|f_2(y) - f(x_2)| < \varepsilon_1\delta$  for every  $y \in V$ . Finally, using the induction hypothesis we choose an infinite subset  $M_2$  of  $\mathbb{N}$  with  $2 < M_2$  satisfying the conclusion of the theorem for the case  $\xi$ ,  $\varepsilon = \varepsilon_1$ ,  $U = V$ , and  $x = x_2$ . The proof for  $s = 2 = r$  is complete.

Let  $s \geq 2$ , and suppose that  $n_1, \dots, n_s, M_1, \dots, M_s$  have been constructed. Then  $n_{s+1} = \min M_s$ . We will construct infinite subsets  $M^1, M^2, \dots, M^{s+1}$  of  $\mathbb{N}$  such that  $M_s \setminus \{n_{s+1}\} = M^1 \supseteq M^2 \supseteq \dots \supseteq M^{s+1}$  and for every  $1 < r \leq s + 1$  there is an open subset  $V$  of  $U$  and  $x_2 \in V$  satisfying (\*\*\*) (iv)–(viii), where we replace “ $s$ ” by “ $s + 1$ ” in (vi) and “ $M_s$ ” by “ $M^r$ ” in (viii). Once this is done we set  $M_{s+1} = M^{s+1}$ .

Let  $1 < r \leq s + 1$ , and suppose  $M^{r-1}$  is defined. Using the property of  $(f_n)$  we can find  $x_2 \in U \cap L$  satisfying (\*) (i)-(iii) for  $t = n_r$ . Hence we have

$$f(x_2) - f(x_1) > (1 - \varepsilon_1)\delta,$$

$$\sum_{1 \leq i < r} |f_{n_i}(x_2) - f(x_1)| < \varepsilon_1\delta, \quad \sum_{r \leq i \leq s+1} |f_{n_i}(x_2) - f(x_2)| < \varepsilon_1\delta.$$

Using the continuity of  $f_{n_1}, \dots, f_{n_{s+1}}$  we can find an open subset  $V$  of  $U$  with  $x_2 \in V$  satisfying (\*\*\*) (v) and (\*\*\*) (vi) with “ $s$ ” replaced by “ $s + 1$ ”. At last by the induction hypothesis we choose  $M^r \subseteq M^{r-1}$  so that (\*\*\*) (viii) holds with “ $M_s$ ” replaced by “ $M^r$ ”.

The sequence  $(f_{n_i})$  satisfies the conclusion of the theorem for the case  $\xi + 1$ . Indeed, let  $1 \leq r_1 < r_2 < t_1 < t_2 < \dots$  be an infinite sequence of integers. We set  $m_i = n_{t_i}$  for every  $i \in \mathbb{N}$ . Then  $m_1 < m_2 < \dots$  and  $\{m_1, m_2, \dots\} \subseteq M_{t_1-1}$ . Hence from (\*\*\*) there exist an open subset  $V$  of  $U$  and  $x_2 \in V$  such that

$$f(x_2) - f(x_1) > (1 - \varepsilon_1)\delta,$$

$$|f_{n_{r_1}}(y) - f(x_1)| < \varepsilon_1\delta, \quad |f_{n_{r_2}}(y) - f(x_2)| < \varepsilon_1\delta \text{ for every } y \in V,$$

$$\lambda \leq u_\xi(f)(x_2) < (1 + \varepsilon_1)M - \delta.$$

Also there exist  $y \in V$  and  $F_2 = \{m_1, m_2, \dots, m_{2k}\} \in \mathcal{A}_\xi$  such that

$$f_{m_{2i}} - f_{m_{2i-1}}(y) > 0 \text{ for all } 1 \leq i \leq k \text{ and } \sum_{i=1}^k f_{m_{2i}} - f_{m_{2i-1}}(y) > (1 - \varepsilon_1)u_\xi(f)(x_2).$$

Set  $F = \{n_{r_1}, n_{r_2}\} \cup F_2 \in \mathcal{A}_{\xi+1}$ . Then

$$f_{n_{r_1}} - f_{n_{r_2}}(y) > f(x_2) - f(x_1) - 2\varepsilon_1\delta > (1 - \varepsilon_1)\delta - 2\varepsilon_1\delta > (1 - 3\varepsilon_1)\delta > 0$$

and

$$f_{n_{r_1}} - f_{n_{r_2}}(y) + \sum_{i=1}^k f_{n_{2i}} - f_{n_{2i-1}}(y) > (1 - 3\varepsilon_1)\delta + (1 - \varepsilon_1)u_\xi(f)(x_2)$$

$$\geq (1 - 3\varepsilon_1)\delta + (1 - \varepsilon_1)\lambda > (1 - 3\varepsilon_1)(\delta + \lambda)$$

$$> (1 - 3\varepsilon_1)(1 - \varepsilon_1)M > (1 - \varepsilon)M.$$

This finishes the proof of the theorem for the case  $\xi + 1$ .

*Case  $\xi$ : limit ordinal.* Suppose the theorem is proved for all ordinal numbers  $a$  with  $a < \xi$ . By the definition of  $u_\xi(f)(x)$  there exist  $x_1 \in U$  and  $a < \xi$  such that:

$$(1 - \varepsilon/2)u_\xi(f)(x) < u_a(f)(x_1) < (1 + \varepsilon/2)u_\xi(f)(x).$$

In particular, if  $\xi = \sum_{i=1}^m \rho_i \omega^{\beta_i}$ , where  $m, \rho_1, \dots, \rho_m$  are positive natural numbers and  $\beta_1 > \beta_2 > \dots > \beta_m > 0$  are countable ordinal numbers, then we can choose  $\mu \in \mathbb{N}$  such that  $a = \beta + \gamma$ , where  $\beta = \sum_{i=1}^{m-1} \rho_i \omega^{\beta_i}$  ( $\beta = 0$  if  $m = 1$ ) and  $\gamma = (\rho_m - 1)\omega^{\beta_m} + \mu\omega^\zeta$  if  $\beta_m = \zeta + 1$  or  $\gamma = (\rho_m - 1)\omega^{\beta_m} + \omega^{\zeta_\mu}$  if  $\beta_m$  is a limit ordinal and  $(\zeta_n)$  is the sequence of ordinal numbers strictly increasing to  $\beta_m$ .

Now, from the inductive hypothesis there exists a subsequence  $(f_{n_i})$  of  $(f_n)$  such that  $2\mu < n_1$  and given  $t_1 < t_2 < \dots$  an infinite sequence of integers there exists  $k \in \mathbb{N}$  and  $y \in U$  such that  $F = \{n_{t_1}, \dots, n_{t_{2k}}\} \in \mathcal{A}_a$ ,



$$f_{n_{2i}} - f_{n_{2i-1}}(y) > 0 \quad \text{for } i = 1, \dots, k$$

and

$$\sum_{i=1}^k f_{n_{2i}} - f_{n_{2i-1}}(y) > (1 - \varepsilon/2)u_a(f)(x_1) > (1 - \varepsilon)u_\xi(f)(x).$$

We claim that  $F \in \mathcal{A}_\xi$ . Indeed, we have that  $2\mu < F$ . If  $\xi = \omega$ , then  $F \in \mathcal{A}_\mu$  and since  $\#F \leq 2\mu$  we have that  $F \in \mathcal{F}_1 = \mathcal{A}_\omega$ . If  $\xi = \omega^{\zeta+1}$ , then  $F \in \mathcal{A}_{\mu\omega^\zeta}$  and since  $F \subseteq F_1 \cup \dots \cup F_\mu$ , where  $F_1 < \dots < F_\mu$  and  $F_i \in \mathcal{F}_\zeta$  for all  $i = 1, \dots, \mu$ , we have that  $F \in \mathcal{F}_{\zeta+1} = \mathcal{A}_\xi$ . If  $\xi = \omega^\beta$  and  $\beta$  is a limit ordinal, then if  $(\beta_n)$  is the sequence of ordinals increasing to  $\beta$ , we have  $F \in \mathcal{F}_{\beta_n}$  and finally  $F \in \mathcal{F}_\beta = \mathcal{A}_\xi$ . Let  $\xi = \rho\omega^\beta$ , where  $\rho \in \mathbb{N}$ ,  $\rho > 1$ , and  $1 \leq \beta < \omega_1$ . Then  $F \in \mathcal{A}_\gamma$ , where  $\gamma = (\rho - 1)\omega^\beta + \gamma_\mu$  with  $\gamma_\mu = \mu\omega^\zeta$  if  $\beta = \zeta + 1$  or  $\gamma_\mu = \omega^{\beta_\mu}$  if  $\beta$  is a limit ordinal. Since  $F \subseteq F_1 \cup \dots \cup F_\rho$ , where  $F_1 \in \mathcal{A}_{\gamma_\mu}$  and  $F_2 < \dots < F_\rho \in \mathcal{F}_\beta$ , it follows, analogously to the previous cases, that  $F_1 \in \mathcal{F}_\beta$  and finally that  $F \in \mathcal{A}_\xi$ . In general, if  $\xi = \sum_{i=1}^m \rho_i\omega^{\beta_i}$  with  $m > 1$ ,  $\rho_1, \dots, \rho_m > 0$ , and  $\beta_1 > \dots > \beta_m > 0$ , then  $F \in \mathcal{A}_{\beta+\gamma}$  and since  $F \subseteq F_1 \cup F_2$ , where  $F_1 \in \mathcal{A}_\gamma$ ,  $F_2 \in \mathcal{A}_\beta$ , and  $F_1 < F_2$ , we have, analogously to the previous cases, that  $F_1 \in \mathcal{A}_\zeta$ , where  $\zeta \in \rho_m\omega^{\beta_m}$  and finally that  $F \in \mathcal{A}_\xi$ . This completes the proof of the theorem.

From the previous theorem we have the main theorem:

**Theorem 9.** *Let  $f$  be a bounded function defined on a compact metric space  $K$ , let  $(f_n)$  be a uniformly bounded sequence of continuous functions converging pointwise to  $f$ , and let  $\xi$  be a countable ordinal number. If  $(f_n)$  has spreading model of order  $\xi$  equivalent to the summing basis of  $c_0$ , then  $u_{\omega^\xi}(f)$  is defined, equivalently  $r_{ND}(f) > \omega^\xi$ .*

*Proof.* Let  $(f_n)$  have spreading model of order  $\xi$   $\delta$ -equivalent (for some  $\delta > 0$ ) to the summing basis of  $c_0$ , and suppose  $u_{\omega^\xi}(f)$  is undefined. Let  $r_{ND}(f) = \omega^\zeta$ , with  $\zeta \leq \xi$ , according to Lemma 4. Hence there exist  $x \in K$  and a countable ordinal number  $a$ , with  $a < \omega^\zeta$ , such that  $2\delta < u_a(f)(x) < \infty$ . We can choose  $\mu \in \mathbb{N}$  such that  $a = \mu\omega^\beta$  if  $\zeta = \beta + 1$  or  $a = \omega^{\beta_\mu}$  if  $\zeta$  is a limit ordinal and  $(\zeta_n)$  is the sequence of ordinal numbers strictly increasing to  $\zeta$ .

From the definition of the families  $\mathcal{F}_\xi$ ,  $1 \leq \xi < \omega_1$ , it is easy to see that for every  $\zeta < \xi < \omega_1$  there exists  $v(\zeta, \xi) \in \mathbb{N}$  such that if  $F \in \mathcal{F}_\zeta$  and  $v(\zeta, \xi) < F$ , then  $F \in \mathcal{F}_\xi$  (see [2]).

Let  $v = \max(v(\zeta, \xi), \mu)$ . According to Theorem 8 there exist  $F \in \mathcal{A}_a$  with  $2v < F = \{n_1, \dots, n_{2k}\}$  ( $k \in \mathbb{N}$ ) and  $y \in K$  such that

$$\sum_{i=1}^k f_{n_{2i}} - f_{n_{2i-1}}(y) > (1/2)u_a(f)(x) > \delta.$$

Since  $2\mu < F$ , we have that  $F \in \mathcal{F}_\zeta$  (see the proof of Theorem 8, case  $\zeta$ : limit ordinal). Consequently, since  $v(\zeta, \xi) < F$ , we have that  $F \in \mathcal{F}_\xi$ . This is a contradiction, because  $(f_n)$  has spreading model of order  $\xi$   $\delta$ -equivalent to the summing basis of  $c_0$ . Hence  $u_{\omega^\xi}(f)$  is defined.

The following two corollaries are already proved in [6]. Here we give a proof using the previous theorem.

**Corollary 10.** *For every compact metric space  $K$  and countable ordinal number  $\xi$  we have  $V_\xi(K) \subseteq \{f \in B_1(K) : r_{ND}(f) > \omega^\xi\}$ .*

*Proof.* This is true according to the previous theorem and Theorem 3.

For the case  $\xi = 1$  the two classes are equal, according to the following:

**Corollary 11.** *Let  $K$  be a compact metric space and  $f$  a function on  $K$  which is not continuous. The following are equivalent:*

- (i)  $f \in B_{1/4}(K)$ ,
- (ii)  $r_{ND}(f) > \omega$ ,
- (iii) *there exists a bounded sequence  $(f_n) \subseteq C(K)$  converging pointwise to  $f$  and generating a spreading model equivalent to the summing basis of  $c_0$ .*

*Proof.* The equivalence of (i) and (iii) is proved in [7] and [5]. According to the previous corollary (i) implies (ii). That (ii) implies (i) is proved in [6].

After these results the following interesting problem remains:

*Problem.* Is it true that for every compact metric space  $K$  and every ordinal number  $\xi < \omega_1$  we have  $V_\xi(K) = \{f \in B_1(K) : r_{ND}(f) > \omega^\xi\}$ ?

For every countable ordinal number  $\xi$  we will construct a Baire-1 function which is not a difference of bounded semicontinuous functions and has rank greater than  $\omega^\xi$ .

**Example 12.** For every countable ordinal  $\xi$ , let  $T_\xi$  be the Tsirelson-like space which is defined by S. Argyros in [2]. For completeness we recall the definition of  $T_\xi$ .

Let  $x: \mathbb{N} \rightarrow \mathbb{R}$  be a finitely supported function. For every  $m \in \mathbb{N}$  set

$$\|x\|_0^\xi = \sup\{|x(p)| : p \in \mathbb{N}\} \text{ and}$$

$$\|x\|_{m+1}^\xi = \max \left\{ \|x\|_m^\xi, \frac{1}{2} \sup \sum_{i=1}^{k-1} \|x|_{p_i, p_{i+1} - 1}\|_m^\xi \text{ for all } (p_1, \dots, p_k) \in \mathcal{B}_\xi \right\},$$

where  $x|_{p, q}$  ( $p \leq q$ ) denotes the restriction of  $x$  on the set  $\{p, p+1, \dots, q\}$  and  $\mathcal{B}_\xi = \mathcal{F}_\xi \cup \{(n, p) : 2 \leq n < p\} \cup \{\emptyset\}$  for all  $1 \leq \xi < \omega_1$ . Finally, define

$$\|x\|^\xi = \liminf_{m \rightarrow \infty} \|x\|_m^\xi$$

$$= \max \left\{ \|x\|_0^\xi, \sup \frac{1}{2} \sum_{i=1}^{k-1} \|x|_{p_i, p_{i+1} - 1}\|^\xi \text{ for } \{p_1, \dots, p_k\} \in \mathcal{B}_\xi \right\}.$$

The space  $T_\xi$  is the completion of the linear space of all finitely supported functions with the norm  $\|\cdot\|^\xi$ . The usual basis  $(e_n)$  is an unconditional basis of  $T_\xi$  and, as proved in [2],  $T_\xi$  is reflexive.

Let  $X_\xi$  be the "Jamesification" of  $T_\xi$  [3]. Let us recall the definition.

For every finitely supported function  $x: \mathbb{N} \rightarrow \mathbb{R}$  define:

$$\|x\|_\xi = \sup \left\{ \left\| \sum_{j=1}^m (S_{n_j} - S_{p_{j-1}})(x) e_{p_j} \right\|^\xi : 1 \leq p_1 \leq n_1 \leq \dots \leq p_m \leq n_m \right\},$$

where  $S_n(x) = \sum_{i=1}^n x(i)$  for every  $n \in \mathbb{N}$ , and  $S_0(x) = 0$ . The space  $X_\xi$  is the completion of the linear space of all finitely supported functions with the norm  $\|\cdot\|_\xi$ .

As shown in [3] the unit vectors  $e_n$ ,  $n \in \mathbb{N}$ , form a boundedly complete normalized basis for  $X_\xi$ . Thus,  $X_\xi$  is isometric to the space  $Y_\xi^*$ , where  $Y_\xi = [e_n^*]_{n=1}^\infty$  and  $(e_n^*)$  is the sequence of biorthogonal functionals of  $(e_n)$ . Furthermore it was shown in [3] that  $Y_\xi$  is quasi-reflexive (of order one) and  $Y_\xi^{**}$  has a basis given by  $\{S, e_1^*, e_2^*, \dots\}$ , where  $S(\sum_{i=1}^\infty a_i e_i) = \sum_{i=1}^\infty a_i$ . Of course  $S_n = \sum_{i=1}^n e_i^*$  for every  $n \in \mathbb{N}$  and  $(S_n)$  converges to  $S$  in the  $w^*$ -topology. Hence  $S$  is a Baire-1 function restricted on  $K = (S_{Y_\xi^*}, w^*)$ .

Since  $c_0$  is not isomorphically embedding into  $Y_\xi$  [3] we have that  $S \notin \text{DBSC}(K)$ . We will prove that  $r_{\text{ND}}(S) > \omega^\xi$ . Let  $x \in K$  and  $F = (n_1, \dots, n_{2k}) \in \mathcal{F}_\xi$  ( $k \in \mathbb{N}$ ). From the definition of the norms and since  $(n_1 + 1, \dots, n_{2k-1} + 1, r) \in \mathcal{F}_\xi$  for  $r \in \mathbb{N}$  with  $r > n_{2k}$  we have

$$1 \geq \|x\|_\xi \geq \left\| \sum_{i=1}^k (S_{n_{2i}} - S_{n_{2i-1}})(x) e_{n_{2i-1}+1} \right\|_\xi \geq \frac{1}{2} \sum_{i=1}^k |S_{n_{2i}}(x) - S_{n_{2i-1}}(x)|.$$

If  $r_{\text{ND}}(S) \leq \omega^\xi$ , then we can find, analogously to the proof of Theorem 9 ( $\delta = 2$ ),  $y \in K$  and  $F = \{n_1, \dots, n_{2k}\} \in \mathcal{F}_\xi$  such that

$$\sum_{i=1}^k |S_{n_{2i}}(y) - S_{n_{2i-1}}(y)| > 2.$$

This is a contradiction; hence,  $r_{\text{ND}}(S) > \omega^\xi$ .

In [9] H. Rosenthal proved the fundamental result that if  $f \notin \text{DBSC}(K)$ , then every bounded sequence  $(f_n)$  in  $C(K)$  converging pointwise to  $f$  has a strongly summing subsequence. In this article we obtain a result, in the same spirit as the above, concerning the classes:

$$\{f \in B_1(K) : r_{\text{ND}}(f) \leq \omega^\xi\} \subseteq B_1(K) \setminus \text{DBSC}(K), \quad 1 \leq \xi < \omega_1.$$

This result requires the following new concept:

**Definition 13.** A sequence  $(x_n)$  in a Banach space is called null-coefficient (n.c.) of order  $\xi$ , where  $\xi$  is a countable ordinal number, if whenever the scalars  $(c_n)$  satisfy

$$\sup \left\{ \left\| \sum_{i=1}^k c_{n_{2i}} (x_{n_{2i}} - x_{n_{2i-1}}) \right\| : (n_1, \dots, n_{2k}) \in \mathcal{F}_\xi \right\} < \infty,$$

the sequence  $(c_n)$  converges to 0.

*Remark.* If a sequence  $(x_n)$  has spreading model of order  $\xi$  equivalent to the summing basis of  $c_0$ , then it is not null-coefficient. Indeed, take  $c_n = 1$  for every  $n \in \mathbb{N}$ .

**Theorem 14.** Let  $K$  be a compact metric space,  $f$  a bounded function on  $K$ ,  $(f_n)$  a bounded sequence of continuous functions on  $K$  converging pointwise to  $f$ , and  $\xi$  a countable ordinal number. If  $r_{\text{ND}}(f) \leq \omega^\xi$ , then  $(f_n)$  is null-coefficient of order  $\xi$ .

*Proof.* Let  $r_{\text{ND}}(f) \leq \omega^\xi$ . Then  $r_{\text{ND}}(f) = \omega^\zeta$  for some ordinal  $\zeta$  with  $\zeta \leq \xi$ , according to Lemma 4. We assume that  $(f_n)$  is not a null-coefficient sequence

of order  $\xi$ . Then there exists a sequence of scalars  $(c_n)$  and  $\varepsilon > 0$  such that

$$\sup \left\{ \left\| \sum_{i=1}^k c_{n_{2i}} (f_{n_{2i}} - f_{n_{2i-1}}) \right\|_{\infty} : (n_1, \dots, n_{2k}) \in \mathcal{F}_{\xi} \right\} \leq 1$$

and  $|c_n| > \varepsilon$  for infinite many  $n$ . Let  $(g_t)$  be a subsequence of  $(f_n)$  with  $g_t = f_{n_t}$  and  $c_{n_t} > \varepsilon$  for every  $t \in \mathbb{N}$  (otherwise set  $-c_n$  instead of  $c_n$ ).

Since  $r_{ND}(f) = \omega^{\zeta}$ , there exist  $x \in K$  and  $a < \omega^{\zeta}$  such that  $2/\varepsilon < u_a(f)(x) < \infty$ . We can choose  $\mu \in \mathbb{N}$  such that  $a = \mu\omega^{\beta}$  if  $\zeta = \beta + 1$  or  $a = \omega^{\zeta_{\mu}}$  if  $\zeta$  is a limit ordinal and  $(\zeta_n)$  is the sequence of ordinal numbers strictly increasing to  $\zeta$  (according to Definition 1).

Let  $v = \max(\mu, v(\zeta, \xi))$  (if  $F \in \mathcal{F}_{\xi}$  and  $v(\zeta, \xi) < F$ , then  $F \in \mathcal{F}_{\xi}$ ). From Theorem 8, there exist  $F \in \mathcal{A}_a$  with  $2v < F = \{n_{t_1}, \dots, n_{t_k}\}$  ( $k \in \mathbb{N}$ ) and  $y \in K$  such that  $g_{t_{2i}} - g_{t_{2i-1}}(y) > 0$  for all  $i = 1, \dots, k$  and

$$\sum_{i=1}^k g_{t_{2i}} - g_{t_{2i-1}}(y) > (1/2)u_a(f)(x) > 1/\varepsilon.$$

Then  $F \in \mathcal{F}_{\xi}$  (see the proof of Theorem 8, case  $\xi$ : limit ordinal) and consequently  $F \in \mathcal{F}_{\xi}$ . Also,

$$\sum_{i=1}^k c_{n_{t_{2i}}} (f_{n_{t_{2i}}} - f_{n_{t_{2i-1}}})(y) > 1.$$

This is a contradiction, since  $(n_{t_1}, \dots, n_{t_k}) \in \mathcal{F}_{\xi}$ . Thus,  $(f_n)$  is null-coefficient of order  $\xi$ .

For the case  $\xi = 1$ , after Corollary 11, we have the following characterization of functions not in  $B_{1/4}(K)$ :

**Theorem 15.** *Let  $K$  be a compact metric space and  $f \in B_1(K) \setminus C(K)$ . Then  $f$  is not in  $B_{1/4}(K)$  if and only if every uniformly bounded sequence of continuous functions on  $K$  converging pointwise to  $f$  is null-coefficient of order 1.*

*Proof.* If  $f \in B_1(K) \setminus B_{1/4}(K)$ , then  $r_{ND}(f) = \omega$  according to Corollary 11. From Theorem 14 we have that every bounded sequence  $(f_n) \subseteq C(K)$  converging pointwise to  $f$  is null-coefficient of order 1. On the other hand, if every bounded sequence of continuous functions on  $K$  converging pointwise to  $f$  is null-coefficient of order 1, then according to the remark there is no bounded sequence  $(f_n)$  in  $C(K)$  converging pointwise to  $f$  with spreading model (of order 1) equivalent to the summing basis of  $c_0$ . From Corollary 11, it follows that  $f \notin B_{1/4}(K)$ .

As a consequence of Theorems 3 and 15 we have the following dichotomy:

**Theorem 16.** *Let  $K$  be a compact metric space and  $f \in B_1(K) \setminus C(K)$ . Then, either there exists a bounded sequence  $(f_n) \subseteq C(K)$  converging pointwise to  $f$  and generating a spreading model equivalent to the summing basis of  $c_0$  or every uniformly bounded sequence of continuous functions converging pointwise to  $f$  is null-coefficient of order 1.*

**Corollary 17.** *Let  $K$  be a compact metric space,  $f \in B_1(K) \setminus C(K)$ , and  $(f_n)$  a bounded sequence in  $C(K)$  converging pointwise to  $f$ . Then either there exists a*

convex block subsequence of  $(f_n)$  generating a spreading model equivalent to the summing basis of  $c_0$  or every convex block subsequence of  $(f_n)$  is null-coefficient of order 1.

*Proof.* If  $f \in B_{1/4}(K) \setminus C(K)$ , then, according to [7] and [5],  $(f_n)$  has a convex block subsequence generating a spreading model equivalent to the summing basis of  $c_0$ . If  $f \notin B_{1/4}(K)$ , then Theorem 15 finishes the proof.

Now we will give the  $c_0$ -spreading model theorem:

**Theorem 18.** *Every weak-Cauchy and non-weakly convergent sequence in a separable Banach space either has a convex block subsequence generating a spreading model equivalent to the summing basis of  $c_0$  or is null-coefficient of order 1 (in fact, every convex block subsequence is null-coefficient of order 1).*

*Proof.* Let  $X$  be a separable Banach space, and let  $K$  denote the unit ball of the dual space  $X^*$  endowed with the weak\*-topology. If  $(x_n)$  is a weak-Cauchy and nonweakly convergent sequence in  $x$ , then let  $x^{**} \in X^{**} \setminus X$  be the weak\*-limit of  $(x_n)$ . The restriction of  $x^{**}$  to  $K$  is in  $B_1(K) \setminus C(K)$ . Theorem 17 finishes the proof.

#### REFERENCES

1. D. Alspach and S. Argyros, *Complexity of weakly null sequences*, *Dissertationes Math.* **321** (1992).
2. S. Argyros, *Banach spaces of the type of Tsirelson* (to appear).
3. S. Bellenot, R. Haydon, and E. Odell, *Quasi-reflexive and tree spaces constructed in the spirit of R. C. James*, *Contemp. Math.*, vol. 85, Amer. Math. Soc., Providence, R.I., 1989, pp. 19–43.
4. C. Bessaga and A. Pelczynski, *On bases and unconditional convergence of series in Banach spaces*, *Studia Math.* **17** (1958), 151–164.
5. V. Farmaki, *On Baire-1/4 functions and spreading models*, *Mathematika* (to appear).
6. V. Farmaki and A. Louveau, *On a classification of functions* (unpublished).
7. R. Haydon, E. Odell, and H. Rosenthal, *On certain classes of Baire-1 functions with applications to Banach space theory*, *Lecture Notes in Math.*, vol. 1470, Springer-Verlag, New York, 1991, pp. 1–35.
8. A. S. Kechris and A. Louveau, *A classification of Baire class 1 functions*, *Trans. Amer. Math. Soc.* **318** (1990), 209–236.
9. H. Rosenthal, *A characterization of Banach spaces containing  $c_0$*  (to appear).

DEPARTMENT OF MATHEMATICS, PANEPISTEMIOPOLIS, 15784, ATHENS, GREECE  
 E-mail address: vgeorgil@atlas.uoa.ariadne-t.gr