

NEW RESULTS IN THE PERTURBATION THEORY OF
MAXIMAL MONOTONE AND M -ACCRETIVE
OPERATORS IN BANACH SPACES

ATHANASSIOS G. KARTSATOS

ABSTRACT. Let X be a real Banach space and G a bounded, open and convex subset of X . The solvability of the fixed point problem $(*) Tx + Cx \ni x$ in $D(T) \cap \overline{G}$ is considered, where $T : X \supset D(T) \rightarrow 2^X$ is a possibly discontinuous m -dissipative operator and $C : \overline{G} \rightarrow X$ is completely continuous. It is assumed that X is uniformly convex, $D(T) \cap G \neq \emptyset$ and $(T + C)(D(T) \cap \partial G) \subset \overline{G}$. A result of Browder, concerning single-valued operators T that are either uniformly continuous or continuous with X^* uniformly convex, is extended to the present case. Browder's method cannot be applied in this setting, even in the single-valued case, because there is no class of permissible homeomorphisms. Let $\Gamma = \{\beta : \mathcal{R}_+ \rightarrow \mathcal{R}_+ ; \beta(r) \rightarrow 0 \text{ as } r \rightarrow \infty\}$. The effect of a weak boundary condition of the type $\langle u + Cx, x \rangle \geq -\beta(\|x\|)\|x\|^2$ on the range of operators $T + C$ is studied for m -accretive and maximal monotone operators T . Here, $\beta \in \Gamma$, $x \in D(T)$ with sufficiently large norm and $u \in Tx$. Various new eigenvalue results are given involving the solvability of $Tx + \lambda Cx \ni 0$ with respect to $(\lambda, x) \in (0, \infty) \times D(T)$. Several results do not require the continuity of the operator C . Four open problems are also given, the solution of which would improve upon certain results of the paper.

1. INTRODUCTION. PRELIMINARIES

The symbol X stands for a real Banach space with norm $\|\cdot\|$ and (normalized) duality mapping J . In what follows, "continuous" means "strongly continuous". We denote by \mathcal{R} , \mathcal{R}_+ the sets $(-\infty, \infty)$, $[0, \infty)$, respectively. The symbols $\text{int}D$, ∂D and \overline{D} denote the strong interior, boundary and closure of the set D , respectively. An operator $T : X \supset D(T) \rightarrow Y$, with Y another real Banach space, is "bounded" if it maps bounded subsets of $D(T)$ onto bounded sets. It is "compact" if it is continuous and maps bounded subsets of $D(T)$ onto relatively compact sets. It is "demicontinuous" if it maps strongly convergent sequences into weakly convergent sequences. It is "completely continuous" if it maps weakly convergent sequences into strongly convergent sequences. For an operator $T : X \rightarrow 2^Y$ we set $D(T) \equiv \{x \in X : Tx \neq \emptyset\}$. We always assume that $D(T) \neq \emptyset$, for every multi-valued operator T considered in this paper. Given $T : X \supset D(T) \rightarrow 2^Y$ and a set $G \subset D(T)$, we

Received by the editors February 7, 1995.

1991 *Mathematics Subject Classification*. Primary 47H17; Secondary 47B44, 47H09, 47H10.

Key words and phrases. m -accretive operator, maximal monotone operator, compact perturbation, compact resolvent, eigenvalues for nonlinear operators, fixed point theory, degree theory.

The results of this paper were announced in a lecture at the International Conference on Nonlinear Differential Equations, Kiev, Ukraine, August 21-27, 1995.

set $TG = \bigcup\{Tx : x \in G\}$. An operator $T : X \supset D(T) \rightarrow 2^X$ is “accretive” if for every $x, y \in D(T)$ there exists $j \in J(x - y)$ such that

$$(1) \quad \langle u - v, j \rangle \geq 0 \text{ for every } u \in Tx, v \in Ty.$$

An accretive operator T is “strongly accretive” if 0 in the right-hand side of (1) is replaced by $\alpha\|x - y\|^2$, where $\alpha > 0$ is a fixed constant. In an obvious way, we define the concept of strong accretiveness on any subset of $D(T)$. An accretive operator T is called “ m -accretive” if $R(T + \lambda I) = X$ for every $\lambda > 0$, where I denotes the identity operator on X . It is called “ m -dissipative” if $-T$ is m -accretive. We denote by $B_r(0)$ the open ball of X (or X^*) with center at zero and radius $r > 0$. For an m -accretive operator T , the “resolvents” $J_\lambda : X \rightarrow D(T)$ of T are defined by $J_\lambda = (I + \lambda T)^{-1}$ for all $\lambda \in (0, \infty)$. J_λ is a non-expansive mapping on X for all $\lambda > 0$. Also the operator $T_\lambda \equiv (1/\lambda)(I - J_\lambda)$ is a globally Lipschitzian mapping with $T_\lambda x \in TJ_\lambda x$, for every $x \in X$. For facts involving accretive operators, and other related concepts, the reader is referred to Barbu [2], Browder [5], Cioranescu [7], Deimling [9], and Lakshmikantham and Leela [31]. For a survey paper on compactness and accretivity, we cite the paper [23].

For a bounded set $\Omega \subset X$, the Kuratowski measure of noncompactness, $\gamma(\Omega)$, is defined by

$$\gamma(\Omega) = \inf\{\epsilon > 0 : \Omega \text{ can be covered by a finite family of sets of diameter } < \epsilon\}.$$

A continuous mapping $g : X \supset D(g) \rightarrow X$ is called “condensing” if, for every non-empty, bounded and non-compact subset Ω of $D(g)$, $g(\Omega)$ is bounded and $\gamma(g(\Omega)) < \gamma(\Omega)$. Browder gave in [5, Theorem 13.23] the following result.

Theorem A. *Let X be a real Banach space and G a bounded, convex, open subset of X . Let $T : \overline{G} \rightarrow X$ and $C : \overline{G} \rightarrow X$ be two mappings with T accretive and C compact. Suppose that either T is locally uniformly continuous or T is continuous and X^* is uniformly convex. Then if $[I - (T + C)]\partial G \subset \overline{G}$ and $(T + C)(\overline{G})$ is closed, there exists $x \in \overline{G}$ such that $(T + C)x = 0$.*

One of our intentions in this paper is to give an analogous result, Theorem 1, for possibly multi-valued and discontinuous m -accretive operators $T : X \supset D(T) \rightarrow 2^X$. We then use this result to obtain a new fixed point theorem, Corollary 1, for a mapping of the type $T + C$, where T is now m -dissipative. It should be mentioned that Browder’s method cannot be applied in our result (even in the single-valued case) because a discontinuous, strongly accretive operator does not belong to a class of permissible homeomorphisms as in Proposition (13.14) of [5]. On the other hand, one could take T to be just a continuous accretive mapping in Theorem A. This is possible because Deimling’s result in [8] says that a continuous strongly accretive mapping maps open sets of its domain onto open sets. It thus belongs to a class of permissible homeomorphisms.

In the proof of Theorem 1 in Section 2, we discovered that we had the ingredients for a new surjectivity result involving a boundary condition of the type

$$\langle u + Cx, j \rangle \geq -\beta(\|x\|)\|x\|^2,$$

for all $x \in D(T)$ with sufficiently large norm, all $u \in Tx$ and some $j \in Jx$. The function $\beta(\rho)$ here tends to zero as $\rho \rightarrow \infty$. This boundary condition can be combined with a growth condition on the sum $T + C$ in order to produce results of the type $\overline{R(T + C)} = X$ and $R(T + C) = X$.

In Section 2 we also examine the existence of certain eigenvalues for the pair (T, C) . These eigenvalues equal $\lambda + 1$, where λ appears in the boundary conditions involving T , C and in the assumption of compactness of the operator $C(\lambda T + I)^{-1}$. Theorem 4 utilizes this eigenvalue situation in order to produce a new result involving the existence of zeros for the sum $T + C$.

Section 3 contains the extension of some results of Section 2 to the case of maximal monotone operators $T : X \supset D(T) \rightarrow 2^{X^*}$. Theorem 6 shows how to extend Theorem 2 above to this setting. Theorem 7 involves the extendability of the main result of the author in [28]. That result was given for m -accretive operators. Our method here involves the use of the Brézis-Crandall-Pazy approximants J_λ , T_λ introduced in [4]. There are some further differences between the proof of the result in [28] and that of Theorem 7. Theorem 9 solves a problem according to which an entire set lies in the range of the sum $T + C$. This interesting result extends a result in [29] and leads to Corollary 6, which is a new surjectivity result for completely continuous perturbations of maximal monotone operators.

Section 4 is devoted to the problem of the existence of positive eigenvalues for the pair (T, C) . Theorem 9 there improves the maximal monotone analogue of Theorem 2.5 in [16], while Theorem 11 improves Theorem 3.3 in [16]. Both results make use of the above approximants from [4] and further demonstrate the usefulness of such a methodology.

In Section 5, we have decided to introduce the mapping $\tilde{J}x \equiv \|x\|^p x$ in the study of nonlinear problems in real Hilbert spaces H . This mapping, whose finite-dimensional versions have been used by a host of authors, was introduced in [30] in order to study certain perturbations of control problems involving accretive operators. In order to demonstrate the effectiveness of this method, we give a result which extends Theorem 2 (Section 2) and Theorem 6 (Section 3). This result, Theorem 12, yields a criterion for the surjectivity of the sum $T + C$ which involves the number p itself. We thus actually have a one-parameter family of surjectivity criteria in Hilbert spaces.

Section 6 contains some discussion on the results of this paper as well as five examples of operators T or C which can be easily used in order to apply several results herein to partial differential equations.

Finally, we have given four open problems, the solution of which would lead to the improvement of related results in this paper.

The results of this paper, as well as a good number of papers in the references, have applications in the control theory with preassigned responses (cf. Kaplan and the author [20] and the author and Mabry [30]) and the construction of methods of lines (cf. [24]). For other recent results of this nature, we refer to the papers by Ding and the author [10-11], Guan [13-14], Guan and the author [15-17], Hirano and Kalinde [19] and the author [22-24].

2. PERTURBATIONS OF m -ACCRETIVE OPERATORS

The following lemma is an extension of a lemma found in [15]. That lemma is more general than a lemma used by the author in [22] and [24]. We need a more localized version of it. In [15], it was assumed that $C : D(T) \rightarrow X$.

Lemma 1. *Let X be uniformly convex. Let $T : X \supset D(T) \rightarrow 2^X$ be m -accretive*

and $C : \overline{D(T)} \supset \overline{G} \rightarrow X$, where G is a convex subset of X . For the inclusions

$$Tx_n + Cx_n + \alpha_n x_n \ni p_n, \quad n = 1, 2, \dots,$$

assume the following:

- (a) $\{x_n\} \subset D(T) \cap \overline{G}$, $x_n \rightarrow x_0 \in \overline{D(T)}$ and $Cx_n \rightarrow Cx_0$;
- (b) $\{\alpha_n\} \subset R_+$ and $\alpha_n \rightarrow 0$;
- (c) $\{p_n\} \subset X$ and $p_n \rightarrow p \in X$.

Then $x_0 \in D(T) \cap \overline{G}$ and $Tx_0 + Cx_0 \ni p$.

Proof. The proof follows from that of Lemma 1 in [15], where there is no set G . All that we need to do in order to state Lemma 1 in [15] is replace G above by $D(T)$. ■

We recall that since X is uniformly convex, $\overline{D(T)}$ is a convex set. This can be found in Barbu [2, Proposition 3.6] and Cioranescu [7, Theorem 1.15, p. 184]. Both authors assumed that X^* is also uniformly convex, but this is not needed in the proof. Actually Reich has shown in [36] that $\overline{D(T)}$ is a convex set even if X^* has only a Fréchet differentiable norm.

Lemma 2. Let $T : X \supset D(T) \rightarrow 2^X$ be m -accretive with $(T + I)^{-1}$ compact. Let $C : \overline{G} \rightarrow X$ be continuous and bounded, where G is a bounded subset of X . Let $\{\alpha_n\}$ be a sequence of positive numbers such that $\alpha_n \rightarrow 0$ as $n \rightarrow \infty$. Let $\{x_n\} \subset D(T) \cap \overline{G}$ and $\{p_n\} \subset X$ be such that $p_n \rightarrow p$ as $n \rightarrow \infty$ and

$$Tx_n + Cx_n + \alpha_n x_n \ni p_n, \quad n = 1, 2, \dots$$

Then $\{x_n\}$ contains a subsequence $\{x_{n_k}\}$ such that $x_{n_k} \rightarrow x_0 \in D(T) \cap \overline{G}$ and

$$Tx_0 + Cx_0 \ni p.$$

Proof. Let $\{x_n\}$, $\{a_n\}$, $\{p_n\}$ be as in the statement of the lemma. Then we have

$$Tx_n + Cx_n + \alpha_n x_n + x_n \ni p_n + x_n,$$

or

$$x_n = (I + T)^{-1} [-(Cx_n + (\alpha_n - 1)x_n - p_n)].$$

Since $(I + T)^{-1}$ is compact and $\{x_n\}$ and C are bounded, we have that $\{x_n\}$ lies in a compact set. As such, it has a convergent subsequence $\{x_{n_k}\}$. Let $x_{n_k} \rightarrow x_0 \in \overline{G}$. Then, for some sequence of terms $v_{n_k} \in Tx_{n_k}$, we have

$$v_{n_k} \rightarrow -Cx_0 + p \text{ as } k \rightarrow \infty.$$

Since T is m -accretive, it is closed. Thus, $x_0 \in D(T)$ and $Tx_0 + Cx_0 \ni p$. ■

Our first theorem is one of the main results in this paper.

Theorem 1. Let G be a bounded, open and convex subset of X . Let $T : X \supset D(T) \rightarrow 2^X$ be m -accretive and (i) $C : \overline{G} \rightarrow X$ compact. Assume that $D(T) \cap G \neq \emptyset$ and that $[I - (T + C)](D(T) \cap \partial G) \subset \overline{G}$. Then $0 \in \overline{R(T + C)}$. Assume that one of the following conditions holds:

- (a) X is uniformly convex and C is completely continuous;
- (b) instead of (i), let J_1 be compact and $C : \overline{G} \rightarrow X$ continuous and bounded.

Then $0 \in R(T + C)$.

Proof. We may assume that $0 \in D(T) \cap G$ and $0 \in T(0)$. In fact, if one of these is not true, we may consider instead the operators \tilde{T} , \tilde{C} defined on $D(\tilde{T})$ and \tilde{G} , respectively, where $D(\tilde{T}) \equiv D(T) - x_0$, $\tilde{G} \equiv G - x_0$, $\tilde{T}x \equiv T(x + x_0) - v_0$ and $\tilde{C}x \equiv C(x + x_0) + v_0$. Here, x_0 is a fixed point in $D(T) \cap G$ and v_0 is a fixed point in Tx_0 . It is easy to see that our assumptions are still satisfied with these arrangements.

We are going to show first that $0 \in R(T + C + \epsilon I)$, i.e., we are going to solve first the inclusion

$$(2) \quad Tx + Cx + \epsilon x \ni 0$$

in $D(T) \cap \overline{G}$, for each $\epsilon > 0$. We fix $\epsilon > 0$ and consider the homotopy inclusion

$$t(T + C)x + [1 - t(1 - \epsilon)]x \ni 0.$$

This inclusion gives rise to the homotopy mapping

$$H(t, x) \equiv [t(T + \epsilon I) + (1 - t)I]^{-1}(-tCx), \quad (t, x) \in [0, 1] \times \overline{G}.$$

We observe that $H(0, x) \equiv 0$. We may assume that $x - H(1, x) = 0$ has no solutions $x \in \partial G$. In fact, if this is not true, then we have a solution of (2) and the proof is finished. The mapping $H(t, \cdot)$ is continuous on \overline{G} and maps \overline{G} onto a relatively compact subset of X , for every $t \in [0, 1]$. The Leray-Schauder degree $d(I - H(t, \cdot), G, 0)$, $t \in [0, 1]$, will be well-defined if we show that (j) $H(\cdot, x)$ is continuous uniformly w.r.t. $x \in \overline{G}$, and (jj) the equation $x - H(t, x) = 0$ has no solutions $x_t \in \partial G$, for any $t \in [0, 1)$. In order to show (j), we set

$$\lambda(t) \equiv t/[\epsilon t + (1 - t)]$$

and we observe that $\lambda(t)$, $t \in [0, 1]$, is a well-defined continuous function. We also observe that

$$H(t, x) = (\lambda(t)T + I)^{-1}(-\lambda(t)Cx) = J_{\lambda(t)}(-\lambda(t)Cx).$$

Since $J_{\lambda(t)}(0) = 0$ and $J_{\lambda(t)}$ is a nonexpansive mapping,

$$\|J_{\lambda(t)}(-\lambda(t)Cx)\| \leq \|\lambda(t)Cx\| = \lambda(t)\|Cx\|.$$

Since $\lambda(0) = 0$, we have the continuity of $J_{\lambda(t)}(-\lambda(t)Cx)$ w.r.t. t at $t = 0$. Since C is bounded, this continuity is uniform w.r.t. $x \in \overline{G}$.

Now, we fix t , $t_0 \in (0, 1]$, $u \in X$ and let $y(t) \equiv J_{\lambda(t)}u$. Since

$$T_{\lambda(t)}u = (1/\lambda(t))(I - J_{\lambda(t)})u,$$

we have

$$y(t) = u - \lambda(t)T_{\lambda(t)}u.$$

Working as in the proof of Theorem 3 in [21], we obtain

$$\|y(t) - y(t_0)\| \leq \frac{2|\lambda(t) - \lambda(t_0)|}{\lambda(t_0)} \|u\|.$$

Consequently,

$$\begin{aligned} & \|J_{\lambda(t)}(-\lambda(t)Cx) - J_{\lambda(t_0)}(-\lambda(t_0)Cx)\| \\ & \leq \|J_{\lambda(t)}(-\lambda(t)Cx) - J_{\lambda(t_0)}(-\lambda(t)Cx)\| \\ & \quad + \|J_{\lambda(t_0)}(-\lambda(t)Cx) - J_{\lambda(t_0)}(-\lambda(t_0)Cx)\| \\ & \leq \frac{2|\lambda(t) - \lambda(t_0)|\lambda(t)}{\lambda(t_0)} \|Cx\| + |\lambda(t) - \lambda(t_0)| \|Cx\| \\ & = \frac{(2\lambda(t) + \lambda(t_0))|\lambda(t) - \lambda(t_0)|}{\lambda(t_0)} \|Cx\|, \end{aligned}$$

which shows the continuity of $H(t, x)$, w.r.t. t at t_0 , uniformly w.r.t. $x \in \overline{G}$. A simple covering argument shows now the desired uniform continuity of $H(t, x)$ w.r.t. $t \in [0, 1]$.

To show (jj), assume that $x_t = H(t, x_t)$, for some $t \in [0, 1)$ and some $x_t \in \partial G$. Since $t = 0$ is impossible, because $x_t \in \partial G$ implies $x_t \neq 0$, we have $t \in (0, 1)$. Also,

$$t(T + C + \epsilon I)x_t + (1 - t)x_t \ni 0.$$

Let $Sx \equiv x - (T + C)x$. Then $x_t \in [t/(1 + \epsilon t)]Sx_t$. By our assumption, $Sx_t \subset \overline{G}$. Since every element of $[t/(1 + \epsilon t)]Sx_t$ is a proper convex combination of $0 \in G$ and some element of Sx_t , we have $[t/(1 + \epsilon t)]Sx_t \subset G$, i.e., a contradiction to the fact that $x_t \in \partial G$. It follows that

$$d(I - H(t, \cdot), G, 0) = d(I - H(0, \cdot), G, 0) = 1$$

and, by the Leray-Schauder theory, the equation $x - H(1, x) = 0$ is solvable in G . This says that the inclusion (2) is solvable in $D(T) \cap G$. Letting x_ϵ denote a solution of (2), and then letting $\epsilon \rightarrow 0$ ($\{x_\epsilon\}$ is bounded), we have that $0 \in \overline{R(T + C)}$.

If we assume, in addition, that X is uniformly convex and C is completely continuous, then C is compact (by the reflexivity of X) and $0 \in R(T + C)$ by Lemma 1. In fact, for some positive sequence $\{\epsilon_n\}$, with $\epsilon_n \downarrow 0$, we may take, in Lemma 1, $x_n = x_{\epsilon_n}$, $\alpha_n = \epsilon_n$ and $p_n = 0$, $n = 1, 2, \dots$. Since $\{x_n\}$ is bounded and X is reflexive, we may assume that $x_n \rightharpoonup x_0 \in \overline{G}$.

On the other hand, in case (b) we observe that the resolvent identity

$$J_\mu x = J_\lambda \left(\frac{\lambda}{\mu} x + \frac{\mu - \lambda}{\mu} J_\mu x \right)$$

implies that J_μ is compact for every $\mu > 0$. Thus, the entire proof of the first conclusion of the theorem goes through with no change. We let $\epsilon_n \downarrow 0$ and apply Lemma 2. The proof is complete. ■

We are now ready for a new fixed point theorem involving completely continuous perturbations of possibly discontinuous m -dissipative operators.

Corollary 1. *Let G be an open, bounded and convex subset of X . Let $T : X \supset D(T) \rightarrow 2^X$ be m -dissipative and $C : \overline{G} \rightarrow X$. Assume that $D(T) \cap G \neq \emptyset$ and $(T + C)(D(T) \cap \partial G) \subset \overline{G}$. Then if (a) or (b) of Theorem 1 holds, the operator $T + C$ has a fixed point in $D(T) \cap \overline{G}$, i.e., there exists $x \in D(T) \cap \overline{G}$ such that $(T + C)x \ni x$.*

Proof. The operator $I - T$ is m -accretive. Also, the operator $I - (I - (T + C)) = T + C$ maps $D(T) \cap \partial G$ into \overline{G} . Theorem 1 (with $I - T$ in place of T and $-C$ in place of C) implies that the operator $I - (T + C)$ has a zero in $D(T) \cap \overline{G}$. This completes the proof. ■

Corollary 1 is an extension of Theorem (13.24) of Browder [5]. The homotopy used in Theorem 1 can be modified so that it can accommodate a new result on the surjectivity of $T + C$ based on an “inner product” condition. The next theorem reflects this situation and extends to the present case a host of surjectivity results in the references. We let Γ denote the family of all functions $\beta : R_+ \rightarrow R_+$ such that $\beta(\rho) \rightarrow 0$ as $\rho \rightarrow \infty$. Sometimes, the functions $\beta(\rho)$ considered in this paper are only defined for all sufficiently large $\rho > 0$. We make no comment on their trivial definition on any initial finite interval in R_+ . For a set $A \subset X$, we set $|A| \equiv \inf\{\|x\| ; x \in A\}$. We denote by $\text{co}G$ the convex hull of the set G .

Theorem 2. *Let $T : X \supset D(T) \rightarrow 2^X$ be m -accretive and (i) $C : D(T) \rightarrow X$ is compact. Assume that there exist $\beta \in \Gamma$ and $Q > 0$ such that: for every $x \in D(T)$ with $\|x\| \geq Q$ and every $v \in Tx$ there exists $j \in Jx$ such that*

$$(3) \quad \langle v + Cx, j \rangle \geq -\beta(\|x\|)\|x\|^2.$$

Then $R(T + C + \epsilon I) = X$, for every $\epsilon \in (0, 1)$. Assume, further, that

$$(*) \quad \liminf_{\substack{\|x\| \rightarrow \infty \\ x \in D(T)}} \frac{|Tx + Cx|}{\|x\|} > 0.$$

Then $\overline{R(T + C)} = X$. Let one of the following conditions hold:

- (a) X is uniformly convex and $C : \overline{D(T)} \rightarrow X$ is completely continuous;
- (b) instead of (i), $C : \overline{D(T)} \rightarrow X$ is continuous and bounded and $(T + I)^{-1}$ is compact.

Then $R(T + C) = X$.

Proof. We are planning to solve first the problem

$$(4) \quad Tx + Cx + \epsilon x \ni s,$$

where s is any (but fixed) point in X . We may assume that, for some $x_0 \in D(T)$, we have $0 \in T(x_0)$. If this is not true, we consider instead the operators $\tilde{T}(x) \equiv Tx - v_0$, $\tilde{C}(x) \equiv Cx + v_0$, for every $x \in D(T)$, where v_0 is any (but fixed) point in Tx_0 . We assume that (i) holds in order to prove the first conclusion of the theorem. In order to solve (4), we consider the homotopy inclusion

$$t(Tx + Cx + \epsilon x) + (1 - t)x \ni ts,$$

or the equation

$$(5) \quad \begin{aligned} u = H(t, u) &\equiv -t [C(t(T + \epsilon I) + (1 - t)I)^{-1}u - s] \\ &= -t [C((t/\lambda(t))T + I)^{-1}([1/\lambda(t)]u) - s], \end{aligned}$$

where we have set $H(0, x) \equiv 0$ and $\lambda(t) \equiv [1 - t(1 - \epsilon)]$. It is easy to see that $\epsilon \leq \lambda(t) \leq 1$, $t \in [0, 1]$. For every solution $u \in X$ of $u - H(1, u) = 0$, we obtain a solution x of (4), where $x = (T + \epsilon I)^{-1}u$. In order to apply the Leray-Schauder degree theory, we shall show first that the mapping $H : [0, 1] \times \overline{B_r(0)} \rightarrow X$ is continuous and such that $H([0, 1], \overline{B_r(0)})$ is relatively compact in X , for every $r > 0$. To this end, we fix $r > 0$ and show that the function

$$y(t, u) \equiv ((t/\lambda(t))T + I)^{-1}([1/\lambda(t)]u)$$

is continuous in $(t, u) \in (0, 1] \times \overline{B_r(0)}$. In fact, setting $\mu(t) \equiv t/\lambda(t)$, $q(t) \equiv 1/\lambda(t)$ and taking into consideration the proof of Theorem 1, we obtain

$$\begin{aligned} \|y(t, u) - y(t_0, v)\| &= \|J_{\mu(t)}q(t)u - J_{\mu(t_0)}q(t_0)v\| \\ &\leq \|J_{\mu(t)}q(t)u - J_{\mu(t_0)}q(t)u\| \\ &\quad + \|J_{\mu(t_0)}q(t)u - J_{\mu(t_0)}q(t_0)v\| \\ &\leq \frac{2|\mu(t) - \mu(t_0)|}{\mu(t_0)} \|q(t)u\| \\ &\quad + \|q(t)u - q(t_0)v\|. \end{aligned}$$

The continuity of $y(t, u)$ on $(0, 1] \times \overline{B_r(0)}$ follows from this inequality because the functions $\mu(t)$, $\lambda(t)$ and $q(t)$ are well-defined, continuous and bounded on the interval $(0, 1]$. Since C is continuous, the continuity of $H(t, u) = -t[Cy(t, u) - s]$ on $(0, 1] \times \overline{B_r(0)}$ has been shown. In order to show the continuity of $H(t, u)$ on the set $\{0\} \times \overline{B_r(0)}$, we observe first that

$$\begin{aligned} \|y(t, u)\| = \|J_{\mu(t)}q(t)u\| &\leq \|J_{\mu(t)}q(t)u - J_{\mu(t)}x_0\| + \|J_{\mu(t)}x_0\| \\ &\leq \|q(t)u - x_0\| + \|x_0\| \\ &\leq r \sup_{t \in [0, 1]} q(t) + 2\|x_0\|. \end{aligned}$$

This implies that the set

$$K = \{y(t, u) : (t, u) \in [0, 1] \times \overline{B_r(0)}\} \subset D(T)$$

is bounded. Hence, from the boundedness of C we obtain that

$$\|H(t, u)\| \leq t \|CJ_{\mu(t)}q(t)u - s\| \rightarrow 0 \text{ as } t \rightarrow 0,$$

uniformly w.r.t. $u \in \overline{B_r(0)}$. We have demonstrated the continuity of the homotopy function $H(t, u)$ on the set $[0, 1] \times \overline{B_r(0)}$.

To show that the set $H([0, 1] \times \overline{B_r(0)})$ is relatively compact, let (t_n, u_n) be a sequence of points in $[0, 1] \times \overline{B_r(0)}$. We are going to show that the sequence of points

$$z(t_n, u_n) \equiv -t_n[Cy(t_n, u_n) - s]$$

contains a convergent subsequence. To this end, we observe first that since $[0, 1]$ is compact, we may assume that $t_n \rightarrow t_0$, for some $t_0 \in [0, 1]$. If $t_0 = 0$, then the boundedness of the sequence $\{Cy(t_n, u_n) - s\}$ implies that $z(t_n, u_n) \rightarrow 0$. Thus, we may assume that $t_0 > 0$ and $t_n > 0$ for all n . We are now going to show the existence of a convergent subsequence of the sequence $\{y(t_n, u_n)\}$. If C is compact, then the continuity of $C : D(T) \rightarrow X$ implies immediately the existence of such a convergent subsequence because, by the continuity of multiplication and translation, the set $-[0, 1](\overline{CK} - s)$ is compact and $z(t_n, u_n)$ lies in this set. If $(T + I)^{-1}$ is compact, then the resolvent identity says that $(\mu(t_0)T + I)^{-1}$ is also compact. From the definition of $y(t_n, u_n)$ we have

$$\mu(t_n)Ty(t_n, u_n) + y(t_n, u_n) \ni q(t_n)u_n,$$

or

$$\mu(t_n)v_n + y(t_n, u_n) = q(t_n)u_n,$$

where $v_n \in Ty(t_n, u_n)$. This implies

$$\mu(t_0)v_n + y(t_n, u_n) = q(t_n)u_n - [\mu(t_n) - \mu(t_0)]v_n,$$

or

$$\begin{aligned} y(t_n, u_n) &= (\mu(t_0)T + I)^{-1} [q(t_n)u_n - [\mu(t_n) - \mu(t_0)]v_n] \\ &= (\mu(t_0)T + I)^{-1} \left[q(t_n)u_n - \frac{[\mu(t_n) - \mu(t_0)][q(t_n)u_n - y(t_n, u_n)]}{\mu(t_n)} \right]. \end{aligned}$$

The compactness of the resolvent $J_{\mu(t_0)}$ and the boundedness of the sequence of points to the right of it above ($\mu(t_n) \rightarrow t_0/\lambda(t_0) > 0$) show that the sequence $\{y(t_n, u_n)\}$ lies in a compact set. It thus has a convergent subsequence. Since C is continuous, we obtain a convergent subsequence of the sequence $\{z(t_n, u_n)\}$.

In order to show that $d(I - H(t, \cdot), B, 0) = \text{const.}$, for some open ball $B \subset X$, we are going to show first that all possible solutions u of the equation $u - H(t, u) = 0$ are bounded by a constant which is independent of the parameter $t \in [0, 1]$. To this end, let us assume that $\{u_m\}$ is a sequence of points in X such that $u_m - H(t_m, u_m) = 0$, for some sequence $\{t_m\} \subset [0, 1]$, and $\|u_m\| \rightarrow \infty$ as $m \rightarrow \infty$. Let

$$x_m = (t_m(T + \epsilon I) + (1 - t_m)I)^{-1}u_m.$$

Then we have $\{x_m\} \subset D(T)$ and

$$t_m(T + \epsilon I)x_m + t_m(Cx_m - s) + (1 - t_m)x_m \ni 0,$$

or

$$(6) \quad t_m v_m + t_m(Cx_m - s) + [1 - t_m(1 - \epsilon)]x_m = 0,$$

for some $v_m \in Tx_m$. We must have $\|x_m\| \rightarrow \infty$ as $m \rightarrow \infty$. In fact, if this is not true, then there exists a subsequence of $\{x_m\}$, denoted again for convenience by $\{x_m\}$, which is bounded. Then the boundedness of the operator C and (6) imply that the sequence $\{t_m v_m\}$ is bounded. Since

$$u_m = t_m v_m + [1 - t_m(1 - \epsilon)]x_m,$$

we have that $\{u_m\}$ is also bounded, i.e., a contradiction.

Using our hypotheses, there exist $\beta \in \Gamma$ and $Q > 0$ such that: for every $m \geq 1$ such that $\|x_m\| \geq Q$ there exists $j \in Jx_m$ such that

$$\begin{aligned} \langle v_m + Cx_m - s, j \rangle &\geq -\langle s, j \rangle - \beta(\|x_m\|)\|x_m\|^2 \\ &\geq -\frac{\|s\|}{\|x_m\|}\|x_m\|^2 - \beta(\|x_m\|)\|x_m\|^2 \\ &= -\tilde{\beta}(\|x_m\|)\|x_m\|^2, \end{aligned}$$

for every sequence $\{v_m\}$ of terms $v_m \in Tx_m$, where $\tilde{\beta}$ is an obvious function in Γ . For such an integer m , we evaluate the above functional $j \in Jx_m$ on both sides of (6) in order to obtain

$$\begin{aligned} \epsilon\|x_m\|^2 &\leq [1 - t_m(1 - \epsilon)]\|x_m\|^2 \\ &\leq -t_m\langle v_m + Cx_m - s, j \rangle \\ &\leq t_m\tilde{\beta}(\|x_m\|)\|x_m\|^2. \end{aligned}$$

We see now that this last inequality holds for all large integers m . It implies

$$\epsilon \leq \tilde{\beta}(\|x_m\|).$$

Since the right-hand side above converges to zero as $m \rightarrow \infty$, we have a contradiction. We have shown that all possible solutions of $x - H(t, x) = 0$ lie in an open ball, say, $B_M(0)$, whose radius $M > 0$ is independent of the parameter t . It follows that $x - H(t, x) \neq 0$, for any $(t, x) \in [0, 1] \times \partial B_M(0)$. Thus, the Leray-Schauder degree $d(I - H(t, \cdot), B_M(0), 0)$ is well-defined and constant for $t \in [0, 1]$. In particular,

$$\begin{aligned} d(I - H(t, \cdot), B_M(0), 0) &= d(I - H(0, \cdot), B_M(0), 0) \\ &= d(I, B_M(0), 0) \\ &= 1 \end{aligned}$$

because $0 \in B_M(0)$. We conclude that the equation $x - H(1, x) = 0$ has a solution $x \in B_M(0)$. This solution satisfies the inclusion (5). It follows that $s \in R(T + C + \epsilon I)$, for every $\epsilon \in (0, 1)$.

For the second part of the proof, we let x_n be a solution to the problem (4) for $\epsilon = 1/n$. We claim that $\{x_n\}$ is a bounded sequence. To see this, assume, without loss of generality, that $\|x_n\| \rightarrow \infty$ as $n \rightarrow \infty$. Then, by our hypothesis, there exists $\alpha > 0$ so that

$$(7) \quad \liminf_{n \rightarrow \infty} \frac{|Tx_n + Cx_n|}{\|x_n\|} \geq \liminf_{\substack{\|x\| \rightarrow \infty \\ x \in D(T)}} \frac{|Tx + Cx|}{\|x\|} = \alpha.$$

However, we know that for some $v_n \in Tx_n$ we have

$$\|v_n + Cx_n\| = \|s - (1/n)x_n\| \leq (1/n)\|x_n\| + \|s\|,$$

which implies

$$\alpha \leq \liminf_{n \rightarrow \infty} \frac{\|v_n + Cx_n\|}{\|x_n\|} \leq \liminf_{n \rightarrow \infty} \left[(1/n) + \frac{\|s\|}{\|x_n\|} \right] = 0,$$

i.e., a contradiction. Since $(1/n)x_n \rightarrow 0$, we have that $s \in \overline{R(T + C)}$.

If (a) is true, then since $\{x_n\}$ is a bounded sequence, we may assume that $x_n \rightarrow x_0 \in \overline{D(T)} \cap \text{co}\{x_n\}$, and then apply Lemma 1 with $G = \text{co}\{x_n\}$.

If (b) is true, then we apply Lemma 2 with G any ball $B_r(0)$ of X containing $\{x_n\}$. ■

We note that (3) is an extremely weak boundary condition and that condition (*) in the statement of Theorem 2 does not generally suffice for $\overline{R(T + C)} = X$ even if $T : X \rightarrow X$ is continuous. In fact, let $Tx \equiv x^m$, for some odd positive integer m , $Cx \equiv |x|^m + |x| + 3$ and $\alpha \in (0, 1)$. Then $T : \mathcal{R} \rightarrow \mathcal{R}$ is m -accretive, $C : \mathcal{R} \rightarrow \mathcal{R}$ is compact and

$$\liminf_{|x| \rightarrow \infty} \frac{|Tx + Cx|}{|x|} \geq 1,$$

but $0 \notin \overline{R(T + C)} = [3, \infty)$. For more uses of boundary conditions like (but different from) (3), the reader is referred to the paper [17].

On the other hand, condition (3) is not generally sufficient for $\overline{R(T + C)} = X$ either. In fact, if $Tx \equiv x^{1/3}$ and $Cx \equiv |x|^{1/3} + 3$, then we have

$$\begin{aligned} \langle Tx + Cx, x \rangle &= (Tx + Cx)x \geq -|x|^{4/3} - 3|x| \\ &= -(1/|x|^{2/3})|x|^2 - (3/|x|)|x|^2 = -\beta_0(|x|)|x|^2, \end{aligned}$$

for all $x \neq 0$, where $\beta_0 \in \Gamma$. However, $0 \notin \overline{R(T + C)} = [3, \infty)$.

It is easy to see that (3) is implied by any other condition which has the same left-hand side while its right-hand side is of the form

$$-K - L\|x\| - \beta(\|x\|)\|x\|^2,$$

for some positive constants K, L . In fact,

$$\begin{aligned} -K - L\|x\| - \beta(\|x\|)\|x\|^2 &= -\frac{K}{\|x\|^2}\|x\|^2 - \frac{L}{\|x\|}\|x\|^2 - \beta(\|x\|)\|x\|^2 \\ &= -\tilde{\beta}(\|x\|)\|x\|^2, \end{aligned}$$

for all $x \in D(T)$ with $\|x\| \geq Q$, where $\tilde{\beta}$ is an obvious function in Γ .

The next result provides a method for proving the existence of certain eigenvalues for the pair (T, C) . It is assumed that CJ_λ is at least condensing, for some $\lambda > 0$, and shown that $\lambda + 1$ is an eigenvalue for (T, C) . A real number λ is called an "eigenvalue" of a pair of operators (T, C) if the equation $\lambda Tx + Cx \ni 0$ is solvable in $D(T) \cap D(C)$.

Theorem 3. Let G be a bounded, open and convex subset of X and let $T : X \supset D(T) \rightarrow 2^X$ be m -accretive with $0 \in D(T) \cap G$ and $0 \in T(0)$. Assume that $C : D(T) \rightarrow X$ is bounded and such that (i) $C(\lambda T + I)^{-1}$ is compact for some $\lambda > 0$. Let $[I - (T + C)](D(T) \cap \partial G) \subset \overline{G}$ and $[(\lambda T + I)(D(T) \cap \partial G)] \cap \text{int}G = \emptyset$. Then $0 \in R((\lambda + 1)T + C)$. Assume that one of the following conditions holds:

(a) X is uniformly convex and $C(\lambda T + I)^{-1}$ is completely continuous;

(b) instead of (i), J_1 is compact and $C(\lambda T + I)^{-1}$ is just condensing.

Then the number $\lambda + 1$ is an eigenvalue for the pair (T, C) .

Proof. We consider the homotopy equation

$$(8) \quad t(T_\lambda + CJ_\lambda)x + (1 + \epsilon)x - tJ_\lambda x = 0.$$

Here, $t \in [0, 1]$, $\epsilon > 0$ and $x \in \overline{U}$, where $U \equiv (\lambda T + I)(D(T) \cap G)$. We are planning to solve first the equation

$$(9) \quad T_\lambda x + CJ_\lambda x + (1 + \epsilon)x - J_\lambda x = 0,$$

which can also be written as

$$(10) \quad (\lambda + 1)T_\lambda x + CJ_\lambda x + \epsilon x = 0$$

because $T_\lambda = (1/\lambda)(I - J_\lambda)$. We write (8) as

$$(11) \quad x = \frac{t(\lambda + 1)}{t + \lambda(1 + \epsilon)}J_\lambda x - \frac{t\lambda}{t + \lambda(1 + \epsilon)}CJ_\lambda x.$$

Since $J_\lambda \equiv (\lambda T + I)^{-1}$ is defined and continuous on the entire space X , the (possibly multi-valued) mapping $V : x \rightarrow (\lambda T + I)x$ maps relatively open (closed) sets of its domain $D(T)$ onto open (closed) subsets of X . Thus, $V(D(T) \cap G)$ is open and $V(D(T) \cap \overline{G})$ is closed in X . This implies, as in [28, Theorem 2], that

$$\overline{(\lambda T + I)(D(T) \cap \overline{G})} \subset (\lambda T + I)(D(T) \cap \overline{G})$$

and

$$(12) \quad \partial[(\lambda T + I)(D(T) \cap G)] \subset (\lambda T + I)(D(T) \cap \partial G),$$

which is a crucial point of our proof. Since the function $t \rightarrow t/(t + a)$, $t \in [0, 1]$, is increasing for every $a > 0$, it is evident that the first term on the right-hand side of (11) is a strict contraction for all $t \in [0, 1]$. In addition, the second term is a compact mapping for all $t \in [0, 1]$. Thus, the right-hand side of (11), denoted by $H(t, x)$, is a condensing mapping w.r.t. $x \in \overline{U}$ for every $t \in [0, 1]$. It is also jointly continuous. As such, it is a homotopy of condensing mappings such that the image of the set \overline{U} by $H(t, \cdot)$ is bounded. Thus, the degree $d(I - H(t, \cdot), U, 0)$ will be well-defined, constant and equal to 1, if the equation $x - H(t, x) = 0$ has no solution x_t on the boundary of the open set U , for any $t \in [0, 1]$. The proof of the existence of such a degree is sketched in Theorem 1 of [28]. Let $x_t \in \partial U$ be a solution of this equation for some $t \in [0, 1]$. Then

$$t(T_\lambda + CJ_\lambda)x_t + (1 + \epsilon)x_t - tJ_\lambda x_t = 0.$$

This implies

$$(13) \quad x_t = \frac{t}{1 + \epsilon} [J_\lambda - (T_\lambda + CJ_\lambda)]x_t.$$

From what we have seen above, since $x_t \in \partial U$, we must have $J_\lambda x_t \in D(T) \cap \partial G$. By our assumption, this implies

$$(14) \quad [J_\lambda - (T_\lambda + CJ_\lambda)]x_t \subset [J_\lambda - (TJ_\lambda + CJ_\lambda)]x_t \subset \overline{G}.$$

Again, since G is open and convex and $0 \in G$, inclusions (13) and (14) imply $x_t \in \text{int}G$. Since we also have

$$x_t \in \partial[(\lambda T + I)(D(T) \cap G)] \subset (\lambda T + I)(D(T) \cap \partial G),$$

we get a contradiction to

$$[(\lambda T + I)(D(T) \cap \partial G)] \cap \text{int}G = \emptyset.$$

It follows that equation (10) has a solution $x_\epsilon \in (\lambda T + I)(D(T) \cap G)$, for every $\epsilon > 0$.

We now observe that (9) implies that the family $\{x_\epsilon\}$ of solutions of (10) obtained as above is bounded. In fact, each x_ϵ satisfies the equation (11) for $t = 1$. Since x_ϵ lies in the set $\overline{(\lambda T + I)(D(T) \cap G)} \subset (\lambda T + I)(D(T) \cap \overline{G})$, we have that $J_\lambda x_\epsilon$ lies in the set $D(T) \cap \overline{G}$. Since this set is bounded and C is also bounded, equation (11) (for $t = 1$) implies that $\{x_\epsilon\}$ lies in a bounded set. Thus, $\epsilon x_\epsilon \rightarrow 0$ and $0 \in \overline{R((\lambda + 1)T + C)}$.

For the second part of the theorem, we let $\epsilon = 1/n$ and $x_n \equiv x_{1/n}$, and observe that since X is reflexive, there exists a subsequence of $\{x_n\}$, denoted again by $\{x_n\}$, such that $x_n \rightarrow x \in X$. Since CJ_λ is completely continuous, we have that $CJ_\lambda x_n \rightarrow CJ_\lambda x$. Thus, we may apply Lemma 1 with T_λ in place of T , CJ_λ in place of C and X in place of $D(T)$ or G .

In the third part of the proof, we observe first that since the operator J_λ is compact, we do not need the term ϵx in the equation (9). Because of this, the homotopy equation (11) must now be replaced by the equation

$$x = \frac{t(\lambda + 1)}{t + \lambda} J_\lambda x - \frac{t\lambda}{t + \lambda} CJ_\lambda x.$$

We observe that the first term of this equation is compact w.r.t. x and the second condensing, for every $t \in [0, 1]$. We may repeat now the previous argument in order to solve the equation (9) with zero in place of ϵ . This immediately implies $((\lambda + 1)TJ_\lambda + CJ_\lambda)x \ni 0$ and finishes the proof. ■

A special, and important, case of Theorem 2 occurs when $G = B_r(0)$, for some $r > 0$. This is the content of Corollary 2.

Corollary 2. *Let $T : X \supset D(T) \rightarrow 2^X$ be m -accretive with $0 \in D(T)$ and $0 \in T(0)$. Assume that $C : D(T) \rightarrow X$ is bounded and such that (i) $C(\lambda T + I)^{-1}$ is compact, for some $\lambda > 0$. Let $[I - (T + C)](D(T) \cap \partial B_r(0)) \subset \overline{B_r(0)}$. Then $0 \in \overline{R((\lambda + 1)T + C)}$. Assume that one of the following conditions holds:*

- (a) X is uniformly convex and $C(\lambda T + I)^{-1}$ is completely continuous;
- (b) instead of (i), J_1 is compact and $C(\lambda T + I)^{-1}$ is just condensing.

Then the number $\lambda + 1$ is an eigenvalue for the pair (T, C) .

Proof. In view of Theorem 3, it suffices to show that

$$(15) \quad [(\lambda T + I)(D(T) \cap \partial B_r(0))] \cap B_r(0) = \emptyset.$$

In fact, if $y \in (\lambda T + I)(D(T) \cap \partial B_r(0))$, then $y = \lambda v + x$, where $v \in Tx$, for some $x \in D(T) \cap \partial B_r(0)$. Since $0 \in T(0)$ and T is accretive, we have, for some $j \in Jx$,

$$\|\lambda v + x\|r = \|\lambda v + x\|\|x\| \geq \langle \lambda v + x, j \rangle = \lambda \langle v, j \rangle + \langle x, j \rangle \geq \|x\|^2 = r^2.$$

This implies that $\|\lambda v + x\| \geq r$ and proves that $y \notin B_r(0)$. ■

Corollary 2 implies the existence of a zero of the operator $T + C$ if we assume that $(T + I)^{-1}$ is compact and $C : \overline{D(T)} \rightarrow X$ is continuous and bounded. In fact, these two assumptions imply that $(\lambda T + I)^{-1}$ and $C(\lambda T + I)^{-1}$ are compact due to the resolvent identity.

Corollary 2'. Let $T : X \supset D(T) \rightarrow 2^X$ be m -accretive with $0 \in D(T)$ and $0 \in T(0)$. Assume that $C : \overline{D(T)} \rightarrow X$ is continuous and bounded. Let $(T + I)^{-1}$ be compact and $[I - (T + C)](D(T) \cap \partial B_r(0)) \subset \overline{B_r(0)}$, for some $r > 0$. Then, for every $\lambda \in \mathcal{R}_+$, $0 \in R((\lambda + 1)T + C)$.

Proof. We know from Corollary 2 that $0 \in R((\lambda + 1)T + C)$, for all $\lambda > 0$. To show this for $\lambda = 0$, we note first that the solvability of $((\lambda + 1)T + C)x \ni 0$, in the proof of Corollary 2, follows from the solvability of $((\lambda + 1)T_\lambda + CJ_\lambda)x \ni 0$, where $x \in (\lambda T + I)(D(T) \cap B_r(0))$. Let $\{\lambda_n\} \subset (0, \infty)$ be such that $\lambda_n \downarrow 0$ and let $x_n \in (\lambda T + I)(D(T) \cap B_r(0))$ be such that

$$T_{\lambda_n}x_n + [1/(\lambda_n + 1)]CJ_{\lambda_n}x_n = 0.$$

Letting $u_n \equiv J_{\lambda_n}x_n \in D(T) \cap B_r(0)$, we have that $\{u_n\}$ is bounded and

$$u_n = (T + I)^{-1}(-[1/(\lambda_n + 1)]Cu_n + u_n).$$

Consequently, due to the boundedness of C and the compactness of $(T + I)^{-1}$, we have that $u_n \rightarrow u_0 \in \overline{D(T) \cap B_r(0)}$. Since, for some $v_n \in Tu_n$, $v_n \rightarrow -Cu_0$, the closedness of T implies that $u_0 \in D(T)$ and $Tu_0 + Cu_0 \ni 0$. ■

It would be very interesting to know anything about the relationship between the boundary condition

$$[I - (T + C)](D(T) \cap \partial B_r(0)) \subset \overline{B_r(0)}$$

and the following “inner product” condition: for every $x \in D(T) \cap \partial B_r(0)$ there exists $j \in Jx$ such that

$$\langle u + Cx, j \rangle \geq 0,$$

for every $u \in Tx$. This last boundary condition and other extensions of it provide the standard criteria for the existence of zeros of the operator $T + C$.

The following variant of Theorem 2 provides conditions for the existence of zeros of $T + C$. It improves Theorem 5 in [29].

Theorem 4. *Let G be a bounded, open and convex subset of X and let $T : X \supset D(T) \rightarrow 2^X$ be m -accretive with $0 \in D(T) \cap G$ and $0 \in T(0)$. Assume that $C : D(T) \rightarrow X$ is bounded and such that (i) $C(\lambda T + I)^{-1}$ is compact for some $\lambda \in (0, \infty)$. Let $\overline{(I - \lambda C)(D(T) \cap \partial G)} \subset \overline{G}$ and $[(\lambda T + I)(D(T) \cap \partial G)] \cap \text{int}G = \emptyset$. Then $0 \in R(T + C)$. Assume that one of the following conditions holds:*

- (a) X is uniformly convex and $C(\lambda T + I)^{-1}$ is completely continuous;
 - (b) instead of (i), J_1 is compact, $C(\lambda T + I)^{-1}$ is just condensing and $\lambda \in (0, 1]$.
- Then $0 \in R(T + C)$.

Proof. We proceed as in Theorem 3, but we are now using the approximate problem

$$(16) \quad (T_\lambda + CJ_\lambda)x + \epsilon x = 0,$$

which can be also written as

$$(17) \quad (1 + \epsilon\lambda)x - J_\lambda x + \lambda CJ_\lambda x = 0.$$

In order to solve (16), we consider the homotopy analogue of (11),

$$(18) \quad x = H(t, x) \equiv \frac{t}{1 + \epsilon\lambda} J_\lambda x - \frac{\lambda t}{1 + \epsilon\lambda} CJ_\lambda x,$$

for $t \in [0, 1]$ and $x \in \overline{(\lambda T + I)G}$. We note that the first term on the right-hand side of (18) is a strict contraction and the second is compact. Thus, $H(t, x)$ is condensing w.r.t. x . It is also jointly continuous. We can thus apply the relevant degree theory to the mapping $I - H(t, \cdot)$, as in the proof of Theorem 3. To this end, we must show that the equation $x - H(t, x) = 0$ has no solution $x \in \partial((\lambda T + I)(D(T) \cap G)) \subset (\lambda T + I)(D(T) \cap \partial G)$, for any $t \in [0, 1]$. In fact, if x_t were such a solution, then we would have

$$x_t = \frac{t}{1 + \epsilon\lambda} (J_\lambda x_t - \lambda CJ_\lambda x_t).$$

We can now argue as in the case of equation (8) in order to obtain a contradiction. Having solved (16) for each $\epsilon > 0$, we have that $0 \in \overline{R(T + C)}$.

The proof under assumption (a) follows as in Theorem 3. It is therefore omitted.

To prove the conclusion under the assumption (b), it suffices to observe that since J_λ is compact, we do not need the term ϵx in the equation (16). In fact, without this term (17) becomes

$$(19) \quad x = J_\lambda x - \lambda CJ_\lambda x,$$

and the homotopy equation (18) is now replaced by

$$(20) \quad x = H(t, x) \equiv tJ_\lambda x - \lambda tCJ_\lambda x.$$

Since, as in the proof of Theorem 1, $J_\lambda x$ is compact and $\lambda \in (0, 1]$, the mapping $H(t, x)$ is condensing w.r.t. x for every $t \in [0, 1]$. If (18) has a solution $x \in (\lambda T + I)(D(T) \cap \partial G)$ for $t = 1$, then we are done. Since it is impossible to have such a solution for $t = 0$, we may assume that $t \in (0, 1)$. The solvability of $(T_\lambda + CJ_\lambda)x = 0$ (hence $(T + C)x \ni 0$) now follows as above. ■

The following corollary sheds more light into the applicability of Theorem 3.

Corollary 3. *Let $T : X \supset D(T) \rightarrow 2^X$ be m -accretive with $0 \in D(T)$ and $0 \in T(0)$. Assume that $C : D(T) \rightarrow X$ is bounded and such that (i) $C(T + I)^{-1}$ is compact. Let*

$$\|x - Cx\| \leq r, \text{ for every } x \in D(T) \cap \partial B_r(0).$$

Then $0 \in \overline{R(T + C)}$. Assume, further, that one of the following conditions holds:

- (a) *X is uniformly convex and $C(T + I)^{-1}$ is completely continuous;*
- (b) *instead of (i), J_1 is compact and $C(T + I)^{-1}$ is just condensing.*

Then $0 \in R(T + C)$.

Proof. We just let $\lambda = 1$ in Theorem 4 and observe that (15) holds as in the proof of Corollary 2. ■

As we mentioned in the introduction, Theorem A is actually true if the operator T is only assumed to be continuous on \overline{G} . Actually, this result is true if we assume that X^* is uniformly convex and T is demicontinuous on \overline{G} , provided that an additional assumption is made to the effect that (i) $(I + T)(\partial G) \cap \text{int}G = \emptyset$. Such an assumption is needed, as in Theorems 3 and 4 for $\lambda = 1$, because the operators J_1, CJ_1 are now defined on the set $\overline{(T + I)G}$. Due to this additional assumption, we must also assume that (ii) $0 \in G$ and $T(0) = 0$. In fact, without (i), we can always assume (ii) without any loss of generality. This was shown in the proof of Theorem 1. However, unlike the continuous case, this setting cannot be handled by Browder's degree theory because, again, strongly accretive demicontinuous mappings do not necessarily belong to classes of permissible homeomorphisms. Theorem 5 below covers this case and is based upon the invariance of domain result of the author in [21]. It is easily seen that the method of proof of Theorem 1 cannot be applied here because the operator $t(T + \epsilon I) + (1 - t)I$ is no longer an onto mapping and the mapping

$$x \rightarrow [t(T + \epsilon I) + (1 - t)I]^{-1} (-tCx)$$

cannot necessarily be defined. The mapping

$$u \rightarrow C [t(T + \epsilon I) + (1 - t)I]^{-1} u$$

cannot necessarily be handled either, because the relevant Leray-Schauder degree function would have the time-dependent domain $\overline{(t(T + \epsilon I) + (1 - t)I)G}$. Such time-dependent domains were considered by Browder [5], where T was at least a continuous mapping. To the best of our knowledge, no Leray-Schauder degree theory has ever been applied to time-dependent domains, as above, arising from strongly discontinuous, or even demicontinuous, operators T . For a degree theory involving demicontinuous operators T mapping a subset of X into X^* and satisfying condition (S_+) (see Section 3), we cite the book of Skrypnik [38].

Theorem 5. *Assume that X^* is uniformly convex. Assume that $T : \overline{G} \rightarrow X$ is demicontinuous, accretive and $C : \overline{G} \rightarrow X$ compact, where G is an open, convex and bounded subset of X with $0 \in G$ and $T(0) = 0$. Let $(T + C)\overline{G}$ be closed, $[(I + T)(\partial G)] \cap \text{int}G = \emptyset$ and $[I - (T + C)](\partial G) \subset \overline{G}$. Then $(T + C)\overline{G} \ni 0$.*

Proof. The proof follows as in Theorem 4. In fact, the proof is based upon a degree theoretic argument involving equation (18) with $\lambda = 1$. The right-hand side of (18) again defines a condensing mapping on the set $\overline{(T + I)G}$, which is the closure of

the open set $(T + I)G$. The set $(T + I)G$ is open by the invariance of domain result of the author in [21]. The solvability of $(T + C)x \ni 0$ follows from the solvability of $(T + C)x + \epsilon x \ni 0$ because the latter says that $0 \in \overline{R(T + C)}$, but we have assumed that $(T + C)\overline{G}$ is a closed set. ■

Condition (a) in Theorem 4 can be used in Theorem 5 to ensure that $R(T + C)$ is a closed set. We cannot use condition (b) of Theorem 4 because the compactness of the resolvent J_1 implies that the set $[J_1(T + I)]G_0 = G_0$ is relatively compact, i.e., the space X is finite-dimensional. The set G_0 here is a sufficiently small open set such that $(T + I)G_0$ is bounded. It is a well-known fact that a demicontinuous accretive mapping is locally bounded whenever the space X^* is assumed to be uniformly convex (cf. [23, Section 3]).

Open Problem 1. *Is Theorem 5 true without the assumption that $[(I + T)(\partial G)] \cap \text{int}G = \emptyset$?* ■

3. PERTURBATIONS OF MAXIMAL MONOTONE OPERATORS

In this section we study perturbation problems

$$(21) \quad Tx + Cx \ni 0,$$

where $T : X \supset D(T) \rightarrow 2^{X^*}$ is now a maximal monotone operator and $C : X \supset D(T) \rightarrow X^*$ is at least bounded. As it is usually assumed in this setting, the space X is reflexive and the spaces X, X^* are locally uniformly convex. Thus, J is now a bicontinuous mapping. If X, X^* are uniformly convex, then $J : X \rightarrow X^*, J^{-1} : X^* \rightarrow X^{**}$ are uniformly continuous on bounded sets. An operator $T : X \supset D(T) \rightarrow 2^{X^*}$ is “monotone” if for every $x, y \in D(T)$ and $u^* \in Tx, v^* \in Ty$ we have

$$\langle u^* - v^*, x - y \rangle \geq 0.$$

A monotone operator is “maximal monotone” if $T + \lambda J$ is surjective for all $\lambda > 0$. An operator $T : X \supset D(T) \rightarrow 2^{X^*}$ is said to be of “type (S) ” on the set $E \subset D(T)$, if for every sequence $\{x_n\} \subset E$ with $x_n \rightarrow x_0 \in X$ and

$$\lim_{n \rightarrow \infty} \langle v_n^*, x_n - x_0 \rangle = 0,$$

for some $v_n^* \in Tx_n$, we have $x_n \rightarrow x_0$. It is said to be of “type (S_+) ” on the set $E \subset D(T)$ if for every sequence $\{x_n\} \subset E$ with $x_n \rightarrow x_0 \in X$ and

$$\limsup_{n \rightarrow \infty} \langle v_n^*, x_n - x_0 \rangle \leq 0,$$

for some $v_n^* \in Tx_n$, we have $x_n \rightarrow x_0$. It is well-known that, under our assumptions on the spaces X, X^* , the duality mapping J is of type (S_+) on X .

For fundamental properties of monotone operators, and other related concepts, the reader is referred to Barbu [2], Browder [5], Cioranescu [7], Pascali and Sburlan [35] and Zeidler [41]. We start this section with an extension of Theorem 2.

Theorem 6. Let $T : X \supset D(T) \rightarrow 2^{X^*}$ be maximal monotone and (i) $C : D(T) \rightarrow X^*$ compact. Assume that there exist $\beta \in \Gamma$ and $Q > 0$ such that: for every $x \in D(T)$ with $\|x\| \geq Q$ and every $v^* \in Tx$ we have

$$(22) \quad \langle v^* + Cx, x \rangle \geq -\beta(\|x\|)\|x\|^2.$$

Then $R(T + C + \epsilon J) = X^*$, for every $\epsilon \in (0, 1)$. Assume, further, that

$$(***) \quad \liminf_{\substack{\|x\| \rightarrow \infty \\ x \in D(T)}} \frac{|Tx + Cx|}{\|x\|} > 0.$$

Then $\overline{R(T + C)} = X^*$. Let one of the following conditions hold:

- (a) $C : \overline{D(T)} \rightarrow X^*$ is completely continuous;
- (b) instead of (i), $C : \overline{D(T)} \rightarrow X^*$ is bounded and continuous, and $(T + J)^{-1}$ is compact.

Then $R(T + C) = X^*$.

Proof. As in the proof of Theorem 2, we consider the homotopy inclusion

$$t(Tx + Cx + \epsilon J)x + (1 - t)Jx \ni ts,$$

where $t \in [0, 1]$, $\epsilon \in (0, 1)$ and $s \in X^*$. This inclusion can be written as

$$(23) \quad u^* = H(t, u^*) \equiv -t[C(t(T + \epsilon J) + (1 - t)J)^{-1}u^* - s],$$

where

$$x = (t(T + \epsilon J) + (1 - t)J)^{-1}u^*.$$

We remark here that $T_t \equiv t(T + \epsilon J) + (1 - t)J : D(T) \rightarrow 2^{X^*}$ is a surjective maximal monotone operator with inverse T_t^{-1} which is defined, bounded and continuous on all of X^* . The maximal monotonicity/surjectivity of T_t can be seen from

$$T_t \equiv tT + [1 - t(1 - \epsilon)]J,$$

because $1 - t(1 - \epsilon) \geq \epsilon$, $t \in [0, 1]$. The continuity of the inverse T_t^{-1} follows from Cioranescu [7, Proposition 3.10]. To see its boundedness, fix $t \in (0, 1]$. If $t = 0$, the boundedness of T_t^{-1} is obvious. Fix $x_0 \in D(T)$, $v_0^* \in Tx_0$, and let $\{y_n^*\} \subset X^*$ be a bounded sequence. Let

$$x_n = (t(T + \epsilon J) + (1 - t)J)^{-1}y_n^*.$$

Then, for some $v_n^* \in Tx_n$,

$$(24) \quad tv_n^* + [1 - t(1 - \epsilon)]Jx_n = y_n^*,$$

or

$$[t/\lambda(t)]v_n^* + Jx_n = [1/\lambda(t)]y_n^*,$$

where $\lambda(t) \equiv 1 - t(1 - \epsilon)$, $t \in [0, 1]$. Now, we evaluate (24) at the point $x_n - x_0$ to get

$$[t/\lambda(t)]\langle v_n^*, x_n - x_0 \rangle + \langle Jx_n^*, x_n - x_0 \rangle = [1/\lambda(t)]\langle y_n^*, x_n - x_0 \rangle,$$

or

$$\begin{aligned} & [t/\lambda(t)]\langle v_n^* - v_0^*, x_n - x_0 \rangle + [t/\lambda(t)]\langle v_0^*, x_n - x_0 \rangle \\ & \quad + \langle Jx_n - Jx_0, x_n - x_0 \rangle + \langle Jx_0, x_n - x_0 \rangle \\ & = [1/\lambda(t)]\langle y_n^*, x_n - x_0 \rangle. \end{aligned}$$

Using the monotonicity of T and the fact that

$$\langle Jx_n - Jx_0, x_n - x_0 \rangle \geq (\|x_n\| - \|x_0\|)^2,$$

we obtain

$$(\|x_n\| - \|x_0\|)^2 \leq [t/\lambda(t)]\|v_0^*\| + \|Jx_0\| + [1/\lambda(t)]\|y_n^*\| \|x_n - x_0\|,$$

which shows the boundedness of the sequence $\{x_n\}$. Hence, T_t^{-1} is bounded.

We can further rewrite (23) as follows:

$$\begin{aligned} (25) \quad u^* = H(t, u^*) &= -t [C(tT + (1 - t(1 - \epsilon))J)^{-1}u^* - s] \\ &= -t [C([t/(1 - t(1 - \epsilon))]T + J)^{-1}[1/(1 - t(1 - \epsilon))]u^* - s] \\ &= -t [C((t/\lambda(t))T + J)^{-1}[(1/\lambda(t))u^*] - s], \end{aligned}$$

where $H(0, u^*) \equiv 0$. The function $H(t, \cdot)$ is compact, for every $t \in [0, 1]$, under any one of the assumptions made on the operators C , $(T + J)^{-1}$. In fact, it was shown in [17] (see, also, the resolvent identity in the comments following the proof of Theorem 8 below) that the compactness of the resolvent $(T + J)^{-1}$ implies the compactness of every resolvent $(kT + J)^{-1}$, $k > 0$. We are going to show, later in this proof, that the set

$$S \equiv \overline{H([0, 1], \overline{B_r(0)})}$$

is compact, for every $r > 0$.

In order to show the continuity of $H(t, u^*)$ on any set $[0, 1] \times \overline{B_r(0)}$, $r > 0$, we set $Q \equiv B_q(0) \subset X^*$ and observe that the set

$$K = \bigcup_{t \in [0, 1]} \{((t/\lambda(t))T + J)^{-1}\overline{Q}\}$$

is bounded. In fact, let $\{x_n\} \subset K$. Then there exist $\{t_n\} \subset [0, 1]$ and $\{u_n^*\} \subset \overline{Q}$ such that

$$x_n = ((t_n/\lambda(t_n))T + J)^{-1}u_n^*.$$

We have, for $q(t) \equiv t/\lambda(t)$ and some $v_n^* \in Tx_n$,

$$q(t_n)v_n^* + Jx_n = u_n^*.$$

This implies

$$\begin{aligned} & q(t_n)\langle v_n^* - v_0^*, x_n - x_0 \rangle + q(t_n)\langle v_0^*, x_n - x_0 \rangle \\ & \quad + \langle Jx_n - Jx_0, x_n - x_0 \rangle + \langle Jx_0, x_n - x_0 \rangle \\ & = \langle u_n^*, x_n - x_0 \rangle, \end{aligned}$$

where $x_0 \in D(T)$ and $v_0^* \in Tx_0$. This inequality implies

$$(\|x_n\| - \|x_0\|)^2 \leq [q(t_n)\|v_0^*\| + \|Jx_0\| + \|u_n^*\|]\|x_n - x_0\|,$$

and the boundedness of $\{x_n\}$, with a bound depending only on $\|v_0^*\|$, $\|x_0\|$ and q , follows from the boundedness of $\{u_n^*\}$.

From the boundedness of the set K we have immediately that the set

$$L \equiv \bigcup_{t \in [0,1]} \left\{ (q(t)T + J)^{-1} \left[[1/\lambda(t)]\overline{B_r(0)} \right] \right\}$$

is bounded, for each $r > 0$, for there exists a ball $B_q(0)$ such that

$$\bigcup_{t \in [0,1]} \{ [1/\lambda(t)]B_r(0) \} \subset B_q(0), \quad t \in [0, 1].$$

Consequently, by the boundedness of C , the set

$$M \equiv \bigcup_{t \in [0,1]} \left\{ C(q(t)T + J)^{-1} \left[[1/\lambda(t)]\overline{B_r(0)} \right] \right\} = CL$$

is bounded. Letting $\mu(r)$ be an upper bound for the set M , we have

$$\|H(t, u^*)\| \leq t[\mu(r) + \|s\|], \quad t \in [0, 1],$$

which shows the continuity of the function $H(\cdot, u^*)$ at $t = 0$, uniformly w.r.t. $u^* \in \overline{B_r(0)}$. It also shows the continuity of $H(t, u^*)$ at any point $(0, u^*)$, $u^* \in \overline{B_r(0)}$.

To see the continuity of $H(t, u^*)$ at any point $(t_0, u_0^*) \in (0, 1] \times \overline{B_r(0)}$, we fix such a point and let $(t_n, u_n^*) \in [0, 1] \times \overline{B_r(0)}$ be such that $t_n \rightarrow t_0$ and $u_n^* \rightarrow u_0^*$ as $n \rightarrow \infty$. Naturally, we may assume that $t_n > 0$, $n = 1, 2, \dots$. Let

$$(26) \quad \begin{aligned} v_n^* &\equiv H(t_n, u_n^*) = t_n [C(q(t_n)T + J)^{-1}(p(t_n)u_n^*) - s], \\ v_0^* &\equiv H(t_0, u_0^*) = t_0 [C(q(t_0)T + J)^{-1}(p(t_0)u_0^*) - s], \end{aligned}$$

where $p(t) = 1/\lambda(t)$, $t \in (0, 1]$. Let

$$\begin{aligned} x_n &\equiv (q(t_n)T + J)^{-1}(p(t_n)u_n^*) \\ x_0 &\equiv (q(t_0)T + J)^{-1}(p(t_0)u_0^*). \end{aligned}$$

We have, for some $z_n^* \in Tx_n$, $z_0^* \in Tx_0$,

$$\begin{aligned} q(t_n)z_n^* + Jx_n &= p(t_n)u_n^*, \\ q(t_0)z_0^* + Jx_0 &= p(t_0)u_0^*, \end{aligned}$$

which leads to

$$(27) \quad \begin{aligned} &\langle q(t_n)z_n^* - q(t_n)z_0^*, x_n - x_0 \rangle + [q(t_n) - q(t_0)]\langle z_0^*, x_n - x_0 \rangle \\ &\quad + \langle Jx_n - Jx_0, x_n - x_0 \rangle \\ &= \langle p(t_n)u_n^* - p(t_0)u_0^*, x_n - x_0 \rangle. \end{aligned}$$

We note that $\{x_n\}$ is a bounded sequence because the sequence $\{p(t_n)u_n^*\}$ is bounded and the sequence $\{x_n\}$ lies in the set L above, which is also bounded. Letting ν denote an upper bound for $\{x_n\}$ and using (27), we obtain

$$\begin{aligned} \langle Jx_n - Jx_0, x_n - x_0 \rangle &\leq |q(t_n) - q(t_0)| \|z_0^*\| \|x_n - x_0\| \\ &\quad + \|p(t_n)u_n^* - p(t_0)u_0^*\| \|x_n - x_0\| \\ &\leq [|q(t_n) - q(t_0)| \|z_0^*\| + \|p(t_n)u_n^* - p(t_0)u_0^*\|] (\nu + \|x_0\|). \end{aligned}$$

It follows that

$$(28) \quad \limsup_{n \rightarrow \infty} \langle Jx_n - Jx_0, x_n - x_0 \rangle \leq 0.$$

Since J is of type (S_+) , we have $x_n \rightarrow x_0$. Using this property in

$$\begin{aligned} \|v_n^* - v_0^*\| &\leq \|t_n Cx_n - t_0 Cx_0\| + |t_n - t_0| \|s\| \\ &\leq |t_n - t_0| \|Cx_n\| + t_0 \|Cx_n - Cx_0\| + |t_n - t_0| \|s\|, \end{aligned}$$

along with the boundedness and the continuity of C , we obtain the desired continuity property of $H(t, u^*)$.

We now show the compactness of the set S . If the operator $C : D(T) \rightarrow X^*$ is compact, then the boundedness of the set L implies that the set CL is relatively compact. This implies that the set $CL - s$ is also relatively compact. Letting $U \equiv [0, 1] \cdot \overline{CL - s}$, we see that U is a compact set because it is the image of the set $[0, 1] \times \overline{CL - s}$ by the operation of scalar multiplication. Since $S \subset U$, we have the compactness of the set S . If, on the other hand, $C : \overline{D(T)} \rightarrow X^*$ is continuous and bounded and the resolvent $(T + J)^{-1}$ is compact, then the compactness of the set S follows exactly as in the corresponding part of the proof of Theorem 2. It is therefore omitted. The degree function $d(I - H(t, \cdot), B_r(0), 0)$ is well-defined and the rest of the proof follows the corresponding proof steps of Theorem 2. We only make a comment for the case (a). In case (a) we use the fact that $\overline{D(T)}$ is a convex set. Thus, whenever $\{x_n\} \subset D(T)$ with $x_n \rightarrow x_0 \in \overline{D(T)}$, then $Cx_n \rightarrow Cx_0$. Moreover, $v_n^* \rightarrow -Cx_0 + s \in Tx_0$, where $v_n^* \in Tx_n$ satisfies

$$v_n^* + Cx_n + (1/n)Jx_n = s. \blacksquare$$

In order to be able to use homotopies as in Theorems 3 and 4, but for maximal monotone operators, we need the Yosida approximants for maximal monotone mappings introduced by Brézis, Crandall and Pazy [4] (cf. also Barbu [2, p. 41] and Pascali and Sburlan [35, p. 128]). Given a maximal monotone operator $T : X \supset D(T) \rightarrow 2^{X^*}$, $x \in X$ and $\lambda > 0$, there exists a unique $x_\lambda \in D(T)$ such that

$$(29) \quad J(x_\lambda - x) + \lambda T x_\lambda \ni 0.$$

We set

$$(30) \quad J_\lambda x = x_\lambda \text{ and } T_\lambda x = \frac{1}{\lambda} J(x - x_\lambda).$$

We know that $J_\lambda : X \rightarrow X$, $T_\lambda : X \rightarrow X^*$ are single-valued and bounded operators. We also know that both operators are demicontinuous with T_λ maximal monotone. In addition, it is easy to see that $T_\lambda x \in T J_\lambda x$, $x \in X$. The demicontinuity of J_λ can be found in the proof of Proposition 1.1 of Barbu [2]. The rest of these properties can also be found in [2], [4] and [35].

Since it is generally easier to check the compactness of $(\lambda T + J)^{-1}$ than that of J_λ , we establish the following lemma.

Lemma 3. For a maximal monotone operator $T : X \supset D(T) \rightarrow 2^{X^*}$, the resolvent J_λ is compact if and only if $(\lambda T + J)^{-1}$ is compact. Consequently, if J_λ is compact for some $\lambda > 0$, the same is true for all $\lambda \in (0, \infty)$.

Proof. Given $x \in X$ and $\lambda > 0$ such that (29) holds, we have

$$x \in x_\lambda + \lambda J^{-1} T x_\lambda.$$

This gives us the following representation for J_λ :

$$J_\lambda x = (I + \lambda J^{-1} T)^{-1} x, \quad x \in X.$$

We show first that if $(\lambda T + J)^{-1}$ is compact, then so is J_λ . To this end, let $(\lambda T + J)^{-1}$ be compact, for some $\lambda > 0$, let $\{x_n\}$ be a bounded sequence in X , and let

$$y_n \equiv J_\lambda x_n = (I + \lambda J^{-1} T)^{-1} x_n.$$

Then

$$\lambda J^{-1} T y_n + y_n \ni x_n,$$

or

$$\lambda T y_n + J y_n \ni J y_n + J(x_n - y_n).$$

This produces the representation

$$y_n = (\lambda T + J)^{-1} (J y_n + J(x_n - y_n)).$$

Since the sequences $\{x_n\}$, $\{y_n\}$ and J are bounded and $(\lambda T + J)^{-1}$ is compact, we have that $\{y_n\}$ lies in a compact set. Thus, J_λ maps bounded sets into relatively compact sets. We now show that J_λ is also continuous independently of the fact that it maps bounded sets onto relatively compact sets. Let $\{x_n\} \subset X$ be such that $x_n \rightarrow x_0$. Then

$$y_n \equiv J_\lambda x_n \rightarrow y_0 \equiv J_\lambda x_0.$$

We have

$$J(y_n - x_n) + \lambda T y_n \ni 0, \quad J(y_0 - x_0) + \lambda T y_0 \ni 0.$$

Thus, for some $v_n^* \in T y_n$, $v_0^* \in T y_0$,

$$\begin{aligned} & \langle J(y_n - x_n) - J(y_0 - x_0), (y_n - x_n) - (y_0 - x_0) \rangle \\ &= \langle J(y_n - x_n) - J(y_0 - x_0), (y_n - y_0) - (x_n - x_0) \rangle \\ &= \langle J(y_n - x_n) - J(y_0 - x_0), y_n - y_0 \rangle \\ &\quad - \langle J(y_n - x_n) - J(y_0 - x_0), x_n - x_0 \rangle \\ &\leq -\lambda \langle v_n^* - v_0^*, y_n - y_0 \rangle + \|J(y_n - x_n) - J(y_0 - x_0)\| \|x_n - x_0\| \\ &\leq \|J(y_n - x_n) - J(y_0 - x_0)\| \|x_n - x_0\|. \end{aligned}$$

This says that

$$\limsup_{n \rightarrow \infty} \langle J(y_n - x_n) - J(y_0 - x_0), (y_n - x_n) - (y_0 - x_0) \rangle \leq 0.$$

Since

$$\lim_{n \rightarrow \infty} \langle J(y_0 - x_0), (y_n - x_n) - (y_0 - x_0) \rangle = 0,$$

because $y_n - x_n \rightarrow y_0 - x_0$, we obtain

$$\limsup_{n \rightarrow \infty} \langle J(y_n - x_n), (y_n - x_n) - (y_0 - x_0) \rangle \leq 0.$$

Since J is of type (S_+) , we have $y_n - x_n \rightarrow y_0 - x_0$, i.e., $y_n \rightarrow y_0$.

Conversely, let J_λ be compact, for some $\lambda > 0$. We already know that the operator $(\lambda T + J)^{-1}$ is continuous. To show its compactness, let $\{x_n^*\} \subset X^*$ be bounded and let

$$x_n = (\lambda T + J)^{-1} x_n^*.$$

Then

$$\lambda T x_n + J x_n \ni x_n^*,$$

or

$$x_n + \lambda J^{-1} T x_n \ni x_n + J^{-1}(x_n^* - J x_n),$$

which produces the representation

$$x_n = (I + \lambda J^{-1} T)^{-1}(x_n + J^{-1}(x_n^* - J x_n)).$$

It follows that since $\{x_n\}$, $\{x_n^*\}$, J and J^{-1} are bounded and $(I + \lambda J^{-1} T)^{-1}$ is compact, we have that $\{x_n\}$ lies in a compact set. Thus, $(\lambda T + J)^{-1}$ is compact.

Guan and the author have shown in [15] that the compactness of $(\lambda T + J)^{-1}$, for some $\lambda > 0$, implies the compactness of $(\mu T + J)^{-1}$ for every $\mu > 0$. In view of the above equivalence, we have that if J_λ is compact, for some $\lambda > 0$, then J_μ is compact for all $\mu > 0$. ■

Using the resolvents J_λ , we can now establish Theorem 7 below, which extends to the maximal monotone case the main result of the author in [28]. All the results in that reference are for m -accretive operators.

Theorem 7. *Let $T : X \supset D(T) \rightarrow 2^{X^*}$ be maximal monotone and $C : D(T) \rightarrow X^*$. Let $(T + J)^{-1}$ be compact. Let $G \subset X$ be open, bounded and such that, for some $z \in D(T) \cap G$ and some $v^* \in Tz$,*

$$(31) \quad Tx - v^* \not\supset 0 \text{ and } \langle u^* + Cx, x - z \rangle > 0, \quad (x, u^*) \in (D(T) \cap \partial G) \times Tx.$$

Assume, further, that the operator $C(\lambda T + J)^{-1}$ is compact, where λ is a fixed positive constant, and the set $C(D(T) \cap \bar{G})$ is bounded. Then $0 \in (T + C)(D(T) \cap \bar{G})$.

Proof. We are planning to solve the perturbed problem

$$(32) \quad T J_\lambda x + C J_\lambda x \ni 0.$$

Using the fact that $T_\lambda x \in T J_\lambda x$, it suffices to solve instead of (32) the equation

$$(33) \quad T_\lambda x + C J_\lambda x = 0.$$

Using the definition of T_λ from (30), we see that (33) is equivalent to

$$(34) \quad x = J_\lambda x - \lambda J^{-1} C J_\lambda x.$$

We note that the operator $-\lambda J^{-1} C J_\lambda$ is compact because it is the composition of the compact mapping $C J_\lambda$ with the homeomorphism $-\lambda J^{-1}$. The compactness of $C J_\lambda$ follows from the compactness of $C(\lambda T + J)^{-1}$ and the fact that

$$C J_\lambda x = C(\lambda T + J)^{-1} [J J_\lambda x + J(x - J_\lambda x)],$$

which follows from the representation of y_n in the proof of Lemma 3. Since J_λ is compact by Lemma 3, the right-hand side of (34) defines a compact mapping.

We show first that we may assume that $z = 0 \in D(T) \cap G$ and $0 \in T(0)$. In fact, if this is not true, we consider the new operators \tilde{T} , \tilde{C} defined by

$$\tilde{T}x \equiv T(x + z) - v^*, \quad \tilde{C}x \equiv C(x + z) + v^*, \quad x \in \tilde{D}(T),$$

where v^* is as in the statement of the theorem and $D(\tilde{T}) \equiv D(T) - z$. We also set $\tilde{G} \equiv G - z$. It is easy to see that the operator \tilde{T} is maximal monotone on $\tilde{D}(T)$. To show the compactness of the resolvent

$$\tilde{S}_1 \equiv (J + \tilde{T})^{-1},$$

and hence any resolvent $(J + \lambda \tilde{T})^{-1}$, we note first that \tilde{S}_1 is a continuous mapping, by the maximal monotonicity of the operator \tilde{T} . Let $\{y_n^*\}$ be a bounded sequence in X^* and let $x_n = \tilde{S}_1 y_n^*$. Then

$$y_n^* = Jx_n + \tilde{v}_n^* = Jx_n + v_n^* - v^*, \quad n = 1, 2, \dots,$$

where $\tilde{v}_n^* \in \tilde{T}x_n$ and $v_n^* \in T(x_n + z)$. Thus,

$$J(x_n + z) + T(x_n + z) \ni y_n^* + J(x_n + z) - Jx_n + v^*,$$

or

$$x_n = (J + T)^{-1} [y_n^* + J(x_n + z) - Jx_n + v^*] - z.$$

We can now invoke the boundedness of the sequences $\{x_n\}$, $\{y_n^*\}$ and the duality mapping J , as well as the compactness of the operator $(J + T)^{-1}$, in order to conclude that $\{x_n\}$ lies in a compact set. This proves the compactness of the operator \tilde{S}_1 .

We now show that the operator $-\lambda J^{-1} \tilde{C} (I + \lambda J^{-1} \tilde{T})^{-1}$ is compact. To this end, we seek to express this operator in terms of the operator $C(I + \lambda T)^{-1}$. In fact, letting

$$\tilde{J}_\lambda \equiv (I + \lambda J^{-1} \tilde{T})^{-1}$$

and

$$x \equiv \tilde{J}_\lambda y = (I + \lambda J^{-1} \tilde{T})^{-1} y \in D(\tilde{T}),$$

for some $y \in X$, we have that there exists $w \in D(T)$ such that $x = w - z$ and

$$y \in x + \lambda \tilde{J}^{-1} \tilde{T}x = x + \lambda J^{-1} (T(x + z) - v^*) = w - z + \lambda J^{-1} [T(w) - v^*].$$

Thus,

$$J(y + z - w) \in \lambda(Tw - v^*)$$

and

$$J^{-1}(J(y + z - w) + \lambda v^*) \in \lambda J^{-1}Tw.$$

Consequently,

$$w = (I + \lambda J^{-1}T)^{-1} (w + J^{-1}(J(y + z - w) + \lambda v^*)).$$

Letting $x = w - z$ and using the above representation of w , we obtain

$$\begin{aligned} x &= \tilde{J}_\lambda y = (I + \lambda J^{-1}T)^{-1} [x + z + J^{-1}(J(y - x) + \lambda v^*)] - z \\ (35) \quad &= J_\lambda \left[\tilde{J}_\lambda y + z + J^{-1} \left[J(I - \tilde{J}_\lambda)y + \lambda v^* \right] \right] - z \\ &= J_\lambda Qy - z, \end{aligned}$$

where Qy is the expression in the brackets to the right of the first J_λ in (35). Since

$$\tilde{C}\tilde{J}_\lambda y = C(\tilde{J}_\lambda y + z) + v^* = CJ_\lambda Qy + v^*$$

and Q is continuous and bounded, we obtain that $-\lambda J^{-1}\tilde{C}\tilde{J}_\lambda$ is still a compact operator. To see that the second condition in (31) is satisfied with $z = 0$, it suffices to observe that

$$\langle (w^* - v^*) + (C(u + z) + v^*), u \rangle \geq 0,$$

for every $u \in D(\tilde{T}) \cap \partial\tilde{G}$ and every $w^* \in T(u + z)$. We note that the first condition (31) becomes now

$$0 \notin \tilde{T}x, \quad x \in D(\tilde{T}) \cap \partial\tilde{G}.$$

Finally, to show that the set $\tilde{C}(D(\tilde{T}) \cap \overline{\tilde{G}})$ is bounded, we note that

$$\begin{aligned} \tilde{C}(D(\tilde{T}) \cap \overline{\tilde{G}}) &= \tilde{C}[(D(T) - z) \cap (\overline{G - z})] \\ &= C [((D(T) - z) \cap (\overline{G - z})) + z] + v^* \\ &= C(D(T) \cap \overline{G}) + v^*. \end{aligned}$$

We have shown that it suffices to prove the theorem with $z = 0$ and $0 \in T(0)$.

Now, we let

$$U = (I + \lambda J^{-1}T)(D(T) \cap G)$$

and consider the homotopy functions

$$H_1(t, x) \equiv x - J_\lambda x + t\lambda J^{-1}CJ_\lambda x, \quad (t, x) \in [0, 1] \times \overline{U},$$

and

$$H_2(t, x) \equiv x - tJ_\lambda x, \quad (t, x) \in [0, 1] \times \overline{U}.$$

Since the operator J_λ is a continuous mapping on all of X , its inverse, $I + \lambda J^{-1}T$, is a set-valued mapping that maps relatively open (closed) sets of its domain $D(T)$

onto open (closed) sets of the space X . Because of this, the set U is open and the set

$$(I + \lambda J^{-1}T)(D(T) \cap \overline{G})$$

is closed. Thus, we have

$$\begin{aligned} & \overline{(I + \lambda J^{-1}T)(D(T) \cap \overline{G})} \\ &= (I + \lambda J^{-1}T)(D(T) \cap \overline{G}) \\ &= (I + \lambda J^{-1}T)(D(T) \cap G) \cup (I + \lambda J^{-1}T)(D(T) \cap \partial G) \\ &\supset \overline{(I + \lambda J^{-1}T)(D(T) \cap G)} \\ &= (I + \lambda J^{-1}T)(D(T) \cap G) \cup \partial((I + \lambda J^{-1}T)(D(T) \cap G)), \end{aligned}$$

which implies that

$$(I + \lambda J^{-1}T)(D(T) \cap \partial G) \supset \partial((I + \lambda J^{-1}T)(D(T) \cap G)).$$

Since

$$\overline{(I + \lambda J^{-1}T)(D(T) \cap G)} \subset (I + \lambda J^{-1}T)(D(T) \cap \overline{G}),$$

the homotopies $H_1(t, y)$, $H_2(t, y)$ are well-defined. They also are compact displacements of the identity in y . Since J_λ maps the set \overline{U} onto a bounded subset of $D(T) \cap \overline{G}$ and $C(D(T) \cap \overline{G})$ is relatively compact, the operators $J_\lambda - t\lambda J^{-1}CJ_\lambda$ and tJ_λ have relatively compact range on \overline{U} . We may thus apply the Leray-Schauder degree function d_{LS} (cf. Nagumo [34]) to the mappings $H_1(t, \cdot)$, $H_2(t, \cdot)$. We are going to show first that

$$(36) \quad d_{LS}(H_1(1, \cdot), U, 0) = d_{LS}(H_2(1, \cdot), U, 0),$$

and then prove that

$$(37) \quad d_{LS}(H_2(1, \cdot), U, 0) = d_{LS}(I, U, 0) = 1,$$

the last equality holding because $0 \in U$. As we shall see below, to show our first assertion, it suffices to prove that the equation $H_1(t, x) = 0$ has no solution x on ∂U . Assume that the contrary is true. Then there exists $x_t \in \partial U$ such that

$$(38) \quad x_t = J_\lambda x_t - t\lambda J^{-1}CJ_\lambda x_t.$$

We may assume that $t \in [0, 1)$. In fact, if $t = 1$, then we have a solution x_1 of (34) which provides a solution to our problem (32). Equality (38) can also be written as

$$T_\lambda x_t + tCJ_\lambda x_t = 0.$$

This says that if $y_t \equiv J_\lambda x_t$, then there exists $w_t^* \in Ty_t$ such that

$$w_t^* + tCy_t = 0.$$

Since $y_t \in D(T) \cap \partial G$ and $Tx \not\equiv 0$ for $x \in D(T) \cap \partial G$, we have that $t \in (0, 1)$. Hence,

$$(39) \quad \langle w_t^* + tCy_t, y_t \rangle = \langle w_t^*, y_t \rangle + t \langle Cy_t, y_t \rangle = 0.$$

We consider the straight line segment $l(u) = au + b$, $u \in [0, 1]$, where $a \equiv \langle Cy_t, y_t \rangle$ and $b \equiv \langle w_t^*, y_t \rangle$. We know that $l(0) \geq 0$, because $0 \in T(0)$ and T is monotone. We also know that $l(1) > 0$, because of our assumed boundary condition. It follows that $l(u) > 0$, for every $u \in (0, 1)$. This contradicts (39). Hence

$$d_{LS}(H_1(1, \cdot), U, 0) = d_{LS}(H_1(0, \cdot), U, 0) = d_{LS}(H_2(1, \cdot), U, 0).$$

In order to prove (37), we are going to show that the equation $H_2(t, x) = 0$ has no solution $x \in \partial U$. Let x_t be such a solution for some $t \in [0, 1]$. Obviously, $t \neq 0$ because $x_t \in \partial U$, while $0 \in U$ and U is an open set. Also, $t \neq 1$. In fact, if $t = 1$, then $x_t - J_\lambda x_t = 0$ implies

$$0 = \frac{1}{\lambda} J(x_t - J_\lambda x_t) = T_\lambda x_t \in T(J_\lambda x_t) \subset T(D(T) \cap \partial G),$$

i.e., a contradiction to our assumption. Thus, $t \in (0, 1)$ and

$$x_t = tJ_\lambda x_t.$$

Letting $y_t \equiv J_\lambda x_t$, we see that

$$y_t + \lambda J^{-1} T y_t \ni t y_t.$$

This implies

$$\lambda T y_t \ni -(1 - t) J y_t,$$

which yields, for some $w_t \in T y_t$,

$$0 \leq \lambda \langle w_t^*, y_t \rangle = -(1 - t) \langle J y_t, y_t \rangle = -(1 - t) \|y_t\|^2 < 0,$$

the last inequality holding because $t \in (0, 1)$ and $y_t \in D(T) \cap \partial G$. It follows that (37) is true. This says that $H_1(1, x)$ is solvable in $x \in U$. The proof is complete. ■

If $z = 0$ in Theorem 7, we obtain the following simpler corollary.

Corollary 4. *Let $T : X \supset D(T) \rightarrow 2^{X^*}$ be maximal monotone, with $0 \in D(T)$ and $0 \in T(0)$, and $C : D(T) \rightarrow X^*$. Let $(T + J)^{-1}$ be compact. Let $G \subset X$ be open, bounded and such that $0 \in G$ and*

$$Tx \not\equiv 0 \text{ and } \langle u^* + Cx, x \rangle > 0, \quad (x, u^*) \in (D(T) \cap \partial G) \times Tx.$$

Assume, further, that the operator $C(\lambda T + J)^{-1}$ is compact, where λ is a fixed positive constant, and the set $C(D(T) \cap \overline{G})$ is bounded. Then $0 \in (T + C)(D(T) \cap \overline{G})$. ■

For condensing mappings CJ_λ , we have the following version of Theorem 7.

Theorem 8. *Let $T : X \supset D(T) \rightarrow 2^{X^*}$ be maximal monotone, with $0 \in D(T)$ and $0 \in T(0)$. Assume that the duality mapping J^{-1} is Lipschitzian with Lipschitz constant $K > 0$, and let J_1 be compact. Let $G \subset X$ be open, bounded and such that $0 \in G$ and*

$$Tx \not\equiv 0 \text{ and } \langle u^* + Cx, x \rangle > 0, \quad (x, u^*) \in (D(T) \cap \partial G) \times Tx.$$

Assume that the operator $C(I + \lambda J^{-1}T)^{-1}$ is condensing, where λ is a fixed constant in $(0, 1/K)$, and the set $C(D(T) \cap \overline{G})$ is bounded. Then $0 \in (T + C)(D(T) \cap \overline{G})$.

Proof. The proof follows as in Theorem 7. However, the operator $-\lambda J^{-1}CJ_\lambda$ is now condensing, while the operator J_λ is still compact. Since

$$J_\lambda \overline{U} \subset D(T) \cap \overline{G}$$

and

$$(CJ_\lambda)(\overline{U}) \subset C(D(T) \cap \overline{G}),$$

we have that the operators J_λ and $-\lambda J^{-1}CJ_\lambda$ have bounded ranges on \overline{U} . As a consequence of this fact, we are allowed to use a degree theory for the mappings $H_1(t, x)$, $H_2(t, x)$ which can be found in Theorem 1 of [28]. That theorem contains a sketch of the proof of the existence of such a degree theory. ■

It is of importance to note that the operator C in Theorems 7 and 8 is not assumed to be continuous. This is why we are precluded from using homotopies like the ones used by the author in [22-24], the author and Liu [29] and Guan and the author [15-17]. The study of operators C which are not assumed to satisfy any continuity property was initiated recently by Hirano and Kalinde [19]. For an application of this setting, we refer to that paper as well as the paper of the author [28].

The compactness of the operator CJ_λ , in Theorem 7 and Corollary 4, implies the compactness of every operator CJ_μ , $\mu > 0$. To see this, we write J_μ in terms of J_λ in the following way. Let $x = J_\mu y$, for some $y \in X$. Then we have $x + \mu J^{-1}Tx \ni y$ and

$$(1/\mu)x + J^{-1}Tx \ni (1/\mu)y.$$

Hence

$$x + (\lambda/\mu)x + \lambda J^{-1}Tx \ni x + (\lambda/\mu)y,$$

or

$$x = J_\mu y = J_\lambda \left(\frac{\lambda}{\mu}y + \frac{\mu - \lambda}{\mu}x \right) = J_\lambda \left(\frac{\lambda}{\mu}y + \frac{\mu - \lambda}{\mu}J_\mu y \right).$$

This is a “resolvent identity” for the mappings J_λ . Since J_μ is a compact mapping, this identity shows that if CJ_λ is compact, then so is CJ_μ , for every $\mu > 0$.

Due to the above comments, we could have used J_1 in Theorem 7, and its proof, instead of J_λ . However, we need the proof of Theorem 7, as it stands, for the proof of Theorem 8 which imitates it.

Calvert and Gupta have shown in [6, Lemma 1.1] that the normalized duality mapping $J : L^p(\Omega) \rightarrow L^q(\Omega)$ satisfies a Lipschitz condition, provided that $p \in [2, \infty)$. Thus, Theorem 8 applies to the space $X = L^q(\Omega)$. Here, Ω is a bounded domain in \mathcal{R}^n and $(1/p) + (1/q) = 1$.

Open Problem 2. *Is Theorem 7 true without the assumption that $Tx - v^* \not\equiv 0$, for every $x \in D(T) \cap \partial G$? ■*

We are now going to study the effect of another boundary condition on the range of the sum of two operators. This condition implies condition (22) of Theorem 6 and is accompanied by a growth condition on the sum $T + C$ which is weaker than (***) . Because of these facts, we have the immediate solvability of $(T + C + \epsilon J)x \ni p$, for all $\epsilon \in (0, 1)$, but we do not have the surjectivity of the operator $T + C + \epsilon J$ as a consequence of Theorem 6.

Theorem 9. Let $T : X \supset D(T) \rightarrow 2^{X^*}$ be maximal monotone and (a) $C : D(T) \rightarrow X^*$ compact. Let $S \subset X^*$. Assume that for every $s \in S$ there exist three positive constants $a = a(s)$, $q = q(s)$, $Q = Q(s)$ such that

$$a > 1 + \frac{\|s\|}{q}$$

and the following two conditions are satisfied:

- (i) $\|u^* + Cx\| \geq aq$, for every $x \in D(T)$ with $\|x\| \geq Q$, $u^* \in Tx$;
- (ii) we have

$$\langle u^* + Cx - s, x \rangle \geq -q\|x\|, \text{ for every } x \in D(T) \text{ with } \|x\| \geq Q$$

and every $u^* \in Tx$.

Then $S \subset \overline{R(T + C)}$. If, moreover, $C : \overline{D(T)} \rightarrow X^*$ is completely continuous, then $S \subset R(T + C)$. This is also true if instead of (a) it is assumed that $C : \overline{D(T)} \rightarrow X^*$ is continuous and bounded, and $(J + T)^{-1}$ is compact.

Proof. We consider the approximate problem

$$(40) \quad Tx + Cx + [(q + \epsilon)/Q]Jx \ni s,$$

where s is a given point in the set S and $\epsilon \in (0, 1)$ is chosen so that

$$a > 1 + \frac{\|s\| + \epsilon}{q}.$$

We note that in our assumptions we may use any number greater than Q , instead of Q itself. Thus, we may assume that Q is large enough so that $Q > q + \epsilon$. Consequently, the solvability of the problem (40), for any $s \in X^*$, follows from Theorem 6 because, by (ii),

$$\langle u^* + Cx, x \rangle \geq -[(\|s\| + q)/\|x\|]\|x\|^2 = -\beta(\|x\|)\|x\|^2,$$

for every $x \in D(T)$ with $\|x\| \geq Q$, $u^* \in Tx$, where

$$\beta(\rho) \equiv (\|s\| + q)/\rho \rightarrow 0 \text{ as } \rho \rightarrow \infty.$$

To show that every solution x of (40) has to lie in the ball $B_Q(0)$, assume that the contrary is true, and let x solve (40) with $\|x\| \geq Q$. Then, for some $u^* \in Tx$,

$$\begin{aligned} 0 &= \langle u^* + Cx - s, x \rangle + [(q + \epsilon)/Q]\langle Jx, x \rangle \\ &\geq -q\|x\| + [(q + \epsilon)/Q]\|x\|^2 \\ &\geq -q\|x\| + [(q + \epsilon)/Q]Q\|x\| \\ &= \epsilon\|x\| \\ &> 0, \end{aligned}$$

i.e., a contradiction. Let $\{Q_n\}$ be a sequence of numbers in $(0, \infty)$ such that $Q_n \geq Q$, $Q_n \rightarrow \infty$ as $n \rightarrow \infty$, and there exists a solution x_n of the problem

$$Tx + Cx + [(q + \epsilon)/Q_n]Jx \ni s$$

with $x_n \in B_{Q_n}(0)$. This is possible by what was shown above. Let us assume that $\|x_n\| \geq Q$, for some n . Then, for some $u_n^* \in Tx_n$, we have

$$\begin{aligned} 0 &= \|u_n^* + Cx_n + [(q + \epsilon)/Q_n]Jx_n - s\| \\ &\geq \|u_n^* + Cx_n\| - [|(q + \epsilon)/Q_n| \|Jx_n\| + \|s\|] \\ &> aq - [|(q + \epsilon)/Q_n| Q_n + \|s\|] \\ &= aq - (q + \epsilon + \|s\|) \\ &= \left[a - \left(1 + \frac{\|s\| + \epsilon}{q} \right) \right] q \\ &> 0, \end{aligned}$$

i.e., a contradiction. Since $\{Jx_n\}$ is a bounded sequence, we have immediately that $s \in \overline{R(T + C)}$ and $S \subset \overline{R(T + C)}$. The rest of the proof follows exactly as in Theorem 6. It is therefore omitted. ■

A special case of the assumptions of Theorem 9 allows for a generally better conclusion. Namely, in the proof of Theorem 9, $s \in \overline{(T + C)(D(T) \cap B_{Q(s)}(0))}$. This case is contained in the following result.

Theorem 9'. *Let $T : X \supset D(T) \rightarrow 2^{X^*}$ be maximal monotone with $0 \in D(T)$ and (a) $C : D(T) \rightarrow X^*$ compact. Let $S \subset X^*$. Assume that for every $s \in S$ there exist three positive constants $a = a(s)$, $q = q(s)$, $Q = Q(s)$ such that*

$$a > 1 + \frac{\|s\|}{q}$$

and the following two conditions are satisfied:

- (i) $\|u^* + Cx\| \geq aq$, for every $x \in D(T)$ with $\|x\| \geq Q$, $u^* \in Tx$;
- (ii) we have

$$\langle Cx + v^* - s, x \rangle \geq -q\|x\|, \text{ for every } x \in D(T) \text{ with } \|x\| \geq Q$$

and every $u^* \in Tx$.

Then, for every $s \in S$, $s \in \overline{(T + C)(D(T) \cap B_{Q(s)}(0))}$. If, moreover, C is completely continuous, then, for every $s \in S$, $s \in \overline{(T + C)(D(T) \cap \overline{B_{Q(s)}(0)})}$. This is also true if instead of (a) it is assumed that C is continuous and bounded, and $(J + T)^{-1}$ is compact.

Proof. Unlike the proof of Theorem 9, we no longer need to impose a lower bound on the parameter Q . In fact, we now use the homotopy equation

$$u^* = H(t, u^*) \equiv -t \left[\tilde{C}(\tilde{T} + (q/Q)J)^{-1}u^* - s \right],$$

where $\tilde{T}x \equiv Tx - v^*$ and $\tilde{C}x \equiv Cx + v^*$. The rest of the proof follows as in Theorem 14 of [29]. It is therefore omitted. ■

We are now ready for the following maximal monotone analogue of Theorem 14 in [29].

Corollary 5. Let $T : X \supset D(T) \rightarrow 2^{X^*}$ be maximal monotone and (a) $C : D(T) \rightarrow X^*$ compact. Assume that $0 \in D(T)$ and there exist three positive constants a_1, b, r and $v^* \in T(0)$ such that

$$a_1 > 2 + \frac{r}{\|v^*\| + r}$$

and the following conditions are satisfied.

- (i) $\|u^* + Cx\| \geq a_1(\|v^*\| + r)$, for every $x \in D(T)$ with $\|x\| \geq b, u^* \in Tx$;
- (ii) $\langle Cx, x \rangle \geq -(\|v^*\| + r)\|x\|$, for every $x \in D(T)$ with $\|x\| \geq b$.

Then $\overline{B_r(0)} \subset \overline{(T+C)(D(T) \cap B_b(0))}$. If, moreover, $C : \overline{D(T)} \rightarrow X^*$ is completely continuous, then $\overline{B_r(0)} \subset \overline{(T+C)(D(T) \cap B_b(0))}$. This is also true if instead of (a) it is assumed that $C : \overline{D(T)} \rightarrow X^*$ is continuous and bounded, and $(J + T)^{-1}$ is compact.

Proof. In Theorem 9', we take $S = \overline{B_r(0)}, q = 2(\|v^*\| + r), Q = b$, and $a = a_1/2$. We see here that the parameters q, Q and a do not depend on the particular point $s \in S$. For every $u^* \in Tx$ with $\|x\| \geq Q$ and every $s \in S$ we have

$$\begin{aligned} \langle u^* + Cx - s, x \rangle &= \langle u^* - v^* + Cx + v^* - s, x \rangle \\ &= \langle u^* - v^*, x \rangle + \langle Cx, x \rangle + \langle v^* - s, x \rangle \\ &\geq -(\|v^*\| + r)\|x\| - (\|v^*\| + \|s\|)\|x\| \\ &\geq -2(\|v^*\| + r)\|x\|. \end{aligned}$$

Thus,

$$a = a_1/2 > 1 + \frac{r}{2(\|v^*\| + r)} \geq 1 + \frac{\|s\|}{q}. \blacksquare$$

Finally, we have a new surjectivity result for perturbations of maximal monotone operators. For a set A , we set $|A| \equiv \inf\{\|x\| : x \in A\}$.

Corollary 6. Let $T : X \supset D(T) \rightarrow 2^{X^*}$ be maximal monotone with $0 \in D(T)$ and (a) $C : D(T) \rightarrow X^*$ compact. Assume that the following two conditions hold:

- (i)
$$\lim_{\substack{\|x\| \rightarrow \infty \\ x \in D(T)}} |Tx + Cx| = \infty;$$
- (ii) there exists a constant $d > 0$ such that $\langle Cx, x \rangle \geq -d\|x\|$, for every $x \in D(T)$ with sufficiently large norm.

Then $\overline{R(T+C)} = X^*$. If, moreover, $C : \overline{D(T)} \rightarrow X^*$ is completely continuous, then $R(T+C) = X^*$. This is also true if instead of (a) it is assumed that $C : \overline{D(T)} \rightarrow X^*$ is continuous and bounded, $(J + T)^{-1}$ is compact.

Proof. Given a ball $S \equiv \overline{B_r(0)} \subset X^*$, for some $r > 0$, we let $s \in S, v^* \in T(0)$ and $u^* \in Tx$, where $x \in D(T)$ has a sufficiently large norm. We have

$$\begin{aligned} \langle u^* - v^* + Cx + v^* - s, x \rangle &\geq -(d + \|v^*\| + \|s\|)\|x\| \\ &\geq -(d + \|v^*\| + r)\|x\| \\ &= -q\|x\|, \end{aligned}$$

where q is an obvious positive constant. Since $|Tx + Cx| \geq aq$, for all large numbers $a > 0$ and all large $\|x\|$, we obtain that $\overline{B_r(0)} \subset \overline{R(T+C)}$. Since $r > 0$ is arbitrary, we have that $\overline{R(T+C)} = X^*$. The rest of the proof follows as before. It is therefore omitted. \blacksquare

4. EIGENVALUES FOR PERTURBATIONS OF MAXIMAL MONOTONE OPERATORS

The Yosida resolvents $J_\lambda = (I + \lambda J^{-1}T)^{-1}$ may also be used in the solvability of nonlinear eigenvalue problems involving perturbations of maximal monotone operators. Since such resolvents were not used in [16], we find it instructive to establish some results in this direction. Again, X is a real reflexive Banach space and X, X^* are locally uniformly convex. We also assume that the space X is infinite dimensional. We do this because of Lemma A below, which can be found in [16].

Lemma A. *Let $K \subset X^*$ be compact and assume that $\|x^*\| \geq \alpha, x^* \in K$, where α is a positive constant. Then there exists $y_0^* \in \partial B_1(0)$ such that $\{ty_0^* : t \geq \alpha\} \cap K = \emptyset$.*

The next result is an interesting extension of the maximal monotone analogue of Theorem 2.5 in [16].

Theorem 10. *Let $G \subset X$ be open, bounded, and let $T : X \supset D(T) \rightarrow 2^{X^*}$ be maximal monotone and such that $0 \in D(T) \cap G, 0 \in T(0)$ and $0 \notin T(J_\mu(\partial G))$, for some $\mu > 0$. Assume that $(J + T)^{-1}$ is compact. Let $C : D(T) \rightarrow X^*$ be such that CJ_1 is compact and $\|CJ_\mu x\| \geq \alpha, x \in \partial G$, where α is a positive constant. Then there exists $(\lambda_0, x_0) \in (0, \infty) \times D(T)$ such that $Cx_0 \neq 0$ and $Tx_0 - \lambda_0 Cx_0 \ni 0$.*

Proof. We consider first the problem

$$(41) \quad T_\mu x - \lambda C J_\mu x = 0, \quad (\lambda, x) \in (0, \infty) \times \partial G.$$

This problem can be written as

$$(42) \quad H(\lambda, x) \equiv x - J_\mu x - \lambda \mu J^{-1} C J_\mu x = 0, \quad (\lambda, x) \in (0, \infty) \times \partial G.$$

We know that J_μ is compact, by Lemma 3, and that the operator CJ_μ is also compact by the comments preceding the statement of Theorem 9. In order to solve (42) in $(\lambda, x) \in (0, \infty) \times \partial G$, we show first that there exists $\lambda_0 \in (0, \infty)$ such that

$$(43) \quad x - J_\mu x - \mu J^{-1}(\lambda_0 C J_\mu x + \eta y_0^*) \neq 0, \quad (\eta, x) \in (0, \infty) \times \partial G,$$

where y_0^* is as in Lemma A with $K \equiv \overline{-(CJ_\mu)(\partial G)}$. The operator

$$x \rightarrow J_\mu x + \mu J^{-1}(\lambda_0 C J_\mu x + \eta y_0^*)$$

is compact for every $\eta \in [0, \infty)$. The set K is compact because ∂G is bounded and the operator CJ_μ is compact.

Let us assume that there is no $\lambda_0 > 0$ such that (43) holds. Then there exist sequences $\{\lambda_m\} \subset (0, \infty), \{\eta_m\} \subset (0, \infty)$ and $\{x_m\} \subset \partial G$ such that $\lambda_m \rightarrow \infty$ and

$$x_m - J_\mu x_m - \mu J^{-1}(\lambda_m C J_\mu x_m + \eta_m y_0^*) = 0, \quad m = 1, 2, \dots$$

Thus,

$$\frac{1}{\mu} J(x_m - J_\mu x_m) - (\lambda_m C J_\mu x_m + \eta_m y_0^*) = 0, \quad m = 1, 2, \dots,$$

or

$$T_\mu x_m - (\lambda_m C J_\mu x_m + \eta_m y_0^*) = 0.$$

Since $\{T_\mu x_m\}$ is a bounded sequence, we obtain

$$\lim_{m \rightarrow \infty} [CJ_\mu x_m + (\eta_m/\lambda_m)y_0^*] = 0.$$

Since $\|CJ_\mu x_m\| \geq \alpha$ for all m and $\{CJ_\mu x_m\}$ is bounded, we may assume that $\{\eta_m/\lambda_m\}$ converges to some number $\nu \geq 0$. Since CJ_μ is compact, we may also assume that $CJ_\mu x_m \rightarrow u^*$ with $u^* = -\nu y_0^*$. Since

$$\alpha \leq \|u^*\| = \nu \|y_0^*\| = \nu$$

and $\nu y_0^* \in \overline{-(CJ_\mu)(\partial G)}$, we have a contradiction to the choice of y_0^* . Consequently, (43) is true. If (42) holds for $\lambda = \lambda_0$ and some $x \in \partial G$, then we have the solvability of the problem (41). Otherwise, (43) holds for all $\eta \geq 0$. Assuming that the latter is the case, we show that the Leray-Schauder degree $d(H(\lambda_0, \cdot), G, 0) = 0$. To this end, let $d(H(\lambda_0, \cdot), G, 0) \neq 0$ and consider the homotopy

$$H_{1,m}(t, x) \equiv x - J_\mu x - \mu J^{-1}(\lambda_0 CJ_\mu x + tm y_0^*), t \in [0, 1], x \in \overline{G}, m = 1, 2, \dots$$

We know that $H_{1,m}(t, \partial G) \not\equiv 0, t \in [0, 1]$. It follows that the Leray-Schauder degree $d(H_{1,m}(t, \cdot), G, 0)$ is well-defined, and we have

$$d(H_{1,m}(1, \cdot), G, 0) = d(H_{1,m}(0, \cdot), G, 0) = d(H(\lambda_0, \cdot), G, 0) \neq 0.$$

This implies that there exists a sequence $\{x_m\} \subset G$ such that $H_{1,m}(1, x_m) = 0$ or

$$T_\mu x_m - \lambda_0 CJ_\mu x_m = m y_0^*.$$

This is a contradiction to the boundedness of the sequence

$$\{(T_\mu - \lambda_0 CJ_\mu)x_m\}.$$

It follows that $d(H(\lambda_0, \cdot), G, 0) = 0$.

We now show that $d(H(0, \cdot), G, 0) = 1$. Consider the new homotopy

$$H_2(t, x) \equiv x - tJ_\mu x, (t, x) \in [0, 1] \times \overline{G}.$$

Assume that x_t is a solution of $H_2(t, x) = 0$ with $x_t \in \partial G$. Let

$$u_t \equiv J_\mu x_t = (I + \mu J^{-1}T)^{-1}x_t \in D(T).$$

Then there exists $v_t^* \in Tu_t$ such that

$$u_t + \mu J^{-1}v_t^* = x_t$$

and

$$v_t^* = -\frac{1}{\mu}(1-t)Ju_t.$$

Since $0 \in T(0)$, we have

$$0 \leq \langle v_t^*, u_t \rangle = -\frac{1}{\mu}(1-t)\langle Ju_t, u_t \rangle = -\frac{1}{\mu}(1-t)\|u_t\|^2 \leq 0.$$

Thus, either $t = 1$ or $u_t = 0$. In either case, $0 = v_t^* \in Tu_t = TJ_\mu x_t$. Since $x_t \in \partial G$, we have a contradiction. It follows that the equation $H_2(t, x) = 0$ has no solutions $x \in \partial G$, for any value of $t \in [0, 1]$. Thus, the degree $d_2(H(t, \cdot), G, 0)$ is well-defined, constant and equal to 1 because $d(H_2(t, \cdot), G, 0) = d(H_2(0, \cdot), G, 0) = 1$. Consequently, there exists $(\lambda_0, x) \in (0, \infty) \times \partial G$ such that $H(\lambda_0, x) = 0$, or

$$T_\mu x - \lambda_0 CJ_\mu x = 0.$$

Letting $x_0 = J_\mu x$, we have $\|Cx_0\| \geq \alpha > 0$ and $Tx_0 - \lambda_0 Cx_0 \ni 0$. This completes the proof. ■

We give Corollary 7 below because we want to comment on the boundary condition imposed upon the operator C in a special case.

Corollary 7. *Let the assumptions of Theorem 10 be satisfied, but with the operator $C : \overline{D(T)} \rightarrow X^*$ and the compactness of the operator CJ_1 replaced by “ C is continuous.” Then there exists $(\lambda_0, x_0) \in (0, \infty) \times D(T)$ such that $Cx_0 \neq 0$ and $Tx_0 - \lambda_0 Cx_0 \ni 0$.*

The condition $\|CJ_\mu x\| \geq \alpha$, $x \in \partial G$, will hold in Corollary 7 if we assume instead that $Cx \neq 0$, $x \in \overline{D(T)}$. In fact, let us assume that the latter is true and that

$$\inf_{x \in \partial G} \{\|CJ_\mu x\|\} = 0.$$

Then there exists a sequence $\{x_m\} \subset \partial G$ such that $CJ_\mu x_m \rightarrow 0$ as $m \rightarrow \infty$. Since J_μ is compact, we may (and do) assume that $J_\mu x_m \rightarrow u_0 \in \overline{D(T)}$. Since $C : \overline{D(T)} \rightarrow X^*$ is continuous, $CJ_\mu x_m \rightarrow Cu_0 = 0$, i.e., a contradiction.

If

$$L \equiv \sup_{x \in \partial G} \{\|x\|\},$$

it suffices to assume, in the appropriate places in Theorem 10 and Corollary 7, that $0 \notin T(D(T) \cap \overline{B_{2L}(0)})$ and $\|Cx\| \geq \alpha$, $x \in D(T) \cap \overline{B_{2L}(0)}$. In fact, on page 43 of Barbu's book [2] we find

$$\|J_\mu x - x\|^2 \leq \|x - u\| \|J_\mu x - x\| + \mu \|x - u\| \|v\| + \mu \|v\| \|x - J_\mu x\|,$$

for any $u \in D(T)$ and $v \in Tu$. Since $0 \in T(0)$, we may take $u = 0$, $v = 0$ to obtain $\|J_\mu x - x\| \leq \|x\|$, which says that $\|J_\mu x\| \leq 2\|x\| \leq 2L$, $x \in \partial G$. Consequently, $J_\mu(\partial G) \subset \overline{B_{2L}(0)}$. In particular, if $G = B_r(0)$, then

$$T(J_\mu(\partial G)) \subset T(D(T) \cap \overline{B_{2r}(0)}) \quad \text{and} \quad C(J_\mu(\partial G)) \subset C(D(T) \cap \overline{B_{2r}(0)}).$$

This assumption on the operator C requires it to be bounded below away from zero only on the ball $\overline{B_{2L}(0)}$, and is considerably weaker than the one above involving the entire set $\overline{D(T)}$. This is particularly important when the operator T is densely defined.

If the set $T(D(T) \cap \overline{G})$ is bounded, then we can have an eigenvalue result under the more natural assumption $\|Cx\| \geq \alpha$, $x \in D(T) \cap \partial G$.

Theorem 11. *Let $G \subset X$ be open and bounded. Let $T : X \supset D(T) \rightarrow 2^{X^*}$ be maximal monotone with $0 \in D(T) \cap G$, $0 \in T(0)$ and $0 \notin T(D(T) \cap \partial G)$. Assume that the set $T(D(T) \cap \overline{G})$ is bounded. Assume, further, that $C : D(T) \rightarrow X^*$ is such that CJ_1 is compact, $(J+T)^{-1}$ is compact, and there exists $\alpha > 0$ such that $\|Cx\| \geq \alpha$, $x \in D(T) \cap \partial G$. Then for every $\mu > 0$ there exists $(\lambda_0, x_0) \in (0, \infty) \times (D(T) \cap \partial G)$ such that $Tx_0 - \lambda_0 Cx_0 \ni 0$.*

Proof. We follow, generally, the proof of Theorem 10. The mapping $J_\mu \equiv (I + \mu J^{-1}T)^{-1} : X \rightarrow D(T)$ is continuous. This was shown in the proof of Lemma 3. Thus, its inverse image, $I + \mu J^{-1}T$, maps relatively open subsets of its domain $D(T)$ onto open sets in X . Consequently, the set $(I + \mu J^{-1}T)(D(T) \cap G)$ is open and the set $(I + \mu J^{-1}T)(D(T) \cap \overline{G})$ is closed in X . Let

$$U \equiv (I + \mu J^{-1}T)(D(T) \cap G).$$

Then, as before,

$$\partial U \subset (I + \mu J^{-1}T)(D(T) \cap \partial G)$$

and

$$\overline{U} \subset (I + \mu J^{-1}T)(D(T) \cap \overline{G}).$$

In order to solve (42) with $x \in \partial U$, we observe first that the operator

$$CJ_\mu : \overline{U} \rightarrow X^*$$

is compact. We also observe that if $x \in \partial U$, then

$$J_\mu x \in J_\mu(I + \mu J^{-1}T)(D(T) \cap \partial G) = D(T) \cap \partial G.$$

This equality is implied by the existence of the single-valued inverse J_μ . Thus,

$$\|CJ_\mu x\| \geq \alpha.$$

We set

$$K \equiv \overline{-CJ_\mu \partial U}.$$

We let $\{\lambda_m\}$, $\{\eta_m\}$, $\{x_m\}$ be as in the proof of Theorem 10, but with

$$\{x_m\} \subset \partial U \subset (I + \mu J^{-1}T)(D(T) \cap \partial G).$$

Since J^{-1} is a bounded mapping and the set $D(T) \cap \overline{G}$ is bounded, we have that the set $(I + \mu J^{-1}T)(D(T) \cap \partial G)$ is bounded. This says that $\{x_m\}$ is a bounded sequence. Moreover, $\{T_\mu x_m\}$ is bounded because of the boundedness of the operator T_μ . Continuing as in Theorem 10, we obtain that $d(H_2(0, \cdot), U, 0) = d(I, U, 0) = 1$, which implies that there exists

$$(\lambda_0, x) \in (0, \infty) \times \partial((I + \mu J^{-1}T)(D(T) \cap G)) \subset (0, \infty) \times ((I + \mu J^{-1}T)(D(T) \cap \partial G))$$

such that

$$T_\mu x - \lambda_0 CJ_\mu x = 0.$$

Letting $x_0 = J_\mu x$, we have our conclusion. ■

Open Problem 3. *Is Theorem 10 true with $\|Cx\| \geq \alpha$, $x \in \partial G$, instead of $\|CJ_\mu x\| \geq \alpha$, $x \in \partial G$?*

5. MORE GENERAL APPROXIMATING INCLUSIONS IN HILBERT SPACES

A basic fact in the theory of maximal monotone and m -accretive operators is that the approximating problems that are usually employed are of the “first power” with respect to x . By this, we mean that we add either ϵJx , in the monotone case, or ϵx , in the accretive case, to the sum of the operators involved in order to solve an approximating problem which is easier to handle than the original one. One of the reasons for doing this is that we want to exploit the properties of resolvents $(\lambda T + I)^{-1}$, in the accretive case, and $(\lambda T + J)^{-1}$ in the monotone case. It is therefore natural to ask whether other “small” approximating terms could actually produce more general results than the ones heretofore obtained. We call the attention of the

reader to the fact that we are not discussing here the introduction of K -monotone mappings, which aims, in part, at a unified theory of accretive and monotone operators. We are actually referring to the fact that most of the “inner-product” and “norm” criteria established until now are tied to the fact that the small terms of the respective stabilizing problems are of the first power with respect to x . Attempts with other small perturbing terms were made by various authors in their efforts to show that the solutions of the stabilizing (or approximating) problems converge to the solution of the original problem. A good number of such references are listed in the paper of Alber [1]. Alber himself considered the problem of approximating a solution of an equation of the type

$$Tx = f,$$

where $T : X \rightarrow X^*$ is maximal monotone and $f \in X^*$, by a solution x_ϵ of the stabilizing problem

$$Tx + \epsilon Ux = f.$$

The interesting feature of that paper is that the operator U is given by $Ux \equiv \text{grad}\|x\|^\gamma$, for some $\gamma > 1$. Alber showed that $x_\epsilon \rightarrow x_0$, where x_0 solves the above equation, $Tx_0 = f$, and

$$\|x_0\| = \inf\{\|x\| : Tx = f\}.$$

It is our intention here to introduce the mapping

$$(44) \quad \tilde{J}x \equiv \|x\|^p x, \quad x \in H,$$

in the perturbation theory of nonlinear operators in real Hilbert spaces H . Here, p is a fixed positive constant. This mapping is a continuous, maximal monotone operator satisfying the coercivity condition

$$(45) \quad \langle \tilde{J}x - \tilde{J}y, x - y \rangle \geq \frac{1}{2^p} \|x - y\|^{p+2}, \quad x, y \in H.$$

Actually, the constant $1/2^p$ is best possible (cf. [30]). Consequently, the mapping \tilde{J} is a homeomorphism. Its inverse is given, for $y \neq 0$, by

$$\tilde{J}^{-1}y = \frac{y}{\|y\|^{p/(p+1)}}, \quad y \in H,$$

with

$$\|\tilde{J}^{-1}y\| = \|y\|^{1/(p+1)}.$$

Using this mapping \tilde{J} , we can define the Yosida approximants, \tilde{J}_λ , \tilde{T}_λ , exactly as before, but with \tilde{J} in place of J . Thus, we have

$$\tilde{J}(\tilde{J}_\lambda x - x) + \lambda \tilde{T}_\lambda x = 0, \quad x \in X.$$

Again, $\tilde{J}_\lambda : X \rightarrow D(T)$, $\tilde{T}_\lambda : X \rightarrow X^*$, and it is easy to see that

$$\tilde{J}_\lambda x = (I + \lambda^{1/(p+1)} \tilde{J}^{-1}T)^{-1}, \quad \tilde{T}_\lambda x = (1/\lambda) \tilde{J}(x - \tilde{J}_\lambda x).$$

Both mappings $\tilde{J}_\lambda, \tilde{T}_\lambda$ are continuous and bounded. Moreover, the mapping \tilde{J} is odd and positively homogeneous of degree $p + 1$. In order to obtain a resolvent identity for the resolvents \tilde{J}_λ , we observe that

$$\begin{aligned} \tilde{J}_\lambda^{-1} \tilde{J}_\mu x &= (I + \lambda^{1/(p+1)} \tilde{J}^{-1} T) \tilde{J}_\mu x \\ &= \tilde{J}_\mu x + \lambda^{1/(p+1)} \tilde{J}^{-1} T \tilde{J}_\mu x \\ &\supset \tilde{J}_\mu x + \lambda^{1/(p+1)} \tilde{J}^{-1} \tilde{T}_\mu x \\ &= \tilde{J}_\mu x + \lambda^{1/(p+1)} \tilde{J}^{-1} \left[\frac{1}{\mu} \tilde{J}(x - \tilde{J}_\mu x) \right] \\ &= \tilde{J}_\mu x + \frac{\lambda^{1/(p+1)}}{\mu^{1/(p+1)}} \tilde{J}^{-1} \tilde{J}(x - \tilde{J}_\mu x), \\ &= \tilde{J}_\mu x + \frac{\lambda^{1/(p+1)}}{\mu^{1/(p+1)}} (x - \tilde{J}_\mu x), \end{aligned}$$

i.e.,

$$(46) \quad \tilde{J}_\mu x = \tilde{J}_\lambda \left[\frac{\lambda^{1/(p+1)}}{\mu^{1/(p+1)}} x + \frac{\mu^{1/(p+1)} - \lambda^{1/(p+1)}}{\mu^{1/(p+1)}} \tilde{J}_\mu x \right].$$

This is the desired resolvent identity. It implies, among other things, the compactness of \tilde{J}_μ , for all $\mu > 0$, if we know the compactness of the resolvent for just one $\lambda > 0$.

Another important and useful fact about the mapping \tilde{J} is that the mapping $(\lambda T + \tilde{J})^{-1}$ is continuous for all $\lambda > 0$. Actually, it is Hölder continuous. In fact, from

$$\begin{aligned} \|(\lambda u + \tilde{J}x) - (\lambda v + \tilde{J}y)\| \|x - y\| &\geq \langle (\lambda u + \tilde{J}x) - (\lambda v + \tilde{J}y), x - y \rangle \\ &\geq \frac{1}{2^p} \|x - y\|^{p+2}, \end{aligned}$$

for any $x, y \in D(T)$ and any $u \in Tx, v \in Ty$, we obtain

$$\|(\lambda T + \tilde{J})^{-1} u - (\lambda T + \tilde{J})^{-1} v\| \leq 2^{p/(p+1)} \|u - v\|^{1/(p+1)}, \quad u, v \in H.$$

In order to demonstrate the usefulness of the mapping \tilde{J} in the perturbation theory of monotone operators in H , we shall establish a surjectivity criterion, in the spirit of Theorems 2 and 6, whose proof utilizes the function \tilde{J} .

Theorem 12. *Let $T : H \supset D(T) \rightarrow 2^H$ be maximal monotone and (a) $C : D(T) \rightarrow H$ compact. Assume that there exist $\beta \in \Gamma$ and $Q > 0$ such that: for every $x \in D(T)$ with $\|x\| \geq Q$ and every $v \in Tx$ we have*

$$(47) \quad \langle v + Cx, x \rangle \geq -\beta(\|x\|) \|x\|^{p+2},$$

where p is a nonnegative constant. Then $R(T + C + \epsilon \tilde{J}) = H$, for every $\epsilon \in (0, 1)$. Assume, further, that

$$(48) \quad \liminf_{\substack{\|x\| \rightarrow \infty \\ x \in D(T)}} \frac{|Tx + Cx|}{\|x\|^{p+1}} > 0.$$

Then $\overline{R(T+C)} = H$. Let one of the following conditions hold:

(b) $C : \overline{D(T)} \rightarrow X$ is completely continuous;

(c) instead of (a), $C : \overline{D(T)} \rightarrow H$ is continuous and bounded and $(T + \tilde{J})^{-1}$ is compact.

Then $R(T+C) = H$.

Proof. We consider the stabilizing problem

$$Tx + Cx + \epsilon \tilde{J}x \ni s,$$

where s is a fixed point in H , and the relevant homotopy inclusion

$$t(Tx + Cx + \epsilon \tilde{J}) + (1-t)\tilde{J}x \ni ts.$$

We may assume that $p > 0$. The case $p = 0$ is covered by Theorem 6. We only sketch the proof because of the similarity between the mappings J and \tilde{J} in terms of being homeomorphisms, of type (S_+) , etc. Also, the spaces H and $H^* \equiv H$ are uniformly convex. We note that in the two occasions where the expression $(\|x_n\| - \|x_0\|)^2$ guaranteed the boundedness of the sequence $\{x_n\}$, in the proof of Theorem 6, we must now consider the alternate expression $(1/2^p)\|x_n - x_0\|^{p+2}$, which gives us the same result. The mapping $H(t, u)$ in Theorem 6 is now replaced by the same mapping, but with \tilde{J} in place of J . It is again true that $H(t, u)$ is a homotopy of compact transformations and that the degree $d(I - H(t, \cdot), B, 0)$ is well-defined and equals 1, as in the proof of Theorem 6, because all solutions of $u - H(t, u) = 0$ are bounded by a constant which is independent of the parameter $t \in [0, 1]$. To see the latter, we let $\{t_m\}$, $\{u_m\}$, $\{x_m\}$ be as in the corresponding part of the proof of Theorem 2 with $\|u_m\| \rightarrow \infty$ and $\|x_m\| \rightarrow \infty$. Using our hypotheses, there exist $\beta \in \Gamma$ and $Q > 0$ such that: for every $m \geq 1$ such that $\|x_m\| \geq Q$, we have

$$\begin{aligned} \langle v_m + Cx_m - s, x_m \rangle &\geq -\langle s, x_m \rangle - \beta(\|x_m\|)\|x_m\|^{p+2} \\ &\geq -\frac{\|s\|}{\|x_m\|^{p+1}}\|x_m\|^{p+2} - \beta(\|x_m\|)\|x_m\|^{p+2} \\ &= -\tilde{\beta}(\|x_m\|)\|x_m\|^{p+2}, \end{aligned}$$

for every sequence $\{v_m\}$ of terms $v_m \in Tx_m$, where $\tilde{\beta}$ is an obvious function in Γ . For such an integer m , we obtain

$$\begin{aligned} \langle \epsilon \tilde{J}x_m, x_m \rangle &= \epsilon \|x_m\|^{p+2} \leq [1 - t_m(1 - \epsilon)]\|x_m\|^{p+2} \\ &\leq -t_m \langle v_m + Cx_m - s, x_m \rangle \\ &\leq t_m \tilde{\beta}(\|x_m\|)\|x_m\|^{p+2}. \end{aligned}$$

This inequality holds for all large integers m . It implies

$$\epsilon \leq \tilde{\beta}(\|x_m\|).$$

Since the right-hand side above converges to zero as $m \rightarrow \infty$, we have a contradiction. We omit the rest of the proof, which follows from that of Theorem 2. ■

Naturally, a result like Theorem 12 most probably holds for maximal monotone operators $T : X \supset D(T) \rightarrow 2^{X^*}$. In its proof, we should make use of the mapping U defined above. It is obvious that the mapping \tilde{J} is a homeomorphism in a general Banach space X as well. It is thus natural to ask whether a theorem like Theorem 12 actually holds for m -accretive operators $T : X \supset D(T) \rightarrow 2^X$. This would certainly be true if the mapping \tilde{J} has the right coercivity properties in a general Banach space.

Open Problem 4. *Let X be a real Banach space and p a positive constant. Prove or disprove the following statement. There is a positive constant $a(p)$ such that: for every $x, y \in X$ there exists $j \in J(x - y)$ such that*

$$\langle \|x\|^p x - \|y\|^p y, j \rangle \geq a(p) \| \|x\| - \|y\| \|^{p+2}. \blacksquare$$

Although the use of the mapping \tilde{J} seems rather promising in applications, it might not work with problems involving “ranges of sums.” In fact, the usual applications of this theory to problems of the type $\text{int}(R(T) + R(C)) \subset \text{int}R(T + C)$ make use of the uniform boundedness principle in an argument that works, to the best of our knowledge, only if $p = 0$ in the definition of \tilde{J} (for some recent results, see [17] and [24]).

6. DISCUSSION. POSSIBLE APPLICATIONS

The transition from the solution of the problem $(T + C)x + \epsilon x \ni 0$ to a solution of the problem $(T + C)x \ni 0$ may also be achieved via a coercivity-type condition on the operator T . The same remark applies to the maximal monotone case.

Theorem 1 does have a maximal monotone analogue. We state it for the sake of completeness and for future reference.

Theorem 13. *Let G be a bounded, open and convex subset of X . Let $T : X \supset D(T) \rightarrow 2^{X^*}$ be maximal monotone and (i) $C : \bar{G} \rightarrow X^*$ compact. Assume that $D(T) \cap G \neq \emptyset$ and that $[J - (T + C)](D(T) \cap \partial G) \subset J\bar{G}$. Then $0 \in \overline{R(T + C)}$. Assume that one of the following conditions holds:*

- (a) C is completely continuous;
- (b) instead of (i), let J_1 be compact and $C : \bar{G} \rightarrow X^*$ continuous and bounded. Then $0 \in R(T + C)$.

Proof. As in the proof of Theorem 6, we now consider the homotopy inclusion

$$t(T + C + \epsilon J)x + (1 - t)Jx \ni 0.$$

The argument involving the operator S , in the proof of Theorem 1, is now replaced by an argument involving the operator

$$Sx \equiv J^{-1}[Jx - (T + C)x]$$

which maps ∂G into \bar{G} . In fact, in order to obtain the contradiction involving the point $x_t \in \partial G$, we would now have to argue again with

$$x_t \in \frac{t}{1 + \epsilon t} Sx_t. \blacksquare$$

Theorem 7 is actually a special case of a more general principle which is contained in Theorem 14 below. As we mentioned in [28], in the case of m -accretive operators, such a result, although fruitful, might not be directly applicable in such a degree of generality. We give its statement here because it has an interesting consequence, namely Corollary 8. In Corollary 8 we impose no assumption of boundedness on the operator C . To avoid unnecessary complications, we assume that $0 \in D(T)$ and $0 \in T(0)$.

Theorem 14. *Let $T : X \supset D(T) \rightarrow 2^{X^*}$ be maximal monotone with $(T + J)^{-1}$ compact, $0 \in D(T)$ and $0 \in T(0)$. Let $C : X \supset D(T) \rightarrow X^*$ be such that $C(\lambda T + J)^{-1} : X \rightarrow X^*$ is compact, for some $\lambda \in (0, \infty)$. Assume that there exists an open set $B \subset X$ such that $0 \in B$, $0 \notin (I + \lambda J^{-1}T)^{-1}(\partial B)$, the set $C(I + \lambda J^{-1}T)^{-1}(\overline{B})$ is bounded and*

$$Tx \not\ni 0 \text{ and } \langle u^* + Cx, x \rangle \geq 0,$$

for all $x \in (I + \lambda J^{-1}T)^{-1}(\partial B)$, $u^* \in Tx$. Then $0 \in R(T + C)$.

Proof. The proof follows the steps of the proof of Theorem 7. The main difference is that the set

$$U = (I + \lambda J^{-1}T)^{-1}(D(T) \cap G)$$

has now been replaced by the set B . ■

Corollary 8. *Let $T : X \supset D(T) \rightarrow 2^{X^*}$ be maximal monotone with $(T + J)^{-1}$ compact. Let $C : X \supset D(T) \rightarrow X^*$ be such that $C(I + T)^{-1} : X \rightarrow X^*$ is compact. Assume that there exist a positive constant b and $z \in D(T)$ such that $\|z\| < b$, Tz is bounded and (31) is satisfied for all $x \in D(T)$, $u^* \in Tx$ with $\max\{\|x\|, \|u^*\|\} > b$. Then $0 \in R(T + C)$.*

Proof. The proof follows exactly as in Corollary 1 of [28]. It is therefore omitted. ■

The compactness of the resolvents of a time-dependent m -accretive operator T plays an important role in the compactness of the evolution operator generated by T . For more details on this subject, the reader is referred to the paper [26] and several of the references therein.

We are now going to exhibit various examples of partial differential as well as Nemytskii-type operators which are susceptible to the applicability of the results in this paper. We are looking for Nemytskii-type operators which are either compact or continuous and/or bounded. We are also looking for maximal monotone (m -accretive) operators which have compact resolvents or compact inverses defined on all of X^* (X). In these examples, Ω denotes an open, bounded and connected subset of \mathcal{R}^n with sufficiently smooth boundary. Also, several relations hold a.e. on Ω or on $\partial\Omega$. This will be assumed without further mention. The following example was used by the author in [22].

Example 1. Consider the problem

$$(E1) \quad -\Delta\rho(u(x)) + g(x, u(x)) = p(x), \text{ a.e. } x \in \Omega.$$

We assume the following statements.

- (i) $\rho \in C(\mathcal{R}) \cap C^1(\mathcal{R} \setminus \{0\})$ is nondecreasing and such that $\rho(0) = 0$ and, for some constants $K > 0$, $\alpha \geq 1$,

$$\rho'(t) \geq K|t|^{\alpha-1}, \quad t \in \mathcal{R} \setminus \{0\};$$

(ii) $g : \Omega \times \mathcal{R} \rightarrow \mathcal{R}$ is continuous and such that

$$|g(x, u)| \leq q(x) + q_1|u|,$$

where $q : \Omega \rightarrow \mathcal{R}_+$ is in $L^1(\Omega)$ and q_1 is a positive constant;

(iii) $p \in L^1(\Omega)$ is fixed.

We let $T : L^1(\Omega) \supset D(T) \rightarrow L^1(\Omega)$ be defined by $(Tu)(x) \equiv -\Delta\rho(u(x))$, where $D(T) = \{u \in L^1(\Omega) ; \rho(u) \in W_0^{1,1}(\Omega), \Delta\rho(u) \in L^1(\Omega)\}$. The operator C is defined by $(Cu)(x) = g(x, u(x))$. Bénéilan showed in [3] that T is m -accretive and $\overline{D(T)} = L^1(\Omega)$, while Vrabie showed in [40, Lemma 2.6.2] that T generates a compact semigroup on $\overline{D(T)}$. The latter implies that $(T+I)^{-1}$ is a compact operator on $L^1(\Omega)$. Vainberg's Theorem 19.1 in [39] says that the operator C is continuous and bounded on all of $L^1(\Omega)$. ■

From Massabo and Stuart [33] (cf. also Ding and the author [11]), we have the following example.

Example 2. Consider the problem

$$(E2) \quad -\Delta u(x) + q(x)u(x) + g(x, u(x), \nabla u(x)) = 0, \quad x \in \mathcal{R}^n,$$

where $n > 2$, under the following assumptions.

(1) $q : \mathcal{R}^n \rightarrow \mathcal{R}$ is continuous and

$$0 < \inf_{x \in \mathcal{R}^n} q(x) \leq \sup_{x \in \mathcal{R}^n} q(x) < +\infty.$$

(2) $g : \mathcal{R}^{2n+1} \rightarrow \mathcal{R}$ is continuous and satisfies the following two conditions.

(2a) There exist constants $p \in [1, n/(n-2))$, $c \in \mathcal{R}_+$ and a continuous function $g \in L^2(\mathcal{R}^n)$ such that

$$|g(x, \eta)| \leq g(x) + c\|\eta\|^p, \quad (x, \eta) \in \mathcal{R}^n \times \mathcal{R}^{n+1},$$

where $\|\eta\|$ denotes the Euclidean norm of η .

(2b) For every $\epsilon > 0$ there exist constants $p = p(\epsilon) \in [1, n/(n-2))$ and $l = l(\epsilon) \geq 0$ such that

$$|g(x, 0) - g(x, \eta)| \leq \epsilon\|\eta\|^p$$

for every $x \in \mathcal{R}^n$ with $\|x\| \geq l$ and all $\eta \in \mathcal{R}^{n+1}$.

The operators $T : W^{2,2}(\mathcal{R}^n) \rightarrow L^2(\mathcal{R}^n)$ and $C : W^{2,2}(\mathcal{R}^n) \rightarrow L^2(\mathcal{R}^n)$ are defined by $(Tu)(x) \equiv -\Delta u(x) + q(x)u(x)$ and $(Cu)(x) \equiv g(x, u(x), \nabla u(x))$, respectively. The operator T is self-adjoint, m -accretive, strongly accretive, and such that $T^{-1} : L^2(\mathcal{R}^n) \rightarrow W^{2,2}(\mathcal{R}^n)$ is a Q^{-1} -set-contraction. Here,

$$Q \equiv \inf \sigma_e(T) \in (0, \infty),$$

where $\sigma_e(T)$ is the essential spectrum of T . As Massabo and Stuart have shown in [33], the operator C is compact. ■

From the Hirano and Kalinde paper [19] (cf. also the paper of the author [28]), we have Example 3 below.

Example 3. Consider the nonlinear elliptic boundary value problem

$$(E3) \quad \begin{cases} -\sum_{i=1}^n \frac{\partial}{\partial x_i} a_i \left(\frac{\partial u}{\partial x_i} \right) + g(x, u(x), \nabla u(x)) = p(x), & x \in \Omega, \\ u(x) = 0, & x \in \partial\Omega, \end{cases}$$

where $p \in X = L^2(\Omega)$. We assume that $a_i \in C^1(\mathcal{R})$ and $0 < \inf_{t \in \mathcal{R}} a_i'(t) \leq \sup_{t \in \mathcal{R}} a_i'(t) < \infty$, $i = 1, 2, \dots, n$. We define the operator T by

$$Tu = -\sum_{i=1}^n \frac{\partial}{\partial x_i} a_i \left(\frac{\partial u}{\partial x_i} \right), \quad u \in D(T) \equiv H^2(\Omega) \cap H_0^1(\Omega).$$

The operator T is m -accretive (or maximal monotone) in $X = L^2(\Omega)$. Let $g \in C(\Omega \times \mathcal{R} \times \mathcal{R}^n, \mathcal{R})$ be such that there exist positive constants c_1, c_2 such that

$$(g1) \quad |g(x, s, t)| \leq c_1 |s| + c_2, \quad (x, s, t) \in \Omega \times \mathcal{R} \times \mathcal{R}^n.$$

Define the operator C by

$$(Cu)(x) = g(x, u(x), \nabla u(x)), \quad (u, x) \in D(T) \times \Omega.$$

Then (g1) implies that $C : D(T) \rightarrow X$ is bounded. We also have that the mapping $(I + T)^{-1}$ maps bounded sets of $L^2(\Omega)$ into bounded sets of $H^2(\Omega)$. Recalling that $H^2(\Omega)$ is compactly embedded in $H^1(\Omega)$, we find that $C(I + T)^{-1}$ is a compact mapping on X . ■

Example 4 can be found in Gupta and Hess [18] and Guan and the author [17].

Example 4. Let γ be a maximal monotone graph in \mathcal{R}^2 with $0 \in \gamma(0)$ and let

$$\mathcal{A} = -\sum_{i,j=1}^n \frac{\partial}{\partial x_i} \left(a_{ij} \frac{\partial}{\partial x_j} \right)$$

be a uniformly elliptic operator with coefficients a_{ij} lying in the space $C^1(\bar{\Omega}, \mathcal{R})$ for every $i, j = 1, 2, \dots, n$. With this operator we associate the maximal monotone operator T in $L^2(\Omega)$ defined by

$$D(T) = \left\{ u \in H^2(\Omega) : -\frac{\partial u}{\partial n_\alpha} \in \gamma(u) \text{ on } \partial\Omega \right\},$$

$$Tu = \mathcal{A}u, \quad u \in D(T).$$

Here, $\frac{\partial u}{\partial n_\alpha}$ is the outward normal derivative given by

$$\frac{\partial u}{\partial n_\alpha} = \sum_{i,j=1}^n a_{ij} \frac{\partial u}{\partial x_j} \cos(\bar{n}, x_i),$$

where \bar{n} is the outward normal to $\partial\Omega$.

For the operator C , we assume that

$$(Cu)(x) = g_1(u(x)) + g_2(x, u(x)),$$

where g_1 is as in Gupta and Hess [18], $\beta \in \Gamma$ is continuous and g_2 satisfies the following condition:

- (i) $g_2 : \bar{\Omega} \times \mathcal{R} \rightarrow \mathcal{R}$ is continuous and there exist positive constants c, d such that

$$|g_2(x, t)| \leq c + d|t|, \quad (x, t) \in \Omega \times \mathcal{R};$$

We define the operators $C_1(u), C_2(u)$ by $(C_1u)(x) = g_1(u(x))$ and $(C_2u)(x) = g_2(x, u(x))$, $x \in \Omega, u \in L^2(\Omega)$, respectively.

As in Gupta and Hess [18], the operator $T_1 \equiv T + C_1 : L^2(\Omega) \supset D(T) \rightarrow 2^{L^2(\Omega)}$ is maximal monotone. Also, the operator $C_2 : L^2(\Omega) \rightarrow L^2(\Omega)$ is continuous and bounded, by well-known facts on Nemytskii operators. Gupta and Hess showed in [18] that the operator T_1 is actually boundedly inversely compact, i.e., for every bounded $Q \subset X$ and every bounded $Q^* \subset X^*$ the set $Q \cap T_1^{-1}(Q^*)$ is relatively compact in X . This property was then used in [18] to show that the operator $(T_1 + I)^{-1}$ maps bounded sets into relatively compact sets. However, since this operator is also continuous, we have that $(T_1 + I)^{-1}$ is a compact resolvent. ■

From Fitzpatrick and Petryshyn [12], we have the following example of a compact operator on a Sobolev space.

Example 5. Let N denote the set of natural numbers and, for some $m \in N$, let S_m denote the cardinality of the set

$$\{\alpha = (\alpha_1, \alpha_2, \dots, \alpha_n) \in (N \cup \{0\})^n : |\alpha| \leq m\},$$

where $|\alpha| = \alpha_1 + \alpha_2 + \dots + \alpha_n$. For $\xi \in \mathcal{R}^{S_m}$, we let

$$|\xi| = \left(\sum_{|\alpha| \leq m} |\xi_\alpha|^{p-1} \right)^{1/(p-1)},$$

where $p \in (1, \infty)$ is fixed. Let

- (i) $g(x, \xi) : \Omega \times \mathcal{R}^{S_m} \rightarrow \mathcal{R}$ satisfy the Carathéodory conditions and be such that

$$|g(x, \xi)| \leq k_1|\xi|^{p-1} + |h_1(x)|,$$

where $k_1 \geq 0$ and $h_1 \in L^q(\Omega)$ ($(1/p) + (1/q) = 1$).

Let X be a closed subspace of $W^{m,p}(\Omega)$ which contains the space $W_0^{m,p}(\Omega)$. For every $u, v \in X$, we define the generalized Dirichlet form $b(u, v)$ as follows:

$$b(u, v) = \langle g(\cdot, \xi(u)), v \rangle,$$

where, for $u \in L^p(\Omega), v \in L^q(\Omega)$,

$$\langle u, v \rangle = \int_{\Omega} u(x)v(x)dx.$$

Here, $\xi(u) : \Omega \rightarrow \mathcal{R}^{S_m}$ is defined by

$$\xi(u)(x) = \{D^\alpha u(x) : |\alpha| \leq m \text{ and } x \in \Omega\},$$

where, for $D_i \equiv \partial/\partial x_i$, $i = 1, 2, \dots, n$, $D^\alpha \equiv D_1^{\alpha_1} D_2^{\alpha_2} \dots D_n^{\alpha_n}$. We define the operator $C : X \rightarrow X^*$ by

$$b(u, v) = \langle Cu, v \rangle, \quad u, v \in X.$$

Fitzpatrick and Petryshyn have shown in [12] that C is a compact operator. ■

The author wishes to express his thanks to Professor Igor V. Skrypnik, Ukrainian Academy of Sciences, for his hospitality and interest during the author's stay in Kiev.

REFERENCES

1. Y. I. Alber, *On the solution of nonlinear equations with monotone operators in a Banach space*, Siberian Math. J. **16** (1975), 3-11. MR **51**:6512
2. V. Barbu, *Nonlinear Semigroups and Differential Equations in Banach Spaces*, Noordhoff Int. Publ., Leyden (The Netherlands), 1975. MR **52**:11666
3. P. Bénéilan, *Équations d'Évolution dans un Espace de Banach Quelconque et Applications*, Thèse, Orsay, 1972.
4. H. Brézis, M. G. Crandall and A. Pazy, *Perturbations of nonlinear maximal monotone sets in Banach space*, Comm. Pure Appl Math. **23** (1970), 123-144. MR **41**:2454
5. F. Browder, *Nonlinear Operators and Nonlinear Equations of Evolution in Banach Spaces*, Proc. Symp. Pure Appl. Math., **18**, Part 2, Amer. Math. Soc., Providence, 1976. MR **53**:8982
6. B. D. Calvert and C. P. Gupta, *Nonlinear elliptic boundary value problems in L^p -spaces and sums of ranges of accretive operators*, Nonlinear Anal. **2** (1978), 1-26. MR **80i**:35083
7. I. Cioranescu, *Geometry of Banach Spaces, Duality Mappings and Nonlinear Problems*, Kluwer Acad. Publ., Boston, 1990. MR **91m**:46021
8. K. Deimling, *Zeros of accretive operators*, Manuscr. Math. **13** (1974), 365-374. MR **50**:3030
9. K. Deimling, *Nonlinear Functional Analysis*, Springer-Verlag, New York, 1985. MR **86j**:47001
10. Z. Ding and A. G. Kartsatos, *Nonzero solutions of nonlinear equations involving compact perturbations of accretive operators in Banach spaces*, Nonlinear Anal. **25** (1995), 1333-1342.
11. Z. Ding and A. G. Kartsatos, *P -Regular mappings and alternative results for perturbations of m -accretive operators in Banach spaces*, Topol. Meth. Nonl. Anal. (to appear).
12. P. M. Fitzpatrick and W. V. Petryshyn, *On the nonlinear eigenvalue problem $T(u) = \lambda C(u)$ involving noncompact abstract and differential operators*, Boll. Un. Mat. Ital. B (5) **15** (1978), 80-107. MR **80b**:47083
13. Z. Guan, *Ranges of operators of monotone type in Banach spaces*, J. Math. Anal. Appl. **174** (1993), 256-264. MR **95b**:47068
14. Z. Guan, *Solvability of semilinear equations with compact perturbations of operators of monotone type*, Proc. Amer. Math. Soc. **121** (1994), 93-102. MR **94g**:47080
15. Z. Guan and A. G. Kartsatos, *Solvability of nonlinear equations with coercivity generated by compact perturbations of m -accretive operators in Banach spaces*, Houston J. Math. **21** (1995), 149-188. CMP 95:12
16. Z. Guan and A. G. Kartsatos, *On the eigenvalue problem for perturbations of nonlinear accretive and monotone operators in Banach spaces*, Nonlinear Anal. (to appear).
17. Z. Guan and A. G. Kartsatos, *Ranges of perturbed maximal monotone and m -accretive operators in Banach spaces*, Trans. Amer. Math. Soc. **347** (1995), 2403-2435. MR **95i**:47096
18. C. P. Gupta and P. Hess, *Existence theorems for nonlinear noncoercive operator equations and nonlinear elliptic boundary value problems*, J. Diff. Eq. **22** (1976), 305-313. MR **57**:13600
19. N. Hirano and A. K. Kalinde, *On perturbations of m -accretive operators in Banach spaces*, Proc. Amer. Math. Soc. (to appear).
20. D. R. Kaplan and A. G. Kartsatos, *Ranges of sums and control of nonlinear evolutions with preassigned responses*, J. Optim. Theory Appl. **81** (1994), 121-141. CMP 94:12

21. A. G. Kartsatos, *Zeros of demicontinuous accretive operators in reflexive Banach spaces*, Integral Equations **8** (1985), 175-184. MR **86j**:47078
22. A. G. Kartsatos, *On compact perturbations and compact resolvents of nonlinear m -accretive operators in Banach spaces*, Proc. Amer. Math. Soc. **119** (1993), 1189-1199. MR **94c**:47076
23. A. G. Kartsatos, *Recent results involving compact perturbations and compact resolvents of accretive operators in Banach spaces*, Proceedings of the First World Congress of Nonlinear Analysts, Tampa, Florida, 1992, Walter De Gruyter, New York, (1995), pp. 2197-2222.
24. A. G. Kartsatos, *Sets in the ranges of sums for perturbations of nonlinear m -accretive operators in Banach spaces*, Proc. Amer. Math. Soc. **123** (1995), 145-156. MR **95c**:47072
25. A. G. Kartsatos, *On the construction of methods of lines for functional evolutions in general Banach spaces*, Nonlinear Anal. **25** (1995), 1321-1331.
26. A. G. Kartsatos, *A compact evolution operator generated by a time-dependent m -accretive operator in a general Banach space*, Math. Ann. **302** (1995), 473-487. CMP 95:15
27. A. G. Kartsatos, *Sets in the ranges of nonlinear accretive operators in Banach spaces*, Studia Math. **114** (1995), 261-273. CMP 95:14
28. A. G. Kartsatos, *On the perturbation theory of m -accretive operators in Banach spaces*, Proc. Amer. Math. Soc. (to appear).
29. A. G. Kartsatos and X. Liu, *Nonlinear equations involving compact perturbations of m -accretive operators in Banach spaces*, Nonlinear Anal. **24** (1995), 469-492. CMP 95:7
30. A. G. Kartsatos and R. D. Mabry, *Controlling the space with pre-assigned responses*, J. Optim. Theory Appl. **54** (1987), 517-540. MR **88j**:49024
31. V. Lakshmikantham and S. Leela, *Nonlinear Differential Equations in Abstract Spaces*, Pergamon Press, Oxford, 1981. MR **82i**:34072
32. N. G. Lloyd, *Degree Theory*, Cambridge Univ. Press, Cambridge, 1978. MR **58**:12558
33. I. Massabo and C. A. Stuart, *Positive eigenvectors of k -set contractions*, Nonlinear Anal. **3** (1979), 35-44. MR **80b**:47073
34. M. Nagumo, *Degree of mapping in convex linear topological spaces*, Amer. J. Math. **73** (1951), 497-511. MR **13**:150b
35. D. Pascali and S. Sburlan, *Nonlinear Mappings of Monotone Type*, Sijthoff and Noordhoff Intern. Publ., Alphen aan den Rijn, 1978. MR **80g**:47056
36. S. Reich, *Extension problems for accretive sets in Banach spaces*, J. Funct. Anal. **26** (1977), 378-395. MR **57**:17393
37. E. H. Rothe, *Introduction to Various Aspects of Degree Theory in Banach Spaces*, Math. Surveys and Monographs, **23**, A.M.S., Providence, 1986. MR **87m**:47145
38. I. V. Skrypnik, *Methods for Analysis of Nonlinear Elliptic Boundary Value Problems*, Transl. Math. Monographs, **139**, A.M.S., Providence, 1994. MR **95i**:35109
39. M. M. Vainberg, *Variational Methods for the Study of Nonlinear Operators*, Holden-Day, Inc., San Francisco, 1964. MR **31**:638
40. I. I. Vrabie, *Compactness Methods in Nonlinear Evolutions*, Longman Sci. Tech., London, 1987. MR **90f**:47101
41. E. Zeidler, *Nonlinear Functional Analysis and its Applications*, **II/B**, Springer-Verlag, New York, 1990. MR **91b**:47002

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTH FLORIDA, TAMPA, FLORIDA 33620-5700
E-mail address: hermes@gauss.math.usf.edu