

COMMUTING TOEPLITZ OPERATORS WITH PLURIHARMONIC SYMBOLS

DECHAO ZHENG

ABSTRACT. By making use of \mathcal{M} -harmonic function theory, we characterize commuting Toeplitz operators with bounded pluriharmonic symbols on the Bergman space of the unit ball or on the Hardy space of the unit sphere in n -dimensional complex space.

1. INTRODUCTION

Let $dA(z)$ denote the Lebesgue volume measure on the open unit ball B_n of C^n , normalized so that the measure of B_n equals 1. The Bergman space $L_a^2(B_n)$ is the Hilbert space consisting of holomorphic functions on B_n that are also in $L^2(B_n, dA)$.

For f in $L^\infty(B_n)$, the Toeplitz operator T_f with symbol f is defined by $T_f(h) = P(fh)$, where P is the orthogonal projection from $L^2(B_n, dA)$ onto $L_a^2(B_n, dA)$, called the Bergman projection.

Let $d\sigma$ be the surface area measure on the unit sphere S_n . For $1 \leq p < \infty$, $H^p(S_n)$ is the Banach space of holomorphic functions on B_n with the norm defined by

$$\|f\|_p = \text{Sup}\left\{ \left| \int_{S_n} |f(rz)|^p d\sigma(z) \right|^{1/p}; 0 < r < 1 \right\}.$$

For f in $L^\infty(S_n, d\sigma)$, the Toeplitz operator T_f with symbol f is defined by $T_f(h) = S(fh)$, where S is the orthogonal projection from $L^2(S_n, d\sigma)$ onto the Hardy space $H^2(S_n)$, called the Hardy projection. Here we use the same notation T_f to denote the Toeplitz operator on both $L_a^2(B_n)$ and $H^2(S_n)$.

In this paper we study commuting Toeplitz operators on the Bergman space $L^2(B_n)$ of the unit ball B_n or on the Hardy space $H^2(S_n)$ of the unit sphere S_n , using harmonic function theory on the unit ball B_n .

In the study of one complex variable, a real valued function f on the unit disk is harmonic if $\Delta f = 0$, where

$$\Delta = 4 \frac{\partial^2}{\partial z \partial \bar{z}}$$

is the Laplacian in C . It is well-known that f is harmonic in B_1 if and only if f is the real part of a holomorphic function.

Received by the editors June 30, 1995 and, in revised form, July 15, 1996.

1991 *Mathematics Subject Classification*. Primary 47B35.

Key words and phrases. Toeplitz operator, Bergman space, Hardy space, pluriharmonic function, \mathcal{M} -harmonic function.

Supported in part by the National Science Foundation.

In addition to the usual Laplacian on B_1 , there is also the Laplace-Beltrami operator $\tilde{\Delta}$ which is defined in terms of the Bergman kernel function of B_1 . This operator is given by

$$\tilde{\Delta} = 2(1 - |z|^2)^2 \frac{\partial^2}{\partial z \partial \bar{z}}.$$

A real valued function f is \mathcal{M} -harmonic if $\tilde{\Delta}f = 0$. It is clear that $\tilde{\Delta}f = 0$ if and only if $\Delta f = 0$. So f is harmonic if and only if f is \mathcal{M} -harmonic.

When one considers B_n , $n > 1$, there are three concepts of “harmonic”, namely harmonic, \mathcal{M} -harmonic and pluriharmonic, and as a general rule, these are all different. The three definitions are as follows:

(1) f is harmonic in B_n if $\Delta f = 0$ in B_n , where

$$\Delta = 4 \sum_{j=1}^n \frac{\partial^2}{\partial z_j \partial \bar{z}_j}$$

is the usual Laplacian induced by the Euclidean metric on B_n .

(2) f is pluriharmonic if

$$\frac{\partial^2 f}{\partial z_i \partial \bar{z}_j} = 0$$

for all i, j .

(3) f is \mathcal{M} -harmonic if

$$\tilde{\Delta}f = 0,$$

where $\tilde{\Delta}$ is the Laplace-Beltrami operator induced by the Bergman metric on B_n .

A pluriharmonic function in the unit ball is the sum of a holomorphic function and the conjugate of a holomorphic function. But the mean value property does not completely characterize pluriharmonic functions. Instead it completely characterizes \mathcal{M} -harmonic functions. So function theory is complicated and rich on the unit ball.

Although in this paper we are concerned only with Toeplitz operators with pluriharmonic symbols, we will need to study \mathcal{M} -harmonic functions. By making use of \mathcal{M} -harmonic function theory, we completely characterize commuting Toeplitz operators with bounded pluriharmonic symbols either on the Bergman space $L_a^2(B_n)$ or on the Hardy space $H^2(S_n)$.

This paper is arranged as follows. Section 2 contains statements of the main results. Section 3 contains preliminaries on function theory on B_n . Section 4 contains lemmas which will be used in Section 5. Section 5 contains results on \mathcal{M} -harmonic functions, which are the key in the proof of the main results. Section 6 contains proofs of our main results.

The author would like to thank S. Axler for pointing out Theorem 2.2, and he is grateful to J. Pincus for encouragement.

2. MAIN RESULTS

We state our main results in this section. We postpone the proofs of these theorems until Section 6.

Theorem 2.1. *Suppose that φ and ψ are bounded pluriharmonic functions on the unit ball. Then*

$$T_\varphi T_\psi = T_\psi T_\varphi$$

on the Bergman space $L^2_a(B_n)$ if and only if φ and ψ satisfy one of the following conditions:

- (1) Both φ and ψ are holomorphic on B_n .
- (2) Both $\overline{\varphi}$ and $\overline{\psi}$ are holomorphic on B_n .
- (3) Either φ or ψ is constant on B_n .
- (4) There is a nonzero constant b such that $\varphi - b\psi$ is constant on B_n .

A special case of Theorem 2.1 on the unit disk was first proved by Axler and Gorkin [AxG], and the author [Zh1]. Axler and Čučković proved Theorem 2.1 in the case that $n = 1$. Some partial results on commuting Toeplitz operators were obtained on the Bergman spaces of the unit ball in [CL] and [Zh2].

Theorem 2.1 does not hold if “pluriharmonic” is replaced by “measurable” or even “continuous”. For example, if φ and ψ are bounded measurable radial functions (meaning $\varphi(z) = \varphi(\|z\|)$ and $\psi(z) = \psi(\|z\|)$ for all $z \in B_n$) on the unit ball B_n , then $T_\varphi T_\psi = T_\psi T_\varphi$. This example is based on a example on the unit disk in [AxC].

Theorem 2.2. *Suppose that φ and ψ are bounded pluriharmonic functions on the unit ball. Then*

$$T_\varphi T_\psi = T_\psi T_\varphi$$

on the Hardy space $H^2(S_n)$ if and only if φ and ψ satisfy one of the following conditions:

- (1) Both φ and ψ are holomorphic on B_n .
- (2) Both $\overline{\varphi}$ and $\overline{\psi}$ are holomorphic on B_n .
- (3) Either φ or ψ is constant on B_n .
- (4) There is a nonzero constant b such that $\varphi - b\psi$ is constant on B_n .

Brown and Halmos [BrH] proved Theorem 2.2 when $n = 1$. They examined the matrix products of Toeplitz operators on the Hardy space. On the Bergman space or the Hardy space $H^2(S_n)$, $n > 1$, Toeplitz operators do not have nice matrix representations. Thus function theory plays a large role in this paper.

The difficult part of the proof of Theorems 2.1 and 2.2 is to show that the conditions in these theorems are necessary for $T_\varphi T_\psi = T_\psi T_\varphi$. We reduce it to the following question about \mathcal{M} -harmonic functions:

Question 2.3. If f, g, h , and k are holomorphic functions in B_n , when is $f\overline{g} - h\overline{k}$ \mathcal{M} -harmonic?

This question is very subtle. For a pair of functions f and g (that is, $h = k = 0$), Ahern and Rudin [AhR] answered the question completely. And in [Zh2] a different method was used to study the question for a pair of functions. In this paper, we apply the same method as in [Zh2] to answer Question 2.3 completely when these functions are in $H^{2n}(S_n)$.

3. PRELIMINARIES ON FUNCTION THEORY

This section contains results on the function theory of the unit ball, which will be needed later. A good reference for this material is Rudin’s book [Rud].

For z in B_n , let φ_z be the Möbius transformation from B_n onto B_n defined by

$$\varphi_z(w) = \frac{z - P_z w - (1 - \|z\|^2)^{1/2} Q_z w}{1 - (w, z)},$$

where P_z is the orthogonal projection from C^n onto the subspace of C^n spanned by z , and $Q_z = I - P_z$. φ_z is in the group $Aut(B_n)$ of all biholomorphic maps of B_n onto B_n . In fact, each $\varphi \in Aut(B_n)$ has a unique representation $\varphi = U \circ \varphi_z$ for some $z \in B_n$ and some $n \times n$ unitary matrix U .

For any z in B_n , the pointwise evaluation of functions in $L_a^2(B_n)$ at z is a bounded functional. So there is a function K_z in $L_a^2(B_n)$ such that

$$f(z) = (f, K_z)$$

for all f in $L_a^2(B_n)$. K_z is called the Bergman reproducing kernel. Indeed, we have

$$K_z(w) = \frac{1}{(1 - (w, z))^{n+1}}.$$

For convenience we let k_z denote the normalized Bergman kernel $\frac{K_z(w)}{K_z(z)}$. As $K_z(z) > 0$ for any z in B_n , B_n admits an Hermitian metric, the so-called Bergman metric, given on B_n by

$$g_{ij}(z) = \frac{\partial^2}{\partial z_i \partial \bar{z}_j} \log K_z(z)$$

for z in B_n . The Laplace-Beltrami operator for the Bergman metric of B_n is given by

$$\tilde{\Delta} = c_n(1 - \|z\|^2) \left[\sum_k \frac{\partial^2}{\partial z_k \partial \bar{z}_k} - \sum_{ij} z_i \bar{z}_j \frac{\partial^2}{\partial z_i \partial \bar{z}_j} \right]$$

for some positive constant c_n .

For any f in $C^2(B_n)$, the gradient of f is given by

$$\nabla f(w) = \left(\frac{\partial f}{\partial z_1}(w), \dots, \frac{\partial f}{\partial z_n}(w) \right)^T.$$

A calculation in [Rud] shows that a function u is \mathcal{M} -harmonic on B_n if and only if

$$(3.1) \quad \sum_{i=1}^n \left(\frac{\partial}{\partial z_i} \frac{\partial}{\partial \bar{z}_i} u \right)(z) = \sum_{i,j=1}^n z_i \bar{z}_j \left(\frac{\partial}{\partial z_i} \frac{\partial}{\partial \bar{z}_j} u \right)(z)$$

for all $z \in B_n$. It is clear that all holomorphic functions on B_n are \mathcal{M} -harmonic, as are the pluriharmonic ones. Since each $\varphi \in Aut(B_n)$ is an isometry of B_n in the Bergman metric, the space of all \mathcal{M} -harmonic functions is invariant under the action of $Aut(B_n)$.

The bounded \mathcal{M} -harmonic functions are characterized by the so-called invariant volume mean value property, i.e., a function f bounded on B_n is \mathcal{M} -harmonic if and only if

$$f(\psi(0)) = \int_B f \circ \psi dV$$

for every $\psi \in Aut(B_n)$ ([AFR], [Eng]).

Let \mathcal{U} be the group of all $n \times n$ unitary matrices. Clearly, \mathcal{U} is a compact group. It is well-known that there is a Haar measure $d\mathcal{U}$ on \mathcal{U} . If f is a measurable function on B_n , the identity

$$(3.2) \quad \int_S f d\sigma = \int_{\mathcal{U}} f(\mathcal{U}\eta) d\mathcal{U}$$

holds for any $\eta \in S_n$.

To identify a holomorphic function in the Hardy space $H^2(S_n)$, we need the following formula:

$$(3.3) \quad \int_{S_n} |(\eta)^\alpha|^2 d\sigma(\eta) = \frac{(n-1)!\alpha!}{(n-1+|\alpha|)!}$$

for every multi-index α .

4. SOME LEMMAS

In this section we prove several lemmas which will be used in Section 5. First we need to introduce some concepts and notations.

Let $H(B_n)$ denote the space of holomorphic functions in B_n . $H(B_n)$ is a linear space. We say that functions $\{f_i\}_{i=1}^n \subset H(B_n)$ are linearly independent if there is no nonzero vector $(c_1, \dots, c_n)^T \in C^n$ such that $\sum_{i=1}^n c_i f_i(z) = 0$ for all $z \in B_n$.

For a vector $c = (c_1, \dots, c_n)^T \in C^n$ and $f \in H(B_n)$, define

$$\frac{\partial f}{\partial c} = \sum_{i=1}^n \bar{c}_i \frac{\partial f}{\partial z_i}.$$

A polynomial $p(z)$ in C^n is said to be homogeneous of degree s if $p(tz) = t^s p(z)$. If $f(z)$ is holomorphic in a neighborhood of the origin in C^n , then the power series of f can be written in the grouped form

$$(4.1) \quad f(z) = \sum_{s=0}^{\infty} f_s(z),$$

where $f_s(z)$ is a homogeneous polynomial of degree s . Equation (4.1) is called the homogeneous expansion of f . By Euler's theorem,

$$\sum_{i=1}^n z_i \frac{\partial f_s}{\partial z_i}(z) = s f_s(z).$$

We will often use this fact in this and the next section.

Lemma 4.2. *For any $f \in H(B_n)$, there is an $n \times n$ unitary matrix U such that for some $1 \leq k \leq n$, the functions $\frac{\partial(f \circ U)}{\partial z_1}, \dots, \frac{\partial(f \circ U)}{\partial z_k}$ are linearly independent and $\frac{\partial(f \circ U)}{\partial z_i} = 0$ for all $i > k$.*

Proof. Let $\mathcal{Z} = \{c \in C^n : \frac{\partial f}{\partial c} = 0\}$. \mathcal{Z} is a subspace of C^n . Then there is an orthonormal basis $\{e_1, \dots, e_n\}$ of C^n such that for some $1 \leq k \leq n$, $\{e_{k+1}, \dots, e_n\}$ is also an orthonormal basis of \mathcal{Z} . Let $U = \overline{(e_1, \dots, e_n)}$. Then U is a unitary matrix. On the other hand, by the chain rule, letting $w = Uz$, we have

$$\frac{\partial(f \circ U)}{\partial z_i} = \frac{\partial f}{\partial e_i}(w).$$

Note that for $i > k$, e_i is in \mathcal{Z} , and so

$$\frac{\partial(f \circ U)}{\partial z_i} = 0$$

for all $i > k$.

To complete the proof of this lemma, we need to show that $\frac{\partial(f \circ U)}{\partial z_1}, \dots, \frac{\partial(f \circ U)}{\partial z_k}$ are linearly independent. Suppose that there are constants c_i so that

$$\sum_{i=1}^k c_i \frac{\partial(f \circ U)}{\partial z_i} = 0.$$

Then by the chain rule again, we obtain

$$\sum_{i=1}^k c_i e_i^*(\nabla f)(Uz) = 0,$$

and so $\sum_{i=1}^k \bar{c}_i e_i \in \mathcal{Z}$. But e_i is orthogonal to \mathcal{Z} for $i = 1, \dots, k$. Thus we conclude that $c_i = 0$ for $i = 1, \dots, k$, completing the proof. \square

Lemma 4.3. *Suppose that $c = (c_1, \dots, c_n)$ is in the closure of B_n . Then the equation*

$$(4.4) \quad \sum_{i=1}^n z_i \frac{\partial f}{\partial z_i} = \sum_{i=1}^n c_i \frac{\partial f}{\partial z_i}$$

does not have any non-constant solutions in $H^{2n}(S_n)$.

Proof. Suppose that Equation (4.4) has a solution f in $H^{2n}(S_n)$. We consider two cases, $c \in B_n$ and $c \in S_n$. In each case we will show that f is constant.

Case 1. In this case assume c is in B_n . Making the change of variables $w = z - c$, and using the chain rule, we have

$$\sum_{i=1}^n w_i \frac{\partial f}{\partial w_i}(w + c) = 0.$$

Let $g(w) = f(w + c)$; then the above equation becomes

$$\sum_{i=1}^n w_i \frac{\partial g}{\partial w_i}(w) = 0.$$

Since f is holomorphic on the open neighborhood B_n of c , g is holomorphic on the neighborhood $B_n - c$ of 0 and satisfies

$$\sum_{i=1}^n w_i \frac{\partial g}{\partial w_i}(w) = 0$$

for all $w \in B_n - c$. Clearly, the above equation holds on $rB_n \subset B_n - c$ for some $0 < r < 1$. Applying Euler's theorem to the homogeneous expansion of g yields

$$\sum_{s=0}^{\infty} s g_s(w) = 0$$

for w in rB_n . Then $g_s(w) = 0$ for $s > 0$, and so g is constant on rB_n . Thus we conclude that f is constant on B_n .

Case 2. Now we assume c is in S_n , $\|c\| = 1$. It follows from linear algebra that there is an $n \times n$ unitary matrix U such that

$$c^T(U)^T = (\|c\|, 0, \dots, 0).$$

Use the chain rule again to get

$$(Uz)^T \nabla(f \circ U^*)(Uz) = z^T \nabla f(z)$$

and

$$\begin{aligned} \|c\| \frac{\partial(f \circ U^*)}{\partial z_1}(Uz) &= (\|c\|, 0, \dots, 0) \nabla(f \circ U^*)(Uz) \\ &= (\|c\|, 0, \dots, 0)(U^*)^T \nabla f(z) = c^T \nabla f(z). \end{aligned}$$

Let $g = f \circ U^*$. Clearly, g is in $H^{2n}(S_n)$. Combining Equation (4.4) with the above equations gives

$$(4.5) \quad \sum_{i=1}^n z_i \frac{\partial g}{\partial z_i} = \frac{\partial g}{\partial z_1}.$$

Write g as a homogeneous expansion

$$g(z) = \sum_{t=0}^{\infty} \sum_{s=0}^{\infty} g_{t,s}(z_2, \dots, z_n) z_1^t,$$

where $g_{t,s}(z_2, \dots, z_n)$ is homogeneous of degree s in z_2, \dots, z_n . Putting the above homogeneous expansion into Equation (4.5) and then applying Euler's theorem, we get

$$\sum_{t=0}^{\infty} \sum_{s=0}^{\infty} (s+t) g_{t,s}(z_2, \dots, z_n) z_1^t = \sum_{t=0}^{\infty} \sum_{s=0}^{\infty} t g_{t,s}(z_2, \dots, z_n) z_1^{t-1}.$$

Comparing the coefficients of powers of z_1 in the above equation gives

$$(s+t) g_{t,s}(z_2, \dots, z_n) = (t+1) g_{t+1,s}(z_2, \dots, z_n).$$

This implies the following reduction formula for t :

$$g_{t+1,s}(z_2, \dots, z_n) = \frac{s+t}{t+1} g_{t,s}(z_2, \dots, z_n).$$

Iterating the formula gives

$$g_{t,s}(z_2, \dots, z_n) = \frac{(s+t-1)!}{t!s!} g_{0,s}(z_2, \dots, z_n).$$

Thus g is expressed as

$$\begin{aligned} g(z) &= \sum_{t=0}^{\infty} \sum_{s=0}^{\infty} \frac{(s+t-1)!}{t!s!} g_{0,s}(z_2, \dots, z_n) z_1^t \\ &= \sum_{s=0}^{\infty} \left(\sum_{t=0}^{\infty} \frac{(s+t-1)!}{t!s!} z_1^t \right) g_{0,s}(z_2, \dots, z_n) \\ &= \sum_{s=0}^{\infty} \frac{g_{0,s}(z_2, \dots, z_n)}{(1-z_1)^s}. \end{aligned}$$

For the sake of simplicity, let g_s be $g_{0,s}$. Noting that $g_s(z_2, \dots, z_n)$ is a homogeneous polynomial of degree s , write it as

$$g_s(z_2, \dots, z_n) = \sum_{|\alpha|=s} a_{\alpha s} z^{(0,\alpha)}.$$

So it is easy to verify that

$$\|f\|_2^2 = \sum_{s=0}^{\infty} \sum_{|\alpha|=s} \sum_{t=0}^{\infty} \left(\frac{(s+t-1)!}{t!s!}\right)^2 |a_{\alpha s}|^2 \|z^{(t,\alpha)}\|_2^2.$$

The formula (3.3) gives

$$(4.6) \quad \|g\|_2^2 = \sum_{s=0}^{\infty} \sum_{|\alpha|=s} \sum_{t=0}^{\infty} \left(\frac{(s+t-1)!}{t!s!}\right)^2 \frac{(n-1)!\alpha!t!}{(n-1+s+t)!} |a_{\alpha s}|^2.$$

Since for t very large,

$$\left(\frac{(s+t-1)!}{t!}\right)^2 \frac{\alpha!t!}{(n-1+s+t)!}$$

is asymptotically equivalent to

$$\frac{1}{t^{n-s+1}},$$

in order that the series (4.6) converge, $n-s+1$ must be greater than one. Therefore

$$g(z) = \sum_{s \leq s_0} \frac{f_s(z_2, \dots, z_n)}{(1-z_1)^s}$$

for some $0 \leq s_0 < n$. It is easy to see that the term with the highest degree of $\frac{1}{(1-z_1)}$ in g^n is

$$\frac{g_{s_0}^n(z_2, \dots, z_k)}{(1-z_1)^{s_0 n}}.$$

On the other hand, g^n is in $H^2(S_n)$. Thus $s_0 n < n$, and so $s_0 = 0$. This means that g is constant. Noting that $g = f \circ U^*$, we get that f is constant, completing the proof. \square

Lemma 4.7. *For each j , the equation*

$$(4.8) \quad \frac{\partial}{\partial z_j} \left(\sum_{i=1}^n z_i \frac{\partial f}{\partial z_i} \right) \overline{\left(\sum_{i=1}^n w_i \frac{\partial g}{\partial w_i} \right)} = \frac{\partial}{\partial z_j} \sum_{i=1}^n \frac{\partial f}{\partial z_i} \overline{\frac{\partial g}{\partial w_i}}$$

does not have a solution in $H^{2n}(B_n)$ unless either $\frac{\partial f}{\partial z_j}$ is zero or g is constant for any fixed j .

Proof. Assume that g is not constant. Let f be a solution in $H^{2n}(S_n)$ of Equation (4.8). We will show that $\frac{\partial f}{\partial z_j}$ is constant. Since g is not constant, for some $w \in B_n$,

$$\sum_{i=1}^n w_i \frac{\partial g}{\partial w_i} \neq 0.$$

For a fixed w , solving for $\frac{\partial}{\partial z_j}(\sum_{i=1}^n z_i \frac{\partial f}{\partial z_i})$ in (4.8) gives that for some vector $b \in C^n$,

$$\frac{\partial}{\partial z_j}(\sum_{i=1}^n z_i \frac{\partial f}{\partial z_i}) = b^* \nabla \frac{\partial f}{\partial z_j}.$$

Let $\mathcal{W} = \{c \in C^n : \frac{\partial^2 f}{\partial c \partial z_j} = 0\}$. \mathcal{W} is a subspace of C^n . So there is a unique vector $a \in C^n$ such that

$$(4.9) \quad \frac{\partial}{\partial z_j}(\sum_{i=1}^n z_i \frac{\partial f}{\partial z_i}) = a^* \nabla \frac{\partial f}{\partial z_j}$$

where a is orthogonal to \mathcal{W} . Putting (4.9) and (4.8) together gives

$$a^* \nabla \frac{\partial f}{\partial z_j}(\sum_{i=1}^n w_i \frac{\partial g}{\partial w_i}) = \nabla g(w)^* \nabla \frac{\partial f}{\partial z_j}.$$

Thus

$$a \sum_{i=1}^n w_i \frac{\partial g}{\partial w_i} - \nabla g(w)$$

is in \mathcal{W} for any w in the unit ball B_n . Noting that a is orthogonal to \mathcal{W} , we obtain

$$\|a\|^2 \sum_{i=1}^n w_i \frac{\partial g}{\partial w_i} - a^* \nabla g(w) = 0.$$

By Lemma 4.3, $\frac{a}{\|a\|^2}$ is not in the closure of B_n , and so a is in B_n .

Now we write (4.9) as

$$\frac{\partial}{\partial z_j} \sum_{i=1}^n (z_i - \bar{a}_i) \frac{\partial f}{\partial z_i} = 0.$$

Making the change of variables $w = z - \bar{a}$ gives

$$\frac{\partial}{\partial w_j} \sum_{i=1}^n w_i \frac{\partial f}{\partial w_i}(w + \bar{a}) = 0.$$

Let $F(w) = f(w + \bar{a})$; then the above equation becomes

$$\frac{\partial}{\partial w_j} \sum_{i=1}^n w_i \frac{\partial F}{\partial w_i}(w) = 0.$$

Since f is holomorphic on the neighborhood B_n of \bar{a} , for some $0 < r < 1$, F is holomorphic on the neighborhood $rB_n \subset B_n - \bar{a}$ of 0 and satisfies

$$\frac{\partial}{\partial w_j} \sum_{i=1}^n w_i \frac{\partial F}{\partial w_i}(w) = 0$$

on rB_n . Applying Euler's theorem to the homogeneous expansion of F yields

$$\frac{\partial}{\partial w_j} \sum_{s=0}^{\infty} sF_s(w) = 0.$$

Thus $\frac{\partial F_s}{\partial w_j}(w) = 0$ on rB_n for $s > 0$, and so $\frac{\partial F}{\partial w_j}(w) = 0$. This implies that $\frac{\partial f}{\partial z_j} = 0$, completing the proof. \square

5. \mathcal{M} -HARMONIC FUNCTIONS

In this section we will characterize functions $f, g, h,$ and k in $H^{2n}(S_n)$ such that $f\bar{g} - h\bar{k}$ is \mathcal{M} -harmonic. In [AhR], an explicit description is given of all pairs of holomorphic functions f and g in B_n for which the product $f\bar{g}$ is \mathcal{M} -harmonic. Throughout this section we assume that $f, g, h,$ and k are holomorphic on the unit ball B_n .

Lemma 5.1. *$f\bar{g} - h\bar{k}$ is \mathcal{M} -harmonic if and only if*

$$\begin{aligned}
 & \left(\sum_{i=1}^n z_i \frac{\partial f}{\partial z_i}\right) \overline{\left(\sum_{i=1}^n w_i \frac{\partial g}{\partial w_i}\right)} - \left(\sum_{i=1}^n z_i \frac{\partial h}{\partial z_i}\right) \overline{\left(\sum_{i=1}^n w_i \frac{\partial k}{\partial w_i}\right)} \\
 (5.2) \quad & = \left(\sum_{i=1}^n \frac{\partial f}{\partial z_i} \overline{\frac{\partial g}{\partial w_i}} - \sum_{i=1}^n \frac{\partial h}{\partial z_i} \overline{\frac{\partial k}{\partial w_i}}\right)
 \end{aligned}$$

for z and w in B_n .

Proof. As pointed out in Section 3, u is \mathcal{M} -harmonic if and only if u satisfies Equation (3.1). When $u = f\bar{g} - h\bar{k}$, then Equation (3.1) becomes

$$\begin{aligned}
 & \left(\sum_{i=1}^n z_i \frac{\partial f}{\partial z_i}\right) \overline{\left(\sum_{i=1}^n z_i \frac{\partial g}{\partial z_i}\right)} - \left(\sum_{i=1}^n z_i \frac{\partial h}{\partial z_i}\right) \overline{\left(\sum_{i=1}^n z_i \frac{\partial k}{\partial z_i}\right)} \\
 (5.3) \quad & = \left(\sum_{i=1}^n \frac{\partial f}{\partial z_i} \overline{\frac{\partial g}{\partial z_i}} - \sum_{i=1}^n \frac{\partial h}{\partial z_i} \overline{\frac{\partial k}{\partial z_i}}\right).
 \end{aligned}$$

Hence Equation (5.3) is a special case of (5.2) when $z = w$. So if f, g, h and k satisfy Equation (5.2), then $f\bar{g} - h\bar{k}$ is \mathcal{M} -harmonic.

Conversely, we will show that Equation (5.3) gives Equation (5.2). Let the homogeneous expansions of $f, g, h,$ and k be

$$f(z) = \sum_{s=1}^{\infty} f_s(z), \quad g(z) = \sum_{s=1}^{\infty} g_s(z)$$

and

$$h(z) = \sum_{s=1}^{\infty} h_s(z), \quad k(z) = \sum_{s=1}^{\infty} k_s(z).$$

Euler’s theorem implies that

$$\sum_{i=1}^n z_i \frac{\partial f_s}{\partial z_i} = s f_s(z).$$

So Equation (5.3) becomes

$$\sum_{s,t=1}^{\infty} st(f_s(z)\overline{g_t(z)} - h_s(z)\overline{k_t(z)}) = \sum_{s,t=0}^{\infty} \sum_{i=1}^n \left(\frac{\partial f_s}{\partial z_i} \overline{\frac{\partial g_t}{\partial z_i}} - \frac{\partial h_s}{\partial z_i} \overline{\frac{\partial k_t}{\partial z_i}}\right).$$

Let $z = \xi w$ for ξ in the unit disk and $w \in B_n$. Note that $f_s, g_s, h_s,$ and k_s are homogeneous. An easy computation yields

$$\begin{aligned} & \sum_{s,t=1}^{\infty} st \xi^s \bar{\xi}^t (f_s(w) \overline{g_t(w)} - h_s(w) \overline{k_t(w)}) \\ &= \sum_{s,t=0}^{\infty} \sum_{i=1}^n \xi^{s-1} \bar{\xi}^{t-1} \left(\frac{\partial f_s}{\partial w_i} \overline{\frac{\partial g_t}{\partial w_i}} - \frac{\partial h_s}{\partial w_i} \overline{\frac{\partial k_t}{\partial w_i}} \right). \end{aligned}$$

Comparing the coefficients of $\xi^s \bar{\xi}^t$ in the above equation yields

$$st(f_s(w) \overline{g_t(w)} - h_s(w) \overline{k_t(w)}) = \sum_{i=1}^n \left(\frac{\partial f_{s+1}}{\partial w_i} \overline{\frac{\partial g_{t+1}}{\partial w_i}} - \frac{\partial h_{s+1}}{\partial w_i} \overline{\frac{\partial k_{t+1}}{\partial w_i}} \right)$$

for any s and t . Applying $\overline{D^\alpha}$ both sides of the above equation with respect to w for $|\alpha| = t$ gives

(5.4)

$$st(f_s(w) \overline{D^\alpha g_t(w)} - h_s(w) \overline{D^\alpha k_t(w)}) = \sum_{i=1}^n \left(\frac{\partial f_{s+1}}{\partial w_i} \overline{D^\alpha \frac{\partial g_{t+1}}{\partial w_i}} - \frac{\partial h_{s+1}}{\partial w_i} \overline{D^\alpha \frac{\partial k_{t+1}}{\partial w_i}} \right).$$

But for any homogeneous polynomial P of degree t

$$P(w) = \sum_{|\alpha|=t} \frac{1}{\alpha!} [D^\alpha P] w^\alpha.$$

Multiplying both sides of Equation (5.4) by $\frac{\overline{w^\alpha}}{\alpha!}$, and then taking the sum for α with $|\alpha| = t$, we get

$$st(f_s(w) \overline{g_t(z)} - h_s(w) \overline{k_t(z)}) = \sum_{i=1}^n \left(\frac{\partial f_{s+1}}{\partial w_i} \overline{\frac{\partial g_{t+1}}{\partial z_i}} - \frac{\partial h_{s+1}}{\partial w_i} \overline{\frac{\partial k_{t+1}}{\partial z_i}} \right).$$

Taking the sum with respect to s and t gives Equation (5.2) for any z and w in B_n , completing the proof. \square

Lemma 5.5. *Suppose that neither g nor k is constant and $f\bar{g} - h\bar{k}$ is \mathcal{M} -harmonic. For any vector $a \in C^n$ and $a \neq 0$, if $\frac{\partial f}{\partial a} = 0$, then $\frac{\partial h}{\partial a} = 0$.*

Proof. First we will show that $\frac{\partial f}{\partial z_1} = 0$ implies $\frac{\partial h}{\partial z_1} = 0$. Clearly, $\frac{\partial f}{\partial z_1} = 0$ implies $\frac{\partial^2 f}{\partial z_1 \partial z_j} = 0$ for all j . Differentiating both sides of Equation (5.2) with respect to z_1 , we have

$$\frac{\partial}{\partial z_1} \left(\sum_{i=1}^n z_i \frac{\partial h}{\partial z_i} \right) \overline{\left(\sum_{i=1}^n w_i \frac{\partial k}{\partial w_i} \right)} = \frac{\partial}{\partial z_1} \sum_{i=1}^n \frac{\partial h}{\partial z_i} \overline{\frac{\partial k}{\partial w_i}}.$$

Noting that k is not constant, by Lemma 4.7, we obtain $\frac{\partial h}{\partial z_1} = 0$.

Next we reduce the general case to the above case. Assume that $\frac{\partial f}{\partial a} = 0$ for some vector $a \neq 0$. For $e = (1, 0, \dots, 0)^* \in C^n$, let U be a unitary matrix such that $\overline{U}e = a$. By the chain rule, we have

$$\frac{\partial(f \circ U)}{\partial z_1} = \frac{\partial f}{\partial a}(Uz).$$

On the other hand, the space of \mathcal{M} -harmonic functions is invariant under the action of unitary matrices. So $f\bar{g} - h\bar{k}$ implies $f \circ U \bar{g} \circ \bar{U} - h \circ U \bar{k} \circ \bar{U}$ is \mathcal{M} -harmonic. As shown above, $\frac{\partial(f \circ U)}{\partial z_1} = 0$ implies $\frac{\partial(h \circ U)}{\partial z_1} = 0$.

By the chain rule again, we have

$$\frac{\partial(h \circ U)}{\partial z_1} = \frac{\partial h}{\partial a}(Uz),$$

which means $\frac{\partial h}{\partial a} = 0$. This completes the proof. \square

If we assume that none of f , g , h and k is constant and $f\bar{g} - h\bar{k}$ is \mathcal{M} -harmonic, Lemma 5.5 tells us that any linear relation among derivatives of f gives the same linear relation among derivatives of h . This fact is also valid for g and k . We will often use this fact in the proof of Theorem 5.6. The following theorem is the main result in this section. It is the key to the proof of Theorems 2.1 and 2.2.

Theorem 5.6. *Suppose that f , g , h , and k are in $H^{2n}(B_n)$. Then $f\bar{g} - h\bar{k}$ is \mathcal{M} -harmonic on the unit ball if and only if f , g , h , and k satisfy one of the following conditions:*

- (1) Both f and h are constants.
- (2) Both f and k are constants.
- (3) Both g and k are constants.
- (4) Both g and h are constants.
- (5) There is a nonzero constant b such that $bg - k$ and $f - \bar{b}h$ are constants.

Proof. If f , g , h and k satisfy one of the conditions (1)–(5) in this theorem, one easily sees that they satisfy Equation (5.2). Hence, by Lemma 5.1, $f\bar{g} - h\bar{k}$ is \mathcal{M} -harmonic.

Conversely, assume that $f\bar{g} - h\bar{k}$ is \mathcal{M} -harmonic. By Lemma 5.1, it is sufficient to show that any solutions in $H^{2n}(B_n)$ of Equation (5.2) must satisfy one of the conditions in this theorem.

If one of f , g , h and k is constant, then (5.2) becomes an equation involving a pair of functions. In this case, it follows from a result in [AhR] and [Zh2] that one of the conditions (1)–(4) holds.

Now we may assume that none of f , g , h and k is constant. We will show that these functions satisfy condition (5) in this theorem. Note that Equation (5.2) is a linear equation in $\frac{\partial f}{\partial z_1}, \dots, \frac{\partial f}{\partial z_n}, \sum_{i=1}^n z_i \frac{\partial f}{\partial z_i}, \frac{\partial k}{\partial z_1}, \dots, \frac{\partial k}{\partial z_n}$, and $\sum_{i=1}^n z_i \frac{\partial k}{\partial z_i}$ for a fixed $w \in B_n$, and a linear equation in $\frac{\partial g}{\partial w_1}, \dots, \frac{\partial g}{\partial w_n}, \sum_{i=1}^n w_i \frac{\partial g}{\partial w_i}, \frac{\partial h}{\partial w_1}, \dots, \frac{\partial h}{\partial w_n}$, and $\sum_{i=1}^n w_i \frac{\partial h}{\partial w_i}$ for a fixed $z \in B_n$. We will use these equations to get condition (5).

First reduce the general case to a special case in which the derivatives of f , g , h and k satisfy the following conditions.

- (a) $\frac{\partial f}{\partial z_1}, \dots, \frac{\partial f}{\partial z_r}$ are linearly independent.
- (b) $\frac{\partial g}{\partial w_1}, \dots, \frac{\partial g}{\partial w_r}$ are linearly independent.
- (c) $\frac{\partial h}{\partial z_1}, \dots, \frac{\partial h}{\partial z_r}$ are linearly independent.
- (d) $\frac{\partial k}{\partial w_1}, \dots, \frac{\partial k}{\partial w_r}$ are linearly independent.
- (e) Either $\frac{\partial f}{\partial z_i} = \frac{\partial h}{\partial z_i} = 0$ or $\frac{\partial g}{\partial w_i} = \frac{\partial k}{\partial w_i} = 0$ for $i > r > 0$.

By Lemma 4.2, there is a unitary matrix V such that $\frac{\partial(f \circ V)}{\partial z_1}, \dots, \frac{\partial(f \circ V)}{\partial z_{r_1}}$ are linearly independent and $\frac{\partial(f \circ V)}{\partial z_i} = 0$ for $i > r_1$. By Lemma 5.5, $\frac{\partial(h \circ V)}{\partial z_1}, \dots, \frac{\partial(h \circ V)}{\partial z_{r_1}}$ are linearly independent and $\frac{\partial(h \circ V)}{\partial z_i} = 0$ for $i > r_1$.

If $\frac{\partial(g \circ V)}{\partial w_1}, \dots, \frac{\partial(g \circ V)}{\partial w_{r_1}}$ are linearly independent, then by Lemma 5.5, we have that $\frac{\partial(k \circ V)}{\partial w_1}, \dots, \frac{\partial(k \circ V)}{\partial w_{r_1}}$ are also linearly independent. So $(f \circ V), (g \circ V), (h \circ V)$ and $(k \circ V)$ satisfy conditions (a)–(e) and $f \circ V \overline{g \circ V} - h \circ V \overline{k \circ V}$ satisfies Equation (5.2).

If $\frac{\partial(g \circ V)}{\partial w_1}, \dots, \frac{\partial(g \circ V)}{\partial w_{r_1}}$ are linearly dependent, let

$$L = \{a \in C^{r_1} : \sum_{i=1}^{r_1} a_i \frac{\partial(g \circ V)}{\partial w_i} = 0\}.$$

Using the same argument as in the proof of Lemma 4.2, we see that there is an r_1 by r_1 unitary matrix U_2 such that $\{\frac{\partial(g \circ V \circ U_1)}{\partial w_1}, \dots, \frac{\partial(g \circ V \circ U_1)}{\partial w_r}\}$ are linearly independent and $\frac{\partial(g \circ V \circ U_1)}{\partial w_i} = 0$ for $i = r + 1$ to r_1 , where $U_1 = U_2 \oplus I$, and for some integer $0 \leq r \leq r_1$. Also by the chain rule, one easily checks that $\frac{\partial(f \circ V \circ U_1)}{\partial z_1}, \dots, \frac{\partial(f \circ V \circ U_1)}{\partial z_r}$ are linearly independent. Then $f \circ V \circ U_1$ and $g \circ V \circ U_1$ satisfy conditions (a), (b) and (e) as required in the above special case. By Lemma 5.5, $h \circ V \circ U_1$ and $k \circ V \circ U_1$ also satisfy conditions (c), (d) and (e). Moreover, $f \circ V \circ U_1 \overline{g \circ V \circ U_1} - h \circ V \circ U_1 \overline{k \circ V \circ U_1}$ satisfies Equation (5.2), as desired.

Next we turn to solving Equation (5.2) with conditions (a)–(e). Under conditions (a)–(e), Equation (5.2) becomes

$$\begin{aligned} & \left(\sum_{i=1}^n z_i \frac{\partial f}{\partial z_i}\right) \overline{\left(\sum_{i=1}^n w_i \frac{\partial g}{\partial w_i}\right)} - \left(\sum_{i=1}^n z_i \frac{\partial h}{\partial z_i}\right) \overline{\left(\sum_{i=1}^n w_i \frac{\partial k}{\partial w_i}\right)} \\ (5.7) \quad & = \sum_{i=1}^r \left(\frac{\partial f}{\partial z_i} \overline{\frac{\partial g}{\partial w_i}} - \frac{\partial h}{\partial z_i} \overline{\frac{\partial k}{\partial w_i}}\right). \end{aligned}$$

The main idea will be to use the linear independence of the derivatives of f, g, h and k to solve Equation (5.7). Let

$$\mathcal{H} = \{c \in C^r : c^* \nabla_r f \in \text{span}\left\{\frac{\partial h}{\partial z_1}, \dots, \frac{\partial h}{\partial z_r}\right\}\},$$

where $\nabla_r f = \left(\frac{\partial f}{\partial z_1}, \dots, \frac{\partial f}{\partial z_r}\right)^T$. We consider three cases.

Case 1. $\dim \mathcal{H} = r$. In this case there is a matrix A such that

$$\nabla_r f = A \nabla_r h.$$

Jordan theory in linear algebra implies that there is a unitary matrix U such that

$$U^T A (U^T)^* = J_1 \oplus \dots \oplus J_k,$$

where J_i is an s_i by s_i matrix

$$\begin{pmatrix} c_i & 0 & 0 & \dots & 0 & 0 \\ 1 & c_i & 0 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & 0 \\ 0 & 0 & 0 & \dots & 1 & c_i \end{pmatrix}$$

and $\sum_{i=1}^k s_i = r$. For the sake of simplicity, we still use f, g, h , and k to denote respectively $f \circ U, g \circ U, h \circ U$ and $k \circ U$. Using the above Jordan form of A and the chain rule, we have the following equations:

$$\frac{\partial f}{\partial z_1} = c_1 \frac{\partial h}{\partial z_1}, \frac{\partial f}{\partial z_2} = c_1 \frac{\partial h}{\partial z_2} + \frac{\partial h}{\partial z_1}, \dots$$

Taking the derivative of both sides of Equation (5.7) with respect to z_1 , along with the first equation above, we obtain the following equation:

$$\left(\frac{\partial}{\partial z_1} \sum_{i=1}^n z_i \frac{\partial h}{\partial z_i}\right) \overline{\left(\sum_{i=1}^n w_i \frac{\partial(c_1 g - l)}{\partial w_i}\right)} = \frac{\partial}{\partial z_1} \sum_{i=1}^n \frac{\partial h}{\partial z_i} \overline{\frac{\partial(c_1 g - k)}{\partial w_i}}.$$

By Lemma 4.7, $c_1 g - k$ is constant, so $\frac{\partial k}{\partial w_i} = c_1 \frac{\partial g}{\partial w_i}$. Also c_1 is zero because k is not constant. Replacing $\frac{\partial k}{\partial w_i}$ by $c_1 \frac{\partial g}{\partial w_i}$ in Equation (5.7), we have

$$\frac{\partial}{\partial z_1} \left(\sum_{i=1}^n z_i \frac{\partial(f - \overline{c_1} h)}{\partial z_i}\right) \overline{\left(\sum_{i=1}^n w_i \frac{\partial g}{\partial w_i}\right)} = \frac{\partial}{\partial z_1} \sum_{i=1}^n \frac{\partial(f - \overline{c_1} h)}{\partial z_i} \overline{\frac{\partial g}{\partial w_i}}.$$

By Lemma 4.7 again, $f - \overline{c_1} h$ is constant. So f, g, h , and k satisfy condition (5).

Case 2. $\dim \mathcal{H} < r - 1$. In this case, using the same argument as in the proof of Lemma 4.2, we may assume that $\{\frac{\partial f}{\partial z_1}, \dots, \frac{\partial f}{\partial z_j}\}$ are linearly independent of $\{\frac{\partial h}{\partial z_1}, \dots, \frac{\partial h}{\partial z_r}\}$ and $\{\frac{\partial f}{\partial z_{j+1}}, \dots, \frac{\partial f}{\partial z_r}\}$ are in the span $\{\frac{\partial h}{\partial z_1}, \dots, \frac{\partial h}{\partial z_r}\}$ for some $j \geq 2$. Then by Equation (5.7), there are vectors a, a', c, c' and a matrix B such that

$$\sum_{i=1}^n z_i \frac{\partial f}{\partial z_i} = a^* \nabla_j f + c^* \nabla_r h,$$

$$\sum_{i=1}^n z_i \frac{\partial h}{\partial z_i} = (a')^* \nabla_j f + (c')^* \nabla_r h,$$

and

$$\left(\frac{\partial f}{\partial z_{j+1}}, \dots, \frac{\partial f}{\partial z_n}\right)^T = (\nabla_r h)^T B^*.$$

Combining Equation (5.7) with the above equations, and using the fact that $\frac{\partial f}{\partial z_1}, \dots, \frac{\partial f}{\partial z_j}, \frac{\partial h}{\partial z_1}, \dots, \frac{\partial h}{\partial z_r}$ are linearly independent, we obtain

$$a \sum_{i=1}^n w_i \frac{\partial g}{\partial w_i} - a' \sum_{i=1}^n w_i \frac{\partial k}{\partial w_i} = \nabla_j g$$

and

$$c \sum_{i=1}^n w_i \frac{\partial g}{\partial w_i} - c' \sum_{i=1}^n w_i \frac{\partial k}{\partial w_i} = B \left(\frac{\partial g}{\partial w_{j+1}}, \dots, \frac{\partial g}{\partial w_n}\right)^T - \nabla_r k.$$

Since $\frac{\partial g}{\partial w_1}, \dots, \frac{\partial g}{\partial w_j}$ are linearly independent, the first system of equations above gives that $\sum_{i=1}^n w_i \frac{\partial g}{\partial w_i}$ and $\sum_{i=1}^n w_i \frac{\partial k}{\partial w_i}$ are in the span $\{\frac{\partial g}{\partial w_1}, \dots, \frac{\partial g}{\partial w_r}\}$. The second system of equations above gives that $\{\frac{\partial k}{\partial w_1}, \dots, \frac{\partial k}{\partial w_r}\}$ are in the span $\{\frac{\partial g}{\partial w_1}, \dots, \frac{\partial g}{\partial w_r}\}$. So this case is reduced to Case 1 if we replace f and h by g and k respectively.

Case 3. $\dim \mathcal{H} = r - 1$. Using the same argument as in the proof of Lemma 4.2, we may assume that $\frac{\partial f}{\partial z_1}, \frac{\partial h}{\partial z_1}, \dots, \frac{\partial h}{\partial z_r}$ are linearly independent, and there is an $(r - 1) \times r$ matrix A such that

$$(5.8) \quad \left(\frac{\partial f}{\partial z_2}, \dots, \frac{\partial f}{\partial z_r}\right)^T = A \nabla_r h.$$

If there is a constant b such that $bg - k$ is constant, then $\frac{\partial k}{\partial w_i} = b \frac{\partial g}{\partial w_i}$. For each $1 \leq j \leq n$, differentiating (5.7) with respect to z_j gives

$$\frac{\partial}{\partial z_j} \left(\sum_{i=1}^n z_i \frac{\partial(f - bh)}{\partial z_i}\right) \overline{\left(\sum_{i=1}^n w_i \frac{\partial g}{\partial w_i}\right)} = \frac{\partial}{\partial z_j} \sum_{i=1}^n \frac{\partial(f - bh)}{\partial z_i} \overline{\frac{\partial g}{\partial w_i}}.$$

By Lemma 4.7, $\frac{\partial(f - bh)}{\partial z_j} = 0$ for j and so $f - bh$ is constant. Thus f, g, h and k satisfy condition (5).

Suppose that there is no constant b such that $bg - k$ is constant. We will derive a contradiction. By Equation (5.7), there are constants c_1, c_2 and vectors a, a' such that

$$(5.9) \quad \sum_{i=1}^n z_i \frac{\partial f}{\partial z_i} = c_1 \frac{\partial f}{\partial z_1} + a^* \nabla_r h$$

and

$$(5.10) \quad \sum_{i=1}^n z_i \frac{\partial h}{\partial z_i} = c_2 \frac{\partial f}{\partial z_1} + (a')^* \nabla_r h.$$

Extend the holomorphic function f on B_n to a holomorphic function \tilde{f} on B_{n+1} by

$$\tilde{f}(z_1, \dots, z_n, z_{n+1}) = f(z_1, \dots, z_n).$$

By Lemma 5.1, $\tilde{f}\tilde{g} - \tilde{h}\tilde{k}$ is \mathcal{M} -harmonic on B_{n+1} .

For c in the unit disk, define

$$\varphi_c(z_1, \dots, z_n, z_{n+1}) = \left(\frac{z_1 s}{1 - \bar{c}z_{n+1}}, \dots, \frac{z_n s}{1 - \bar{c}z_{n+1}}, \frac{c - z_{n+1}}{1 - \bar{c}z_{n+1}}\right),$$

where $s = (1 - |c|^2)^{1/2}$; then $\varphi_c \in \text{Aut}(B_{n+1})$. Note that the space of \mathcal{M} -harmonic functions is invariant under the action of $\text{Aut}(B_{n+1})$. Then

$$\tilde{f} \circ \varphi_c \overline{\tilde{g} \circ \varphi_c} - \tilde{h} \circ \varphi_c \overline{\tilde{k} \circ \varphi_c}$$

is \mathcal{M} -harmonic on B_{n+1} .

By the chain rule, we have

$$(5.11) \quad \frac{\partial(\tilde{f} \circ \varphi_c)}{\partial z_{n+1}} = \sum_{i=1}^n \frac{\bar{c} s z_i}{(1 - \bar{c}z_{n+1})^2} \frac{\partial \tilde{f}}{\partial z_i} \circ \varphi_c,$$

and

$$(5.12) \quad \frac{\partial(\tilde{f} \circ \varphi_c)}{\partial z_j} = \frac{s}{(1 - \bar{c}z_{n+1})} \frac{\partial \tilde{f}}{\partial z_j} \circ \varphi_c,$$

for $j < n + 1$. Thus (5.11) gives

$$(5.13) \quad \frac{\partial(\tilde{f} \circ \varphi_c)}{\partial z_{n+1}} = \frac{c}{(1 - \bar{c}z_{n+1})} \sum_{i=1}^n (\varphi_c)_i \frac{\partial \tilde{f}}{\partial z_i} ((\varphi_c(z))_i, \dots, (\varphi_c(z))_n).$$

Since $\sum_{i=1}^n z_i \frac{\partial f}{\partial z_i} = c_1 \frac{\partial f}{\partial z_1} + a^* \nabla_r h$, we have

$$\begin{aligned} & \sum_{i=1}^n (\varphi_c)_i \frac{\partial f}{\partial z_i} ((\varphi_c(z))_i, \dots, (\varphi_c(z))_n) \\ &= c_1 \frac{\partial \tilde{f}}{\partial z_1} \circ \varphi_c + a^* \nabla_r h((\varphi_c(z))_i, \dots, (\varphi_c(z))_n). \end{aligned}$$

Then (5.13) combined with (5.12) gives

$$(5.14) \quad \frac{\partial(\tilde{f} \circ \varphi_c)}{\partial z_{n+1}} = \frac{\bar{c}}{s} c_1 \frac{\partial(\tilde{f} \circ \varphi_c)}{\partial z_1} + \frac{\bar{c}}{s} a^* \nabla_r(\tilde{h} \circ \varphi_c).$$

Similarly we can show that

$$(5.15) \quad \frac{\partial(\tilde{h} \circ \varphi_c)}{\partial z_{n+1}} = \frac{\bar{c}}{s} c_2 \frac{\partial(\tilde{f} \circ \varphi_c)}{\partial z_1} + \frac{\bar{c}}{s} (a')^* \nabla_r(\tilde{h} \circ \varphi_c).$$

(i) If $c_2 \neq 0$, let $F = \tilde{f} \circ \varphi_c$, $G = \tilde{g} \circ \varphi_c$, $H = \tilde{h} \circ \varphi_c$ and $K = \tilde{k} \circ \varphi_c$; then (5.14) and (5.15) give that $\frac{\partial F}{\partial z_i}$ is in the span $\{\frac{\partial H}{\partial z_1}, \dots, \frac{\partial H}{\partial z_r}, \frac{\partial H}{\partial z_{n+1}}\}$ for $i = 1, \dots, r$, and $n + 1$. Since $F\bar{G} - H\bar{K}$ is \mathcal{M} -harmonic on B_{n+1} , by Lemma 5.1, we have

$$\begin{aligned} & \left[\sum_{i=1}^{n+1} z_i \frac{\partial F}{\partial z_i} \right] \overline{\left[\sum_{i=1}^{n+1} w_i \frac{\partial G}{\partial w_i} \right]} - \left[\sum_{i=1}^{n+1} z_i \frac{\partial H}{\partial z_i} \right] \overline{\left[\sum_{i=1}^{n+1} w_i \frac{\partial K}{\partial w_i} \right]} \\ &= \sum_{i=1}^r \left(\frac{\partial F}{\partial z_i} \overline{\frac{\partial G}{\partial w_i}} - \frac{\partial H}{\partial z_i} \overline{\frac{\partial K}{\partial w_i}} \right) - \left(\frac{\partial F}{\partial z_{n+1}} \overline{\frac{\partial G}{\partial w_{n+1}}} - \frac{\partial H}{\partial z_{n+1}} \overline{\frac{\partial K}{\partial w_{n+1}}} \right). \end{aligned}$$

because

$$\sum_{i=r+1}^n \left(\frac{\partial F}{\partial z_i} \overline{\frac{\partial G}{\partial w_i}} - \frac{\partial H}{\partial z_i} \overline{\frac{\partial K}{\partial w_i}} \right) = 0.$$

Thus this case is reduced to Case 1. Hence F, G, H and K satisfy condition (5) of this theorem. So do f, g, h , and k . But this contradicts that there is no constant b such that $bg - k$ is constant.

(ii) If $c_2 = 0$, then (5.10) becomes

$$\sum_{i=1}^n z_i \frac{\partial h}{\partial z_i} = (a')^* \nabla_r h.$$

Replacing $\sum_{i=1}^n z_i \frac{\partial h}{\partial z_i}$ in Equation (5.7) by the right hand side of the above equation gives

$$(5.16) \quad \bar{c}_1 \sum_{i=1}^n w_i \frac{\partial g}{\partial w_i} = \frac{\partial g}{\partial w_1}$$

and

$$(5.17) \quad a \sum_{i=1}^n w_i \frac{\partial g}{\partial w_i} - a' \sum_{i=1}^n w_i \frac{\partial k}{\partial w_i} = A^* \left(\frac{\partial g}{\partial w_2}, \dots, \frac{\partial g}{\partial w_r} \right)^T - \nabla_r k.$$

These equations come from linear independence of $\frac{\partial f}{\partial z_1}, \frac{\partial h}{\partial z_1}, \dots, \frac{\partial h}{\partial z_r}$.

By the chain rule, (5.16) gives

$$\frac{\partial(\tilde{g} \circ \varphi_c)}{\partial w_{n+1}} = \frac{\bar{c}}{s} \frac{\partial(\tilde{g} \circ \varphi_c)}{\partial w_1},$$

which means

$$\overline{c_1} \frac{\partial G}{\partial w_{n+1}} = \frac{\overline{c}}{s} \frac{\partial G}{\partial w_1},$$

Since neither F nor H is constant, Lemma 5.5 implies

$$\overline{c_1} \frac{\partial K}{\partial w_{n+1}} = \frac{\overline{c}}{s} \frac{\partial K}{\partial w_1},$$

and so

$$\overline{c_1} \frac{\partial(\tilde{k} \circ \varphi_c)}{\partial w_{n+1}} = \frac{\overline{c}}{s} \frac{\partial(\tilde{k} \circ \varphi_c)}{\partial w_1}.$$

Then by the chain rule again, we obtain

$$\overline{c_1} \sum_{i=1}^n w_i \frac{\partial k}{\partial w_i} = \frac{\partial k}{\partial w_1}.$$

Putting the above equation, (5.15) and (5.17) together gives

$$\left(\frac{a}{\overline{c_1}}, -A^*\right) \nabla_r g = \begin{pmatrix} \frac{a'_1}{\overline{c_1}} - 1 & 0 & 0 & \cdots & 0 \\ \frac{a'_2}{\overline{c_1}} & -1 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \frac{a'_r}{\overline{c_1}} & 0 & 0 & \cdots & -1 \end{pmatrix} \nabla_r k.$$

If $\frac{a'_1}{\overline{c_1}} - 1 \neq 0$, the system of equations above gives that $\frac{\partial k}{\partial w_i}$ is in the space spanned by $\{\frac{\partial g}{\partial w_1}, \dots, \frac{\partial g}{\partial w_r}\}$, so this case is reduced to Case 1 for g and k . This contradicts that there is no constant b such that $bg - k$ is constant.

If $\frac{a'_1}{\overline{c_1}} - 1 = 0$, the system of equations above gives

$$\frac{a_1}{\overline{c_1}} \frac{\partial g}{\partial w_1} - \frac{a_{21}}{\overline{c_1}} \frac{\partial g}{\partial w_2} - \cdots - \frac{a_{r1}}{\overline{c_1}} \frac{\partial g}{\partial w_r} = 0.$$

Note that $\{\frac{\partial g}{\partial w_1}, \dots, \frac{\partial g}{\partial w_r}\}$ are linearly independent. Then $a_{i1} = 0$ for $i = 2, \dots, r$. By (5.8), we have

$$\left(\frac{\partial f}{\partial z_2}, \dots, \frac{\partial f}{\partial z_r}\right)^T = (0, A_{(r-1) \times (r-1)}) \nabla_r h.$$

So

$$\left(\frac{\partial f}{\partial z_2}, \dots, \frac{\partial f}{\partial z_r}\right)^T = A_{(r-1) \times (r-1)} \left(\frac{\partial h}{\partial z_2}, \dots, \frac{\partial h}{\partial z_r}\right)^T.$$

Without loss of generality, we may assume that

$$A = J_1 \oplus \cdots \oplus J_{k_1}$$

where J_i is the s_i by s_i matrix

$$\begin{pmatrix} c_i & 0 & 0 & \cdots & 0 & 0 \\ 1 & c_i & 0 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & 0 \\ 0 & 0 & 0 & \cdots & 1 & c_i \end{pmatrix}$$

and $\sum_{i=1}^{k_1} s_i = r - 1$. Using the Jordan form of A , we get

$$\frac{\partial f}{\partial z_2} = c_1 \frac{\partial h}{\partial z_2}.$$

Taking the derivative of both sides of Equation (5.7) with respect to z_2 , and using the above equation, we have

$$\frac{\partial}{\partial z_2} \left(\sum_{i=1}^n z_i \frac{\partial h}{\partial z_i} \right) \overline{\left(\sum_{i=1}^r w_i \frac{\partial(\overline{c_1 g - k})}{\partial w_i} \right)} = \frac{\partial}{\partial z_2} \sum_{i=1}^n \frac{\partial h}{\partial z_i} \overline{\frac{\partial(\overline{c_1 g - k})}{\partial w_i}}.$$

By Lemma 4.7, $\overline{c_1 g - k}$ is constant. This contradicts that there is no constant b such that $bg - k$ is constant. This completes the proof. \square

To end this section, we restate Theorem 5.6 as follows.

Corollary 5.18. *Suppose that f, g, h and k are in $H^{2n}(B_n)$. Then $f\overline{g} - h\overline{k}$ is \mathcal{M} -harmonic if and only if it is pluriharmonic.*

6. PROOFS OF MAIN RESULTS

In this section we will present the proof of the main results. We begin with a simple fact about bounded pluriharmonic functions on B_n .

Proposition 6.1. *If f is a bounded pluriharmonic function on B_n , then there are functions f_1 and f_2 in $H^p(S_n)$ for all $p > 1$ such that $f = f_1 + \overline{f_2}$.*

Proof. Without loss of generality, we consider just the case when f is real-valued. Because f is pluriharmonic, there is a holomorphic function f_1 on B_n such that $f = f_1 + \overline{f_1}$. We need to show that f_1 is in $H^p(S_n)$ for all $p > 1$. It is easy to check that $S(f) = f_1 + \overline{f_1}(0)$, where S is the Hardy projection. It is well-known that the Hardy projection S is bounded on $L^p(S_n)$ for any $p > 1$. Note that f is in $L^\infty(B_n)$. Then f_1 is in $H^p(S_n)$ for any $p > 1$. This completes the proof. \square

The following theorem easily follows from recent result of Ahern, Flores and Rudin [AFR] for $n < 12$. The result here is valid for all n . It was also obtained by a different method in [CL]. Our method extends the method in [AxC] and [Zh1] directly.

Theorem 6.2. *Suppose that f, g, h and k are in $H^2(S_n)$ with the property*

$$(6.3) \quad \int_{B_n} (f\overline{g} - h\overline{k}) \circ \psi(w) |k_z(w)|^2 dA(w) = (f\overline{g} - h\overline{k}) \circ \psi(z)$$

for all $\psi \in \text{Aut}(B_n)$. Then $(f\overline{g} - h\overline{k})$ is \mathcal{M} -harmonic on the unit ball B_n .

Proof. Let $d\mathcal{U}$ be the Haar measure on the group \mathcal{U} of $n \times n$ unitary matrices. Define

$$G(z) = \int_{\mathcal{U}} (f\overline{g} - h\overline{k})(Uz) d\mathcal{U}$$

for z in B_n . For any f in $H^2(S)$, write f as a power series

$$f(z) = \sum_{\alpha} a_{\alpha}(f) z^{\alpha}$$

with

$$\sum_{\alpha} |a_{\alpha}(f)|^2 I(\alpha) < +\infty,$$

where $I(\alpha) = \|z^{\alpha}\|_{H^2(S)}^2$.

As \mathcal{U} is a subgroup of $Aut(B_n)$, for any $U \in \mathcal{U}$, letting $\psi = U$ in (6.3), we have

$$\int_{B_n} (f\bar{g} - h\bar{k}) \circ U(w) |k_z(w)|^2 dA(w) = (f\bar{g} - h\bar{k}) \circ U(z).$$

Integrating both sides of the above equation with respect to U over \mathcal{U} gives

$$(6.4) \quad \int_B G(w) |k_z(w)|^2 dA(w) = G(z).$$

By the formula (3.2), we have

$$\begin{aligned} G(z) &= \int_S (f\bar{g} - h\bar{k})(\|z\|\xi) d\sigma(\xi) \\ &= \sum_{\alpha} (a_{\alpha}(f)\overline{a_{\alpha}(g)} - a_{\alpha}(h)\overline{a_{\alpha}(k)}) I(\alpha) \|z^{\alpha}\|^{2|\alpha|}, \end{aligned}$$

where $|\alpha| = \sum_{i=1}^n \alpha_i$.

Since f, g, h and k are in $H^2(S_n)$, $G(z)$ is continuous on \bar{B}_n . The mean value theorem tells us that $\tilde{\Delta}G = 0$. In addition, $G(z)$ is constant on S_n . It follows from the maximal principle that $G(z)$ is constant on B_n . Thus

$$(6.5) \quad \sum_{|\alpha|=1} [a_{\alpha}(f)\overline{a_{\alpha}(g)} - a_{\alpha}(h)\overline{a_{\alpha}(k)}] = 0.$$

For convenience we let $a_i(f)$ denote $a_{(0, \dots, 0, 1, 0, \dots, 0)}(f)$. Then Equation (6.5) becomes

$$\sum_i [a_i(f)\overline{a_i(g)} - a_i(h)\overline{a_i(k)}] = 0.$$

By the chain rule, $a_i(f) = \frac{\partial f}{\partial z_i}(0)$. Letting $a_i(f)(w) = \frac{\partial(f \circ \varphi_w)}{\partial z_i}(0)$, similarly we get that

$$(6.6) \quad \sum_i [a_i(f)(z)\overline{a_i(g)(z)} - a_i(h)(z)\overline{a_i(k)(z)}] = 0.$$

The above equation comes from the fact that the space of \mathcal{M} -harmonic functions is invariant under the action of $Aut(B_n)$.

Calculating $a_i(f)(w)$ by the chain rule, we obtain

$$(6.7) \quad a(f)(z) = s(1-s)P_{\bar{z}}\nabla_z f - s\nabla_z f,$$

where $s = (1 - \|z\|^2)^{1/2}$ and

$$a(f)(z) = \begin{pmatrix} a_1(f)(z) \\ \vdots \\ a_n(f)(z) \end{pmatrix}.$$

Replacing $a_i(f)(z)$ in (6.6) by the right hand side of (6.7) gives

$$\begin{aligned} 0 &= [s(1-s)P_{\bar{z}}\nabla_z g - s\nabla_z g]^* [s(1-s)P_{\bar{z}}\nabla_z f - s\nabla_z f] \\ &\quad - [s(1-s)P_{\bar{z}}\nabla_z k - s\nabla_z k]^* [s(1-s)P_{\bar{z}}\nabla_z h - s\nabla_z h]. \end{aligned}$$

Simplifying the above equation yields

$$\begin{aligned}
 & -s^2(1-s^2)\langle \bar{z}, \nabla g \rangle \langle \nabla f, \bar{z} \rangle + s^2(1-s^2)\langle \nabla f, \nabla g \rangle \\
 & = -s^2(1-s^2)\langle \bar{z}, \nabla k \rangle \langle \nabla h, \bar{z} \rangle + s^2(1-s^2)\langle \nabla h, \nabla k \rangle.
 \end{aligned}$$

Multiplying both sides of the above equation by $\frac{1}{s^2(1-s^2)}$, we obtain

$$\begin{aligned}
 & \langle \bar{z}, \nabla g \rangle \langle \nabla f, \bar{z} \rangle - \langle \bar{z}, \nabla k \rangle \langle \nabla h, \bar{z} \rangle \\
 & = \langle \nabla f, \nabla g \rangle - \langle \nabla h, \nabla k \rangle.
 \end{aligned}$$

We conclude that

$$\begin{aligned}
 & \left(\sum_{j=1}^n z_j \frac{\partial f}{\partial z_j}\right) \left(\sum_{i=1}^n \overline{z_i} \frac{\partial g}{\partial z_i}\right) - \left(\sum_{j=1}^n z_j \frac{\partial h}{\partial z_j}\right) \left(\sum_i \overline{z_i} \frac{\partial k}{\partial z_i}\right) \\
 & = \sum_{j=1}^n \frac{\partial f}{\partial z_j} \overline{\frac{\partial g}{\partial z_j}} - \sum_{j=1}^n \frac{\partial h}{\partial z_j} \overline{\frac{\partial k}{\partial z_j}},
 \end{aligned}$$

which is Equation (3.1). So $f\bar{g} - h\bar{k}$ is \mathcal{M} -harmonic. This completes the proof. \square

Lemma 6.8. *Let $f, g, h,$ and k be holomorphic on the unit ball B_n . Suppose that $\varphi = f + \bar{k}$ and $\psi = h + \bar{g}$. Then φ and ψ satisfy one of conditions (1)–(4) in Theorem 2.1 if and only if $f, g, h,$ and k satisfy one of conditions (1)–(5) in Theorem 5.6.*

The proof of Lemma 6.8 is easy. We leave it as an exercise to readers.

For each $z \in B_n$, define an operator U_z on the Bergman space $L_a^2(B_n)$ by

$$U_z u = u \circ \varphi_z k_z,$$

where φ_z is the Möbius transformation and k_z is the normalized reproducing kernel. If V is unitary matrix in C^n , define an operator U_V on $L_a^2(B_n)$ by $U_V u = u \circ V$. A simple computation shows that U_z and U_V are a unitary operators from $L_a^2(B_n)$ onto $L_a^2(B_n)$. So for each element τ in $Aut(B_n)$, we can associate a unitary operator U_τ on $L_a^2(B_n)$ with τ by $U_\tau = U_V U_z$ if $\tau = U \circ \varphi_z$. Immediately we see that

$$U_\tau T_f U_\tau^* = T_{f \circ \tau}.$$

Hence

$$(6.9) \quad U_\tau (T_\varphi T_\psi - T_\psi T_\varphi) U_\tau^* = (T_{\varphi \circ \tau} T_{\psi \circ \tau} - T_{\psi \circ \tau} T_{\varphi \circ \tau}).$$

We need the following lemma in the proof of Theorem 2.1.

Lemma 6.10. *Let $f, g, h,$ and k be holomorphic on the unit ball B_n . Suppose that $\varphi = f + \bar{k}$ and $\psi = h + \bar{g}$ are bounded on B_n . Then*

$$\begin{aligned}
 & \langle (T_{\varphi \circ \tau} T_{\psi \circ \tau} - T_{\psi \circ \tau} T_{\varphi \circ \tau}) k_z, k_z \rangle \\
 (6.11) \quad & = - \int_{B_n} (f \circ \tau(w) \overline{g \circ \tau(w)} - h \circ \tau(w) \overline{k \circ \tau(w)}) |k_z(w)|^2 dA(w) \\
 & \quad + (f \circ \tau(z) \overline{g \circ \tau(z)} - h \circ \tau(z) \overline{k \circ \tau(z)})
 \end{aligned}$$

for each $\tau \in Aut(B_n)$.

Proof. Suppose that

$$(6.12) \quad \varphi = f + \bar{k} \text{ and } \psi = h + \bar{g}.$$

Since

$$T_\varphi 1 = P(\varphi) = P(f + \bar{k}) = f + \overline{k(0)},$$

we have

$$\begin{aligned} \langle T_\psi T_\varphi 1, 1 \rangle &= \langle T_\varphi 1, T_\psi 1 \rangle = \langle T_\varphi 1, \bar{\psi} \rangle = \langle (f + \overline{k(0)}), \bar{\psi} \rangle = \langle (\varphi + \overline{k(0)} - \bar{k}), \bar{\psi} \rangle \\ &= \langle \varphi, \bar{\psi} \rangle + \langle (\overline{k(0)} - \bar{k}), \bar{\psi} \rangle = \langle \psi \varphi, 1 \rangle + \overline{k(0)} h(0) - \langle \bar{k} h, 1 \rangle. \end{aligned}$$

Interchanging φ and ψ in the above equation gives

$$\langle T_\varphi T_\psi 1, 1 \rangle = \langle \varphi \psi, 1 \rangle + \overline{g(0)} f(0) - \langle \bar{g} f, 1 \rangle.$$

Hence we obtain

$$(6.13) \quad \langle (T_\varphi T_\psi - T_\psi T_\varphi) 1, 1 \rangle = \overline{g(0)} f(0) - \overline{k(0)} h(0) - [\langle \bar{g} f, 1 \rangle - \langle \bar{k} h, 1 \rangle].$$

For each $\tau \in \text{Aut}(B_n)$, composing both sides of Equation (6.12) with τ expresses each of the functions $\varphi \circ \tau$ and $\psi \circ \tau$ as the sum of a holomorphic function and the conjugate of a holomorphic function:

$$\varphi \circ \tau = f \circ \tau + \overline{k \circ \tau} \text{ and } \psi \circ \tau = h \circ \tau + \overline{g \circ \tau}.$$

Note that $\tau \circ \varphi_z$ is still in $\text{Aut}(B_n)$ for each $\tau \in \text{Aut}(B_n)$ and $z \in B_n$. Replacing each function in (6.13) by its composition with $\tau \circ \varphi_z$ gives

$$\begin{aligned} &\langle (T_{\varphi \circ \tau \circ \varphi_z} T_{\psi \circ \tau \circ \varphi_z} - T_{\psi \circ \tau \circ \varphi_z} T_{\varphi \circ \tau \circ \varphi_z}) 1, 1 \rangle \\ &= - \int_{B_n} (f \circ \tau \circ \varphi_z(w) \overline{g \circ \tau \circ \varphi_z(w)} - h \circ \tau \circ \varphi_z(w) \overline{k \circ \tau \circ \varphi_z(w)}) dA(w) \\ (6.14) \quad &+ (f \circ \tau \circ \varphi_z(0) \overline{g \circ \tau \circ \varphi_z(0)} - h \circ \tau \circ \varphi_z(0) \overline{k \circ \tau \circ \varphi_z(0)}). \end{aligned}$$

On the other hand, $U_z 1 = k_z$. By (6.9), we have

$$(6.15) \quad \langle (T_{\varphi \circ \tau \circ \varphi_z} T_{\psi \circ \tau \circ \varphi_z} - T_{\psi \circ \tau \circ \varphi_z} T_{\varphi \circ \tau \circ \varphi_z}) 1, 1 \rangle = \langle (T_{\varphi \circ \tau} T_{\psi \circ \tau} - T_{\psi \circ \tau} T_{\varphi \circ \tau}) k_z, k_z \rangle.$$

Making the change of variables $\lambda = \varphi_z(w)$, we get

$$\begin{aligned} (6.16) \quad &\int_{B_n} (f \circ \tau \circ \varphi_z(w) \overline{g \circ \tau \circ \varphi_z(w)} - h \circ \tau \circ \varphi_z(w) \overline{k \circ \tau \circ \varphi_z(w)}) dA(w) \\ &= \int_{B_n} (f \circ \tau(\lambda) \overline{g \circ \tau(\lambda)} - h \circ \tau(\lambda) \overline{k \circ \tau(\lambda)}) |k_z(\lambda)|^2 dA(\lambda). \end{aligned}$$

Combining (6.14) with (6.15) and (6.16), we obtain (6.11), as desired. This completes the proof. \square

Now we are ready to prove Theorems 2.1 and 2.2.

Proof of Theorem 2.1. First we prove that if φ and ψ satisfy one of the conditions in Theorem 2.1, then T_φ commutes with T_ψ .

If φ is holomorphic on B_n , then T_φ is the operator on the Bergman space $L_a^2(B_n)$ of multiplication by φ . So if φ and ψ satisfy condition (1), then T_φ commutes with T_ψ . Since the adjoint of T_φ on the Bergman space is $T_{\bar{\varphi}}$, T_φ commutes with T_ψ if φ and ψ satisfy condition (2). If φ and ψ satisfy condition (3), either T_φ or T_ψ is a scalar operator, so T_φ commutes with T_ψ . If φ and ψ satisfy condition (4), let $c = b\psi - \varphi$ be a constant. Thus $T_\varphi = bT_\psi + c$. So T_φ commutes with T_ψ .

To prove the other direction of Theorem 2.1, suppose that T_φ commutes with T_ψ on $L_a^2(B_n)$. We need to show that φ and ψ satisfy one of the conditions in Theorem 2.1. By Proposition 6.1, there are four functions f, g, h , and k in $H^{2n}(S_n)$ such that

$$\varphi = f + \bar{k} \text{ and } \psi = h + \bar{g}.$$

By Lemma 6.8, it suffices to show that f, g, h , and k satisfy one of the conditions in Theorem 5.6. To do this, we need to show that these functions satisfy Equation (6.3) in Theorem 6.2.

Since T_φ commutes with T_ψ , $T_\varphi T_\psi - T_\psi T_\varphi = 0$. Then by (6.9), we have

$$T_{\varphi \circ \tau} T_{\psi \circ \tau} - T_{\psi \circ \tau} T_{\varphi \circ \tau} = 0$$

for each $\tau \in \text{Aut}(B_n)$. Thus it follows from Lemma 6.10 that

$$\begin{aligned} & \int_{B_n} (f \circ \tau(w) \overline{g \circ \tau(w)} - h \circ \tau(w) \overline{k \circ \tau(w)}) |k_z(w)|^2 dA(w) \\ &= f \circ \tau(z) \overline{g \circ \tau(z)} - h \circ \tau(z) \overline{k \circ \tau(z)} \end{aligned}$$

for each $\tau \in \text{Aut}(B_n)$. So f, g, h and k satisfy the condition of Theorem 6.2, as desired. This completes the proof. \square

Proof of Theorem 2.2. The proof of the “if” part is similar to that in the proof of Theorem 2.1, so it is omitted. It remains to prove the “only if” part.

Suppose that T_φ commutes with T_ψ on $L_a^2(B_n)$. We need to show that φ and ψ satisfy one of the conditions in Theorem 2.2. By Proposition 6.1, there are four functions f, g, h , and k in $H^{2n}(S_n)$ such that

$$\varphi = f + \bar{k} \text{ and } \psi = h + \bar{g}.$$

Every function in $L^p(S_n)$ extends to an \mathcal{M} -harmonic function on the unit ball, via the Poisson integral. In fact, functions in the Hardy space correspond to holomorphic functions on the unit ball, and the evaluation at z for any z in B_n

$$f \rightarrow f(z)$$

is a bounded linear functional, so there is a function S_z in $H^2(S_n)$ such that $f(z) = \langle f, S_z \rangle$. It is easy to check that, for any function u in $H^2(S_n)$,

$$T_{\bar{u}}(S_z) = \overline{u(z)} S_z.$$

Then for $z \in B_n$,

$$\begin{aligned}
 (6.17) \quad & \langle T_\psi T_\varphi \frac{S_z}{\|S_z\|}, \frac{S_z}{\|S_z\|} \rangle = \langle T_\varphi \frac{S_z}{\|S_z\|}, T_{\bar{\psi}} \frac{S_z}{\|S_z\|} \rangle = \langle (f + \overline{k(z)}) \frac{S_z}{\|S_z\|}, \bar{\psi} \frac{S_z}{\|S_z\|} \rangle \\
 & = \langle (\varphi + \overline{k(z)} - \bar{k}) \frac{S_z}{\|S_z\|}, \bar{\psi} \frac{S_z}{\|S_z\|} \rangle = \langle \varphi \frac{S_z}{\|S_z\|}, \bar{\psi} \frac{S_z}{\|S_z\|} \rangle + \langle (\overline{k(z)} - \bar{k}) \frac{S_z}{\|S_z\|}, \bar{\psi} \frac{S_z}{\|S_z\|} \rangle \\
 & = \langle \psi \varphi \frac{S_z}{\|S_z\|}, \frac{S_z}{\|S_z\|} \rangle + \overline{k(z)} h(z) - \bar{k} h \frac{S_z}{\|S_z\|}, \frac{S_z}{\|S_z\|} \rangle.
 \end{aligned}$$

Interchanging φ and ψ in the above equations gives

$$(6.18) \quad \langle T_\varphi T_\psi \frac{S_z}{\|S_z\|}, \frac{S_z}{\|S_z\|} \rangle = \langle \varphi \psi \frac{S_z}{\|S_z\|}, \frac{S_z}{\|S_z\|} \rangle + \overline{g(z)} f(z) - \langle \bar{g} f \frac{S_z}{\|S_z\|}, \frac{S_z}{\|S_z\|} \rangle.$$

Combining (6.17) with (6.18) gives

$$\begin{aligned}
 & \langle (T_\varphi T_\psi - T_\psi T_\varphi) \frac{S_z}{\|S_z\|}, \frac{S_z}{\|S_z\|} \rangle \\
 & = - \int_{S_n} (f(w) \overline{g(w)} - h(w) \overline{k(w)}) \frac{|S_z(w)|^2}{\|S_z\|^2} d\sigma(w) + (f(z) \overline{g(z)} - h(z) \overline{k(z)}).
 \end{aligned}$$

Because T_φ commutes with T_ψ , we have

$$\int_{S_n} (f(w) \overline{g(w)} - h(w) \overline{k(w)}) \frac{|S_z(w)|^2}{\|S_z\|^2} d\sigma(w) = f(z) \overline{g(z)} - h(z) \overline{k(z)}.$$

Thus the right hand side of the above equation defines an \mathcal{M} -harmonic function on the unit ball, and so the left hand side must be \mathcal{M} -harmonic on B_n . This implies that $f\bar{g} - h\bar{k}$ is \mathcal{M} -harmonic. By Theorem 5.6, f , g , h , and k satisfy one of the conditions in Theorem 5.6. It follows from Lemma 6.8 that φ and ψ satisfy one of the conditions of Theorem 2.2. This completes the proof. \square

REFERENCES

[AFR] P. Ahern, M. Flores and W. Rudin, An invariant volume-mean-value property, *J. Functional Analysis* 111 (1993), 380-397. MR **94b**:31002
 [AhR] P. Ahern and W. Rudin, \mathcal{M} -harmonic products, *Indag. Math.* 2 (1991), 141-147. MR **92g**:32013
 [AxC] S. Axler and Z. Čučković, Commuting Toeplitz operators with harmonic symbols, *Integral Equations Operator Theory* 14 (1991), 1-12. MR **92f**:47018
 [AxG] S. Axler and P. Gorkin, Algebras on the disk and doubly commuting operators, *Trans. Amer. Math. Soc.* 309 (1988), 711-723. MR **90a**:46133
 [BrH] A. Brown and P.R. Halmos, Algebraic properties of Toeplitz operators, *J. Reine Angew. Math.* 213 (1964), 89-102. MR **28**:3350
 [CL] B. Choe and Y. Lee, Pluriharmonic symbols of commuting Toeplitz operators, *Illinois J. Math.* 37 (1993), 424-436. MR **94i**:47041
 [Eng] M. Engliš, Functions invariant under the Berezin transform, *J. Functional Analysis* 121 (1994), 233-254. MR **95h**:31001
 [Rud] W. Rudin, *Function theory on the unit ball of C^n* , Grundlehren 241, Springer-Verlag, Berlin-Heidelberg-New York, 1980. MR **82i**:32002

- [Zh1] D. Zheng, Hankel operators and Toeplitz operator on the Bergman space, *J. Functional Analysis* 83 (1989), 98-120. MR **91b**:47057
- [Zh2] D. Zheng, Semi-commutators of Toeplitz operators on the Bergman space, *Integral Equations Operator Theory* 25 (1996), 347-372. MR **97e**:47039

DEPARTMENT OF MATHEMATICS, MICHIGAN STATE UNIVERSITY, EAST LANSING, MICHIGAN
48824

Current address: Department of Mathematics, Vanderbilt University, Nashville, Tennessee
37240

E-mail address: zheng@math.vanderbilt.edu