

INFINITE TYPE HOMEOMORPHISMS OF THE CIRCLE AND CONVERGENCE OF FOURIER SERIES

ANTÔNIO ZUMPARO

ABSTRACT. We consider the problem of convergence of Fourier series when we make a change of variable. Under a certain reasonable hypothesis, we give a necessary and sufficient condition for a homeomorphism of the circle to transform absolutely convergent Fourier series into uniformly convergent Fourier series.

1. INTRODUCTION

Let $A(T)$ be the space of all continuous functions of the circle T with absolutely convergent Fourier series, and let $U(T)$ be the space of all continuous functions on the circle T that have uniformly convergent Fourier series. If φ is a homeomorphism of the circle T , we say that φ transports $A(T)$ to $U(T)$ if $f \circ \varphi \in U(T)$ for all $f \in A(T)$.

A great deal of attention has been given to the following question: which homeomorphisms of the circle T transport $A(T)$ to $U(T)$? We say that a homeomorphism φ of the circle is of finite type if there is an integer v with $v \geq 3$ such that: φ is of class C^v and $|\varphi''(t)| + \dots + |\varphi^{(v)}(t)| \neq 0$ for all $t \in \mathbb{R}$. In 1974, R. Kaufman showed that if φ is of finite type, then it transports $A(T)$ to $U(T)$. If φ is an analytic homeomorphism of the circle, it is easy to see that either it is of finite type or $\varphi''(t) = 0$ for all t . In the first case it transports $A(T)$ to $U(T)$ by the result of Kaufman; in the second case φ transports $A(T)$ to $A(T)$, see [1]. But, not every C^∞ homeomorphism of the circle transports $A(T)$ to $U(T)$, see [1] for the counterexample. We will be interested, therefore, in the case when φ is of class C^∞ , but it may have a flat point, i.e. a point $t \in \mathbb{R}$ such that $\varphi^{(k)}(t) = 0$ for all $k = 2, 3, \dots$ (note that a C^∞ homeomorphism with no flat point is of finite type). In a certain class of homeomorphisms of the circle, we will give a necessary and sufficient condition to transport $A(T)$ to $U(T)$.

2. STATEMENT OF THE RESULTS

The theorem of R. Kaufman that was mentioned above states that a homeomorphism φ of the circle transports $A(T)$ to $U(T)$ if it is of finite type. The proof of this fact can be found in [1] and it is based on Lemma 2 below, but we suggest an alternative proof based on a result due to Stein and Wainger. See [2]. We state that result here as a lemma:

Received by the editors November 6, 1994 and, in revised form, October 23, 1996.

1991 *Mathematics Subject Classification*. Primary 42A20; Secondary 26A45.

Key words and phrases. Oscillatory integral, Fourier analysis, Fourier series, homeomorphism of the circle.

Lemma 1 (Stein-Wainger). *Let $p(t)$ be a real polynomial of degree d . Then,*

$$\left| \int_{-r}^r e^{ip(t)} \frac{1}{t} dt \right| \leq 6(2^{d+1}) - 2d - 10 \text{ for all } r > 0.$$

They proved that lemma in a more general form in 1965 and the proof was published five years later [2].

The second lemma we state below was proved by R. Kaufman in 1974, see [1].

Lemma 2 (R. Kaufman). *Let f be a function of class C^k on the interval $[-r, r]$ with $k \geq 2$. Suppose $1 \leq |f^{(k)}(t)| \leq b$ for all $t \in [-r, r]$. Then,*

$$\left| \int_{-r}^r e^{if(t)} \frac{1}{t} dt \right| \leq C(k, b)$$

where $C(k, b)$ is a constant that depends only on k and b .

The fact is that Lemma 2 can be proved from Lemma 1 in a quite simple way. The proof of the second lemma given in [1] does not use Lemma 1 at all. Also, it is not difficult to see that Lemma 1 follows from Lemma 2 if we consider $d \geq 2$. So, they are indeed equivalent results.

The primary tool in dealing with oscillatory integrals as those in the lemmas is the Van der Corput lemma, see [3].

Our purpose is to deal with homeomorphisms of the circle not necessarily of finite type. To be precise, let φ be a homeomorphism of a circle of class C^v , with $v \geq 3$, such that

$$(1) \quad |\varphi''(t)| + |\varphi'''(t)| + \dots + |\varphi^{(v)}(t)| \neq 0 \text{ for all } t \neq 0, t \in [-\pi, \pi].$$

Suppose that there is a neighbourhood of zero, say $(-r, r)$ with $r < \pi$, such that:

$$(2) \quad \varphi \text{ is an odd function on } (-r, r).$$

$$(3) \quad \varphi'(0) = 0 \text{ and } \varphi''(t) > 0 \text{ for all } t \in (0, r].$$

Also, assume that there is a constant θ , with $0 < \theta < 1$ so that

$$(4) \quad \varphi'((1 - \theta)a + \theta b) \leq \frac{1}{2}[\varphi'(a) + \varphi'(b)]$$

for all $a, b \in [0, r)$ with $a \leq b$.

We describe (4) by saying that φ' has uniform bounded doubling time. In particular φ' has what is called bounded doubling time, for if we put $a = 0$ above, we have that $\varphi'(\theta t) \leq \frac{1}{2}\varphi'(t)$ for all $t \in [0, r)$. Also, condition (4) with $\theta = \frac{1}{2}$ means that φ' is a convex function on $[0, r)$; but it does not imply convexity of φ' (see [5] for an example of a homeomorphism φ of the circle that satisfies all the four conditions and φ' is not convex in any interval of the form $(0, r)$, $r > 0$).

We shall prove the following result:

Theorem 1. *Let φ be a homeomorphism of the circle of class C^v , $v \geq 3$. Suppose that φ satisfies (1), (2), (3) and (4). Then, φ transports $A(T)$ to $U(T)$ if and only if there are constants ζ, M and λ with $0 < \zeta < 1$, $M > 0$ and $\lambda > 1$ such that*

$$(5) \quad \frac{\varphi'(b) - \varphi'((1 - \zeta)a + \zeta b)}{\varphi'((1 - \zeta)a + \zeta b) - \varphi'(a)} \leq M$$

whenever $\varphi'((1 - \zeta)a + \zeta b) \leq \lambda\varphi'(a)$, for all $0 < a < b < r$.

An easy way to verify condition (5) is that: suppose $\varphi'(0) = 0$ and $\varphi''(t) > 0$ for all $t \in (0, r)$; then all three conditions below imply condition (5). Moreover, they are related in the following way: $(i) \Rightarrow (ii) \Leftrightarrow (iii)$, where

(i) $\log \circ \varphi'(t)$ is a concave function on $(0, r)$.

(ii) There is ζ , with $0 < \zeta < 1$, such that for all $a, b \in (0, r)$ with $a \leq b$,

$$\varphi'((1 - \zeta)a + \zeta b) \geq \sqrt{\varphi'(a)}\sqrt{\varphi'(b)}.$$

(iii) There are ζ and σ with $0 < \zeta < 1, 0 < \sigma < 1$ such that for all $a, b \in (0, r)$ with $a \leq b$ we have

$$\varphi'((1 - \zeta)a + \zeta b) \geq [\varphi'(a)]^{1-\sigma}[\varphi'(b)]^\sigma.$$

To see that condition (iii) implies (5) we use the mean value theorem to get

$$\begin{aligned} & \frac{\varphi'(b) - \varphi'((1 - \zeta)a + \zeta b)}{\varphi'((1 - \zeta)a + \zeta b) - \varphi'(a)} \\ & \leq \frac{\varphi'((1 - \zeta)a + \zeta b)[\varphi'((1 - \zeta)a + \zeta b)^{\frac{1-\sigma}{\sigma}} - \varphi'(a)^{\frac{1-\sigma}{\sigma}}]}{\varphi'(a)^{\frac{1-\sigma}{\sigma}}[\varphi'((1 - \zeta)a + \zeta b) - \varphi'(a)]} \\ & \leq \frac{1 - \sigma}{\sigma} \left[\frac{\varphi'((1 - \zeta)a + \zeta b)}{\varphi'(a)} \right]^{\frac{1-\sigma}{\sigma}} \quad \text{for } \sigma < \frac{1}{2}. \end{aligned}$$

Also, assuming (iii), we can prove by induction that

$$\varphi'((1 - \zeta)^k a + [1 - (1 - \zeta)^k]b) \geq [\varphi'(a)]^{(1-\sigma)^k} [\varphi'(b)]^{1-(1-\sigma)^k} \quad \text{for all } k \in \mathbb{N}.$$

So, (iii) implies (ii). Condition (ii) does not imply (i), see [5].

Using the theorem we can see that homeomorphisms of the circle as $\varphi'(t) = e^{-1/t}$ or $\varphi'(t) = e^{-1/t^2}$ for $t \in (0, 1/4)$, transport $A(T)$ to $U(T)$.

In [5] there is an example of a C^∞ homeomorphism φ of the circle that satisfies conditions (1), (2), (3) and (4) but does not satisfy condition (5).

3. PROOF OF THE THEOREM

The space $A(T)$ is a Banach space with the norm

$$\| f \|_{A(T)} = \sum_{n \in \mathbb{Z}} | \hat{f}_n |, \quad \text{where } \hat{f}_n = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(t) e^{-int} dt, n \in \mathbb{Z},$$

are the Fourier coefficients of f . Also, $U(T)$ is a Banach space with the norm

$$\| f \|_{U(T)} = \sup \{ | S_m(f, x) | ; x \in [-\pi, \pi], m = 0, 1, 2, \dots \},$$

where $S_m(f, x) = \sum_{-m}^m \hat{f}_n e^{inx}$ are the partial sums of the Fourier series of f .

A homeomorphism of the circle φ of class $C^k, k \geq 1$, transports $A(T)$ to $U(T)$ if and only if there is a constant C , that does not depend on m, n or x , such that

$$(6) \quad \left| \int_{-\pi}^{\pi} e^{in\varphi(t+x)} D_m(t) dt \right| \leq C$$

for all $m = 0, 1, 2, \dots, n \in \mathbb{Z}$ and $x \in [-\pi, \pi]$,

where $D_m(t)$ is the Dirichlet Kernel, i.e.

$$D_m(t) = \frac{\sin(m + \frac{1}{2})t}{\sin(\frac{t}{2})} = \frac{2 \sin mt}{t} + \mathcal{O}(1)$$

on any compact subset of $(-2\pi, 2\pi)$.

Let $(-\tau, \tau)$ be a small neighbourhood of zero and suppose that $\tau \leq |x| \leq \pi$. By (1), there is $\delta > 0$ such that: if $\tau \leq |x| \leq \pi$, then, for some k , depending on x , with $2 \leq k \leq v$ we have $|\varphi^{(k)}(t+x)| \geq \delta$ for all t with $|t| \leq \delta$. Therefore, as in reference [1], the integral in (6) is bounded for $\tau \leq |x| \leq \pi$. So, φ transports $A(T)$ to $U(T)$ if and only if

$$(7) \quad \left| \int_{-\pi}^{\pi} e^{in\varphi(t)} \frac{\sin m(t-x)}{t-x} dt \right| \leq C$$

for all $n \in \mathbb{Z}, m = 0, 1, 2, \dots$ and $x \in (-\tau, \tau)$,

where τ is any positive number less than or equal to $\frac{\pi}{2}$ and C is a constant that does not depend on n, m and x . (The number τ will be chosen conveniently and it will depend on the constant θ .) Let τ be a number such that $\tau < \frac{r}{2} < \frac{\pi}{2}$. So,

$$\left| \int_{-\pi}^{\pi} e^{in\varphi(t)} \frac{\sin m(t-x)}{t-x} dt \right| \leq \left| \int_{-r}^r e^{in\varphi(t)} \frac{\sin m(t-x)}{t-x} dt \right| + \frac{4\pi}{r}$$

for all $n \in \mathbb{Z}, m = 0, 1, 2, \dots$ and for all $x \in (-\tau, \tau)$.

Also, since φ is odd on $(-r, r)$, then, if we prove that the integral

$$(8) \quad \int_0^r e^{in\varphi(t)} \frac{\sin m(t+x)}{t+x} dt$$

is bounded for all $x \in [0, \tau), n > 0$ and $m > 0$, we conclude that (7) holds if and only if the integral

$$(9) \quad \int_0^r e^{in\varphi(t)} \frac{\sin m(t-x)}{t-x} dt$$

is bounded for all $x \in (0, \tau), n > 0$ and $m > 0$.

The boundedness of the integral (8) was proved by Nagel, Vance, Wainger and Weinberg in the second paragraph of [4]. The proof does not use the full force of condition (4); it is based on the bounded doubling time only.

The main part of the theorem is the question of boundedness of the integral (9) for $x \in (0, \tau)$ and $n > 0, m > 0$.

To get boundedness of the integral (9) we have to deal with two major problems: the first one is that the derivative of $mt - n\varphi(t)$ may vanish somewhere between zero and r ; and in that case we do not have enough oscillation around the vanishing point. The second one is that we have a bad singularity at the point x , and around that point, oscillation will not be enough to bound the integral. But, condition (5) will tell us whether there is cancellation around the point x or not. So, to estimate the integral (9) we deal simultaneously with oscillation and cancellation.

Put $f(t) = mt - n\varphi(t)$ and suppose that $x \leq \frac{1}{m}$. Assume first that $f'(r) = m - n\varphi'(r) \geq 0$. Then, we have two possibilities: the first one is $\theta r \leq \frac{2}{m}$. If so, (9) is bounded by $\frac{2}{\theta}$. The second one is $\theta r > \frac{2}{m}$, and in that case we divide the integral

in three parts: over $[0, \frac{2}{m}]$ it is bounded by 2, over $[\theta r, r]$ by $\frac{2}{\theta}$ and over $[\frac{2}{m}, \theta r]$ we have

$$\begin{aligned} & \left| \int_{2/m}^{\theta r} e^{in\varphi(t)} \frac{\sin m(t-x)}{t-x} dt \right| \\ & \leq \left| \int_{2/m}^{\theta r} e^{i(mt+n\varphi(t))} \frac{1}{t-x} dt \right| + \left| \int_{2/m}^{\theta r} e^{if(t)} \frac{1}{t-x} dt \right| \\ & \leq \frac{4}{m(\frac{2}{m}-x)} + \frac{4}{[m-n\varphi'(\theta r)](\frac{2}{m}-x)} \leq 4 + \frac{4m}{[m-n\frac{1}{2}\varphi'(r)]} \leq 12. \end{aligned}$$

We used the Van der Corput lemma, the bounded doubling time, the monotonicity of φ' and the assumption $m-n\varphi'(r) \geq 0$ to obtain the above estimates. We have not used the full force of condition (4). Now, let's assume that $f'(r) = m-n\varphi'(r) < 0$. Then, there is a number ξ with $0 < \xi < r$ such that $f'(\xi) = 0$. This means that $\varphi'(\xi) = \frac{m}{n}$. As before, we have two possibilities: $\theta\xi \leq \frac{2}{m}$ or $\frac{2}{m} < \theta\xi$. If $\theta\xi \leq \frac{2}{m}$, then $\xi < \frac{1}{\theta}\xi \leq \frac{1}{\theta^2}\frac{2}{m}$. We can assume that $\frac{1}{\theta^2}\frac{2}{m} < r$, because otherwise, (9) is bounded by $mr \leq \frac{2}{\theta^2}$. So, assuming $\frac{2}{\theta^2 m} < r$ we divide the integral in two parts: over $[0, \frac{2}{\theta^2 m}]$ it is bounded by $\frac{2}{\theta^2}$, and over $[\frac{2}{\theta^2 m}, r]$ we have

$$\begin{aligned} & \left| \int_{2/\theta^2 m}^r e^{in\varphi(t)} \frac{\sin m(t-x)}{t-x} dt \right| \\ & \leq \left| \int_{2/\theta^2 m}^r e^{i(mt+n\varphi(t))} \frac{1}{t-x} dt \right| + \left| \int_{2/\theta^2 m}^r e^{if(t)} \frac{1}{t-x} dt \right| \\ & \leq \frac{4}{m(\frac{2}{\theta^2 m}-x)} + \frac{4}{[n\varphi'(\frac{2}{\theta^2 m})-m](\frac{2}{\theta^2 m}-x)} \\ & \leq 4 + \frac{4m}{[n\varphi'(\frac{\xi}{\theta})-m]} \leq 4 + \frac{4m}{2n\varphi'(\xi)-m} = 8. \end{aligned}$$

If $\frac{2}{m} < \theta\xi$, then we do the following: the integral is bounded by 2 over the interval $[0, \frac{2}{m}]$ and

$$\begin{aligned} & \left| \int_{2/m}^{\theta\xi} e^{in\varphi(t)} \frac{\sin m(t-x)}{t-x} dt \right| \leq 4 + \left| \int_{2/m}^{\theta\xi} e^{if(t)} \frac{1}{t-x} dt \right| \\ & \leq 4 + \frac{4}{[m-n\varphi'(\theta\xi)](\frac{2}{m}-x)} \leq 4 + \frac{4m}{[m-\frac{1}{2}n\varphi'(\xi)]} = 12. \end{aligned}$$

Now, if $\frac{\xi}{\theta} \geq r$, then

$$\left| \int_{\theta\xi}^r e^{in\varphi(t)} \frac{\sin m(t-x)}{t-x} dt \right| \leq \frac{r}{\theta\xi-x} \leq \frac{\frac{\xi}{\theta}}{\frac{1}{2}\theta\xi} = \frac{2}{\theta^2}.$$

So, we may assume $\frac{\xi}{\theta} < r$. In that case we have

$$\left| \int_{\theta\xi}^{\xi/\theta} e^{in\varphi(t)} \frac{\sin m(t-x)}{t-x} dt \right| \leq \frac{\frac{\xi}{\theta}}{\theta\xi-x} \leq \frac{2}{\theta^2}$$

and

$$\begin{aligned} \left| \int_{\xi/\theta}^r e^{in\varphi(t)} \frac{\sin m(t-x)}{t-x} dt \right| &\leq 4 + \frac{4}{[n\varphi'(\frac{\xi}{\theta}) - m](\frac{\xi}{\theta} - x)} \\ &\leq 4 + \frac{4m}{2n\varphi'(\xi) - m} = 8. \end{aligned}$$

Again we used the Van der Corput lemma and the bounded doubling time to obtain the above estimates. Hence, the integral (9) is bounded if $x \leq \frac{1}{m}$.

Let's consider now the case when $x > \frac{1}{m}$. We have that (7) holds if and only if

$$(10) \quad \left| \int_0^{x-\frac{1}{m}} e^{if(t)} \frac{1}{t-x} dt + \int_{x+\frac{1}{m}}^r e^{if(t)} \frac{1}{t-x} dt \right| \leq C$$

for all $x \in (0, \tau), m, n \in \mathbb{N}$ with $\frac{1}{m} < x$. (C is of course a constant that does not depend on x, m and n .) So, we will prove that (10) holds if and only if the homeomorphism φ satisfies condition (5).

Let's prove first that condition (5) is necessary. For this, let $(\zeta_k)_{k \in \mathbb{N}}$ and $(\lambda_k)_{k \in \mathbb{N}}$ be sequences such that,

$$0 < \zeta_k < 1, \lambda_k > 1 \text{ for all } k \in \mathbb{N} \text{ and } \zeta_k \rightarrow 1, \lambda_k \rightarrow 1.$$

Suppose that φ does not satisfy condition (5). Then, for each $k \in \mathbb{N}$ there exist numbers a_k and b_k , with $0 < a_k < b_k < \frac{1}{k}r$, so that

$$(11) \quad \frac{\varphi'(b_k) - \varphi'(x_k)}{\varphi'(x_k) - \varphi'(a_k)} > \frac{\zeta_k}{1 - \zeta_k}$$

and

$$(12) \quad \varphi'(x_k) \leq \lambda_k \varphi'(a_k)$$

where $x_k = (1 - \zeta_k)a_k + \zeta_k b_k$. Put

$$\begin{aligned} \epsilon_k &= \zeta_k(b_k - a_k), \\ \epsilon'_k &= (1 - \zeta_k)(b_k - a_k), \\ \nu_k &= \frac{1}{\epsilon_k[\varphi'(x_k) - \varphi'(a_k)]}, \\ \mu_k &= \varphi'(x_k)\nu_k. \end{aligned}$$

Let m_k be the smallest integer greater than μ_k and n_k be the greatest integer less than ν_k . So, by (11), for k sufficiently large we have

$$(13) \quad \frac{1}{2} \leq \epsilon_k[m_k - n_k\varphi'(x_k - \epsilon_k)] \leq \frac{3}{2}$$

and

$$(14) \quad \epsilon'_k[n_k\varphi'(x_k + \epsilon'_k) - m_k] > \frac{1}{2}.$$

Inequality (14) implies that there is $\bar{\epsilon}_k$ with $0 < \bar{\epsilon}_k < \epsilon'_k$ such that

$$(15) \quad \bar{\epsilon}_k[n_k\varphi'(x_k + \bar{\epsilon}_k) - m_k] = \frac{1}{2}.$$

Also, by (12), $\epsilon_k m_k \rightarrow +\infty$ and since $x_k > \epsilon_k$, then $x_k > \frac{1}{m_k}$. Since $x_k \rightarrow 0$, then $0 < x_k < \tau$ for k large enough.

Suppose that $\bar{\epsilon}_k \leq \frac{1}{m_k}$ for an infinite number of indices k .

Let $f_k(t) = m_k t - n_k \varphi(t)$. Then, by (15) and the Van der Corput lemma we have that

$$\left| \int_{x_k + \frac{1}{m_k}}^r e^{if_k(t)} \frac{1}{t - x_k} dt \right| \leq \frac{4m_k}{|f'_k(x_k + \bar{\varepsilon}_k)|} \leq \frac{4}{\bar{\varepsilon}_k |f'_k(x_k + \bar{\varepsilon}_k)|} = 8,$$

for an infinite number of indices k . By (13) and the Van der Corput lemma,

$$\left| \int_0^{x_k - \varepsilon_k} e^{if_k(t)} \frac{1}{t - x_k} dt \right| \leq \frac{4}{\varepsilon_k f'_k(x_k - \varepsilon_k)} \leq 8.$$

Now, using (13),

$$\left| \int_{x_k - \varepsilon_k}^{x_k - \frac{1}{m_k}} [e^{if_k(t)} - e^{if_k(x_k)}] \frac{1}{t - x_k} dt \right| \leq \varepsilon_k f'_k(x_k - \varepsilon_k) \leq \frac{3}{2}$$

and

$$\left| \int_{x_k - \varepsilon_k}^{x_k - \frac{1}{m_k}} e^{if_k(x_k)} \frac{1}{t - x_k} dt \right| = \int_{1/m_k}^{\varepsilon_k} \frac{1}{t} dt = \log(\varepsilon_k m_k).$$

Hence, we conclude that (10) does not hold. Suppose now that $\bar{\varepsilon}_k > \frac{1}{m_k}$ for all k large enough. Then, by (13) and (15),

$$\left| \int_{x_k + \bar{\varepsilon}_k}^r e^{if_k(t)} \frac{1}{t - x_k} dt \right| \leq \frac{4}{\bar{\varepsilon}_k |f'_k(x_k + \bar{\varepsilon}_k)|} = 8$$

and

$$\left| \int_0^{x_k - \varepsilon_k} e^{if_k(t)} \frac{1}{t - x_k} dt \right| \leq \frac{4}{\varepsilon_k f'_k(x_k - \varepsilon_k)} \leq 8,$$

for all k large enough. Finally, by (13) and (15),

$$\left| \int_{x_k - \varepsilon_k}^{x_k - \frac{1}{m_k}} (e^{if_k(t)} - e^{if_k(x_k)}) \frac{1}{t - x_k} dt \right| \leq \varepsilon_k f'_k(x_k - \varepsilon_k) \leq \frac{3}{2}$$

and

$$\left| \int_{x_k + \frac{1}{m_k}}^{x_k + \bar{\varepsilon}_k} (e^{if_k(t)} - e^{if_k(x_k)}) \frac{1}{t - x_k} dt \right| \leq \varepsilon_k f'_k(x_k - \varepsilon_k) \leq \frac{3}{2}.$$

Hence, since

$$\begin{aligned} & \left| \int_{x_k - \varepsilon_k}^{x_k - \frac{1}{m_k}} e^{if_k(x_k)} \frac{1}{t - x_k} dt + \int_{x_k + \frac{1}{m_k}}^{x_k + \bar{\varepsilon}_k} e^{if_k(x_k)} \frac{1}{t - x_k} dt \right| \\ &= \left| \int_{-\varepsilon_k}^{-\frac{1}{m_k}} \frac{1}{t} dt + \int_{1/m_k}^{\bar{\varepsilon}_k} \frac{1}{t} dt \right| = \log\left(\frac{\varepsilon_k}{\bar{\varepsilon}_k}\right) \geq \log\left(\frac{\varepsilon_k}{\varepsilon'_k}\right) = \log\left(\frac{\zeta_k}{1 - \zeta_k}\right), \end{aligned}$$

we conclude that (10) does not hold. Therefore, condition (5) is necessary.

Now, let's prove sufficiency. So, assuming that φ satisfies condition (5), we have to show that (10) is true. For this, let $x \in (0, \tau), n, m \in \mathbb{N}$ with $x > \frac{1}{m}, m > 0, n > 0$. We said before that the number τ would be chosen conveniently. So, let $\tau < \frac{1}{2}\theta r$. (The reason for that choice will be clear in the proof that follows. It is not quite necessary, but it simplifies the proof a lot.) Put $f(t) = mt - n\varphi(t)$ as before and suppose that $f'(r) = m - n\varphi'(r) \geq 0$. Since $x + \frac{1}{m} < 2x < 2\tau < \theta r$, then using the bounded doubling time, the monotonicity of φ' , and the Van der Corput lemma we have that

$$\begin{aligned} \left| \int_0^{x-\frac{1}{m}} e^{if(t)} \frac{1}{t-x} dt \right| &\leq \frac{4m}{f'(x-\frac{1}{m})} \leq \frac{4m}{f'(\theta r)} \leq \frac{4m}{m-n\frac{1}{2}\varphi'(r)} \leq 8, \\ \left| \int_{x+\frac{1}{m}}^{\theta r} e^{if(t)} \frac{1}{t-x} dt \right| &\leq \frac{4m}{f'(\theta r)} \leq 8 \end{aligned}$$

and

$$\left| \int_{\theta r}^r e^{if(t)} \frac{1}{t-x} dt \right| \leq \frac{r}{\theta r - x} \leq \frac{2}{\theta}.$$

Hence, (10) holds if $f'(r) = m - n\varphi'(r) \geq 0$ and $\frac{1}{m} < x < \tau$.

Suppose now that $f'(r) = m - n\varphi'(r) < 0$. So, there exists a number ξ with $0 < \xi < r$ such that $f'(\xi) = m - n\varphi'(\xi) = 0$. Since $m x > 1$, then there exists a number ε with $0 < \varepsilon < x$ such that $\varepsilon(m - n\varphi'(x - \varepsilon)) = 1 - \frac{1}{\lambda}$. We have two cases to consider: one is when $x \leq \xi$ and the other is when $\xi < x$. We will treat each one of them separately. From now on we have to deal simultaneously with oscillation and cancellation. Roughly, the problem is that: when we approach near x , the singularity is so bad that even if there is oscillation at x , this is not enough to guarantee boundedness. So, we have to use the cancellation we have around x . In order to have an effective cancellation we need to use condition (5). On the other hand, when we approach ξ , we don't have oscillation since $f'(\xi) = 0$. So, to deal with the integral around ξ we have to use condition (4). An important factor will be how far (with respect to ε) is ξ from x .

Assume that $x \leq \xi$. Call $\alpha = \xi - x$. The arguments we will use depend on the relation between ε, α and m . Suppose we have the following situation: $0 < \frac{1}{m} < \varepsilon < \frac{1}{2}\theta\alpha$. Let's choose a number A such that $\alpha + \varepsilon = \theta(A + \alpha + \varepsilon)$. We see that $A = \frac{1-\theta}{\theta}(\alpha + \varepsilon)$ is that number. We are going to divide the integral in several parts and prove boundedness of each one separately:

$$\begin{aligned} \left| \int_0^{x-\varepsilon} e^{if(t)} \frac{1}{t-x} dt \right| &\leq \frac{4}{\varepsilon f'(x-\varepsilon)} = \frac{4\lambda}{\lambda-1}, \\ \left| \int_{x-\varepsilon}^{x-\frac{1}{m}} e^{if(t)} \frac{1}{t-x} dt + \int_{x+\frac{1}{m}}^{x+\varepsilon} e^{if(t)} \frac{1}{t-x} dt \right| \\ &= \left| \int_{x-\varepsilon}^{x-\frac{1}{m}} (e^{if(t)} - e^{if(x)}) \frac{1}{t-x} dt + \int_{x+\frac{1}{m}}^{x+\varepsilon} (e^{if(t)} - e^{if(x)}) \frac{1}{t-x} dt \right| \\ &\leq 2\varepsilon f'(x-\varepsilon) = \frac{2(\lambda-1)}{\lambda}. \end{aligned}$$

To estimate the integral from $x + \varepsilon$ to $x + \frac{1}{2}\theta\alpha$ we use condition (4). This condition assures enough oscillation in that interval. We claim that

$$(16) \quad m - n\varphi'(x + \frac{1}{2}\theta\alpha) \geq \frac{1}{2}[m - n\varphi'(x - \varepsilon)].$$

To prove (16) we use condition (4): since $\varepsilon < \frac{1}{2}\theta\alpha$, then

$$x + \frac{1}{2}\theta\alpha \leq (1 - \theta)(x - \varepsilon) + \theta\xi;$$

this implies that

$$\varphi'(x + \frac{1}{2}\theta\alpha) \leq \varphi'((1 - \theta)(x - \varepsilon) + \theta\xi) \leq \frac{1}{2}(\varphi'(x - \varepsilon) + \varphi'(\xi))$$

and (16) follows. Therefore, by (16)

$$\left| \int_{x+\varepsilon}^{x+\frac{1}{2}\theta\alpha} e^{if(t)} \frac{1}{t-x} dt \right| \leq \frac{4}{\varepsilon f'(x + \frac{1}{2}\theta\alpha)} \leq \frac{8}{\varepsilon f'(x - \varepsilon)} = \frac{8\lambda}{\lambda - 1}.$$

Now, we have to pass through the point ξ where we don't have oscillation at all. But since we are away from x , things become much simpler:

$$\left| \int_{x+\frac{1}{2}\theta\alpha}^{\xi+A} e^{if(t)} \frac{1}{t-x} dt \right| \leq \log \left(\frac{\xi + A - x}{\frac{1}{2}\theta\alpha} \right) \leq \log \left(\frac{2 + \theta(1 - \theta)}{\theta^2} \right).$$

Note that we can assume that $\xi + A < r$, because otherwise,

$$\left| \int_{x+\frac{1}{2}\theta\alpha}^r e^{if(t)} \frac{1}{t-x} dt \right| \leq \log \left(\frac{r-x}{\frac{1}{2}\theta\alpha} \right) \leq \log \left(\frac{\xi + A - x}{\frac{1}{2}\theta\alpha} \right)$$

and we are done. Finally, to evaluate the integral from $\xi + A$ to r we use again condition (4). This condition, as before, assures enough oscillation in that interval. We claim that

$$(17) \quad n\varphi'(\xi + A) - m \geq m - n\varphi'(x - \varepsilon).$$

By the choice of A , we have that $\xi = (1 - \theta)(x - \varepsilon) + \theta(\xi + A)$. So,

$$\varphi'(\xi) \leq \frac{1}{2}(\varphi'(x - \varepsilon) + \varphi'(\xi + A))$$

and (17) follows. Using (17) now we have

$$\begin{aligned} \left| \int_{\xi+A}^r e^{if(t)} \frac{1}{t-x} dt \right| &\leq \frac{4}{(\xi + A - x) |f'(\xi + A)|} = \frac{4}{(\alpha + A) |f'(\xi + A)|} \\ &\leq \frac{4}{\varepsilon f'(x - \varepsilon)} = \frac{4\lambda}{\lambda - 1} \end{aligned}$$

since $\alpha + A > \alpha > \varepsilon$.

Let's see now another possible situation: suppose that $0 < \varepsilon \leq \frac{1}{m} < \frac{1}{2}\theta\alpha$. Then,

$$\begin{aligned} \left| \int_0^{x-\frac{1}{m}} e^{if(t)} \frac{1}{t-x} dt \right| &\leq \frac{4m}{f'(x - \frac{1}{m})} \leq \frac{4}{\varepsilon f'(x - \varepsilon)} = \frac{4\lambda}{\lambda - 1}, \\ \left| \int_{x+\frac{1}{m}}^{x+\frac{1}{2}\theta\alpha} e^{if(t)} \frac{1}{t-x} dt \right| &\leq \frac{4m}{f'(x + \frac{1}{2}\theta\alpha)} \leq \frac{8}{\varepsilon f'(x - \varepsilon)} = \frac{8\lambda}{\lambda - 1}, \end{aligned}$$

by (16).

From $x + \frac{1}{2}\theta\alpha$ to $\xi + A$ and from $\xi + A$ to r there is no change.

If $0 < \varepsilon < \frac{1}{2}\theta\alpha \leq \frac{1}{m}$, then

$$\left| \int_0^{x-\frac{1}{m}} e^{if(t)} \frac{1}{t-x} dt \right| \leq \frac{4m}{f'(x-\frac{1}{m})} \leq \frac{4}{\varepsilon f'(x-\varepsilon)} = \frac{4\lambda}{\lambda-1}.$$

If $x + \frac{1}{m} < \xi + A$, then

$$\begin{aligned} \left| \int_{x+\frac{1}{m}}^{\xi+A} e^{if(t)} \frac{1}{t-x} dt \right| &\leq \log\left(\frac{\xi+A-x}{\frac{1}{m}}\right) \leq \log\left(\frac{\alpha+A}{\frac{1}{2}\theta\alpha}\right) \\ &\leq \log\left(\frac{2+\theta(1-\theta)}{\theta^2}\right). \end{aligned}$$

Again we can assume $\xi + A < r$. Note that $x + \frac{1}{m} < r$ because $\frac{1}{m} < x < \tau < \frac{r}{2}$.

$$\left| \int_{\xi+A}^r e^{if(t)} \frac{1}{t-x} dt \right| \leq \frac{4\lambda}{\lambda-1} \quad \text{as before.}$$

If $\xi + A \leq x + \frac{1}{m}$, then

$$\begin{aligned} \left| \int_{x+\frac{1}{m}}^r e^{if(t)} \frac{1}{t-x} dt \right| &\leq \frac{4m}{|f'(x+\frac{1}{m})|} \leq \frac{4}{\varepsilon |f'(\xi+A)|} \\ &\leq \frac{4}{\varepsilon f'(x-\varepsilon)} = \frac{4\lambda}{\lambda-1}, \end{aligned}$$

by (17).

The next situation is $\frac{1}{2}\theta\alpha \leq \varepsilon$ and $\frac{1}{m} \geq \varepsilon$. Assuming this we have

$$\left| \int_0^{x-\frac{1}{m}} e^{if(t)} \frac{1}{t-x} dt \right| \leq \frac{4m}{f'(x-\frac{1}{m})} \leq \frac{4}{\varepsilon f'(x-\varepsilon)} = \frac{4\lambda}{\lambda-1}.$$

If $x + \frac{1}{m} < \xi + A$, then

$$\left| \int_{x+\frac{1}{m}}^{\xi+A} e^{if(t)} \frac{1}{t-x} dt \right| \leq \log\left(\frac{\alpha+A}{\frac{1}{m}}\right) \leq \log\left(\frac{2+\theta(1-\theta)}{\theta^2}\right).$$

As before, we can assume $\xi + A < r$, so

$$\left| \int_{\xi+A}^r e^{if(t)} \frac{1}{t-x} dt \right| \leq \frac{4}{(\alpha+A)|f'(\xi+A)|} \leq \frac{4}{\varepsilon f'(x-\varepsilon)} = \frac{4\lambda}{\lambda-1},$$

by (17).

If $\xi + A \leq x + \frac{1}{m}$, then, by (17), the integral from $x + \frac{1}{m}$ to r is bounded by $\frac{4\lambda}{\lambda-1}$ as in the preceding situation.

Now suppose that $\frac{1}{2}\theta\alpha \leq \varepsilon$ and $\frac{1}{m} < \varepsilon$. Let B be a number such that $\alpha + \varepsilon = \zeta(B + \alpha + \varepsilon)$. We see that $B = \frac{1-\zeta}{\zeta}(\alpha + \varepsilon)$ is that number. Since we can assume that $\theta < \zeta$, then $B < A$. Indeed we will assume $\theta < \frac{1}{2}$ and $\zeta > \frac{1}{2}$. So, we have that $x + \frac{1}{m} < \xi + A$, because $A = \frac{1-\theta}{\theta}(\alpha + \varepsilon) > \alpha + \varepsilon \geq \varepsilon > \frac{1}{m}$. But, it may happen that $\xi + B \leq x + \frac{1}{m}$, because α can be even zero. Let's see first this special case,

i.e. if $\frac{1}{2}\theta\alpha \leq \varepsilon$ and $\alpha + B \leq \frac{1}{m} < \varepsilon$. We do the following:

$$\begin{aligned} \left| \int_0^{x-\varepsilon} e^{if(t)} \frac{1}{t-x} dt \right| &\leq \frac{4}{\varepsilon f'(x-\varepsilon)} = \frac{4\lambda}{\lambda-1}, \\ \left| \int_{x-\varepsilon}^{x-\frac{1}{m}} e^{if(t)} \frac{1}{t-x} dt \right| &\leq m\varepsilon \leq \frac{\varepsilon}{\alpha+B} \leq \frac{\zeta}{1-\zeta}, \\ \left| \int_{x+\frac{1}{m}}^{\xi+A} e^{if(t)} \frac{1}{t-x} dt \right| &\leq \log\left(\frac{\alpha+A}{\frac{1}{m}}\right) \leq \log\left(\frac{\alpha+A}{\alpha+B}\right) \\ &\leq \log\left[\left(\frac{1-\theta}{\theta}\right)\left(\frac{\zeta}{1-\zeta}\right)\right]. \end{aligned}$$

(As before, we can assume $\xi + A < r$. Indeed, since $\frac{1}{2}\theta\alpha \leq \varepsilon$, then $\xi + A < r$ if x is sufficiently small.)

$$\left| \int_{\xi+A}^r e^{if(t)} \frac{1}{t-x} dt \right| \leq \frac{4}{(\alpha+A) |f'(\xi+A)|} \leq \frac{4}{\varepsilon f'(x-\varepsilon)} = \frac{4\lambda}{\lambda-1},$$

by (17).

Let's see now the case that was left over, i.e. we assume $\frac{1}{2}\theta\alpha \leq \varepsilon$, $\frac{1}{m} < \varepsilon$ and $\frac{1}{m} < \alpha + B$. We claim that

$$(18) \quad n\varphi'(\xi + B) - m \leq M[m - n\varphi'(x - \varepsilon)].$$

To prove (18) we use condition (5). By the choice of B we have that $\xi = (1 - \zeta)(x - \varepsilon) + \zeta(\xi + B)$. Also, since $\varepsilon f'(x - \varepsilon) = \frac{\lambda-1}{\lambda}$ and $\varepsilon m > 1$, then $\varphi'(\xi) < \lambda\varphi'(x - \varepsilon)$. Hence, (18) follows from (5). Now, let's see how to get boundedness of the integral in that last situation. From zero to $x - \varepsilon$ is quite easy and we have $\left| \int_0^{x-\varepsilon} e^{if(t)} \frac{1}{t-x} dt \right| \leq \frac{4}{\varepsilon f'(x-\varepsilon)} = \frac{4\lambda}{\lambda-1}$. (We recall again here that we are free to choose the number τ . So, let's choose τ sufficiently small in order to have $\xi + B < r$. This procedure is not quite necessary, but it simplifies a little bit that part of the proof. Note that it is possible to choose such τ , for in that situation we have

$$\begin{aligned} \xi + B = x + \alpha + \frac{1-\zeta}{\zeta}(\alpha + \varepsilon) &\leq x + \frac{2}{\theta}\varepsilon + \frac{1-\zeta}{\zeta}\left(\frac{2}{\theta}\varepsilon + \varepsilon\right) \\ &\leq x + \frac{2}{\theta}x + \frac{1-\zeta}{\zeta}\left(\frac{2}{\theta}x + x\right) \\ &\leq \left(\frac{4}{\theta} + 2\right)x. \end{aligned}$$

So, $\tau < \frac{1}{6}\theta r$ will do the job.)

$$\left| \int_{\xi+B}^{\xi+A} e^{if(t)} \frac{1}{t-x} dt \right| \leq \log\left(\frac{\alpha+A}{\alpha+B}\right) \leq \log\left[\left(\frac{1-\theta}{\theta}\right)\left(\frac{\zeta}{1-\zeta}\right)\right].$$

We can also assume that $\xi + A < r$, because if not, we get boundedness of the integral over the interval $[\xi + B, r]$ as we did above. So,

$$\left| \int_{\xi+A}^r e^{if(t)} \frac{1}{t-x} dt \right| = \frac{4}{(\alpha+A) |f'(\xi+A)|} \leq \frac{4}{\varepsilon f'(x-\varepsilon)} = \frac{4\lambda}{\lambda-1},$$

by (17).

It remains to prove boundedness over $[x - \varepsilon, x - \frac{1}{m}] \cup [x + \frac{1}{m}, \xi + B]$. We will use cancellation around x to do this. The cancellation depends on the inequality (18) as well as on the assumption that $\varepsilon \geq \frac{1}{2}\theta\alpha$. We have

$$\begin{aligned} & \left| \int_{x-\varepsilon}^{x-\frac{1}{m}} e^{if(t)} \frac{1}{t-x} dt + \int_{x+\frac{1}{m}}^{\xi+B} e^{if(t)} \frac{1}{t-x} dt \right| \\ &= \left| \int_{x-\varepsilon}^{x-\frac{1}{m}} (e^{if(t)} - e^{if(x)}) \frac{1}{t-x} dt + \int_{x+\frac{1}{m}}^{\xi+B} (e^{if(t)} - e^{if(x)}) \frac{1}{t-x} dt \right. \\ & \quad \left. + \int_{x-\varepsilon}^{x-\frac{1}{m}} e^{if(x)} \frac{1}{t-x} dt + \int_{x+\frac{1}{m}}^{\xi+B} e^{if(x)} \frac{1}{t-x} dt \right| \\ &\leq \varepsilon f'(x-\varepsilon) + M f'(x-\varepsilon)(\alpha+B) + \left| \int_{1/m}^{\alpha+B} \frac{1}{t} dt - \int_{1/m}^{\varepsilon} \frac{1}{t} dt \right| \\ &\leq \varepsilon f'(x-\varepsilon) + M \varepsilon f'(x-\varepsilon) \left[\frac{2}{\theta} + \frac{1-\zeta}{\zeta} \left(\frac{2}{\theta} + 1 \right) \right] + \left| \log \left(\frac{\alpha+B}{\varepsilon} \right) \right| \\ &= \frac{\lambda-1}{\lambda} + M \left(\frac{\lambda-1}{\lambda} \right) \left[\frac{2}{\theta} + \frac{1-\zeta}{\zeta} \left(\frac{2}{\theta} + 1 \right) \right] + \left| \log \left(\frac{\alpha+B}{\varepsilon} \right) \right|, \end{aligned}$$

by (18).

Since $\frac{\alpha+B}{\varepsilon} \geq \frac{B}{\varepsilon} = \frac{(\frac{1-\zeta}{\zeta})(\alpha+\varepsilon)}{\varepsilon} \geq \frac{1-\zeta}{\zeta}$ and $\frac{\alpha+B}{\varepsilon} < \frac{\frac{2}{\theta}\varepsilon + (\frac{1-\zeta}{\zeta})(\frac{2}{\theta}\varepsilon + \varepsilon)}{\varepsilon} \leq \frac{4+\theta}{\theta}$, then $\left| \log \left(\frac{\alpha+B}{\varepsilon} \right) \right| \leq \max \left\{ \left| \log \left(\frac{1-\zeta}{\zeta} \right) \right|, \log \left(\frac{4+\theta}{\theta} \right) \right\}$. This concludes the case $x \leq \xi$.

Assume now that $\xi < x$. Call $\beta = x - \xi$. This case is quite different from the previous one, and it requires different arguments. We will introduce a new number, say δ , and that number controls the amount of oscillation and the amount of cancellation that we have. The control is done in the following sense: when δ becomes small, this means that we lose cancellation but we gain oscillation; and when δ becomes large this means that we lose oscillation, but we gain cancellation.

Let δ be a positive number such that $\delta[n\varphi'(\xi + \frac{1+\zeta}{2}\beta) - m] = \lambda - 1$. Since $f'(x - \varepsilon) > 0$, then $x - \varepsilon < \xi$. So, $\varepsilon > \beta$. Let's consider first the following situation. Suppose that $0 < \frac{1}{m} < \delta < \frac{1-\zeta}{2}\beta$ and $\varepsilon \leq \frac{1+\zeta}{1-\zeta}\beta$. We claim that

$$(19) \quad n\varphi'(\xi + \frac{1+\zeta}{2}\beta) - m \leq (M+1)[n\varphi'(\xi + \frac{1+\zeta}{2}\beta) - m].$$

To prove (19) we use condition (5). (Note that we can choose τ sufficiently small in order to have $\xi + \frac{1+\zeta}{2}\beta < r$.) We have that $\xi + \frac{1+\zeta}{2}\beta = (1-\zeta)\xi + \zeta(\xi + \frac{1+\zeta}{2}\beta)$. Since $\delta | f'(\xi + \frac{1+\zeta}{2}\beta) | = \lambda - 1$ and $\delta m > 1$, then $\varphi'(\xi + \frac{1+\zeta}{2}\beta) < \lambda\varphi'(\xi)$. Hence, (19) follows from (5). Let's work with the integral now. We divide it in several parts.

$$\left| \int_0^{x-\varepsilon} e^{if(t)} \frac{1}{t-x} dt \right| \leq \frac{4}{\varepsilon f'(x-\varepsilon)} = \frac{4\lambda}{\lambda-1},$$

$$\left| \int_{x-\varepsilon}^{\xi+\frac{1+\zeta}{2}\beta} e^{if(t)} \frac{1}{t-x} dt \right| \leq \log \left(\frac{\varepsilon}{\frac{1-\zeta}{2}\beta} \right) \leq \log \left(\frac{2(1+\zeta)}{(1-\zeta)^2} \right),$$

$$\left| \int_{\xi+\frac{1+\zeta}{2}\beta}^{x-\delta} e^{if(t)} \frac{1}{t-x} dt \right| \leq \frac{4}{\delta |f'(\xi+\frac{1+\zeta}{2}\beta)|} = \frac{4}{\lambda-1},$$

and

$$\left| \int_{x+\delta}^r e^{if(t)} \frac{1}{t-x} dt \right| \leq \frac{4}{\delta |f'(x+\delta)|} \leq \frac{4}{\delta |f'(\xi+\frac{1+\zeta}{2}\beta)|} = \frac{4}{\lambda-1}.$$

To prove boundedness around x we use the small amount of cancellation that we have there:

$$\begin{aligned} & \left| \int_{x-\delta}^{x-\frac{1}{m}} e^{if(t)} \frac{1}{t-x} dt + \int_{x-\frac{1}{m}}^{x+\delta} e^{if(t)} \frac{1}{t-x} dt \right| \\ &= \left| \int_{x-\delta}^{x-\frac{1}{m}} (e^{if(t)} - e^{if(x)}) \frac{1}{t-x} dt + \int_{x+\frac{1}{m}}^{x+\delta} (e^{if(t)} - e^{if(x)}) \frac{1}{t-x} dt \right| \\ &\leq 2\delta [n\varphi'(\xi + \frac{1+\zeta}{2\zeta}\beta) - m] \\ &\leq 2\delta(M+1)[n\varphi'(\xi + \frac{1+\zeta}{2}\beta) - m] = 2(M+1)(\lambda-1), \end{aligned}$$

by (19).

Another possible situation is that: $0 < \delta \leq \frac{1}{m} < \frac{1-\zeta}{2}\beta$ and $\varepsilon \leq \frac{1+\zeta}{1-\zeta}\beta$. This case is better than the previous one because the small δ here means that we have a large amount of oscillation, and $\frac{1}{m} \geq \delta$ means that $\frac{1}{m}$ is large enough to put us away from the bad singularity at x . Hence, as we shall see, we don't need conditions (4) and (5) in this case. We have:

$$\begin{aligned} & \left| \int_0^{x-\varepsilon} e^{if(t)} \frac{1}{t-x} dt \right| \leq \frac{4}{\varepsilon f'(x-\varepsilon)} = \frac{4\lambda}{\lambda-1}, \\ & \left| \int_{x-\varepsilon}^{\xi+\frac{1+\zeta}{2}\beta} e^{if(t)} \frac{1}{t-x} dt \right| \leq \log \left(\frac{\varepsilon}{\frac{1-\zeta}{2}\beta} \right) \leq \log \left(\frac{2(1+\zeta)}{(1-\zeta)^2} \right), \\ & \left| \int_{\xi+\frac{1+\zeta}{2}\beta}^{x-\frac{1}{m}} e^{if(t)} \frac{1}{t-x} dt \right| \leq \frac{4m}{|f'(\xi+\frac{1+\zeta}{2}\beta)|} \\ & \leq \frac{4}{\delta |f'(\xi+\frac{1+\zeta}{2}\beta)|} = \frac{4}{\lambda-1}, \\ & \left| \int_{x+\frac{1}{m}}^r e^{if(t)} \frac{1}{t-x} dt \right| \leq \frac{4m}{|f'(x+\frac{1}{m})|} \\ & \leq \frac{4}{\delta |f'(\xi+\frac{1+\zeta}{2}\beta)|} = \frac{4}{\lambda-1}. \end{aligned}$$

The next situation is: $0 < \delta < \frac{1-\zeta}{2}\beta \leq \frac{1}{m}$ and $\varepsilon \leq \frac{1+\zeta}{1-\zeta}\beta$. In that case we do the following: if $x - \frac{1}{m} \leq x - \varepsilon$, then

$$\begin{aligned} \left| \int_0^{x-\frac{1}{m}} e^{if(t)} \frac{1}{t-x} dt \right| &\leq \frac{4m}{f'(x-\frac{1}{m})} \leq \frac{4}{\varepsilon f'(x-\varepsilon)} = \frac{4\lambda}{\lambda-1}, \\ \left| \int_{x+\frac{1}{m}}^r e^{if(t)} \frac{1}{t-x} dt \right| &\leq \frac{4m}{|f'(x+\frac{1}{m})|} \leq \frac{4}{\delta |f'(\xi + \frac{1+\zeta}{2}\beta)|} = \frac{4}{\lambda-1}. \end{aligned}$$

If $x - \varepsilon < x - \frac{1}{m}$, then

$$\begin{aligned} \left| \int_0^{x-\varepsilon} e^{if(t)} \frac{1}{t-x} dt \right| &\leq \frac{4}{\varepsilon f'(x-\varepsilon)} = \frac{4\lambda}{\lambda-1}, \\ \left| \int_{x-\varepsilon}^{x-\frac{1}{m}} e^{if(t)} \frac{1}{t-x} dt \right| &\leq \log(\varepsilon m) \leq \log\left(\frac{2(1+\zeta)}{(1-\zeta)^2}\right), \\ \left| \int_{x+\frac{1}{m}}^r e^{if(t)} \frac{1}{t-x} dt \right| &\leq \frac{4m}{|f'(x+\frac{1}{m})|} \\ &\leq \frac{4}{\delta |f'(\xi + \frac{1+\zeta}{2}\beta)|} = \frac{4}{\lambda-1}. \end{aligned}$$

Suppose now that $\frac{1}{m} < \frac{1-\zeta}{2}\beta \leq \delta$ and $\varepsilon \leq \frac{1+\zeta}{1-\zeta}\beta$. In that situation we have to use conditions (4) and (5) because the large δ means that we don't have enough oscillation, and the small $\frac{1}{m}$ means that we are too close to the singularity at x . But, on the other hand, the large δ means that we have much cancellation around x , and we will use that fact to compensate the lack of oscillation in this case.

$$\begin{aligned} \left| \int_0^{x-\varepsilon} e^{if(t)} \frac{1}{t-x} dt \right| &\leq \frac{4}{\varepsilon f'(x-\varepsilon)} = \frac{4\lambda}{\lambda-1}, \\ \left| \int_{x-\varepsilon}^{\xi+\frac{1+\zeta}{2}\beta} e^{if(t)} \frac{1}{t-x} dt \right| &\leq \log\left(\frac{\varepsilon}{\frac{1-\zeta}{2}\beta}\right) \\ &\leq \log\left(\frac{2(1+\zeta)}{(1-\zeta)^2}\right), \\ \left| \int_{\xi+\frac{1+\zeta}{2}\beta}^{x-\frac{1}{m}} e^{if(t)} \frac{1}{t-x} dt + \int_{x+\frac{1}{m}}^{x+\frac{1-\zeta}{2}\beta} e^{if(t)} \frac{1}{t-x} dt \right| \\ &= \left| \int_{\xi+\frac{1+\zeta}{2}\beta}^{x-\frac{1}{m}} (e^{if(t)} - e^{if(x)}) \frac{1}{t-x} dt + \int_{x+\frac{1}{m}}^{x+\frac{1-\zeta}{2}\beta} (e^{if(t)} - e^{if(x)}) \frac{1}{t-x} dt \right| \\ &\leq 2\delta \left| f'(\xi + \frac{1+\zeta}{2}\beta) \right| \\ &\leq 2\delta(M+1) \left| f'(\xi + \frac{1+\zeta}{2}\beta) \right| = 2(M+1)(\lambda-1), \end{aligned}$$

by (19).

We still have to prove boundedness of the integral over the interval $[x + \frac{1-\zeta}{2}\beta, r]$. Since δ is large, then we may not have enough oscillation at the point $x + \frac{1-\zeta}{2}\beta$.

So, we cannot just integrate from $x + \frac{1-\zeta}{2}\beta$ to r using the Van der Corput lemma. We have to use condition (4).

Let A' be a number such that $\varepsilon - \beta = \theta(A' + \varepsilon - \beta)$. We can see that $A' = \frac{1-\theta}{\theta}(\varepsilon - \beta)$ is that number. We claim that

$$(20) \quad \begin{aligned} n\varphi'(x + \frac{1-\zeta}{2}\beta + A') - m &\geq n\varphi'(\xi + A') - m, \\ n\varphi'(\xi + A') - m &\geq m - n\varphi'(x - \varepsilon). \end{aligned}$$

To prove (20) we use (4). By the choice of A' we have that $\xi = (1 - \theta)(x - \varepsilon) + \theta(\xi + A')$. Since $\xi + A' < x + \frac{1-\zeta}{2}\beta + A'$, (20) follows from (4).

We will use inequality (20) to show boundedness over the interval $[x + \frac{1-\zeta}{2}\beta, r]$. We divide the interval from $x + \frac{1-\zeta}{2}\beta$ to $x + \frac{1-\zeta}{2}\beta + A'$ and from $x + \frac{1-\zeta}{2}\beta + A'$ to r . (Note that we can choose τ sufficiently small in order to have $x + \frac{1-\zeta}{2}\beta + A' < r$, but again this is not quite necessary because if not we just integrate from $x + \frac{1-\zeta}{2}\beta$ to r as we have done before.) We have

$$\left| \int_{x + \frac{1-\zeta}{2}\beta}^{x + \frac{1-\zeta}{2}\beta + A'} e^{if(t)} \frac{1}{t - x} dt \right| \leq \log \left(\frac{\frac{1-\zeta}{2}\beta + A'}{\frac{1-\zeta}{2}\beta} \right) \leq \log \left(\frac{\frac{1-\zeta}{2} + \frac{1-\theta}{\theta} \left(\frac{1+\zeta}{1-\zeta} \right)}{\frac{1-\zeta}{2}} \right)$$

and

$$\begin{aligned} \left| \int_{x + \frac{1-\zeta}{2}\beta + A'}^r e^{if(t)} \frac{1}{t - x} dt \right| &\leq \frac{4}{|f'(x + \frac{1-\zeta}{2}\beta + A')| \left(\frac{1-\zeta}{2}\beta + A' \right)} \\ &\leq \frac{4}{\left(\frac{1-\zeta}{2}\beta \right) f'(x - \varepsilon)}, \end{aligned}$$

by (20).

But now, since $\varepsilon \leq \frac{1+\zeta}{1-\zeta}\beta$, then $\frac{1-\zeta}{2}\beta \geq \left(\frac{1-\zeta}{2} \right) \left(\frac{1-\zeta}{1+\zeta} \right) \varepsilon$. Hence,

$$\begin{aligned} \left(\frac{1-\zeta}{2}\beta \right) f'(x - \varepsilon) &\geq \left(\frac{1-\zeta}{2} \right) \left(\frac{1-\zeta}{1+\zeta} \right) \varepsilon f'(x - \varepsilon) \\ &= \left(\frac{\lambda - 1}{\lambda} \right) \left(\frac{1-\zeta}{2} \right) \left(\frac{1-\zeta}{1+\zeta} \right) \end{aligned}$$

and we are done. (Note that in order to bound the above integral over $[x + \frac{1-\zeta}{2}\beta, r]$ we could have worked with the point $x + \varepsilon + A'$ instead of $x + \frac{1-\zeta}{2}\beta + A'$.)

The next situation is the following. Assume that $\frac{1-\zeta}{2}\beta \leq \frac{1}{m}$, $\frac{1-\zeta}{2}\beta \leq \delta$ and $\varepsilon \leq \frac{1+\zeta}{1-\zeta}\beta$. This case is a little better than the previous one because the large $\frac{1}{m}$ means that we are far away from the singularity at x . But since δ can be still large, we might not have enough oscillation. So we still have to use condition (4). If $x - \frac{1}{m} \leq x - \varepsilon$, then

$$\left| \int_0^{x - \frac{1}{m}} e^{if(t)} \frac{1}{t - x} dt \right| \leq \frac{4m}{f'(x - \varepsilon)} \leq \frac{4}{\varepsilon f'(x - \varepsilon)} = \frac{4\lambda}{\lambda - 1}.$$

If $x - \frac{1}{m} > x - \varepsilon$, then

$$\left| \int_0^{x-\varepsilon} e^{if(t)} \frac{1}{t-x} dt \right| \leq \frac{4}{\varepsilon f'(x-\varepsilon)} = \frac{4\lambda}{\lambda-1}$$

and

$$\left| \int_{x-\varepsilon}^{x-\frac{1}{m}} e^{if(t)} \frac{1}{t-x} dt \right| \leq \log(\varepsilon m) \leq \log\left(\frac{2(1+\zeta)}{(1-\zeta)^2}\right).$$

This proves boundedness over the interval $[0, x - \frac{1}{m}]$. To work out the integral over $[x + \frac{1}{m}, r]$ we do the following:

$$\left| \int_{x+\frac{1}{m}}^{x+\frac{1}{m}+A'} e^{if(t)} \frac{1}{t-x} dt \right| \leq \log\left(\frac{\frac{1}{m} + A'}{\frac{1}{m}}\right) \leq \left(1 + \frac{(\frac{1-\theta}{\theta})(\frac{1+\zeta}{1-\zeta})}{(\frac{1-\zeta}{2})}\right)$$

and

$$\begin{aligned} \left| \int_{x+\frac{1}{m}+A'}^r e^{if(t)} \frac{1}{t-x} dt \right| &\leq \frac{4m}{|f'(x + \frac{1}{m} + A')|} \\ &\leq \frac{4}{\frac{1-\zeta}{2}\beta |f'(x + \frac{1-\zeta}{2}\beta + A')|} \\ &\leq \frac{4}{(\frac{1-\zeta}{2})(\frac{1-\zeta}{1+\zeta})\varepsilon f'(x-\varepsilon)} = \frac{4}{(1-\frac{1}{\lambda})(\frac{1-\zeta}{2})(\frac{1-\zeta}{1+\zeta})}, \end{aligned}$$

by (20).

So far we studied all possible situations with $\varepsilon \leq \frac{1+\zeta}{1-\zeta}\beta$. We still have to analyse the problem when $\varepsilon > \frac{1+\zeta}{1-\zeta}\beta$. For this, let B' be a number such that $\varepsilon - \beta = \zeta(B' + \varepsilon - \beta)$. We see that $B' = \frac{1-\zeta}{\zeta}(\varepsilon - \beta)$ is that number.

Suppose we have the following situation : $\frac{1}{m} < B' - \beta$ and $\varepsilon > \frac{1+\zeta}{1-\zeta}\beta$. (Note that $B' > 2\beta$ since $\varepsilon > \frac{1+\zeta}{1-\zeta}\beta$.) Since $\frac{1}{m} < B' - \beta$, then $x + \frac{1}{m} < \xi + B'$. Also, $B' - \beta \leq B' + \frac{1-\zeta}{\zeta}(\varepsilon - \beta) \leq \varepsilon - \beta < \varepsilon$. So, $\frac{1}{m} < B' - \beta$ implies that $\frac{1}{m} < \varepsilon$. We claim that

$$(21) \quad n\varphi'(\xi + B') - m \leq M[m - n\varphi'(x - \varepsilon)].$$

To prove (21) we use condition (5). By the choice of B' we have that $\xi = (1 - \zeta)(x - \varepsilon) + \zeta(\xi + B')$. Since $\varepsilon f'(x - \varepsilon) = \frac{\lambda-1}{\lambda}$ and $\varepsilon m > 1$, $\varphi'(\xi) \leq \lambda\varphi'(x - \varepsilon)$. Hence, (21) follows from (5).

Now, let's estimate the integral. We have

$$\begin{aligned} \left| \int_0^{x-\varepsilon} e^{if(t)} \frac{1}{t-x} dt \right| &\leq \frac{4}{\varepsilon f'(x-\varepsilon)} = \frac{4\lambda}{\lambda-1}, \\ \left| \int_{x-\varepsilon}^{x-\frac{1}{m}} e^{if(t)} \frac{1}{t-x} dt + \int_{x+\frac{1}{m}}^{\xi+B'} e^{if(t)} \frac{1}{t-x} dt \right| \\ &\leq M\varepsilon f'(x-\varepsilon) + M(B' - \beta)f'(x-\varepsilon) + \left| \int_{1/m}^{B'-\beta} \frac{1}{t} dt - \int_{1/m}^\varepsilon \frac{1}{t} dt \right| \end{aligned}$$

$$\leq 2M\varepsilon f'(x - \varepsilon) + \log\left(\frac{\varepsilon}{B' - \beta}\right) \leq 2M\left(\frac{\lambda - 1}{\lambda}\right) + \log\left(\frac{1 + \zeta}{1 - \zeta}\right),$$

by (21).

$$\begin{aligned} \left| \int_{\xi + B'}^{\xi + A'} e^{if(t)} \frac{1}{t - x} dt \right| &\leq \log\left(\frac{A' - \beta}{B' - \beta}\right) \\ &\leq \log\left(\frac{2A'}{B'}\right) = \log\left[\left(\frac{1 - \theta}{\theta}\right)\left(\frac{2\zeta}{1 - \zeta}\right)\right]. \\ \left| \int_{\xi + A'}^r e^{if(t)} \frac{1}{t - x} dt \right| &\leq \frac{4}{(A' - \beta) |f'(\xi + A')|} \leq \frac{4}{(B' - \beta)f'(x - \varepsilon)} \\ &\leq \frac{8}{B'f'(x - \varepsilon)} \leq \frac{4(1 + \zeta)\lambda}{(\lambda - 1)(1 - \zeta)}, \end{aligned}$$

by (20).

Finally, the last situation is that: $B' - \beta \leq \frac{1}{m}$ and $\varepsilon > \frac{1 + \zeta}{1 - \zeta}\beta$. If $x - \frac{1}{m} \leq x - \varepsilon$, then

$$\left| \int_0^{x - \frac{1}{m}} e^{if(t)} \frac{1}{t - x} dt \right| \leq \frac{4m}{f'(x - \frac{1}{m})} \leq \frac{4}{\varepsilon f'(x - \varepsilon)} = \frac{4\lambda}{\lambda - 1}.$$

If $x - \frac{1}{m} > x - \varepsilon$, then

$$\left| \int_0^{x - \varepsilon} e^{if(t)} \frac{1}{t - x} dt \right| \leq \frac{4}{\varepsilon f'(x - \varepsilon)} = \frac{4\lambda}{\lambda - 1},$$

and

$$\begin{aligned} \left| \int_{x - \varepsilon}^{x - \frac{1}{m}} e^{if(t)} \frac{1}{t - x} dt \right| &\leq \log(\varepsilon m) \leq \log\left(\frac{\varepsilon}{B' - \beta}\right) \\ &\leq \log\left(\frac{1 + \zeta}{1 - \zeta}\right). \end{aligned}$$

This proves boundedness of the integral over $[0, x - \frac{1}{m}]$. To estimate the integral on $[x + \frac{1}{m}, r]$ we do as follows: if $x + \frac{1}{m} \geq \xi + A'$, then

$$\begin{aligned} \left| \int_{x + \frac{1}{m}}^r e^{if(t)} \frac{1}{t - x} dt \right| &\leq \frac{4m}{|f'(x + \frac{1}{m})|} \leq \frac{4}{(B' - \beta) |f'(\xi + A')|} \\ &\leq \frac{4}{\left(\frac{1 - \zeta}{1 + \zeta}\right) \varepsilon f'(x - \varepsilon)} = \frac{4\lambda(1 + \zeta)}{(\lambda - 1)(1 - \zeta)}, \end{aligned}$$

by (20).

If $x + \frac{1}{m} < \xi + A'$, then

$$\begin{aligned} \left| \int_{x+\frac{1}{m}}^{\xi+A'} e^{if(t)} \frac{1}{t-x} dt \right| &\leq \log \left(\frac{A' - \beta}{\frac{1}{m}} \right) \leq \log \left(\frac{A'}{B' - \beta} \right) \\ &\leq \log \left(\frac{2A'}{B'} \right) = \log \left(\frac{2\zeta(1-\theta)}{\theta(1-\zeta)} \right), \end{aligned}$$

and

$$\begin{aligned} \left| \int_{\xi+A'}^r e^{if(t)} \frac{1}{t-x} dt \right| &\leq \frac{4}{(A' - \beta) |f'(\xi + A')|} \leq \frac{8}{B' f'(x - \varepsilon)} \\ &\leq \frac{4}{\left(\frac{1-\zeta}{1+\zeta}\right) \varepsilon f'(x - \varepsilon)} = \frac{4\lambda(1+\zeta)}{(\lambda-1)(1-\zeta)}, \end{aligned}$$

by (20).

This concludes the proof of the theorem.

REFERENCES

1. Kahane, J. P. - Quatre Leçons sur les Homéomorphismes du Cercle et les Series de Fourier. Topics in Modern Harmonic Analysis, Proceedings of a seminar held in Torino and Milano, May - June 1982, vol. II, Roma, 1983. MR **85k**:42012
2. Stein, E. M. and Wainger, S. - The Estimation of an Integral Arising in Multiplier Transformations, *Studia Math.* **35** (1970), 101 - 104. MR **42**:904
3. Stein, E. M. - Oscillatory Integral in Fourier Analysis, Beijing Lectures in Harmonic Analysis, edited by E. M. Stein, Princeton University Press. (1986), 307 - 355. MR **88g**:42022
4. Nagel, A., Vance, J., Wainger, S. and Weinberg, D. - Hilbert Transforms for Convex Curves, *Duke Mathematical Journal*. Vol. 50 (1983), 735 - 744. MR **85a**:42025
5. Santos Zumpano, A. - Homeomorphisms Of The Circle And Fourier Series, Ph.D thesis - University of Wisconsin - Madison, May 1987.

DEPARTAMENTO DE MATEMATICA, UNIVERSIDADE FEDERAL DE MINAS GERAIS, CAIXA POSTAL 702, 30.123-970 BELO HORIZONTE MG, BRAZIL
E-mail address: zumpano@mat.ufmg.br